

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Emulsifying Properties of Hydrolyzed Sunflower Lecithins by Phospholipases A₂ of Different Sources

D. M. Cabezas¹, R. Madoery²,
B. W. K. Diehl³ and M. C. Tomás¹

¹*Centro de Investigación y Desarrollo en Criotecnología de Alimentos (CIDCA) – CCT La Plata – CONICET - Facultad de Ciencias Exactas (FCE), Universidad Nacional de La Plata (FCE - UNLP), La Plata*

²*Cátedra de Química Orgánica – Facultad de Ciencias Agrarias, Universidad Nacional de Córdoba (FCA - UNC), Córdoba*

³*Spectral Service GmbH Laboratorium für Auftragsanalytik, Cologne*
^{1,2}*Argentina*
³*Germany*

1. Introduction

Lecithins are a mixture of acetone insoluble phospholipids, containing mainly phosphatidylcholine (PC), phosphatidylethanolamine (PE), phosphatidylinositol (PI), minor compounds such as phosphatidic acid (PA), and other minor substances such as carbohydrates and triglycerides (Schneider, 1989). The production of sunflower oil, in Argentina, is of utmost importance from an economic point of view (Franco, 2008). In this country, sunflower lecithin might represent an alternative to soybean lecithin because it is considered a non-GMO product, which is in accordance with the preference of some consumers.

The introduction of changes in the original concentration of these phospholipids, by chemical or enzymatic modification of their structure can lead to obtain lecithins with different physicochemical and functional properties, with respect to native lecithin (van Nieuwenhuyzen & Tomás, 2008). The modification processes usually applied on native lecithins are the fractionation with ethanol (Sosada, 1993; Wu & Wang, 2004; Cabezas et al., 2009a, 2009b) and the enzymatic hydrolysis (Schmitt & Heirman, 2007; Cabezas et al., 2011a). Native and modified lecithins are used in a wide range of industrial applications: nutritional, pharmaceutical applications, food, cosmetics, etc. (Prosise, 1985; Wendel, 2000). In the food industry, lecithin represents a multifunctional additive in the manufacture of chocolate, bakery and instant products, margarines, and mayonnaise, due to the characteristics of its phospholipids (van Nieuwenhuyzen, 1981).

In particular, enzymatic hydrolyzed lecithin may present technological and commercial advantages over native lecithins: (1) enhanced O/W emulsifying property; (2) increased emulsion stability under acid conditions and in the coexistence with salts; (3) improved

capability to bind proteins and starch; (4) excellent mold- or pan-releasing property. Consequently, the demand for lysolecithins was increasing in recent years (Hirai et al., 1998; Erickson, 2008).

The main application of lecithin at the food industry is associated with its role as emulsifier agent for dispersions or emulsions (Hernández & Quezada, 2008). Emulsions are thermodynamically unstable systems from a physicochemical point of view. In virtue of that, it is important to characterize and know their behaviour against different destabilization processes (flocculation, coalescence, creaming, etc.) (McClements, 1999).

Enzymatic hydrolysis is carried out mainly by two groups of enzymes: phospholipases and lipases (Mustranta et al., 1995). Phospholipases A₂ catalyze the hydrolysis of the ester bond in the sn-2 position of glycerophospholipids, producing free fatty acids and the corresponding lysophospholipid. Advances in biotechnology and certain requirements of consumers (*kosher* or *halal* foods) have influenced the development of the production of microbial enzymes (bacteria, fungi, yeasts) which could be substitute of the traditionally obtained from porcine pancreas (Minchiotti, 2006; Cabezas et al., 2011b).

The aim of this work was analyze the emulsifying activity of sunflower lysolecithins obtained by phospholipases A₂ from diverse sources: bacterial (LysoMax PLA₂, Danisco) and porcine pancreas (Lecitase 10L, Novo Nordisk) in O/W systems. This study seeks to contribute to the oil industry with useful information for rescaling of the mentioned hydrolysis process, with the aim of increasing the aggregated value of sunflower lecithins.

2. Materials and methods

2.1 Materials

Native sunflower lecithin was used as starting material, and was provided by a local oil industry (Vicentin S.A.I.C.). Enzymatic hydrolysis processes were carried out using a porcine pancreatic PLA₂ (Lecitase 10L, Novo Nordisk) and a microbial PLA₂ (*Streptomyces violaceoruber*, LysoMax PLA₂, Danisco). All solvents used were of analytical grade.

The sunflower lecithin used as starting material presented the following composition: 43.1% phospholipids (16.5% PI, 16.2% PC, 5.3% PE, and 5.1% minor phospholipids), 33.4% oil, and 23.5% of other compounds (glycolipids, complex carbohydrates).

2.2 Enzymatic hydrolysis process

Enzymatic hydrolysis was carried out in a thermostated reactor at laboratory scale, using 27 g of native sunflower lecithin and 18 ml of 0.4 M CaCl₂. Initial pH was adjusted to 7 or 9 by adding 4 N NaOH solution. Then, the resulting mixture was set to the optimal temperature of each phospholipase, i.e. 60 °C for porcine pancreatic PLA₂ and 50 °C for microbial PLA₂, which were incorporated in a concentration of 2.0% ml lipase per 100 g lecithin. Next, continuous agitation (50 rpm) was applied during 5 h. Evolution of hydrolysis process was followed by measuring pH, using a pH meter for solid samples (840049 Puncture Tip, Saen S.R.L.). Products of enzymatic hydrolysis were subjected to a sudden decrease in temperature to stop the process of hydrolysis and then deoiled using acetone, according to AOCS Official Method Ja 4-46, procedures 1-5 (Cabezas et al., 2011a). After that, samples were stored at 0 °C. The hydrolysis process was carried out in duplicate.

Also, native sunflower lecithin was deoiled with acetone obtaining the deoiled sunflower lecithin (DSL). DSL was used as a sample control. Deoiling procedure was performed by duplicate.

2.3 Phospholipid composition

2.3.1 Sample preparation

100 mg of each hydrolyzed sample were diluted in 1 ml of deuterated chloroform, 1 ml of methanol and 1 ml of Cs-EDTA (pH 8). The organic layer was separated after 15 min shaking, and analyzed by ^{31}P NMR (Cabezas et al., 2009a).

2.3.2 Quantitative ^{31}P NMR analysis

Quantitative ^{31}P NMR analysis was carried out in a Bruker Avance 600 MHz automatic spectrometer using triphenyl phosphate as internal standard (Spectral Service GmbH, Köln, Germany) (Diehl, 1997; 2001; 2008). Phospholipid content of samples obtained under different conditions of enzymatic hydrolysis, was determined by this spectroscopic technique.

2.4 Oil-in-water (O/W) emulsions preparation

Commercial sunflower oil was used to prepare oil-in-water (O/W) emulsions with a formulation of 30:70 (wt/wt) according to Pan et al., 2004. Emulsions were prepared at room temperature in an Ultra-Turrax T25 homogenizer using S 25 N-10 G dispersing tool (7.5 mm rotor diameter) at 10,000 rpm for 1 min, with the addition of the different modified sunflower lecithins in a range of 0.1–2.0% (wt/wt). This process was carried out in triplicate for each case.

2.5 Optical characterization of emulsions

The backscattering of light was measured using a QuickScan Vertical Scan Analyzer (Coulter Corp., Miami, FL). The backscattering of monochromatic light ($\lambda = 850$ nm) of the emulsions was determined as a function of the height of the sample tube (ca. 65 mm) in order to quantify the rate of the different destabilization processes during the first 90 min. This methodology allowed to discriminate between particle migration (sedimentation, creaming) and particle size variation (flocculation, coalescence) processes (Pan et al., 2002). The basis of the vertical scan analyzer profiles has been exhaustively studied by Mengual (Mengual et al., 1999).

2.6 Particle size measurements

Particle size distribution, and De Brouckere (D[4,3]) and Sauter (D[3,2]) mean diameters of particles of the emulsions were determined with a particle size analyzer (Malvern Mastersizer 2000E, Malvern Instruments Ltd., Worcestershire, U.K.). Samples were diluted in the water bath of the dispersion system (Hydro 2000MU), which is a laser diffraction based particle size analyzer (Márquez & Wagner, 2010). This determination was carried out in triplicate for each case.

2.7 Statistical analysis

Data were evaluated by analysis of variance (ANOVA) using the software Systat® 12.0 (Systat, 2007). For this purpose, differences were considered significant at $p < 0.05$.

	DSL	SHLM9	SHLM7	SHLP9	SHLP7
PC	36.7	7.4	6.7	< 0.1	< 0.1
1-LPC	< 0.1	< 0.1	< 0.1	< 0.1	< 0.1
2-LPC	1.5	39.2	39.4	46.1	45.5
PI	35.3	11.2	12.3	< 0.1	< 0.1
LPI	< 0.1	18.6	15.5	26.5	27.9
PE	15.1	3.7	6.4	1.0	0.7
LPE	< 0.1	8.8	9.0	10.7	13.0
APE	1.8	< 0.1	< 0.1	< 0.1	< 0.1
PA	5.2	3.1	3.4	2.5	2.0
LPA	< 0.1	1.0	1.0	5.1	4.1
Other	4.5	7.1	6.4	8.0	6.7
PL / 100g lecithin	62.8	40.6	45.9	38.7	44.4

Table 1. Phospholipid (PL) composition of modified sunflower lecithins by ^{31}P NMR^a

3. Results and discussion


3.1 Compositional analysis of modified lecithins

The phospholipid composition of different modified sunflower lecithins obtained in this chapter is shown in Table 1. The sunflower hydrolyzed lecithins with a porcine pancreatic (SHLP) and microbial (SHLM) phospholipase A_2 , at different levels of pH (7, 9), recorded a marked difference regarding the phospholipid composition in relation to the native sunflower lecithin (DSL). The hydrolyzed lecithins presented a high concentration of major lysophospholipids (> 64.9 mol LPL/ mol of total PL) compared to the native sunflower lecithin ($\approx 1.5\%$), showing the efficiency of the enzymatic hydrolysis processes. In particular, the pancreatic PLA_2 (SHLP7, SHLP9) produced a higher hydrolysis degree of the main phospholipids in comparison with the microbial phospholipase (SHLM7, SHLM9).

PC presented a high degree of hydrolysis in all performed conditions on the hydrolysis processes. These results can be correlated with the ones described by Penci, 2010. In that work, it was reported that phosphatidylcholine is the phospholipid with higher tendency to be hydrolyzed when using a porcine pancreatic PLA_2 with a very low amount of sunflower lecithins (1 mg). In this way, the residual PC and PI concentration in the hydrolyzed lecithin was lower than the detection limit of the ^{31}P NMR equipment when the pancreatic porcine PLA_2 was used.

3.2 Optical characterization of O/W emulsions

Stability of the different O/W emulsions (30:70 wt/wt) was studied recording the backscattering (BS) profiles as a function of the cell length and time, by a vertical scan analyzer (QuickScan). For instance, Figure 1 shows two typical profiles obtained for emulsions with the addition of 0.1% of DSL and SHLP9.



^a Mean values are shown ($n = 3$). The coefficient of variation was lower than 5%

Fig. 1. Backscattering (%BS) profiles of O/W emulsions (30:70 wt/wt) with the addition of: (a) DSL, 0.1%; (b) SHLP, 0.1%

The creaming destabilization process (i.e. migration of oil particles to the upper portion of the tube) is evidenced by a decrease of %BS values at the bottom of the tube. The QuickScan profiles corresponding to the zone I (10-20 mm) showed an increase of the emulsion stability against the creaming process, as a function of increasing concentration of different modified lecithins (Fig. 2). In particular, the hydrolyzed lecithins (SHLP and SHLM) generated a high stability in O/W emulsions than DSL, over the studied range of concentration. Moreover, O/W emulsions with 0.1-0.5% of DSL showed a sharp decrease of %BS in the Zone I.

The tube zone between 40-45 mm (Zone II) is characterized by the accumulation of oil droplets after the creaming process (cream phase); Figure 3 shows the %BS values vs. time in Zone II. Emulsions formulated with hydrolyzed lecithins presented higher %BS values than those obtained using DSL, for all concentrations studied. The higher levels of %BS and the greater stability of these emulsions would be associated with the formation of dense cream phases with a lower proportion of continuous phase inside (Palazolo, 2006). However, emulsions with 0.1 -0.5% of DSL did not allow the formation of the cream phase. These results are related to the rapid decrease of %BS and the formation of an oil layer in the upper part of the tube (Fig. 1a) suggesting the occurrence of a cream phase destabilization by coalescence (Pan et al., 2002).




Fig. 2. Backscattering (%BS) values of O/W emulsions (30:70 wt/wt) with the addition of different modified sunflower lecithins in Zone I (10-20 mm). Mean values ($n = 3$) \pm sd




Fig. 3. Backscattering (%BS) values of O/W emulsions (30:70 wt/wt) with the addition of different modified sunflower lecithins in Zone II (40-45 mm). Mean values ($n = 3$) \pm sd

3.3 Particle size distribution

Particle size distribution in volume and surface of O/W emulsions obtained with different modified lecithins was measured just after emulsification ($t=0$); corresponding results can be seen in Figures 4 and 5, respectively. These distributions presented a bimodal or trimodal character depending on the concentration of aggregated lecithin, and the following particle size populations: (I) particle size $< 4 \mu\text{m}$; (II) particle size between 4 and $30 \mu\text{m}$; (III) particle size $> 30 \mu\text{m}$. In this sense, only SHLP7 showed a trimodal character for all concentrations assayed. It should be noted that the SHLM7 presented a bimodal character for concentration in the range 0.1-0.5%, but with a high percentage of particles of the population II in comparison with DSL.

In order to complete the analysis of particle size distribution, Figure 6 depicts the evolution of De Brouckere ($D [4,3]$) and Sauter ($D [3,2]$) mean diameters as a function of the concentration of the different emulsifiers. Hydrolyzed lecithins generated values of $D [4,3]$ and $D [3,2]$ significantly lower than those corresponding to DSL. These results are correlated with the high stability of the O/W emulsions recorded when using hydrolyzed lecithins, considering the main destabilization processes determined by the corresponding QuickScan profiles (creaming or coalescence). It is worth to note that a high concentration of small particles produces a slow creaming process, according to the Stokes' law (McClements, 1999; Palazolo, 2006).




Fig. 4. Volume particle size distribution for O/W emulsions with the addition of different modified sunflower lecithins. Mean values ($n = 3$)

The hydrophilic-lipophilic balance value (HLB) is often used in connection with the performance of emulsifiers (McClements, 1999). The high concentration of hydrophilic phospholipids (lysophospholipids) presented in the hydrolyzed lecithins (SHLP and SHLM) increase this empirical value. In this sense, according to Carlsson (Carlsson, 2008), these modified lecithins with higher HLB values presented best properties as O/W emulsifying agents.

Also, the phase structure at the interface of the different phospholipids influences the emulsion formation and stability (van Nieuwenhuyzen & Tomás, 2008). LPC and LPE form hexagonal wide spread clusters. These structures have a great importance for the stabilisation of O/W emulsions. This behaviour is in relation to the low mean diameters and the high concentration of small particle populations recorded in emulsions using sunflower hydrolyzed lecithins (Figs. 4-6). However, PE gives reversed hexagonal phase, which are more difficult to arrange at the interface (van Nieuwenhuyzen, 1998). The presence of PE could explain the minor characteristics as emulsifying agent of DSL, and the high mean diameters when was using SHLM in contrast to when using SHLP (Fig. 6).

Taking into account the results presented in Figures 2 to 6, the addition of a concentration between 0.5 and 1.0% of hydrolyzed lecithin (SHLP, SHLM) is enough for covering all droplets surface. High concentrations of this modified lecithin do not show significant differences in the % BS values, nor in the mean particle sizes. On the other hand, concentration levels higher than 0.1% of hydrolyzed lecithins from different phospholipases at different initial pH levels showed similar characteristics in terms of their emulsifying activity. However, DSL presented an improved in the stability of O/W emulsions as a function of increasing concentration.




Fig. 5. Surface particle size distribution for O/W emulsions with the addition of different modified sunflower lecithins. Mean values ($n = 3$)




Fig. 6. De Brouckere (D[4,3]) and Sauter (D[3,2]) mean diameters for O/W emulsions with the addition of different modified sunflower lecithins. Mean values ($n = 3$) \pm sd

4. Conclusions

The best emulsifying properties exhibited by hydrolyzed sunflower lecithins (LHM7, LHP7, LHM9, LHP9) in comparison with the native lecithin could indicate that their application by the local oil industry could revalue this by-product of the refining process of crude sunflower oil.

The use of a microbial phospholipase A₂ gives the possibility to obtain diverse sunflower lysolecithins, which functionality could be applied to the development of foods with *kosher* and *halal* certification.

5. Acknowledgments

This work was supported by grants from Agencia Nacional de Promoción Científica y Tecnológica (ANPCyT), Argentina, PICT 1085 (ANPCyT); Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina, PIP 1735 (CONICET); and Universidad Nacional de La Plata (UNLP), Argentina, 11/X502 (UNLP).

M. C. Tomás is a member of the Career of Scientific and Technological Researcher of Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina. D. M. Cabezas is a recipient of a postdoctoral fellowship from Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). R. Madoery is a Scientific and Technological Researcher, and Professor at Universidad Nacional de Córdoba (UNC). B. W. K. Diehl is Director of Spectral Service GmbH, Cologne, Germany.

Sunflower lecithin and microbial PLA₂ were provided by Vicentin S.A.I.C. and Danisco, respectively. Thorsten Buchen is acknowledged for his technical assistance.


6. References

- Cabezas, D.M., Diehl, B. W. K. & Tomás M.C. (2009a). Effect of processing parameters on sunflower PC enriched fractions extracted with aqueous-ethanol. *European Journal of Lipid Science and Technology*, No 111, pp. 993-1002
- Cabezas, D.M., Diehl, B. W. K. & Tomás M.C. (2009b). Sunflower lecithin: Application of a fractionation process with absolute ethanol. *Journal of the American Oil Chemists' Society*, No 86, pp. 189-196.
- Cabezas, D.M., Madoery, R., Diehl, B.W.K. & Tomás, M.C. (2011a). Application of enzymatic hydrolysis on sunflower lecithin using a pancreatic PLA₂. *Journal of the American Oil Chemists' Society*, No 88, pp. 443-446
- Cabezas, D.M., Madoery, R., Diehl, B.W.K. & Tomás, M.C. (2011b). Enzymatic hydrolysis of sunflower lecithins using a microbial PLA₂. In: Hughes VC (ed) *Sunflowers: cultivation, nutrition, and biodiesel uses*. Nova Science Publishers, New York (USA), Chapter 9, pp. 303-316
- Carlsson, A. (2008). Physical properties of phospholipids. In: Gunstone FD (ed) *Phospholipid technology and applications*. The Oily Press, Bridgwater (UK), Chapter 5, pp. 104
- Diehl, B. W. K. (1997). Multinuclear high-resolution magnetic resonance spectroscopy. In: *Lipid Analysis in Oils and Fats*. Ed. R. J. Hamilton, Blackie Academic & Professional, London (UK), pp. 87-135

- Diehl, B. W. K. (2001). High resolution NMR spectroscopy. *European Journal of Lipid Science and Technology*, No 103, pp. 830–834
- Diehl, B. W. K. (2008). NMR spectroscopy of natural substances. In: Holzgrabe U, Wawer I, Diehl B (eds). *NMR spectroscopy in pharmaceutical analysis*, Chapter 5. Elsevier, Jordan Hill (UK), pp. 194–196
- Erickson, M.C. (2008). Chemistry and function of phospholipids. In: Akoh CC, Min DB (eds) *Food lipids: chemistry, nutrition and biotechnology*, Chapter 2, 3rd edn. CRC Press, Boca Raton, pp. 39–62
- Franco, D. (2008). Informe de Cadenas Alimentarias: Sector Oleaginoso. Ministerio de Economía y Producción Argentina. http://www.alimentosargentinos.gov.ar/0-3/revistas/r_41/cadenas/Oleaginosos_Oleaginosas.htm (online)
- Hernández, E. & Quezada, N. (2008). Uses of phospholipids as functional ingredients. In: F.D. Gunstone (Ed.), *Phospholipid Technology and Applications*, Chapter 4, Bridgwater, The Oily Press
- Hirai, H., Sono, R. & Koh, H. (1998). Process for manufacturing vegetable lysolecithins. European Patent EP0870840
- Márquez, A.L. & Wagner, J.R. (2010). Rheology of double (W/O/W) emulsions prepared with soybean milk and fortified with calcium. *Journal of Texture Studies*, No 41, pp. 651–671
- McClements, D.J. (1999). *Food emulsions: principles, practice and techniques*. CRC Press, Florida (USA)
- Mengual, O., Meunier, G., Cayré, I., Puech, K. & Snabre, P. (1999). Turbiscan MA 2000: multiple light scattering measurement for concentrated emulsion and suspension instability analysis. *Talanta*, No 50, pp. 445–456
- Minchiotti, M.C. (2006). Purificación, caracterización y estudio de propiedades de fosfolipasa A2 presente en soja (*Glycine max*). Doctoral Thesis. Facultad de Ciencias Agropecuarias, Universidad Nacional de Córdoba (UNC), Argentina
- Mustranta, A., Forssell, P. & Poutanen, K. (1995). Comparison of lipases and phospholipases in the hydrolysis of phospholipids. *Process Biochemistry*, No 30, pp. 393–401
- Palazolo, G.G. (2006). Formación y estabilidad de emulsiones O/W preparadas con proteínas nativas y desnaturalizadas de soja. Doctoral Thesis, Universidad Nacional de La Plata, Argentina
- Pan, L.G., Tomás, M.C. & Añón, M.C. (2002). Effect of sunflower lecithins on the stability of water in Oil (W/O) and Oil in Water (O/W) emulsions. *Journal of Surfactants and Detergents*, No 5, pp. 135 – 143
- Pan, L.G., Tomás, M.C. & Añón, M.C. (2004). Oil-in-Water emulsions formulated with sunflower lecithins: vesicle formation and stability. *Journal of the American Oil Chemists' Society*, No 81, pp. 241–244
- Penci, M.C., Constenla, D.T. & Carelli, A.A. (2010). Free-fatty acid profile obtained by enzymatic solvent-free hydrolysis of sunflower and soybean lecithins. *Food Chemistry*, No 120, pp. 332–338
- Prosise, W.E (1985). Commercial lecithin products: Food use of soybean lecithin. In: *Lecithins*. Eds. B. F. Szuhaj, G. List, AOCS Press, Champaign (USA), pp. 163–181
- Schmitt, H. & Heirman, M. (2007). Enzymatic modification of lecithin. US Patent US7189544B2

- Schneider, M. (1989). Fractionation and Purification of Lecithin. In *Lecithins: Sources, Manufacture, & Uses*. Ed. B.F. Szuhaj, AOCS Press, Champaign (USA), pp. 109-130
- Sosada, M. (1993). Optimal conditions for fractionation of rapeseed lecithin with alcohols. *Journal of the American Oil Chemists' Society*, No. 70, pp. 405-410
- Systat, Version 12 (2007). Systat Software Inc.. San José, California (USA).
- van Nieuwenhuyzen W. and Tomás, M. C. (2008). Update on vegetable lecithin and phospholipid technologies. *European Journal of Lipid Science and Technology*, No 110, pp. 472-486
- van Nieuwenhuyzen, W. (1981). The industrial uses of special lecithins. *Journal of the American Oil Chemists' Society*, No 58, pp. 886-888
- van Nieuwenhuyzen, W. (1998). Effects of lecithins and proteins on the stability of emulsions. *Fett/Lipid*, No 100, pp. 282-291
- Wendel, A. (2000). Lecithin: The first 150 years. Part I: From discovery to early commercialization. *Inform*, No 11, pp. 885-892
- Wu Y. & Wang T. (2004). Fractionation of crude soybean lecithin with aqueous ethanol. *Journal of the American Oil Chemists' Society*, No 81, pp. 697-704

IntechOpen



Food Industrial Processes - Methods and Equipment

Edited by Dr. Benjamin Valdez

ISBN 978-953-307-905-9

Hard cover, 418 pages

Publisher InTech

Published online 22, February, 2012

Published in print edition February, 2012

The global food industry has the largest number of demanding and knowledgeable consumers: the world population of seven billion inhabitants, since every person eats! This population requires food products that fulfill the high quality standards established by the food industry organizations. Food shortages threaten human health and are aggravated by the disastrous, extreme climatic events such as floods, droughts, fires, storms connected to climate change, global warming and greenhouse gas emissions that modify the environment and, consequently, the production of foods in the agriculture and husbandry sectors. This collection of articles is a timely contribution to issues relating to the food industry. They were selected for use as a primer, an investigation guide and documentation based on modern, scientific and technical references. This volume is therefore appropriate for use by university researchers and practicing food developers and producers. The control of food processing and production is not only discussed in scientific terms; engineering, economic and financial aspects are also considered for the advantage of food industry managers.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

D. M. Cabezas, R. Madoery, B. W. K. Diehl and M. C. Tomás (2012). Emulsifying Properties of Hydrolized Sunflower Lecithins by Phospholipases A2 of Different Sources, Food Industrial Processes - Methods and Equipment, Dr. Benjamin Valdez (Ed.), ISBN: 978-953-307-905-9, InTech, Available from: <http://www.intechopen.com/books/food-industrial-processes-methods-and-equipment/emulsifying-properties-of-hydrolized-sunflower-lecithins-by-phospholipases-a2-of-different-sources>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen