

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

## 4,800

Open access books available

## 122,000

International authors and editors

## 135M

Downloads

Our authors are among the

## 154

Countries delivered to

## TOP 1%

most cited scientists

## 12.2%

Contributors from top 500 universities

**WEB OF SCIENCE™**Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)

## Probiotics and Oral Health

Harini Priya Vishnu  
 Department of Pedodontics and Preventive Dentistry,  
 Vydehi Institute of Dental Sciences and Research Centre, Bangalore  
 India

### 1. Introduction

*All great thoughts have already been thought; what is necessary is only to try to think them again....*

*Johann Wolfgang*

Man is the most intelligent animal on the earth, as he is the only species gifted with the power of thinking and reasoning. He always manages to come up with solutions for *most problems* on earth. Ironically, most problems on earth are directly or indirectly created by Man himself. That's the reason he is most often referred to as an *intelligent fool*.

*The man-made problem* prevailing across the field of medicine is the development of *antibiotic resistance*, largely contributed by us, the health professionals. The indiscriminate and reckless use of antibiotics has led to the emergence of multi-resistant strains of bacteria. This unfortunate development has led scientists to seek other means of fighting infections. [1]

The good-old but forgotten concept of *bacteriotherapy* seem to offer innovative tools for the treatment of infectious diseases. Hence, rightly quoted "All great thoughts have already been thought; what is necessary is only to try to think them again".

*Bacteriotherapy is the administration of naturally occurring bacteria of human origin as a therapeutic manipulation of the bacterial microenvironment in the patient's body.* [1] The Basic principle is to use good bacteria to compete against pathogenic bacteria. Bacteriotherapy has been studied and tested to control infectious diseases, particularly in the GI tract. Recent studies have shown that this therapeutic method may be used to influence body function in other systems too, beyond the intestine. [1, 2, 3]

### 2. History

It was in the first decade of 1900, the beginning of the 20<sup>th</sup> century when the Ukrainian-born Nobel prize laureate Elie Metchnikoff observed the positive beneficial effect of some bacteria on the human health and suggested that these beneficial bacteria can be used to replace harmful microbes in the body. He reported that Bulgarians lived longer than other population which was due to their consumption of Bulgarian yoghurt which contained lactic acid bacteria.[4] Metchnikoff worked at the Pasteur Institute in Paris and had discovered *Lactobacillus bulgarius*, a strain he later introduced into commercial production of sour-milk products in France and throughout Europe. He devoted the last decade of his life to the study of lactic-acid producing bacteria as a means of increasing human longevity. [5] He developed a theory stating that senility is caused by poisoning of

the body by the products of some of the bacteria for which he proposed a diet containing milk fermented by lactobacilli producing lactic acid to prevent the multiplication of these organisms. [6] The concept of probiotics was thus born and a new field of microbiology came into light.

The first clinical trials were performed in the 1930s on the effect of probiotics on constipation. Ever since then, different microorganisms have been used for their ability to prevent and cure diseases. [7]

In 1994, the WHO deemed probiotics to be the next-most important immune defence system when commonly prescribed antibiotics are rendered useless by antibiotic resistance. The use of probiotics in antibiotic resistance is termed as *microbial interference therapy*. [7]

The literature on the possible role of probiotics on oral and dental health is scarce and the studies on probiotics v/s oral health are still in their cradle. [6]

### 3. What are Probiotics?

The term probiotic means 'for life', was first coined by Lilly and Stillwell. [8]

### 4. Definition

[Adopted by the International Scientific Association for Probiotics and Prebiotics term]

Probiotics are defined as "Live microorganisms, which when administered in adequate amounts, confer a health benefit on the host" (Guarner et al 2005) [9]

### 5. Oral probiotic organisms [7]

The most common probiotic strains belong to the genera *Lactobacillus* and *Bifidobacterium*. Bacterial strains that have been tested for probiotic action in the oral cavity include:

LACTOBACILLI SPECIES	BIFIDOBACTERIUM SPECIES	OTHERS
<ul style="list-style-type: none"> <li>• <i>L. acidophilus</i></li> <li>• <i>L. rhamnosus</i> GG</li> <li>• <i>L. johnsonii</i></li> <li>• <i>L. casei</i></li> <li>• <i>L. rhamnosus</i></li> <li>• <i>L. gasseri</i></li> <li>• <i>L. reuteri</i></li> <li>• <i>L. paracasei</i></li> </ul>	<ul style="list-style-type: none"> <li>• <i>B. bifidum</i></li> <li>• <i>B. longum</i></li> <li>• <i>B. infantis</i></li> <li>• <i>Bifidobacterium</i> DN-173 010</li> </ul>	<ul style="list-style-type: none"> <li>• <i>S. salivarius</i></li> <li>• <i>W. cibaria</i></li> <li>•</li> </ul>

Table 1.

*L. rhamnosus* GG, ATCC 53103 (LGG) is the most widely studied probiotic bacterium. Named after the discoverers, Sherwood Gorbach and Barry Goldin, produces a substance

with potential inhibitory activity against different bacterial species including cariogenic Streptococcus species. [1]

## 6. Vehicles for probiotic administration for oral health purpose

The various means of administration of probiotics for oral health purpose that has been studies are:

- Lozenges
- Tablets
- Cheese
- Yoghurt
- Mouth rinse
- Capsule, Liquid

## 7. Mechanisms of probiotic action on oral health

An essential requirement for an organism to be “an oral probiotic” is its ability to adhere to and colonize surfaces in the oral cavity. Microorganisms generally considered as probiotics may not have oral cavity as their inherent habitat, so their possibility to confer benefit on oral health is then questionable. [6]The suggested mechanisms of probiotic action are drawn entirely from gastrointestinal studies. Since, mouth is the gateway of the GI tract, there is every reason to believe that atleast some probiotic mechanisms may play a role in the oral cavity. [1]

Some of the hypothetical mechanisms of probiotic action in the oral cavity are:

- Lactobacilli play an important role in maintaining the microecological balance in the oral cavity.[11]
- Direct interaction in dental plaque
- Involvement in binding of oral micro-organisms to proteins [interference in formation of acquired pellicle]
- Action on plaque formation and on its complex ecosystem by competing and intervening with bacterial attachments.
- Involvement in metabolism of substrate and production of chemicals that inhibit oral bacteria.[12, 13]

Indirect probiotic actions are also featured such as

- Modulating systemic immune function.
- Effect on local immunity.
- Effect on non-immunologic defense mechanisms.
- Regulation of mucosal permeability.
- Probiotics as an antioxidants and produce antioxidants.
- Prevent plaque formation by neutralizing the free electrons. [14, 15, 16]

Immune inductive sites in the oral cavity are within the diffuse lymphoid aggregates of the Waldeyer's ring. Lingual and pharyngeal tonsils and adenoids contain most of the lymphatic tissue. Dendritic cells in the mucosal surfaces play vital role in antigen presentation and in activating T-cell responses. Depending on the signals from dendritic cells either immune tolerance or active immune response toward a specific antigen may occur. [6] However, more studies investigating the role probiotics on activation of the oral immune inductive sites are required before further conclusions are drawn.




Fig. Possible Mechanisms of Probiotic Action in the Oral Cavity [10]

## 8. Action of probiotics on oral health

### 8.1 Action on organisms associated with dental caries

Several investigations have shown reduction in the number of mutans streptococci in saliva after consumption of various probiotic products, [17 – 25] however such an effect has not been observed in all studies. [26]

Náse et al, 2001 showed that supplementing 1-6 year old children with *L. rhamnosus* for 7 months significantly reduced the risk of dental caries. [17]

Comelli EM et al (2002) studied 23 dairy bacterial strains for the prevention of dental caries and reported that only two strains namely *Streptococcus thermophilus* and *Lactococcus lactis* were able to adhere to saliva-coated hydroxyapatite and were further successfully incorporated into a biofilm similar to the dental plaque. Furthermore, they could grow together with five strains of oral bacterial species commonly found in supragingival plaque. In this system, *Lactobacillus lactis* was able to modulate the growth of the oral bacteria, and in particular to diminish the colonization of *Streptococcus oralis*, *Veillonella dispar*, *Actinomyces naeslundii* and of the cariogenic *Strep.sobrinus*. [27]

Chung et al, 2004 showed that the probiotic strain *L. fermentum* found in the saliva of healthy children significantly inhibited the formation of the insoluble glucan produced by *S. mutans*. It did not affect the multiplication of this pathogenic strain, but it completely inhibited the adherence onto cuvette walls. [28]

Koll-Klais et al (2005) used various lactobacilli strains in their study and stated that 69% of these strains inhibited *S. mutans*, 82% inhibited *P. gingivalis*. [29]

An increase in the number of salivary lactobacilli has also been seen in some studies. [18, 26]

The products containing probiotic LGG bacteria may have beneficial effects on the dental health. The LGG bacteria had been shown not to ferment lactose or sucrose. [30]

Stamatova et al, in their study that was conducted in 2007 stated that *L. rhamnosus* & *Lactobacillus bulgaricus* produced inhibitory effects against *P. gingivalis*, *Fusobacterium nucleatum* & streptococcal species. [31]

Strahnic et al, 2007 conducted a study using probiotic strains *L. salivarius* & *L. fermentum* and both strains showed antagonistic activity on the growth of *S. mutans* and *Streptococcus pneumoniae*. *L. salivarius* was able to survive an environment of low pH as that produced by a high number of *S. mutans*. [32]

N.S.H. Mehanna et al (2009) investigated the effect of plant meswak and some probiotic bacteria on *Streptococcus mutans* and *Porphyromonas gingivalis* isolated from human oral cavity as most common oral pathogenic strains. The results of the study indicated that *Lactobacillus rhamnosus* had a marked decreasing effect on the colonization of both *Streptococcus mutans* and *Porphyromonas gingivalis*. [33]

A recently published review article by Anna Haukioja in 2010 does not give a conclusive statement about the effect of probiotics on dental caries or caries related organisms since the study groups have been relatively small and of fairly shorter duration. [10]

## 8.2 Action on periodontal diseases

Encouraging results have been obtained in studies investigating the role of probiotics for the treatment of various periodontal diseases, gingivitis, plaque levels, and periodontitis.

Reduction in the number of periodontopathogens in the plaque has also been observed. Again, most studies have been fairly short. [10]

Koll-Klais and team (2005) found a prevalence of *Lactobacillus gasseri* and *L. fermentum* in the oral cavity of healthy individuals compared to those with chronic periodontitis. Further to this, the same researchers have found that lactobacilli inhibit the growth of periodontopathogens, demonstrating the influence of lactobacilli in the oral cavity of a healthy individual. [29]

Riccia, et al (2007) recently studied the anti-inflammatory effects of *Lactobacillus brevis* in a group of patients with chronic periodontitis. The treatment, which involved sucking on lozenges containing *L. brevis* over a period of 4 days, led to improvements in the targeted clinical parameters (plaque index, gingival index, bleeding on probing) for all patients. In that study, a significant reduction in salivary levels of prostaglandin E<sub>2</sub> (PGE<sub>2</sub>) and matrix metalloproteinases (MMPs) was also observed. The authors suggested that the beneficial anti-inflammatory effects of *L. brevis* could be attributed to its capacity to prevent the production of nitric oxide and, consequently, the release of PGE<sub>2</sub> and the activation of MMPs induced by the nitric oxide. However, *L. brevis* may also be antagonistic, leading to a reduction in the quantity of plaque and therefore an improvement in the gingival index. [34]

Shimazaki and colleagues (2008) used epidemiological data to assess the relationship between periodontal health and the consumption of dairy products such as cheese, milk and yogurt. The authors found that individuals, particularly nonsmokers, who regularly consumed yogurt or beverages containing lactic acid exhibited lower probing depths and less loss of clinical attachment than individuals who consumed few of these dairy products. A similar effect was not observed with milk or cheese. By controlling the growth of the pathogens responsible for periodontitis, the lactic acid bacteria present in yogurt would be in part responsible for the beneficial effects observed. [35]

Harini PM and Anegundi RT 2010 found that probiotic mouth rinse was effective in reducing plaque accumulation and gingival inflammation in 6-8 year old children. [36]

### 8.3 Halitosis

There are a number of reasons for the onset of halitosis (bad smelling breath) –consumption of particular foods, metabolic disorders and respiratory tract infections – but commonly it is associated with an imbalance of the commensal microflora of the oral cavity. [37]

An unbalanced oral microflora has been associated with the production of malodorous substances called volatile sulphur compounds (VSCs). These are by-products of microbial degradation of proteins, blood, mucins found in saliva, and traces of food retained on oral surfaces. Kazor and team (2003) looked at the species of bacteria found on the tongue of patients suffering from halitosis and compared the findings with subjects who were considered healthy. The species found to be most associated with halitosis were *Atopobium parvulum*, *Eubacterium sulci*, *Fusobacterium periodonticum*. In the same study, *Streptococcus salivarius* was found to be the most prevalent in the healthy subjects, and this is thought to be due to the capability of *S. salivarius* to produce bacteriocins which could contribute to reducing the number of bacteria that produce VSCs. [38]

Probiotics are marketed for the treatment of both mouth and gut associated halitosis.

### 8.4 Probiotics and *Candida albicans*

*C. albicans* is a leading cause of infection in oral cavity; it is particularly common in the elderly and in immunocompromised patients. Hatakka et al (2007) showed a reduced prevalence of *C. albicans* after taking probiotics in cheese containing *L. rhamnosus* GG and *Propionibacterium freudenreichii*. [39]

Results obtained by Koll et al (2008) when assessing the effects of various *Lactobacillus* strains in oral cavity were markedly different; most strains suppressed growth of periodontal pathogens, including *actinomycescomitans* (60 out of 67 tested strains); *Porphyromona gingivalis* (35 out of 42 strains), *P. intermedia* (26 out of 42 strains), and cariogenic *S. mutans* (37 out of 67 strains). No inhibition was found, however, for *C.albicans* growth. [40]

Hasslöf P et al (2010) investigated the ability of a selection of lactobacilli strains, used in commercially available probiotic products, to inhibit growth of oral mutans streptococci and *C. albicans* in vitro by agar overlay method. At concentrations ranging from 10<sup>9</sup> to 10<sup>5</sup> CFU/ml, all lactobacilli strains inhibited the growth of the mutans streptococci completely with the exception of *L. acidophilus* La5 that executed only a slight inhibition of some strains at concentrations corresponding to 10<sup>7</sup> and 10<sup>5</sup> CFU/ml. All the tested lactobacilli strains reduced candida growth but the effect was generally weaker than for mutans streptococci. The two *L. plantarum* strains and *L. reuteri* ATCC 55730 displayed the strongest inhibition on *Candida albicans*. [41]

### 8.5 Probiotics and HIV

Recently it has been postulated that the probiotic bacteria may slow down AIDS progression. Lin Tao and his colleagues (2008) screened hundreds of bacteria taken from the saliva of volunteers. The results showed that some *Lactobacillus* strains had produced proteins capable of binding a particular type of sugar found on HIV envelope, called mannose. The binding of the sugar enables the bacteria to stick to the mucosal lining of the mouth and digestive tract, forming colonization. One strain secreted abundant mannose-binding protein particles into its surroundings, neutralizing HIV by binding to its sugar coating. They also observed that immune cells trapped by lactobacilli formed a clump. This configuration would immobilize any immune cells harboring HIV and prevent them from infecting other cells. [42]

## 9. Residence time of probiotics in oral cavity

Residence time of probiotics in oral cavity after treatment withdrawal was studied by Çağlar et al (2006) A reduced *S. mutans* level was shown after a two-week use of a *L. reuteri*-enriched yogurt; effects were observed during use and for a few days after discontinuation. [43]

A loss of *L. reuteri* colonization was observed by Wolf et al (1995) two months after having discontinued probiotic use. [44] *L. rhamnosus* GG administration and oral cavity colonization was studied by Yli-Knuutila et al in 2006. The authors concluded that permanent colonization in oral cavity was unlikely (although possible in some cases) and suggested the probiotic to be used on a regular basis. [45]

Binding strength of 17 *Lactobacillus* strains and 7 bifidobacteria strains to saliva and oral mucous membrane was variable in different strains, according to a study by Haukioja et al in 2006, such a strength variation caused an increased residence time of probiotic in oral cavity. [46] Latency time of probiotic *S. salivarius* K12, 4 tablets/day for 3 days, was assessed in several oral cavity areas in a 35-day follow-up, by Horz et al (2007) probiotic could be found on oral mucous membrane, tongue and in stimulated saliva for more than 3 weeks, with a gradually reduced *S. salivarius* K12 level being detected beginning 8 days after treatment withdrawal. [47]

The findings of the studies on oral colonization of probiotics did not suggest that a permanent installation can take place. However, one needs to bear in mind that most studies were conducted in adults and it may be questioned if a permanent installation readily can occur in persons with an already established microflora. [1] Therefore, it seems especially important that further research needs to be carried on infants because it is very likely that the chance of a permanent colonization of probiotics increases with a regular exposure from early childhood. [3]

## 10. Conclusion

The oral cavity with its diversity of microbial species has been shown to harbor strains also distinguished as probiotics as such. Further studies identifying resident probiotics in the mouth, and their eventual effect on the oral environment are required.

Probiotics are a new and interesting field of research in oral microbiology and oral medicine. Bacteriotherapy in the form of probiotics seems to be a natural way to maintain health and protect oral tissues from diseases. But, this area of research in relation to oral health is still in



its infancy and further long term randomized controlled studies are required before definite conclusions are drawn regarding their effective action on oral health.

## 11. References

- [1] Meurman JH. (2005). Probiotics: do they have a role in oral medicine and dentistry? *Eur J Oral Sci*, 113: 188-196.
- [2] Noordin K & Kamin S. (2007). The effect of probiotic mouthrinse on plaque and gingival inflammation. *Ann Dent Univ Malaya*, 14: 19-25.
- [3] Twetman S & Steckslen-Blicks C. (2008). Probiotics and oral health effects in children. *Int J of Ped Dent*, 18: 3-10.
- [4] Metchnikoff E. (1907). Lactic acid as inhibitory intestinal putrefaction. In Chalmers Mitchell P, ed. *The prolongation of life: optimistic studies*. London: Heinemann, 161-183.
- [5] *The Columbia Encyclopedia*, (2001). 6th Ed. New York: Columbia University Press. Available at <http://www.bartleby.com/65/me/Metchnik.html>.
- [6] Meurman JH & Stamatova I. (2007) Probiotics: contributions to oral health. *Oral Diseases*, 13: 443-451.
- [7] Parvez S et al. (2006). Probiotics and their fermented food products are beneficial for health. *J of Appl Microbiol*, 100: 1171-1185.
- [8] Lilly DM & Stillwell RH. (1965) Probiotic growth promoting substances produced by microorganisms. *Science*, 147: 747-748.
- [9] FAO/WHO. (2001). Evaluation of health and nutritional properties of powder milk and live lactic acid bacteria. Cordoba, Argentina: Food and Agriculture Organization of the United Nations and World Health Organization Expert Consultation Report, 1-34.
- [10] Haukioja A. (2010). Probiotics and Oral Health. *Eurp J of Dent*, 4: 348-355.
- [11] Koll-Klais P et al. (2006) Oral lactobacilli in chronic periodontitis: species composition and antimicrobial activity. *IADR Congress, Dublin, 13-16 Sept (Abstract 0081)*
- [12] Isolauri E et al. (1993). Lactobacillus casei strain GG reverses increased intestinal permeability induced by cow milk in suckling rats. *Gastroenterology*, 105: 1643-1650.
- [13] Vanderhoof JA et al. (1999) Lactobacillus GG in the prevention of antibiotic associated diarrhea in children. *J Ped*, 135: 564-568.
- [14] Suvarna VC & Bobby VG. (2005). Probiotics in human health - A current assessment. *Current science*, 88: 1744-48
- [15] Izumita D. (2001). A new approach in dentistry. *Clinical and Basic Medical Research on EM-X--A Collection of Research Papers*, 2:77-81.
- [16] Minna KS, et al. (2002). Lactobacillus bacterium during a rapid increase in probiotic use of L. Rhamnosus GG in Finland. *CID*, 35:1155-60.
- [17] Nase L et al. (2001). Effect of long-term consumption of a probiotic bacterium, Lactobacillus rhamnosus GG, in milk on dental caries and caries risk in children. *Caries Res*, 35: 412-420.
- [18] Ahola AJ et al. (2002). Short-term consumption of probiotic-containing cheese and its effects on dental caries risk factors. *Arch Oral Biol*, 47: 799-804.
- [19] Nikawa H, et al. (2004). Lactobacillus reuteri in bovine milk fermented decreases the oral carriage of mutans streptococci. *Int J Food Microbiol*, 95:219-223.

- [20] Caglar E, et al. (2005). Effect of yoghurt with Bifidobacterium DN-173010 on salivary mutans streptococci and lactobacilli in young adults. *Acta Odont Scand*, 63:317-320.
- [21] Caglar E, et al. (2006). Salivary mutans streptococci and lactobacilli levels after ingestion of the probiotic bacterium *Lactobacillus reuteri* ATCC 55730 by straws or tablets. *Acta Odontol Scand*, 64:314-318.
- [22] Caglar E, et al. (2007). Effect of chewing gums containing xylitol or probiotic bacteria on salivary mutans streptococci and lactobacilli. *Clin Oral Investig*, 11:425-429.
- [23] Caglar E, et al. (2008). Short-term effect of ice-cream containing Bifidobacterium lactis Bb-12 on the number of salivary mutans streptococci and lactobacilli. *Acta Odontol Scand*, 66:154-158.
- [24] Caglar E, et al. (2008). A probiotic lozenge administered medical device and its effect on salivary mutans streptococci and lactobacilli. *Int J Paediatr Dent*, 18:35-39.
- [25] Cildir SK, et al. (2009). Reduction of salivary mutans streptococci in orthodontic patients during daily consumption of yoghurt containing probiotic bacteria. *Eur J Orthod*, 31:407-4011.
- [26] Montalto M, et al. (2004). Probiotic treatment increases salivary counts of lactobacilli: a double-blind, randomized, controlled study. *Digestion*. 69:53-56.
- [27] Comelli EM, et al. (2002). Selection of dairy bacterial strains as probiotics for oral health. *Eur J Oral Sci*, 110(3):218-24.
- [28] Chung J, et al. (2004). Isolation and characterization of *Lactobacillus* species inhibiting the formation of *Streptococcus mutans* biofilm. *Oral Microbiol Immunol*, 19(3): 214-6.
- [29] Koll-Klais P, et al. (2005). Oral Lactobacilli in chronic periodontitis and periodontal health: species composition and antimicrobial activity. *Oral Microbiol Immunol*, 20(6): 354-61.
- [30] Meurman J, et al. (1994). Recovery of *Lactobacillus* strain GG (ATCC 53103) from saliva of healthy volunteers after consumption of yoghurt prepared with the bacterium. *Microb Ecol Health Dis*, 7:295-298.
- [31] Stamatova I, et al. (2009). In vitro evaluation of yoghurt starter lactobacilli and *Lactobacillus rhamnosus* GG adhesion to saliva-coated surfaces. *Oral Microbiol Immunol*. 24:218-223.
- [32] Strahinic I, et al. (2007). Molecular and biochemical characterizations of human oral lactobacilli as putative probiotic candidates. *Oral Microbiol Immunol*, 22(2): 111-7.
- [33] N.S.H. Mehanna et al. (2009). Effect of some probiotic strains and meswak plant on certain oral pathogenic strains. *Int J of Academic Res*, 1(2): 128-132.
- [34] Riccia DN, et al. (2007). Anti-inflammatory effects of *Lactobacillus brevis* (CD2) on periodontal disease. *Oral Dis*, 13(4):376-85.
- [35] Shimazaki Y et al. (2008). Intake of dairy products and periodontal disease: the Hisayama Study. *J Periodontol*, 79(1): 131-7.
- [36] Harini PM & Anegundi RT. (2010). Efficacy of a probiotic and chlorhexidine mouth rinses: A short term clinical study. *JISPPD*, 28(3):179-182.
- [37] Scully G & Greenman J. (2008). Halitosis (breath Odor). *Periodontology 2000*, 48: 667-75.
- [38] Kazor CE, et al. (2003). Diversity of bacterial populations on the tongue dorsa of patients with halitosis and healthy patients. *J. Clin. Microbiol*, 41(2): 558-563.

- [39] Hatakka K, et al. (2007). Probiotics reduces the prevalence of oral candida in the elderly- a randomized controlled trial. *J Dent Res*, 86:125-30.
- [40] Kõll P, et al. (2008). Characterization of oral lactobacilli as potential probiotics for oral health. *Oral Microbiol Immunol*, 23:139-47.
- [41] Hasslöf P et al. (2010). Growth inhibition of oral mutans streptococci and candida by commercial probiotic lactobacilli - an in vitro study. *BMC Oral Health*, 10:18.
- [42] Lin T. (2008). Current opinion in HIV and AIDS, 3:599-602.
- [43] Caglar E, et al. (2006). Salivary mutans streptococci and lactobacilli levels after ingestion of the probiotic bacterium *Lactobacillus reuteri* ATCC 55730 by straws or tablets. *Acta Odontol Scand*, 64:314-8.
- [44] Wolf BW, et al. (1995). Safety and tolerance of *Lactobacillus reuteri* in healthy adult male subjects. *Microb Ecol Health Dis*, 8:41-50.
- [45] Yli-Knuuttila H, et al. (2006). Colonization of *Lactobacillus rhamnosus* GG in the oral cavity. *Oral Microbiol Immunol*, 21:129-31.
- [46] Haukioja A, et al. (2006). Oral adhesion and survival of probiotic and other lactobacilli and bifidobacteria in vitro. *Oral Microbiol Immunol*, 21:326-32.
- [47] Horz HP, et al. (2007). Distribution and persistence of probiotic *Streptococcus salivarius* K12 in the human oral cavity as determined by real-time quantitative polymerase chain reaction. *Oral Microbiol Immunol*, 22:126-30.

IntechOpen



## **Oral Health Care - Pediatric, Research, Epidemiology and Clinical Practices**

Edited by Prof. Mandeep Virdi

ISBN 978-953-51-0133-8

Hard cover, 302 pages

**Publisher** InTech

**Published online** 29, February, 2012

**Published in print edition** February, 2012

Oral health care in pediatric dentistry deals with complete oral health, including preventive aspects for children right from their conception to adolescence, encompassing all the spheres of dentistry including various specialties. It also includes planning a preventive program at individual and community levels. The current research interests in oral health care include studies regarding the role of stem cells, tissue culture, and other ground-breaking technologies available to the scientific community in addition to traditional fields such as anatomy, physiology, and pharmaceuticals etc of the oral cavity. Public health and epidemiology in oral health care is about the monitoring of the general oral health of a community, general afflictions they are suffering from, and an overall approach for care and correction of the same. The oral health care-giver undertakes evaluation of conditions affecting individuals for infections, developmental anomalies, habits, etc. and provides corrective action in clinical conditions. The present work is a compendium of articles by internationally renowned and reputed specialists about the current developments in various fields of oral health care.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Harini Priya Vishnu (2012). Probiotics and Oral Health, Oral Health Care - Pediatric, Research, Epidemiology and Clinical Practices, Prof. Mandeep Virdi (Ed.), ISBN: 978-953-51-0133-8, InTech, Available from: <http://www.intechopen.com/books/oral-health-care-pediatric-research-epidemiology-and-clinical-practices/probiotics-oral-health>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen