

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Partial Mapping of the IL-10 Promoter Region: Identification of New SNPs and Association with Tuberculosis Outcome in Brazilians

L.B. Spinassé¹, M.Q.P. Lopes¹, A.B. Miranda², R.L.F. Teixeira¹,
F.C.Q. Mello³, J.R. Lapa e Silva⁴, P.N. Suffys¹ and A.R. Santos¹

¹Laboratory of Molecular Biology Applied to Mycobacteria,
Oswaldo Cruz Institute, Fiocruz, Rio de Janeiro

²Laboratory of computational Biology and Systems,
Oswaldo Cruz Institute, Fiocruz, Rio de Janeiro

³Medical School – Academic TB Program, Hospital Complex HUCFF-IDT,
Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro

⁴Multidisciplinary Research Laboratory, Hospital Complex HUCFF-IDT,
Federal University of Rio de Janeiro (UFRJ), Rio de Janeiro
Brazil

1. Introduction

Tuberculosis (TB) is one of the oldest infectious diseases which affect humankind. According to the World Health Organization (WHO), in 2009 the TB global burden reached 9.4 million of incident cases, 14 million of prevalent cases and 1.3 million deaths. Most cases were in the South-East Asia and Western Pacific regions. Concerning HIV-TB co-infection, 11-13% of the incident cases were HIV-positive, been the African continent accounted for approximately 80% of these cases (World Health Organization and Global Tuberculosis Programme 2010). Several factors are involved in susceptibility to active disease after infection by *Mycobacterium tuberculosis* including environmental and host genetic markers. Today, the search for functional genetic markers in target genes and association studies with different TB outcomes has been common in the literature (Amim et al, 2007, Calado et al, 2006). Several genes encoding different cytokines, receptors and transporters molecules are involved in the host immune response against *M. tuberculosis*, therefore playing crucial roles in TB susceptibility. The interleukin-10 (IL-10), one of the most important Th2 cytokine involved in the immunoregulation. It is a homodimeric molecule of 37kDa consisting of two monomers of 18.5kDa that interacts with the heterodimeric receptor complex to modulate the biological activities of several cells lineage such as T-cells and the myeloid cells, including monocytes, dendritic cells and macrophages. Interleukin 10 inhibits activation of these cells resulting in a reduced production of pro-inflammatory mediators, including different cytokines and chemokines (Groux et al, 2003; David et al, 2008, Lin et al, 2003; Westendorp et al, 1997).

Variability in IL-10 production has a hereditary component of approximately 75% and this variation is mainly due to polymorphisms within the promoter region of the IL-10 gene (Shin et al 2005; Eskdale et al, 1997).

The human IL-10 coding gene spans about 4.7 kb on chromosome 1q31-32 and contains five exons. It is a highly polymorphic gene, with sixty described single nucleotide polymorphisms (SNPs), spreaded along the promoter, coding and intronic regions (Ensembl Genome Browser April 2011, ENSG00000136634). According to the literature data, the genetic diversity of the IL-10 gene expressed by allele frequencies, varies depends on the ethnicity of the studied population (Moraes et al, 2003). Based in the admixture genetic background of Brazilians, the aim of this work was to map the proximal region of the regulatory sequence (promoter region) of the IL-10 coding gene to look for new SNPs and to evaluate the possible association between IL-10 gene polymorphisms and different TB outcomes after infection with *M. tuberculosis* in Brazilians.

2. Material and methods

2.1 Selection of subjects and sample collection

Four hundred and ninety two consanguineously unrelated individuals from Rio de Janeiro were enrolled in this study. They comprised 221 patients with culture-confirmed active TB (who had been treated at the University Hospital Complex: Thoracic Institute/Clementino Fraga University Hospital from Federal University of Rio de Janeiro-UFRJ and 271 close healthy contacts of TB cases, with no previous TB history and with information of Tuberculin Skin Test (TST) response, (TST+ and TST-) from the same Hospital. This study was approved by the ethics committee of Oswaldo Cruz Foundation.

After a written informed consent a volume of 5 mL of venous blood was collected from each volunteer and stored at -20°C. Genomic DNA was isolated from 100 µL of frozen whole blood using the FlexiGene DNA Kit (Qiagen Inc., USA), according to the manufacturer's specifications. After extraction, DNA samples were stored at -20°C.

2.2 IL-10 genotyping

The partial mapping of the promoter region of IL-10 coding gene was performed by PCR amplification followed by direct sequencing of a 1500pb region upstream of the transcription site. Two sets of primers, EF: 5' CTGTGCCTCAGTTTGCTCAC 3', ER: 5'ACTCTGCTGAAGGCATCTCG3', IF 5'GCAATTTGTCCACGTCACCTG 3' and IR 5'TTGGTTGAACATGAACCTTCTG 3' were used for amplification and sequencing of a DNA fragment of 1001pb (fragment 1) and one set, EF 5' TTCCCAGGTAGAGCAACAC3' and ER5'GGCACATGTTTCCACCTCTT3' for amplification and sequencing of a second DNA fragment of 565pb (fragment 2). Primers were design by using the Primers3 software (<http://frodo.wi.mit.edu/cgi-bin/primer3/primer3 www.cgi>).

The PCR-mediated DNA amplification of the fragment 1 was performed from 100 ng of genomic DNA in a reaction mixture containing 200ng of each primer (IL10frag1) EF and (IL10frag1) ER, 0,2mM of dNTPs, 2,5mM MgCl₂ and 1 U *Taq* DNA polymerase (Invitrogen by Life Technologies, USA) The cycling conditions were: initial denaturation at 94°C for 5 min, followed by 35 cycles of 94°C for 1 min., 70°C for 1 min. and 72°C for 1 min with a final

extension at 72°C for 5 min. For amplification of the fragment 2, 100ng of genomic DNA were used in a similar mixture except for the primers (IL10frag2) EF and (IL10Frag2) ER. Amplification conditions were also the same used for fragment 1 except for the annealing temperature of 63,3°C. Sequencing of the amplified fragments was performed in both DNA strands using a combination of the internal and external primers using ABI PRISM Big Dye Terminator v. 3.1 Kit (PE Applied BioSystems), according to the manufacturer's recommendations, on an ABI PRISM 3730 DNA Analyser (PE Applied BioSystems). All singletons and even new/rare mutation identified were confirmed by PCR and re-amplification for three times followed by re-sequencing.

2.3 Computational analysis

The SNPs identification in each individual sample was achieved after alignment with the reference sequences AF418271 and Z30175 for the fragments 1 and 2 (<http://www.ncbi.nlm.nih.gov/GenBank>). We have defined the DNA sequence of the transcription starting site of the human IL-10 gene from the first nucleotide immediately preceding the A of the ATG taken position -1 (Eskdale et al, 1997), through SeqScape v.s2.6 software (Applied Biosystem www.appliedbiosystems.com). Haplotype reconstruction were achieved through the of Software PHASE v.s2.1.1, development by Yongtao Guan and Matthew Stephens from University of Chicago department of human genetics and department of Statistics.

2.4 Statistical analysis

Pair-wise linkage disequilibrium was also tested for the loci studies. The Hard-Weinberg equilibrium using χ^2 test. Statistics were performed in XLSTAT 2008.7 (Addinsoft Software Inc - New York USA). The magnitude of the associations was estimated by odds ratio values. All tests were performed at the 0.05 level of significance. Epi Info version 3.5.1 2008, (Centers for Disease Control and Prevention, USA) regarding the value of 5% ($p < 0.05$) as threshold for statistical significance in the association study of haplotypes between the numbers observed and expected separately in patients and contacts.

3. Results

Two epidemiological model of study were adopted in this work and the study population was stratified according the different models. Expected genotype frequencies were calculated from respective single allele frequencies and were consistent with Hard Weinberg Equilibrium using χ^2 test.

3.1 SNPs profile of the human IL-10 promoter among Brazilian subjects

Using the direct PCR sequencing approach, analysis of the 492 DNA samples enrolled in this study revealed the presence of thirteen SNPs within *IL-10* promoter region, seven of which not reported yet and located at positions (-537G>T, -633G>A, -637G>C, -750C>T, -840A>C, -1162A>G/T and -1189C>T) respectively. The six remaining SNPs found, already deposited in GenBank-Entrez SNP database, were located at positions (-1117G>A, reported as -1082, -886G>A, -854C>T, reported as -819, -692G>A, -627C>A, reported as -592

Locus	Genotype	Subjects (N=492)	Absolute Frequency	Allele Frequency
-1189*	CC	483	0,982	0,0090
	CT	9	0,018	
	(fa) T	9	-	
-1162*	AA	485	0,986	0,0060 (G) 0,0030 (T)
	AG	2	0,004	
	AT	3	0,006	
	GG	2	0,004	
	(fa) T	3	0,006	
	(fa) G	4	0,008	
-1117	GG	55	0,112	0,6780
	GA	207	0,421	
	AA	230	0,467	
	(fa) A	667	0,888	
-886	GG	464	0,943	0,0295
	GA	27	0,055	
	AA	1	0,002	
	(fa) A	29	0,057	
-854	CC	198	0,402	0,3700
	CT	224	0,455	
	TT	70	0,142	
	(fa) T	364	0,597	
-840*	AA	484	0,984	0,0080
	AC	8	0,016	
	(fa) C	8	-	
-750*	CC	473	0,960	0,0213
	CT	17	0,034	
	TT	2	0,004	
	(fa) T	21	0,038	
-692	GG	475	0,965	0,0170
	GA	17	0,035	
	(fa) A	17	-	
-637*	CC	487	0,989	0,0050
	CG	5	0,011	
	(fa) G	5	-	
-633*	GG	490	0,996	0,0020
	GA	2	0,004	
	(fa) A	2	-	
-627	CC	192	0,390	0,3770
	CA	229	0,465	
	AA	71	0,144	
	(fa) A	371	0,610	
-537*	GG	491	0,998	0,0010
	GT	1	0,002	
	(fa) T	1	-	

Locus	Genotype	Subjects (N=492)	Absolute Frequency	Allele Frequency
-464	GG	479	0,973	0,0130
	GT	13	0,026	
	(fa) T	13		

Table 1. Genotype and allele frequencies of SNPs within *IL-10* promoter in Brazilians from Rio de Janeiro.

and -464G>T. Upon genotype and allele frequency analysis, the three more frequent SNPs were the ones located at positions -627C>A, -854C>T and -1117G>A (Table1). Analysis of linkage disequilibrium (LD) showed that only two, SNPs -627C>A and -854C>T out of the thirteen SNPs identified in the total sample were in LD.

3.2 Haplotypes characterization

Haplotypes	-1189	-1162	-1117	-886	-854	-840	-750	-692	-637	-633	-627	-537	-464	Alleles	Frequency
1	C	A	G	G	C	A	C	G	C	G	C	G	G	265	0,270
2	T	A	G	G	C	A	C	G	C	G	C	G	G	2	0,002
3	C	T	G	G	C	A	C	G	C	G	C	G	G	2	0,002
4	C	A	A	G	C	A	C	G	C	G	C	G	G	274	0,280
5	C	A	G	A	C	A	C	G	C	G	C	G	G	28	0,028
6	C	A	A	G	T	A	C	G	C	G	C	G	G	10	0,010
7	C	A	A	G	C	C	C	G	C	G	C	G	G	2	0,002
8	C	A	G	G	C	A	T	G	C	G	C	G	G	6	0,006
9	C	A	A	G	C	A	T	G	C	G	C	G	G	2	0,002
10	C	A	G	G	C	A	C	A	C	G	C	G	G	2	0,002
11	C	A	A	G	C	A	C	A	C	G	C	G	G	2	0,002
12	C	A	G	G	C	A	C	G	G	G	C	G	G	2	0,002
13	C	A	G	G	C	A	C	G	C	G	A	G	G	1	0,001
14	C	A	A	G	C	A	C	G	C	G	A	G	G	11	0,011
15	C	A	G	G	T	A	C	G	C	G	A	G	G	2	0,002
16	C	A	A	G	T	A	C	G	C	G	A	G	G	311	0,316
17	T	A	A	G	T	A	C	G	C	G	A	G	G	7	0,007
18	C	T	A	G	T	A	C	G	C	G	A	G	G	1	0,001
19	C	A	A	G	T	C	C	G	C	G	A	G	G	6	0,006
20	C	A	G	G	C	A	T	G	C	G	A	G	G	1	0,001
21	C	A	A	G	T	A	T	G	C	G	A	G	G	10	0,010
22	C	A	G	G	C	A	C	A	C	G	A	G	G	2	0,002
23	C	A	G	A	C	A	C	A	C	G	A	G	G	1	0,001
24	C	A	A	G	T	A	C	A	C	G	A	G	G	10	0,010
25	C	A	A	G	T	A	C	G	G	G	A	G	G	2	0,002
26	C	A	A	G	T	A	C	G	C	A	A	G	G	2	0,002
27	C	A	G	G	C	A	C	G	C	G	C	T	G	1	0,001
28	C	A	A	G	C	A	C	G	C	G	C	G	T	7	0,007
29	C	A	G	G	C	A	T	G	C	G	C	G	T	2	0,002
30	C	A	A	G	T	A	C	G	C	G	A	G	T	3	0,003
31	C	A	A	G	C	A	C	G	G	G	A	G	T	1	0,001
32	C	G	A	G	C	A	C	G	C	G	C	G	G	6	0,006

Table 2. Haplotypes description and frequencies

3.3 Association of the IL-10 SNPs and TB outcomes

Frequencies of the *IL-10* polymorphisms in patients affected by active tuberculosis (n=221) and healthy controls TST+ (n=271) were compared among stratified groups to evaluate the possible association of the identified SNPs and the outcomes of susceptibility *per ser* to the occurrence of active TB, and disease severity. Table 3 shows that the most studied and well characterized SNPs at positions -1117, -854, -627, (also reported respectively as -1082, -819 and -592), did not show any association with the studied TB outcomes. Additionally, any of the remaining 10 SNPs showed to be associated with TB occurrence at neither genotype nor allele level.

Locus	Genotype	Patients N=221	Controls TST+ N= 126	χ^2	<i>p</i> -valor	OR																																																																																																																																						
-1189	CC	212	126	5,26	0,022	#																																																																																																																																						
	CT	9	0				-1162	AA	217	125	1,16	0,763	#	AG	1	0	AT	2	1	GG	1	0	-1117	GG	24	12	1,20	0,547	#	GA	100	51	AA	97	63	-886	GG	208	123	2,23	0,135	0,39	GA	13	3	-854	CC	89	45	1,67	0,432	#	CT	100	66	TT	32	15	-840	AA	213	126	ND	ND	ND	AC	8	0	-750	CC	221	114	ND	ND	ND	CT	0	10	TT	0	2	-692	GG	215	121	0,41	0,521	1,48	GA	6	5	-637	CC	216	126	ND	ND	ND	CG	5	0	-633	GG	221	125	ND	ND	ND	GA	0	1	-627	CC	85	44	0,95	0,619	#	CA	102	65	AA	34	17	-537	GG	220	126	ND	ND	ND	GT	1	0	-464	GG	211	123	1,02	0,311
-1162	AA	217	125	1,16	0,763	#																																																																																																																																						
	AG	1	0																																																																																																																																									
	AT	2	1																																																																																																																																									
	GG	1	0																																																																																																																																									
-1117	GG	24	12	1,20	0,547	#																																																																																																																																						
	GA	100	51																																																																																																																																									
	AA	97	63																																																																																																																																									
-886	GG	208	123	2,23	0,135	0,39																																																																																																																																						
	GA	13	3																																																																																																																																									
-854	CC	89	45	1,67	0,432	#																																																																																																																																						
	CT	100	66																																																																																																																																									
	TT	32	15																																																																																																																																									
-840	AA	213	126	ND	ND	ND																																																																																																																																						
	AC	8	0																																																																																																																																									
-750	CC	221	114	ND	ND	ND																																																																																																																																						
	CT	0	10																																																																																																																																									
	TT	0	2																																																																																																																																									
-692	GG	215	121	0,41	0,521	1,48																																																																																																																																						
	GA	6	5																																																																																																																																									
-637	CC	216	126	ND	ND	ND																																																																																																																																						
	CG	5	0																																																																																																																																									
-633	GG	221	125	ND	ND	ND																																																																																																																																						
	GA	0	1																																																																																																																																									
-627	CC	85	44	0,95	0,619	#																																																																																																																																						
	CA	102	65																																																																																																																																									
	AA	34	17																																																																																																																																									
-537	GG	220	126	ND	ND	ND																																																																																																																																						
	GT	1	0																																																																																																																																									
-464	GG	211	123	1,02	0,311	0,514																																																																																																																																						
	GT	10	3																																																																																																																																									

Table 3. Genotype distribution of the *IL-10* SNPs among TB patients and healthy controls (TST+)

Locus	Genotype	Pacients N=221	Controls (TST+/TST-) N= 271	χ^2	p-valor	OR																																																																																																																																									
-1189	CC	212	271	ND	ND	ND																																																																																																																																									
	CT	9	0				-1162	AA	217	269	0,621	0,892	#	AG	1	1	AT	2	1	GG	1	1	-1117	GG	24	31	1,699	0,428	#	GA	100	107	AA	97	133	-886	GG	208	256	0,931	0,638	#	GA	13	14	AA	0	1	-854	CC	89	109	0,025	0,988	#	CT	100	124	TT	32	38	-840	AA	213	271	ND	ND	ND	AC	8	0	-750	CC	221	252	ND	ND	ND	CT	0	17	TT	0	2	-692	GG	215	260	0,659	0,417	1,51	GA	6	11	-637	CC	216	271	ND	ND	ND	CG	5	0	-633	GG	221	269	1,637	0,200	#	GA	0	2	-627	CC	85	107	0,299	0,861	#	CA	102	127	AA	34	37	-537	GG	220	271	ND	ND	#	GT	1	0	-464	GG	211	268	5,528	0,019
-1162	AA	217	269	0,621	0,892	#																																																																																																																																									
	AG	1	1																																																																																																																																												
	AT	2	1																																																																																																																																												
	GG	1	1																																																																																																																																												
-1117	GG	24	31	1,699	0,428	#																																																																																																																																									
	GA	100	107																																																																																																																																												
	AA	97	133																																																																																																																																												
-886	GG	208	256	0,931	0,638	#																																																																																																																																									
	GA	13	14																																																																																																																																												
	AA	0	1																																																																																																																																												
-854	CC	89	109	0,025	0,988	#																																																																																																																																									
	CT	100	124																																																																																																																																												
	TT	32	38																																																																																																																																												
-840	AA	213	271	ND	ND	ND																																																																																																																																									
	AC	8	0																																																																																																																																												
-750	CC	221	252	ND	ND	ND																																																																																																																																									
	CT	0	17																																																																																																																																												
	TT	0	2																																																																																																																																												
-692	GG	215	260	0,659	0,417	1,51																																																																																																																																									
	GA	6	11																																																																																																																																												
-637	CC	216	271	ND	ND	ND																																																																																																																																									
	CG	5	0																																																																																																																																												
-633	GG	221	269	1,637	0,200	#																																																																																																																																									
	GA	0	2																																																																																																																																												
-627	CC	85	107	0,299	0,861	#																																																																																																																																									
	CA	102	127																																																																																																																																												
	AA	34	37																																																																																																																																												
-537	GG	220	271	ND	ND	#																																																																																																																																									
	GT	1	0																																																																																																																																												
-464	GG	211	268	5,528	0,019	4,23																																																																																																																																									
	GT	10	3																																																																																																																																												

Table 4. Genotype frequencies of the *IL-10* SNPs among TB patients and healthy controls (TST+/TST-)

Genetic diversity of the *IL-10* gene, mainly in the promoter region has been studied in several populations and many studies have been associated polymorphisms in this region with differential *IL-10* production. Among the sixty already described SNPs, the ones at positions -1117 G>A, -854C>T, -627C>A, also described as (-1082, -819 and -592) are the better characterized and taken as haplotypes are related to *IL-10* production. The allele G at -1117, and haplotypes containing this allele have been associated with high *IL-10* production, while the allele A and the haplotype ATA have been associated with low *IL-10* production. Even so, association studies have been shown conflicting results depending on several factors,

including differences in laboratory experiments and ethnicity of the studied populations. As a result, different kinds of associations with immune response and diseases outcomes are commonly reported. Here, no association was found of any between the already described SNPs within IL-10 promoter with TB. This result corroborates at least in part with previous association study of these well characterized SNPs with leprosy, in which, only the only the -854C>T (-819) showed associated with leprosy susceptibility in Brazilians residents in Rio de Janeiro (Santos et al, 2002). No data concerning TB association are available in Brazil.

Locus	Genótipo	TBP (N=139)	TBE (N=43)	χ^2	p-valor	OR																																																																																																																			
-1189	CC	136	39	4,532	0,033	0,215																																																																																																																			
	CT	3	4				-1162	AA	137	42	3,854	0,278	#	AG	0	1	AT	1	0	-1117	GG	1	0	2,619	0,270	#	GA	19	2	AA	62	21	-886	GG	58	20	1,089	0,297	#	GA	130	42	-854	CC	9	1	1,193	0,551	#	CT	61	16	TT	58	22	-840	AA	20	5	0,010	0,565	0,640	AC	134	42	-692	GG	5	1	ND	ND	ND	GA	134	43	-637	CC	5	0	1,577	0,209	0,290	CG	137	41	-627	CC	2	2	1,802	0,406	#	CA	60	14	AA	59	23	-537	GG	20	6	ND	ND	ND	GT	139	42	-464	GG	0	1	2,274	0,132	0,363	GT	134	39			5
-1162	AA	137	42	3,854	0,278	#																																																																																																																			
	AG	0	1																																																																																																																						
	AT	1	0																																																																																																																						
-1117	GG	1	0	2,619	0,270	#																																																																																																																			
	GA	19	2																																																																																																																						
	AA	62	21																																																																																																																						
-886	GG	58	20	1,089	0,297	#																																																																																																																			
	GA	130	42																																																																																																																						
-854	CC	9	1	1,193	0,551	#																																																																																																																			
	CT	61	16																																																																																																																						
	TT	58	22																																																																																																																						
-840	AA	20	5	0,010	0,565	0,640																																																																																																																			
	AC	134	42																																																																																																																						
-692	GG	5	1	ND	ND	ND																																																																																																																			
	GA	134	43																																																																																																																						
-637	CC	5	0	1,577	0,209	0,290																																																																																																																			
	CG	137	41																																																																																																																						
-627	CC	2	2	1,802	0,406	#																																																																																																																			
	CA	60	14																																																																																																																						
	AA	59	23																																																																																																																						
-537	GG	20	6	ND	ND	ND																																																																																																																			
	GT	139	42																																																																																																																						
-464	GG	0	1	2,274	0,132	0,363																																																																																																																			
	GT	134	39																																																																																																																						
		5	4																																																																																																																						

The influence of demographic characteristics such as gender and age in the studied outcomes were also evaluated, no significant difference was found for gender or age. (data not shown).

Table 5. Genotype distribution of the IL-10 SNPs among patients with pulmonary and extrapulmonary tuberculosis (TBP and TBE)

When genotype association was evaluated by comparison of the frequencies of TB patients against healthy controls without stratification by TST response (TST+/TST-), a significant increased frequency of the heterozygous -464 GT, showed to be associated with susceptibility *per se* to the occurrence of active TB (OR 4,23, *p* value 0,019). An interesting finding was the observation of the SNPs -750C>T and -633G>A, only the group of health

controls and on the contrary, the SNPs at positions -1189C>T, -840A>C, -637C>G and -537G>T, present only in the group of TB patients (Table 4).

Assessment of the possible evaluation of the *IL-10* SNPs found with the outcome of disease severity, were achieved by comparing the frequencies found in the group of pulmonary TB patients (TBP) and the group of extrapulmonary TB patients (TBE). Table 5 shows a significantly higher frequency of the allele variant -1189T in the group of patients with extrapulmonary TB in comparison with pulmonary TB suggesting an association of this allele variant with the outcome of severity. No statistically significant differences between the groups were observed for the other SNPs evaluated.

After haplotyping reconstruction and identification of thirty two different haplotypes, only the three more frequent, respectively, haplotype 1 (CAGGCACGCGCGG), haplotype 4 (CAAGCACGCGCGG) and haplotype 16 (CAAGTACGCGAGG) with frequencies of 27, 28 and 31.6 % respectively were considered for the association study. No haplotype association was found for any of the evaluated outcomes.

4. Conclusions

In this study, using a PCR direct sequencing approach, seven new SNPs and six already described were identified, in a region of one thousand and fifty hundred base pairs, proximal to the transcription starting site of *IL-10* coding gene.

Evaluation of the possible association of the identified SNPs with different TB outcomes by comparison of allelic, genotype and haplotype frequencies between groups showed that none of the new SNPs were associated neither with susceptibility nor with severity. However, an interesting finding is that three of the new SNPs were present only among patients and one, only in the group of controls. Although it can suggest an association with susceptibility *per se* to TB occurrence and resistance to TB respectively, it is not conclusive. The functional role of these SNPs should be elucidated and sample size should also be increased.


5. Acknowledgements

This work was supported by Faperj/Pronex: Proc: E-26/170.0003/2008 and FAPERJ Pensa Rio: E-26/110.288/2007. Afrânio Lineu Kritski and Fernanda C. Q. Mello, and Philip Noel Suffys are supported by CNPq Grant numbers 308786/2005-0, 35 04 77/1995-7, 306702/2007-0 and 312165/2006-4 respectively.

6. References

- Amim, L.H.; Pacheco, A.G.; Fonseca-Costa, J.; Loredó, C.S.; Rabahi, M.F.; Melo, M.H.; Ribeiro, F.C.; Mello, F.C.; Oliveira, M.M.; Lapa e Silva, J.R.; Ottenhoff, T.H.; Kritski, A.L.; Santos, A.R (2007). Role of IFN-gamma +874 T/A single nucleotide polymorphism in the tuberculosis outcome among Brazilians subjects. *Mol Biol Rep* 35(4):563-6, Rio de Janeiro. Brazil.
- Asadullah, K.; Sabat, R.; Wiese, A.; Döcke, W.D; Dieter, W.D; Volk, H.D; Sterry, W. (1999). Interleukin-10 in cutaneous disorders: implications for its pathophysiological importance and therapeutic use. *Arch Dermatol Res*; 291 :628–636, Charité, Germany

- Ates, Ö.; Musselim, B.; Ongen, G.; Topal Sarikaya, A. (2007). Interleukin-10 and Tumor Necrosis Factor α Gene polymorphism in Tuberculosis. *J Clin Immunol*; 9155-2, Istanbul, Turkey.
- Bellamy, R. (2006). Genome-wide approaches to identifying genetic factors in host susceptibility to tuberculosis. *V Microbes Infect*; 1119-23. Middlesbrough, UK.
- Bellamy, R.; Ruwende, C.; Corrah, T.; Mc Adam, KP.; Whittle, HC.; Hill, AV. (1998). Assessment of the interleukin-1 gene cluster and other candidate gene polymorphisms in host susceptibility to tuberculosis. *Tuber Lung Disease*: 79:83-9, Oxford, UK.
- Calado, D.P.; Paixão, T.; Holmberg, D.; Haury, M. (2006). Stochastic monolallelic expression of IL-10 in T Cells. *J Immunol*; 177: 5358-5364, Oeiras, Portugal.
- David, M. Mosser & Xia, Zhang (2008). Interleukin-10: new perspectives on an old cytokine. *Immunological Reviews*; 226: 205-218, College Park, USA.
- de Waal Malefyt, R.; Abrams, J.; Bennett, B.; Figdor, C.G.; de Vries, J.E. (1991). Interleukin 10(IL-10) inhibits cytokine synthesis by human monocytes: an autoregulatory role of IL-10 produced by monocytes. *J Exp Med*; 174:1209, Palo Alto, California, USA.
- Eskdale, J.; Kube, D.; Tesch, H.; Gallagher, G. (1997). Mapping of the human IL-10 gene and further characterization of the 5' flanking sequence. *Immunogenetics*; 46: 120-128, Glasgow, Scotland.
- Groux, H. & Cottrez, F. (2003). The complex role of interleukin-10 in autoimmunity. *Journal of Autoimmunity* 20: 281-285, Nice, France.
- López-Maderuelo, D.; Arnalich, F.; Serantes, R.; González, A.; Codoceo, R.; Madero, R.; Vázquez, J.J.; Montiel, C (2003). Interferon-gamma and interleukin-10 gene polymorphisms in pulmonary tuberculosis. *Am J Respir Crit Care Med*; 167:970-975, Madrid, Spain.
- Lin, MT.; Storer, B.; Martin, P.J.; Tseng, L.H.; Googley, T.; Chen, P.J.; Hansen, J.A. (2003). Relation of an interleukin-10 promoter polymorphism to graft-versus-host disease and survival after hematopoietic-cell transplantation. *N Engl J Med*; 349: 2201-2210, Seattle, USA.
- Opdal, H.; Opstad, A.; Vege, A.; Rognum, T.O (2003). IL-10 Gene Polymorphisms Are Associated With Infectious Cause of Sudden Infant Death. *Human* 64: 1183-1189, Oslo, Norway.
- Peguet-Navarro, J.; Moulon, C.; Caux, C.; Dalbiez-Gauthier, C.; Banchereau, J.; Schmitt, D.; (1994). Interleukin-10 inhibits the primary allogeneic T cell response to human epidermal Langerhans cells. *Eur J Immunol*; 24:884, Lyon, France.
- Pyo, C.W.; Hur, S.S.; Kim, Y.K.; Choi, H.B.; Hong, Y.S.; Kim, D.W.; Kim, C.C.; Kim, H.K.; Kim, T.G. (2003). Polymorphisms of IL-1B, IL-1RN, IL-2, IL-4, IL-6, IL-10, and IFN- γ genes in the Korean population. *Hum Immunol.*; 64:979-89, Seoul, South Korea.
- Santos, A.R.; Suffys, P.N.; Vanderborght, P.R.; Moraes, M.O.; Vieira, L.M.; Cabello, P.H.; Bakker, A.M.; Matos, H.J.; Huizinga, T.W.; Ottenhoff, T.H.; Sampaio, E.P.; Sarno, E.N.; (2002). Role of tumor necrosis factor-alpha and interleukin-10 promoter gene polymorphisms in leprosy. *J Infect Dis*; 1;186(11):1687-91, Rio de Janeiro, Brazil.
- Shin, H.D.; Park, B.L.; Kim, Y.H.; Cheong, H.S.; Lee, I.H.; Park, S.K.; (2005). Common interleukin-10 polymorphisms associated with decreased risk of tuberculosis. *Exp Mol Med*; 37(2):128-32, Seoul, Korea.
- Westendorp, R.G.; Langermans, J.A.; Huizinga, T.W.; Elouali, A.H.; Vervej, C.L.; Boomsma, D.I.; (1997). Genetic influence on cytokine production and fatal meningococcal Disease. *Lancet* ; 349 170-173, Leiden , Netherlands
Site: <http://www.ensembl.org/index.html#ENSG00000136634>



Understanding Tuberculosis - Analyzing the Origin of Mycobacterium Tuberculosis Pathogenicity

Edited by Dr. Pere-Joan Cardona

ISBN 978-953-307-942-4

Hard cover, 560 pages

Publisher InTech

Published online 24, February, 2012

Published in print edition February, 2012

Mycobacterium tuberculosis in an attempt to understand the extent to which the bacilli has adapted itself to the host and to its final target. On the other hand, there is a section in which other specialists discuss how to manipulate this immune response to obtain innovative prophylactic and therapeutic approaches to truncate the intimal co-evolution between Mycobacterium tuberculosis and the Homo sapiens.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

L.B. Spinassé, M.Q.P. Lopes, A.B. Miranda, R.L.F. Teixeira, F.C.Q. Mello, J.R. Lapa e Silva, P.N. Suffys and A.R. Santos (2012). Partial Mapping of the IL-10 Promoter Region: Identification of New SNPs and Association with Tuberculosis Outcome in Brazilians, Understanding Tuberculosis - Analyzing the Origin of Mycobacterium Tuberculosis Pathogenicity, Dr. Pere-Joan Cardona (Ed.), ISBN: 978-953-307-942-4, InTech, Available from: <http://www.intechopen.com/books/understanding-tuberculosis-analyzing-the-origin-of-mycobacterium-tuberculosis-pathogenicity/partial-mapping-of-the-il-10-promoter-region-identification-of-new-snps-and-association-with-tubercu>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen