

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Essential Oils of Umbelliferae (Apiaceae) Family Taxa as Emerging Potent Agents for Mosquito Control

Epameinondas Evergetis¹, Antonios Michaelakis²
and Serkos A. Haroutounian^{1,*}

¹*Chemistry Laboratory, Agricultural University of Athens, Athens*

²*Department of Entomology, Benaki Phytopathological Institute, Athens
Greece*

1. Introduction

Warm-humid areas around the globe constitute the cradle of humanity, providing their inhabitants the most favorable environments for living and agricultural production. In this “Garden of Eden”, which spreads within the globe’s temperate and tropical zones, is also thriving an annoying but dangerous daemon, the mosquito. This little devil constitutes the main vector of malaria and human encephalitis, both infectious diseases that account as major threats of public health (Becker et al., 2003). Recently, these threats have been spread to a broader geographical area, as a consequence of their vectors (*Aedes* sp., *Anopheles* sp. and *Culex* sp.) introduction into metropolitan areas of northern hemisphere, such as Chicago (Tedesco et al., 2010), New York (Peterson et al., 2006) and Paris (Delaunay et al., 2009). Since mosquito breeding habitats in both urban and rural areas are man-made (Imbahale et al., 2010), there are several restrictions limiting the efforts towards the development of an integrated vector management system. To date, the history of evolutions of malaria vector interventions is directly connected with the mosquito control tools development, concerning either environmental modifications/manipulations or their chemical and/or biological control (Kilama, 2009).

In respect the chemical control, a significant milestone was the DichloroDiphenyl-Trichloroethane (DDT) synthesis by Zeidler in 1874. The DDT success was followed by the fast introduction of numerous chlorinated hydrocarbons, which were used in massive amounts for the control of mosquito-borne diseases (Ray, 2010). Despite their efficiency, the use of organochlorines had severe environmental impacts which were publicly (and dramatically) addressed by Carson (1962) in *Silent Spring*, initiating the development of insecticide resistant mosquito populations. These undesirable characteristics, in combination with concerns on public health risks, derived from the organochlorine residues detected in humans and animals, led to their ban in early 70’s. Thus, they were replaced by less persistent chemicals, such as organophosphates, pyrethroids and avermectin derivatives,

*Corresponding Author

substitutes that also display the major disadvantage of resistance development (Alves et al., 2010; Daaboub et al., 2008; Lima et al., 2011).

Recent research trend on mosquito chemical control mainly focuses on currently used compounds, aiming to enhance their potency and circumvent the problems connected with their application. In this respect, the so far developed pyrazole derivatives are quite efficient exhibiting however adverse environmental effects (Stevens et al., 2011), while the corresponding pyrroles display the desirable efficiency (Raghavendra et al., 2011) but adequate research on their environmental side effects is still underway. Amides, such as methazolamide and acetazolamide, were also evaluated as potent mosquito larvicides but were found to display significant bioaccumulation properties (Del Pilar Corena et al., 2006) which discourage their broad use. Finally, various novel pyrimido-quinolone molecules have been developed and assessed as highly toxic for other organisms (Rajanarendar et al., 2010). Ray (2010) recognized that the insecticide treated nets, in connection with the long lasting insecticidal nets, have resurrected the chemical control of malaria's mosquito vector. This may be rationalized considering that their targeted application resolves the problems connected with the environmental impacts of chemical control agents since limit their expansion, availability and environmental penetration.

Despite the numerous efforts and progress achieved, the efficacy of insecticidal nets in malaria prevention still constitutes a hot issue, since depends strongly upon a plethora of additional factors (Killeen & Smith, 2007). In particular, despite efforts (Pennetier et al., 2010) to overcome the recognized for longtime resistance development issues of insecticidal nets, todate these problems have not been resolved (Yadouleton et al., 2011). An additional drawback derives by the combined impact of herbicide application that promotes the cross-resistance to mosquito populations (Boyer et al., 2006; Riaz et al., 2009).

The corresponding biological control has dictated the development of novel-alternative mosquito control tools, including the sterile males technique (Patersson et al., 1968), the genetically modified mosquitoes (Gu et al., 2011; Lavalie-Defaix et al., 2011), the entomopathogenic fungi (Van Breukelen et al., 2011; Kanzok & Jacobs-Lorena, 2006) and bacteria. Among the tools developed the bacterial pathogens application is considered as the most prominent intervention, displaying species selective insecticidal ability (Hayes et al., 2011) which is considered as an efficient means for mosquito control without harmful impacts for the environment (Caquet et al., 2011). Major thresholds limiting the wider application of this technique are related with the induced pathogen introduction among the natural mosquito populations (Hancock et al., 2011) and the threats connected with bioaccumulation and resistance development (Tilquin et al., 2008). In general, the biological control tools are still under development, presenting todate a low degree of maturity for large-scale interventions.

Temephos was considered as one of the most potent-safe insecticides. Its recent exclusion from Annex I of the Directive 98/8/EC resulted in the discontinuation of its application in mosquito control programs by the European, emerging the development and use of new-safer insecticides. Thus, relative research directed towards the discovery-development of novel molecules, capable to control the mosquito populations without exhibiting the disadvantages of synthetic pesticides. In this respect, the plant originated natural compounds constitute a large deposit of such molecules, inherently allowing the retrieval of

various commercially successful molecules like pyrethrins. To date, the search for novel, potent and safer pesticides from this deposit has already provided several candidates, either as pure compounds and/or their extracts. Specifically, various organic acids such as lactic and orthophosphoric acids (Chakraborty et al., 2010), alkaloids (Talontsi et al., 2011) and plant proteins (Chowdhury et al., 2008) have been identified as efficient mosquito control agents. Furthermore, several plants were used as the maternal material to produce bio-products which were applied against mosquitoes with hopeful results (Shaan et al., 2005; Sukumar et al., 1991). On the other hand, the plant derived Essential Oils (EOs) constitute a special category of natural products that exhibit the major advantage -for the mosquito control endeavor- of exhibiting an insect oriented mode of action with low penetrability to the ecosystems that does not affect larger animals. In addition, the natural diversity of their constituents addresses effectively the problem of resistance development (Isman, 2000).

2. Literature review

2.1 Umbelliferae (Apiaceae) family: A source of potent natural agrochemicals

Many EOs originated from diverse plant families have been considered and studied as potential sources of natural agrochemicals. In this respect, previous research results on Umbelliferae (Apiaceae) family plant materials revealed the significant acaricidal activities of butyridenepthalides isolated from *Angelica acutiloba* Kitagawa var. *sugiyame* Hikino (Kwon & Ahn, 2002) and the similar activity of the EO of *Foeniculum vulgare*, attributed to the presence of *p*-anisaldehyde and (+,-)-fenchone in the EO (Lee, 2004). These EOs were practically inactive in fumigant toxicity tests against *Lycoriella mali* though they are known to contain the active monoterpenes α -pinene and β -pinene (Choi et al., 2006), which are common constituents of many Umbelliferae EOs. Methanolic extracts of *Angelica dahurica*, *Cnidium officinale*, and *Foeniculum vulgare* were also tested against the Coleoptera *Lasioderma sericorne*, *Sitophilus oryzae* and *Callosobruchus chinensis* exhibiting a moderate activity only the second extract (Kim et al., 2003a; Kim et al., 2003b). Other EOs of this family screened as inactive against coleoptera were originated from the species *Anethum graveolens* L., *Apium graveolens* Houtt., *Coriandrum sativum* L., *Cuminum cyminum* L. and *Petroselinum sativum* L. (Regnault-Roger & Hamraoui, 1994; Papachristos & Stamopoulos, 2002). On the contrary, the EOs of *Pimpinella anisum* L. and *Cuminum cyminum* L. displayed excellent ovicidal and insecticidal activities against the *Tribolium confusum* du Val and the *Ephesia kuehniella* Zeller (Tunc et al., 2000). In addition, the aqueous extract of *Pimpinella anisum* exhibited good repellent effect against the adults of sweet potato whitefly *Bemisia tabaci* (Ateyyat et al., 2007).

These rather controversial results are not connected with the impressive activities that Umbelliferae EOs were found to exhibit against the Diptera, with the EO of *Ammi visnaga* displaying -among 19 EOs- the most potent ovicidal activity against *Mayetiola destructor* (Lamiri et al., 2001). In addition, tests against *Drosophila melanogaster* of furanocoumarins and pthalides isolated from *Angelica acutiloba* Kitagawa var. *sugiyame* Hikino revealed the hypothesis that the insecticidal properties of the plant extracts are connected with the acetylcholinesterase inhibition (Miyazawa et al., 2004). Finally, alkylpthalides originated from *Cnidium officinale* Makino were tested as extremely effective against *Drosophila melanogaster* (Tsukamoto et al., 2005).

2.2 Umbelliferae (Apiaceae) family: A strong focal point for mosquito control

Table 1 summarizes the test results against various mosquito species reported for all extracts and EOs derived from plants belonging to the Umbelliferae family. Same table also contains the test results of fourteen EOs, which appear herein for the first time. Results indicate that the organic phase of the *Cryoptaenia canadensis* extract is the most active against fourth instars of *Culex pipiens*, leading to the isolation –from the extract- of the acetylated very toxic (LC₅₀ values lower than 10 mg l⁻¹) molecules of falcarinol and falcarindiol (Eckenbach et al., 1999). The larvicidal properties of hexane soluble fraction of *Apium graveolens* seeds -a plant with pleasant aroma- and three isolated compounds (sedanolide, senkyunolide-N, senkyunolide-J) against *Aedes aegypti* mosquitoes highlighted sedanolide as very active (100% mortality at 50 mg l⁻¹, Momin & Nair, 2001). As a consequence, a gel containing 5% of the *Apium graveolens* hexane extract was developed, providing full protection to volunteers from mosquito bites for two hours (Tuetun et al., 2009), while the ethanolic formulations from the same plant also provided protection against *Aedes aegypti*. Another formulation containing the aforementioned hexane extract and 5% vanillin showed strong repellent activities against different mosquito species (Tuetun et al., 2005, see also **Table 1** for details). The crude seed extract had no adverse effects on human volunteers skins when tested for several anti-mosquito properties (Choochote et al., 2004). This plant's EO exhibits potent larvicidal activity against two laboratory-reared mosquito species, the malaria vector *Anopheles dirus* and the vector of dengue *Aedes aegypti* (Pitasawat et al., 2007).

Species	Part used	Mosquito species	Bioactivity	Reference
<i>Ammi visnaga</i>	Seeds	<i>Culex quinquefasciatus</i>	Larvicidal	Pavela, 2008
<i>Anethum graveolens</i>	(not mentioned)	<i>Anopheles stephensi</i> , <i>Aedes aegypti</i> , <i>Culex quinquefasciatus</i>	Larvicidal	Amer and Mehlhorn, 2006
<i>Anethum graveolens</i>	Leaves, twigs	<i>Aedes aegypti</i>	Larvicidal, effects on growth and development	Promsiri et al., 2006
<i>Angelica archangelica</i>	Fruits	<i>Culex quinquefasciatus</i>	Larvicidal	Pavela, 2009
<i>Angelica sylvestris</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Apium graveolens</i>	Seeds	<i>Aedes aegypti</i>	Larvicidal	Momin & Nair, 2001
<i>Apium graveolens</i>	Seeds	<i>Aedes aegypti</i>	Larvicidal, adulticidal, repellent	Choochote, 2004
<i>Apium graveolens</i>	Seeds	<i>Aedes aegypti</i> , <i>Aedes gardnerii</i> , <i>Aedes lineatopennis</i> , <i>Anopheles barbirostris</i> , <i>Armigeres subalbatus</i> , <i>Culex tritaeniorhynchus</i> , <i>Culex gelicus</i> , <i>Culex vishnui</i> group, <i>Mansonia uniformis</i>	Repellent	Tuetun et al., 2005

Species	Part used	Mosquito species	Bioactivity	Reference
<i>Apium graveolens</i>	Seeds	<i>Aedes aegypti</i> , <i>Anopheles ditrus</i>	Larvicidal	Pitasawat et al., 2007
<i>Apium graveolens</i>	Seeds	<i>Aedes</i> , <i>Anopheles</i> , <i>Armigeres</i> , <i>Culex</i> , <i>Mansonia</i>	Repellent	Tuetun et al., 2009
<i>Athamanta densa</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Bupleurum fruticosum</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	Evergetis et al., 2009
<i>Carum carvi</i>	Fruits	<i>Aedes aegypti</i> , <i>Culex quinquefasciatus</i>	Larvicidal	Lee, 2006
<i>Carum carvi</i>	Seeds	<i>Aedes aegypti</i> , <i>Anopheles ditrus</i>	Larvicidal	Pitasawat et al., 2007
<i>Carum ptroselinum</i>	(not mentioned)	<i>Culex pipiens</i>	Larvicidal	Khater and Shalaby, 2008
<i>Chaerophyllum heldreichii</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Conium divaricatum</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Conopodium capillifolium</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	Evergetis et al., 2009
<i>Coriander sativum</i>	Seeds	<i>Ochlerotatus caspius</i>	Larvicidal	Knio et al., 2008
<i>Cryptotaenia canadensis</i>	Fresh foliage, root, fruits	<i>Culex pipiens</i>	Larvicidal	Eckenbach et al., 1999
<i>Daucus carota</i>	Roots	<i>Aedes aegypti</i> , <i>Culex quinquefasciatus</i>	Larvicidal	Lee, 2006
<i>Eleoselinum asclepium</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	Evergetis et al., 2009
<i>Ferula assa-foetida</i>	Stems	<i>Culex quinquefasciatus</i>	Larvicidal	Pavela, 2009
<i>Ferula galbaniflua</i>	(not mentioned)	<i>Anopheles stephensi</i> , <i>Aedes aegypti</i> , <i>Culex quinquefasciatus</i>	Larvicidal	Amer and Mehlhorn, 2006
<i>Ferula lancerottensis</i>	Stems	<i>Culex quinquefasciatus</i>	Larvicidal	Pavela, 2008
<i>Ferulago nodosa</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Foeniculum vulgare</i>	Fruits	<i>Aedes aegypti</i>	Repellent	Kim et al., 2002
<i>Foeniculum vulgare</i>	(not mentioned)	<i>Aedes aegypti</i>	Larvicidal	Orozco & Lentz, 2005
<i>Foeniculum vulgare</i>	Flowers	<i>Culex pipiens</i>	Larvicidal, repellent	Trabousli et al., 2005
<i>Foeniculum vulgare</i>	Fruits	<i>Aedes aegypti</i> , <i>Anopheles ditrus</i>	Larvicidal	Pitasawat et al., 2007
<i>Foeniculum vulgare</i>	Stems, inflorescences, leaves	<i>Culex pipiens</i>	Larvicidal	Manolakou et al., 2009

Species	Part used	Mosquito species	Bioactivity	Reference
<i>Foeniculum vulgare</i>	Leaves	<i>Aedes albopictus</i>	Larvicidal	Conti et al., 2010
<i>Heracleum sphondylium</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	Evergetis et al., 2009
<i>Imperatoria ostruthium</i>	Roots	<i>Culex quinquefasciatus</i>	Larvicidal	Pavela, 2009
<i>Laserpitium pseudomeum</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Oenanthe pimpinelloides</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	Evergetis et al., 2009
<i>Petroselinum crispum</i>	Seeds	<i>Ochlerotatus caspius</i>	Larvicidal	Knio et al., 2008
<i>Peucedanum neumayeri</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Peucedanum officinale</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Pimpinella anisum</i>	Seeds	<i>Anopheles stephensi</i> , <i>Aedes aegypti</i> , <i>Culex quinquefasciatus</i>	Larvicidal, adulticidal, ovicidal, oviposition- deterrent, repellent	Prajapati et al., 2005
<i>Pimpinella anisum</i>	Seeds	<i>Ochlerotatus caspius</i>	Larvicidal	Knio et al., 2008
<i>Pimpinella peregrina</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Pimpinella rigidula</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Pimpinella tragiium</i> ssp <i>tragiium</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Scaligeria cretica</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Seseli montanum</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	Evergetis et al., 2009
<i>Seseli pallasii</i>	Stems	<i>Culex quinquefasciatus</i>	Larvicidal	Pavela, 2009
<i>Seseli parnassicum</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Seseli tortuosum</i>	Stems	<i>Culex quinquefasciatus</i>	Larvicidal	Pavela, 2008
<i>Smyrniium rotundifolium</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Thamnosciadium junceum</i>	Aerial parts	<i>Culex pipiens</i>	Larvicidal	(present study)
<i>Trachyspermum ammi</i>	Seeds	<i>Anopheles stephensi</i>	Larvicidal, oviposition- deterrent, vapor toxicity, repellent	Pandey et al., 2009

Table 1. Reported phytochemicals derived from plants belonging to Apiaceae family against various mosquito species.

Another EO found to possess potent larvicidal, oviposition-deterrent, vapor toxicity and repellent activities against *Aedes aegypti* was isolated from ajowan (*Tachyspermum ammi*, Pandey et al. 2009). *Anethum graveolens* extract exhibited larval toxicity with LC₅₀ values from 27 to 20 mg l⁻¹ (for 24 and 48 hours exposures respectively), while on growth survival and prolongation tests of the various instar larvae of *Aedes aegypti*, the second instar larvae was determined as the more susceptible. The lowest concentration of crude extracts of *Anethum graveolens* used (caused more than 50% larval mortality) was not toxic to guppy fish (*Poecilia reticulata*) at concentrations of 12.5 mg l⁻¹ (Promsiri et al., 2006).

Among all EOs tested for mosquito control, the most potent was derived from *Foeniculum vulgare*, which caused the highest mortality against *Aedes albopictus* (Conti et al., 2010) and moderate against *Anopheles dirus* and *Aedes aegypti* (Pitasawat et al., 2007). Main component of this EO is methyl chavicol (more than 43%), while its methanolic extract (*trans*-anethole chemotype) was moderately active against *Aedes aegypti*, the yellow fever mosquito (Orozco & Lentz, 2005). The hexane fraction from its fruit-derived parts showed 99% repellency against *Aedes aegypti*, while the other fractions (chloroform, ethyl acetate and water: 37, 37 and 17% respectively) were practically inactive (Kim et al., 2002). Repellency and toxicity were also studied against *Culex pipiens* (Trabousli et al., 2005), indicating that the EO of *Foeniculum vulgare* was the most effective, while the repellency assays revealed protection time for almost one hour when applied at concentration of 3%.

Pimpinella anisum L. EO proved to possess equally potent larvicidal and ovicidal activities against *Anopheles stephensi*, *Aedes aegypti*, *Culex quinquefasciatus* and only larvicidal against *Ochlerotatus caspius* (Prajapati et al., 2005; Knio et al., 2008). Similar larvicidal activity results were also observed when the EOs of *Coriander sativum* and *Petroselinum crispum* were tested against *Ochlerotatus caspius* (Knio et al., 2008). The larvicidal tests of EOs of genus *Carum* were performed for *Carum carvi* against *Anopheles dirus* and *Aedes aegypti* and for *Carum petroselinum* against *Culex pipiens* (Lee, 2006; Pitasawat et al., 2007; Khater & Shalaby, 2008). The results were directly similar to those of the EO of *Daucus carota* (against *Anopheles dirus* and *Aedes aegypti*) proving their inability to cause 100% mortality at the lowest concentration (Lee, 2006).

Among the methanolic extracts of 118 Euroasiatic plants, tested for their larvicidal effects against *Culex quinquefasciatus*, the species *Ammi visnaga* and *Seseli pallasii* were determined as two of the most toxic materials tested, with LC₅₀ values lower than 10 mg l⁻¹ (Pavela, 2008, 2009). On the other hand, the extracts of *Angelica archangelica* and *Imperatoria ostruthium* exhibited LC₅₀ values lower than 70 mg l⁻¹, while *Seseli tortuosum* and *Ferula lancerottensis* displayed moderate larvicidal activity (LC₅₀ values around 430 mg l⁻¹). The only inactive Apiaceae plant tested was *Ferula assa-foetida* (LC₅₀ value higher of 1000 mg l⁻¹), with the EO of *Ferula galbaniflua* exhibiting the weakest activity against *Culex quinquefasciatus* and *Anopheles stephensi* (mortality level less than 14% of dead larvae after 48 hours, Amer & Mehlhorn, 2006a). The same authors also reported that *Anopheles stephensi* was the most resistant to dill (*Anethum graveolens*), while the *Culex quinquefasciatus* the more sensitive. Dill was also evaluated for persistency to larvicidal effects under different conditions for 1 month after the preparation of its solutions. In all cases (open, closed, in light or in dark) the EO was active only when was used immediately after preparation (Amer & Mehlhorn, 2006b).

Finally, an interesting result was obtained during the study of several EOs using coupled gas chromatography-electroantennographic detection (GC-EAD), on the hypothesis that compounds can be detected by the antennae of the yellow fever mosquito, *Aedes aegypti*. Thus, cumin aldehyde and cumin alcohol the *Cuminum cyminum* EO components were identified as such molecules. It must be noted that for both components, their EO (cumin oil) was also EAD-active (Campbell et al., 2011)

2.3 Greek Umbelliferae (Apiaceae) plants extract activities against *Culex pipiens* mosquitoes

The larvicidal activity of the EO obtained from the stem of Greek *Foeniculum vulgare* was determined against *Culex pipiens* larvae, while methyl chavicol was determined as its main component (more than 32%). Although the LC₅₀ value of methyl chavicol was more than 80 mg l⁻¹, the respective EO was determined as 2.1-fold more toxic (Manolakou et al., 2009). *Culex pipiens* larvae were also used to test the mosquito control properties of EO from various naturally growing plants throughout Greece, belonging to the following six different Apiaceae family taxa: *Heracleum sphondylium*, *Seseli montanum*, *Conopodium capillifolium*, *Bupleurum fruticosum*, *Oenanthe pimpinelloides*, *Eleoselinum asclepium*. All EOs tested displayed good larvicidal activities with LC₅₀ values ranging from 40.26-96.96 mg l⁻¹ (Evergetis et al., 2009)

As a continuation of our ongoing efforts to exploit the use of natural products for the development of environmentally friendly means for the mosquito population control, our interest was stimulated on the investigation of Umbelliferae (Apiaceae) plants EOs. In this context, we report herein the chemical composition and larvicidal activity results for 14 EOs originated from different taxon obtained during Greek Umbelliferae biodiversity studies (Table 1).

3. Materials and methods

3.1 Plant material

Fourteen different taxa of the Umbelliferae (Apiaceae) family, Apioideae subfamily belonging to seven tribes and twelve different genera have been collected during the present study. Representatives of the Apieae Tribe are *Pimpinella peregrina* L., and 5 Greek endemics, namely *Athamanta densa* Boiss. & Orph., *Pimpinella tragioides* ssp *tragioides* Vill., *Pimpinella rigidula* (Boiss. & Orph.) H. Wolf, *Seseli parnassicum* Boiss. & Heldr. and *Thamnosciadium junceum* (Sibth. & Sm.) Hartvig.; of Smyrnieae tribe *Scaligeria cretica* (Miller) Boiss. and *Smyrniium rotundifolium* Miller; of Angeliceae tribe *Angelica sylvestris* L.; of Scandiceae tribe the Greek endemic *Chaerophyllum heldreichii* Orph. Ex Boiss.; of Peucedaneae tribe *Ferulago nodosa* (L.) Boiss., *Peucedanum neumayeri* (Vis.) Reichenb, *Peucedanum officinale* L., and of Laserpitieae tribe the Greek endemic *Laserpitium pseudomeum* Orph., Heldr. & Sart. Ex Boiss (Pimenov & Leonov, 1993; Tutin et al., 1968).

Full collection details are provided in Table 2. A voucher specimen of each plant is deposited in the herbarium of the Agricultural University of Athens, Athens, Greece.

3.2 Essential oils isolation

The freshly collected plant materials (stems, leaves and flowers) were washed thoroughly, chopped off finely and subjected to steam distillation in a Clevenger-type apparatus, using the Microwave Accelerated Reaction System (MARS 5) at 1400 W for 40 min with 3 L of H₂O in order to obtain their EOs. The resulting oils were dried over anhydrous sodium sulphate and stored at 4 °C. The EO yield of each plant is included in **Table 3**.

Species	Abbreviation	Vegetative Stage	Date	Location
<i>Angelica sylvestris</i> L.	AS	Flowering	05.09.2004	Mt. Parnon, Peloponnisos, forest streams
<i>Athamanta densa</i> Boiss. & Orph. *	AD	Flowering	15.06.2005	Mt. Parnassos, Sterea Hellas, vertical cliffs
<i>Chaerophyllum heldreichii</i> Orph. Ex Boiss. *	CH	Flowering	25.07.2004	Mt. Parnon, Peloponnisos, forest clearings
<i>Ferulago nodosa</i> (L.) Boiss.	FN	Flowering	02.05.2005	Antikyra, Sterea Hellas, olive groves
<i>Laserpitium pseudomeum</i> Orph., Heldr. & Sart. Ex Boiss. *	LP	Flowering	15.07.2004	Mt. Oiti, Sterea Hellas, rocky slopes
<i>Peucedanum neumayeri</i> (Vis.) Reichenb	PN	Flowering	28.08.2004	Mt. Smolikas, Hepiros, forest clearings
<i>Peucedanum officinale</i> L.	PO	Flowering	15.07.2004	Mt. Oiti, Sterea Hellas, rocky slopes
<i>Pimpinella tragiium</i> ssp <i>tragiium</i> Vill. *	PT	Flowering	15.07.2004	Mt. Oiti, Sterea Hellas, rocky slopes
<i>Pimpinella peregrina</i> L.	PP	Flowering	14.05.2005	Iraklio, Is. Crete, olive groves
<i>Pimpinella rigidula</i> (Boiss. & Orph.) H. Wolf *	PR	Flowering	17.08.2004	Molai, Peloponnisos, roadside
<i>Scaligeria cretica</i> (Miller) Boiss.	SC	Flowering	22.05.2005	Vouliagmeni, Sterea Hellas, seaside
<i>Seseli parnassicum</i> Boiss. & Heldr. *	SP	Flowering	15.07.2004	Mt. Oiti, Sterea Hellas, forest clearings
<i>Smyrniium rotundifolium</i> Miller	SR	Flowering	02.05.2005	Distomo, Sterea Hellas, roadside
<i>Thamnosciadium junceum</i> (Sibth. & Sm.) Hartvig *	TJ	Flowering	25.07.2004	Mt. Parnassos, Sterea Hellas, alpic ravine

*=Greek Endemic.

Table 2. Collection data.

Species	Part distilled	Weight of aerial parts (g)	Volume of oil (mL)
<i>Angelica sylvestris</i> L.	Aerial	920	0,5
<i>Athamanta densa</i> Boiss. & Orph. *	Aerial	450	0,5
<i>Chaerophyllum heldreichii</i> Orph. Ex Boiss. *	Aerial	530	0,7
<i>Ferulago nodosa</i> (L.) Boiss.	Aerial	400	0,7
<i>Laserpitium pseudomeum</i> Orph., Heldr. & Sart. Ex Boiss. *	Aerial	270	0,9
<i>Peucedanum neumayeri</i> (Vis.) Reichenb	Aerial	600	0,5
<i>Peucedanum officinale</i> L.	Aerial	180	0,8
<i>Pimpinella tragiium</i> ssp <i>tragiium</i> Vill. *	Aerial	650	1,5
<i>Pimpinella peregrina</i> L.	Aerial	500	0,4
<i>Pimpinella rigidula</i> (Boiss. & Orph.) H. Wolf *	Aerial	235	0,7
<i>Scaligeria cretica</i> (Miller) Boiss.	Aerial	200	0,5
<i>Seseli parnassicum</i> Boiss. & Heldr. *	Aerial	200	0,5
<i>Smyrniium rotundifolium</i> Miller	Aerial	530	0,9
<i>Thamnosciadium junceum</i> (Sibth. & Sm.) Hartvig *	Aerial	600	2,0

*=Greek Endemic.

Table 3. Essential oils yields.

3.3 Gas Chromatography-Mass Spectrometry (GC-MS) analyses

Gas Chromatography (GC). All GC analyses were carried out on a Agilent Technologies 7890A gas chromatograph, fitted with a HP 5MS 30m x 0.25mm x 0.25µm film thickness capillary column and FID. The column temperature was programmed from 60 to 280 °C at a initial rate of 3 °C/min. The injector and detector temperatures were programmed at 230 and 300 °C, respectively. Helium was used as the carrier gas at a flow rate 1 ml/min.

Gas Chromatography-Mass Spectrometry (GC-MS). The GCMS analyses were performed on the same instrument using the Agilent 5957C, VL MS Detector with Triple-Axis Detector system operating in EI mode (equipped with a HP 5MS 30m x 0.25mm x 0.25µm film thickness capillary column), using He (1 ml/min) as the carrier gas. The initial temperature of the column was 60 °C. The column was heated gradually to 280 °C with a 3 °C/min rate. The identification of the compounds was based on comparison of their retention indices (RI) (Van den Dool & Kratz, 1963), obtained using various n-alkanes (C9-C24). Also, their EI-mass spectra were compared with the NIST/NBS and Wiley library spectra and the literature (Adams, 1995; Massada, 1976). Additionally, the identity of the indicated phytochemicals was confirmed by comparison with available authentic samples.

3.4 Mosquito rearing

A colony of the species *Culex pipiens* biotype *molestus* is maintained for more than 25 years in the laboratory of Entomology of the Benaki Phytopathological Institute, Kifissia, Greece. Adult mosquitoes are kept in wooden framed cages (33x33x33 cm) with a 32x32 mesh at 25±2 °C, 80±2% relative humidity and photoperiod of 14:10 (L:D) h. Cotton wicks saturated with 10% sucrose solution are used as food source. Females lay eggs in round, plastic containers (10 cm

diameter x 5 cm depth) filled with 150 ml of tap water. Egg rafts are removed daily and placed in cylindrical enamel pans (with diameter of 35 cm and 10 cm deep), in order to hatch. Larvae are reared under the same conditions of temperature and light and are fed daily with baby fish food (TetraMin, Baby Fish Food) at a concentration of 0.25 g^l⁻¹ of water until pupation. Pupae are then collected and introduced into the adult rearing cages.

3.5 Larvicidal bioassays

Stock solutions of EOs tested were prepared in ethanol and maintained in a freezer as 1% mg^l⁻¹ solutions. They were dissolved in double distilled water to produce solutions of the tested materials in concentrations ranging from 5 to 150 mg^l⁻¹. Prior to biological determinations the toxicity of each EO was evaluated (data not shown).

The larval mortality bioassays were carried out according to the test method for larval susceptibility, proposed by the World Health Organization (WHO, 1981). Twenty 3rd to 4th instar larvae of the species *Culex pipiens* biotype *molestus* were collected from the colony, placed in a glass beaker with 250 ml of aqueous suspension of the tested material at various concentrations and an emulsifier was added in the final test solution (less than 0.05%). Four replicates were made per each concentration and a control treatment with tap water and emulsifier was also included. Beakers with larvae were placed at 25±2 °C, 80±2% relative humidity and photoperiod of 14:10 h (L:D).

3.6 Data analysis

Larvicidal effect was recorded 48 h after treatment. Data obtained from each dose-larvicidal bioassay (total mortality, mg^l⁻¹ concentration in water) were subjected to probit analysis in which probit-transformed mortality was regressed against log₁₀-transformed dose; LC₅₀, LC₉₀ values, and slopes were calculated (SPSS 11.0).

4. Results and discussion

4.1 Phytochemical analysis

Fourteen distinct Umbelliferae taxa (twelve genera) are studied herein, one of which is endemic to Greece (*Thamnosciadium* Hartvig). It must be noted that there are no literature reports and studies on the EOs and their chemical compositions for the material obtained from the plants *Athamanta densa* Boiss. & Orph. (AD), *Chaerophyllum heldreichii* Orph. Ex Boiss. (CH), *Laserpitium pseudomeum* Orph., Heldr. & Sart. Ex Boiss. (LP), *Peucedanum neumayeri* (Vis.) Reichenb (PN), *Pimpinella tragiium* ssp *tragiium* Vill. (PT), *Pimpinella rigidula* (Boiss. & Orph.) H. Wolf (PR), *Scaligeria cretica* (Miller) Boiss. (SC), *Seseli parnassicum* Boiss. & Heldr. (SP) and *Smyrniium rotundifolium* Miller (SR). In addition, the discussion section on the related taxa EOs compositions includes ten (out of twelve) genera studied herein, since there are also no previous reports on the composition of EOs obtained from *Conium* L. and *Thamnosciadium* Hartvig genera.

In total seventy phytochemicals, representing 76.64 to 99.83 % of the respective EOs samples have been identified as their constituents using combined GC and GC/MS analyses and in certain occasions verified by NMR studies. The detailed qualitative and quantitative analytical data of the main constituents of steam volatiles (and their respective retention indices) are presented in **Table 4**.

Components	RI	PN	AS	TJ	SP	PO	CH	LP	PT	SC	PP	PR	FN	SR	AD	Identification
<i>trans</i> -2-hexanal	803												0.96	0.27		a, b
<i>a</i> -pinene	939	21.27	24.65	2.80		2.14	1.67	49.58	1.21	8.76			30.85	0.93	0.46	a, b, c
camphene	954	2.99	3.32			1.72							4.36			a, b, c
sabinene	975	2.76				1.15	71.76	24.73	4.30	13.74			1.96	0.93		a, b, c
β -pinene	979	2.66	1.33					8.51					1.79	8.86		a, b, c
myrcene	991	3.93	4.75	1.52		0.81	1.51	1.82					6.68	11.25	0.92	a, b, c
<i>a</i> -phellandrene	1003	2.53	2.35	3.83									1.33			a, b, c
<i>a</i> -terpinene	1017		3.58										0.32			a, b, c
<i>p</i> -cymene	1025	4.71						0.74								a, b, c
<i>o</i> -cymene	1026			0.91												a, b, c
limonene	1029	4.71		40.75		2.78			1.42	1.43					0.66	a, b, c
β -phellandrene	1030	12.76	42.96		1.42		10.86	6.73					10.20			a, b, c
<i>cis</i> -ocimene	1037	4.78		18.59									2.76	1.66		a, b, c
<i>trans</i> -ocimene	1050			0.82									1.01	5.16		a, b, c
γ -terpinene	1060	32.25				2.54	1.42			1.40			0.41			a, b, c
<i>cis</i> -sabinene hydrate	1070							2.53								a, b, c
terpinolene	1089			12.97												a, b, c
linalool	1097														0.50	a, b, c
<i>trans</i> -limonene oxide	1137			0.46												a, b, c
geijerene	1143								10.23							a, b
<i>a</i> -terpineol	1189						3.35	2.43		0.78						a, b, c
pregeijerene	1287								5.13							a, b
1-bornyl acetate	1289		3.84			81.13							0.52			a, b
2,3,4-trimethyl benzaldehyde	1359			2.16		4.68										a, b
2,3,6-trimethyl benzaldehyde	1371			0.62												a, b
isolekene	1376										0.69					a, b
<i>a</i> -copaene	1377									0.67			0.42		0.26	a, b
β -cubebene	1388													0.86		a, b
β -elemene	1391				10.85					0.50			0.51	2.09	0.33	a, b
aristolene	1407										19.92					a, b
calarene	1411										3.40				0.40	a, b
β -caryophyllene	1419		1.73		2.76				0.92	3.07			2.10			a, b, c
<i>a</i> -bergamontene	1435										62.15					a, b, c
γ -elemene	1437				3.22									0.72		a, b, c
β -humulene	1439				3.30					0.40						a, b, c
β -farnesene	1457									29.27		2.47			1.17	a, b, c
C ₁₄ H ₃₀ O (m/z: 189, 147, 105, 91, 204)	1483														19.80	b
<i>a</i> -amorphene	1484											0.32				a, b
C ₁₄ H ₂₈ O (m/z: 119, 91, 105, 145, 131)	1485								20.39							b
germacrene D	1487	2.55	4.42	0.89	13.02		2.53	0.71	0.87	28.37	0.85		6.42		1.21	a, b, c
β -selinene	1490				5.14						3.78	23.80			3.95	a, b, c
<i>a</i> -selinene	1498					0.95						3.53		5.28		a, b, c
<i>a</i> -zingiberene	1499											7.75				a, b
bicyclogermacrene	1500				6.25					3.51	1.21		4.04			a, b
<i>a</i> -farnesene	1506		1.84													a, b, c
β -bisabolene	1506				2.85						1.18	4.16			12.72	a, b, c
myristicin	1519											6.72			4.32	a, b, c
β -sesquiphellandrene	1523				30.39						1.04	1.80		0.92		a, b, c
δ -cadinene	1524	1.30	3.02							0.38						a, b, c
germacrene B	1561				10.64				19.28					2.13	0.98	a, b, c
spathulenol	1578				1.52					0.46						a, b, c
caryophyllene oxide	1583				1.02					0.61						a, b, c
β -elemenone	1601								1.72							a, b
isofuranogermacrene	1648													1.28		a, b
furanodiene	1649													11.81		a, b
<i>a</i> -bisabolol	1686		2.04													a, b
germacrone	1694								23.33						5.62	a, b
<i>trans</i> -isomyristicin	1721			10.14								7.74				a, b
<i>trans</i> -pseudoisoeugenyl	1774											7.74				a, b
2-methylbutyrate	1783												26.72			a, b
<i>trans</i> -epoxypseudoisoeugenyl	1880													6.42	0.91	a, b
2-methylbutyrate	1889													8.87		a, b
1 β -acetoxyfuranouedism-4(15)-ene	1911													20.72		a, b
C ₁₂ H ₂₅ O ₂ N (m/z: 91, 55, 115, 129, 77)	1923														2.94	b
C ₁₂ H ₂₅ O ₂ N (m/z: 91, 115, 55, 129, 77)	1943														8.58	b
C ₁₃ H ₂₇ O ₂ N (m/z: 91, 115, 55, 129, 159)	2030														12.17	b
<i>n</i> -heneicosane	2100														0.59	a, b, c
tricosane	2300														0.32	a, b, c
pentacosane	2500														0.25	a, b, c
Total		99.20	99.83	96.46	92.38	95.36	94.22	99.20	88.80	93.35	94.22	92.75	76.64	79.83	89.39	

^aComparison of mass spectra with MS libraries and retention times

^bComparison of experimental RI with reported RI

^cComparison with authentic compounds

RI: Retention indices calculated against C₈ to C₂₄ n-alkanes on the HP 5MS column.

Table 4. Chemical constituents of the essential oils tested.

The determined chemical composition of the EO from the aerial part of *Angelica sylvestris* L. (AS) is consistent with the literature reports for EOs obtained from its seeds (Bernard, 2001) and roots (Bernard & Clair, 1997), with α -pinene and β -phellandrene being the major components. Same compounds were reported as the prevailing phytochemicals in the EOs of *A. archangelica* L. *sensu lato* (Bernard, 2001; Nykanen et al., 1991; Bernard & Clair, 1997; Chalcat & Garry, 1997; Nivinskiene et al., 2005), while the EO of *A. glauca* is reported to contain β -phellandrene as major component and only small portions of α -pinene (Aghinotri et al., 2004; Kaul et al., 1996). Other *Angelica* L. taxa, such as *A. sinensis* (Dung et al., 1996; Kim et al., 2006), *A. gigas* (Kim et al., 2006), *A. acutiloba* (Kim et al., 2006), *A. heterocarpa* (Bernard, 2001; Bernard & Clair, 1997) and *A. tenuissima* (Ka et al., 2005) display a completely different, both qualitative and quantitative, EO composition profile.

In addition to α -pinene, which is the main constituent as previously reported by Demetzos et al. (2000), the studied EO of *Ferulago nodosa* (L) Boiss. (FN) was found to contain thirteen new components for the *taxon's* EO. More specifically, the molecules of *trans*-2-hexenal, myrcene, α -phellandrene, α -terpinene, β -phellandrene, *cis*-ocimene, *trans*-ocimene, γ -terpinene, bornyl acetate, β -elemene, β -caryophyllene, germacrene D and bicyclgermacrene were also determined as constituents of this EO. With the exception of *trans*-2-hexenal all the abovementioned compounds have been assayed in the EOs of the following *Ferulago* W.D.J. Koch taxa; *F. asparagifolia* (Baser et al., 2001), *F. phialocarpa* (Masoudi et al., 2004b), *F. macrocolea* (Rustaiyan et al., 2005), *F. galbaniflua* (Rustaiyan et al., 2002a), and *F. thirkeana* (Baser et al., 2002).

The EO of *Peucedanum officinale* L. (PO) is dominated by bornyl acetate, which was previously found only in *P. scoparium* (Masoudi et al., 2004a). It is also characterized by the presence of 2,3,4-trimethyl benzaldehyde, which has not been previously reported as constituent of *Peucedanum* L. EOs. In addition, the EO tested was found to contain five molecules, namely α -pinene, sabinene, myrcene, limonene and β -selinene, never reported in a EO of *P. officinale* (Jaimand et al., 2006). These five phytochemicals are abundant in the general profile of *Peucedanum* L. EOs, as reported for *P. scoparium* (Masoudi et al., 2004a), *P. zenkeri* (Menut et al., 1995), *P. verticillare* (Fraternale et al., 2000), *P. petiolare* (Rustaiyan et al., 2001) and *P. cervariifolium* (Bazgir et al., 2005).

The EOs of *Pimpinella* L. have been thoroughly studied, mainly because the application of their several taxa as culinary herbs and/or spices. Though the EOs of fourteen (14) *taxa* were studied, only one (Tabanca et al., 2005) refers to PP (*Pimpinella peregrina* L.) and none to PT (*Pimpinella tragiium* ssp *tragiium* VIII) and PR (*Pimpinella rigidulla* Boiss. & Orph. H. Wolf). The main constituent of EO of PO is α -bergamontenene, reported so far only for *P. anagodendron* (Velasco-Negueruela et al. 2005) and *P. anisum* (Santos et al., 1998). Two additional components determined herein, β -bisabolene and β -sesquiphellandrene, have not been reported in previous studies for PP but are well documented for *P. anagodendron* (Velasco-Negueruela et al. 2005), *P. junoniae* (Velasco-Negueruela et al. 2003), *P. anisum* (Santos et al., 1998), *P. anisetum* (Baser et al., 1999; Tepe et al., 2006) and *P. tragioides* (Askari & Sefidcon, 2007). New entries, for this genera EO components list, are isodenedene, aristolene, calarene and β -selinene which were also assayed in the EO of PP. On the contrary, the EO of PR is characterized by the complete absence of monoterpenes, advocating previous record of β -selinene and introducing α -amorphene, α -selinene and *trans*-isomyristicin as components of the *Pimpinella* L. EOs. Finally, the EO of PT has only two differences as

compared to the genus EO components, an unidentified component and β -elemenone. In general, its composition is in accordance with the phytochemical profiles reported for the EOs of *P. aromatica* (Baser et al., 1996), *P. serbica* (Ivanic et al., 1983), *P. flabellifolia* (Tepe et al., 2006), *P. aurea* (Tabanca et al., 2005; Assadian et al., 2005), *P. acuminata* (Melkani et al., 2006), *P. barbata* (Fakhari & Sonboli, 2006), *P. rupicola* (Velasco-Negueruela et al. 2005), *P. corymbosa* and *P. puberula* (Tabanca et al., 2005).

Major components of the EO of *Scaligeria cretica* (Miller) Boiss (SC) are *a*-pinene, β -farnesene and germacrene D, which have also been detected in previous studies on the EOs of *Scaligeria* DC. In this respect, the EO of *S. lazica* contains β -farnesene as major and *a*-pinene, germacrene D as minor components (Baser et al., 1993). On the contrary, the EO of *S. tripartite* contains β -farnesene and germacrene D as minor compounds, while *a*-pinene is absent (Tabanca et al., 2007). Compounds assayed herein and never reported before in the EOs of *Scaligeria* DC are *a*-terpineol, β -elemene and β -humulene.

The EOs of *Laserpitium pseudomeum* Orph. Heldr. & Sart Ex Poiss. (LP) contains *a*-pinene, β -pinene, sabinene and β -phellandrene as major components, all well known constituents of the EOs of *Laserpitium* L. Previous literature reports indicated that the EOs of *L. latifolium* contains *a*-pinene and β -pinene as major components (Borg-Karlson et al., 1994), the *L. petrophilum* *a*-pinene and sabinene (Baser et al., 1997), while the molecule of β -phellandrene is present in traces in both EOs. On the contrary, the phytochemical profile of the EO of *L. siler* is completely different containing mainly limonene and perillaldehyde (Chizzola et al., 1999)

The EO composition of *Smyrniium* L. has also been scarcely investigated, since only three *taxa*'s EOs, namely *S. perfoliatum* (Molleken et al., 1998a; Tirillini et al., 1996; Tirillini & Tosi, 1992), *S. cordifolium* (Amiri et al., 2006) and *S. olusatrum* (Molleken et al., 1998b), have been studied to date. The studied of EO of *Smyrniium rotundifolium* Miller (SR) contains 7 major components, with the molecule of *a*-selinene reported for the first time as EO component of *Smyrniium* L.. Other compounds present in large quantities are furanodiene (reported as major constituent in *S. olusatrum*), myrcene, furanoremonophil-1-one, 1 β -acetoxyfuranoeudesm-4(15)-ene, 1 β -acetoxyfurano eudesm-3-ene (detected in *S. olusatrum* and *S. perfoliatum*, Molleken et al., 1998) and germacrene (present in *S. cordifolium*).

The phytochemical profile of *Chaerophyllum* L. EOs was studied previously for *C. macropodium* (Baser et al., 2006), *C. crinitum* (Baser et al., 2006; Nematollahi et al., 2005), *C. macrospermum* (Sefidcon & Abdoli, 2005; Rustaiyan et al., 2002b, Mamedova, 1994), *C. bulbosum sensu lato* (Mamedova & Akhmedova, 1991; Kokkalou & Stefanou, 1989), *C. aksekiense* (Baser et al., 2000b), *C. coloratum* (Vajs et al., 1995), *C. azoricum* (Pedro et al., 1999) and *C. prescottii* (Letchamo et al., 2005). The more significant differentiation among the literature results and the assayed herein EO of *Chaerophyllum heldreichi* Orph. Ex Boiss (CH) comprises the identification for first time of *a*-terpineol as main component of EO of *Chaerophyllum* L..

The EO of *Seseli parnassicum* Boiss. & Heldr. (SP) was found to contain three new compound entries, β -humulene, β -selinene and β -sesquiphellandrene, as compared with the EO of the *Seseli* L. *taxa* (also including the synonymous *Lomatopodium* Fisch. et C.A. Mey *taxa*). The remaining components are in accordance with the EO content of same *taxa* plants, such as *S. montanum* (Evergetis et al., 2009), *S. campestre* and *S. peucedanoides* (Baser et al. 2000a;

Bulatovic et al. 2006) and in *S. buchtormence*. These compounds were also present in the EOs of *S. resinosum* and *S. tortuosum*, obtained from the fruits and not the herbal part of the plants (Dogan et al. 2006). The *L. khorassanicum* and *L. staurophyllum* EOs were assayed to contain mostly aliphatic terpenes, while the corresponding cyclic terpenes were present in smaller amounts compared to EOs of *Seseli* L. (Sedghat et al. 2003; Sefidkon et al. 1997).

Finally, the investigated EO of *Athamanta densa* Boiss. & Orph., contains as major constituents myristicin and various unidentified alkaloids, which account for almost 24 % of its weight. The literature reports of EOs of *Athamanta* L. indicate that they mainly contain either myristicin, such as the EOs of *A. sicula* (Camarda & Di Stefano, 2003), *A. turbitus sensu lato* (Tomic et al., 2009), *A. macedonica* (Verykokidou et al., 1995) and *A. haynaldi* (Zivanovic et al., 1994), or apiole as in *A. sicula* (Camarda & Di Stefano, 2008).

4.2 Larvicidal assays

The investigated EOs were evaluated —for the first time— in respect to their larvicidal activities against 3rd- 4th instar larvae of *Culex pipiens*. The relative results expressed as the respective LC₅₀ and LC₉₀ values are included in **Table 5**. Among the EOs tested only two were rather inactive (AS and PP, displaying LC₅₀ values above 150 mg l⁻¹), while the EOs of SC and SP were moderately active displaying LC₅₀ values above 100 mg l⁻¹ (111.99 and 122.54 mg l⁻¹ respectively).

Essential Oils tested	LC ₅₀ (95% CL) ^a	LC ₉₀ (95% CL) ^a	Slope (±SE)
<i>Athamanta densa</i>	10.15 (9.49-10.73)	15.75 (14.52-17.76)	6.72±0.80
<i>Pimpinella tragiium</i> ssp <i>tragiium</i>	40.13 (32.43-45.95)	71.10 (61.51-91.00)	5.15±0.52 ^b
<i>Pimpinella rigidula</i>	40.31 (34.75-43.64)	60.41 (55.66-70.57)	7.29±1.44 ^b
<i>Thamnosciadium junceum</i>	44.17 (41.52-46.62)	64.42 (59.94-71.28)	7.82±0.86 ^b
<i>Peucedanum neumayeri</i>	47.40 (40.25-54.15)	81.47 (68.63-113.57)	5.44±0.53 ^b
<i>Chaerophyllum heldreichii</i>	53.61 (50.29-56.55)	75.96 (71.53-82.15)	8.46±0.87
<i>Laserpitium pseudomeum</i>	56.73 (53.50-59.60)	79.59 (75.18-85.71)	8.46±0.86
<i>Ferulago nodosa</i>	67.39 (64.17-70.41)	95.59 (89.90-103.94)	8.43±0.84
<i>Smyrniium rotundifolium</i>	80.32 (76.88-84.16)	105.30 (98.33-116.61)	10.89±1.29
<i>Peucedanum officinale</i>	86.46 (82.27-90.30)	125.05 (117.23-136.95)	7.99±0.84
<i>Scaligeria cretica</i>	111.99 (107.86-115.47)	133.83 (128.35-143.21)	6.58±0.73 ^b
<i>Seseli parnassicum</i>	122.54 (115.54-141.06)	167.15 (143.83-268.76)	6.30±0.68
<i>Angelica sylvestris</i>	>150		
<i>Pimpinella peregrina</i>	>150		

^a LC values are expressed in mg l⁻¹ and they are considered significantly different when 95% CL fail to overlap.

^b Since goodness-of-fit test is significant (P<0.05), a heterogeneity factor is used in the calculation of confidence limits (CL)

Table 5. LC₅₀ and LC₉₀ values for the tested essential oils against larvae of *Culex pipiens* biotype *molestus*.

The EO derived from the endemic in Greece plant *Athamanta densa* was determined as the most active since displayed the highest toxicity against mosquito larvae, with LC₅₀ value 10.15 mg l⁻¹. The EO tested contains a series of compounds which were not found in the other EOs tested, such as bisabolene and the unidentified compounds C₁₄H₃₀O, C₁₂H₂₅O₂N and C₁₃H₂₇O₂N, which have to study more thoroughly in order to determine their activities. The remaining EOs (PR, TJ, PT, PN, CH, LP, FN, SR and PO) displayed LC₅₀ values ranging from 40.31 to 86.46 mg l⁻¹. No significant relationship between toxicity and phytochemical content was detected.

5. References

- Adams, R. (1995). Identification of essential oil components by Gas Chromatography / Mass Spectroscopy. Carol Stream, Allured Publishing, Illinois, USA. ISBN 0931710421.
- Aghinotri, V.K., Thappa R.K., Meena, B., Kapaphi, B.K. Saxena, R.K., Qazi, G.N. & Agarwal S.G. (2004). Essential oil composition of aerial parts of *Angelica glauca* growing wild in North-West Himalaya (India). *Phytochemistry*, 65, 2411-2413.
- Alves, S.N., Serro, J.E. & Melo, A.L. (2010). Alterations in the fat body and midgut of *Culex quinquefasciatus* larvae following exposure to different insecticides. *Micron*, 41, 592-597.
- Amer, A. & Mehlhorn, H. (2006a) Larvicidal effects of various essential oils against *Aedes*, *Anopheles*, and *Culex* larvae (Diptera: Culicidae). *Parasitology Research*, 99, 466-472.
- Amer, A. & Mehlhorn, H. (2006b) Persistency of larvicidal effects of plant oil extracts under different storage conditions. *Parasitology Research*, 99, 473-477.
- Amiri, H., Khavari-Nejad R.A., Masoud, S., Chalabian, F. & Rustaiyan, A. (2006). Composition and antimicrobial activity of the essential oil from stems, leaves, fruits and roots of *Smyrniium cordifolium* Boiss. from Iran. *Journal of Essential Oil Research*, 18, 574-577.
- Askari, F. & Sefidkon, F. (2007). Essential oil composition of *Pimpinella tragioides* (Boiss.) Benth et Hook from Iran. *Journal of Essential Oil Research*, 19, 54-56.
- Assadian, F., Masoudi, S., Nematollahi, F., Rustaiyan, A., Larijani, K. & Mazloomifar, H. (2005). Volatile constituents of *Xanthogalum purpurascens* Ave-Lall., *Eryngium caeruleum* M.B. and *Pimpinella aurea* DC. three Umbelliferae herbs growing in Iran. *Journal of Essential Oil Research*, 17, 243-245.
- Ateyyat, M.A., Al-Mazra'awi, M., Abu-Rjai, T. & Shatnawi, M.A. (2009). Aqueous extracts of some medicinal plants are as toxic as Imidacloprid to the sweet potato whitefly, *Bemisia tabaci*. *Journal of Insect Science*, 9:15, 6pp.
- Baser, K.H.C. Demirci, B., Demirci, F., Hashimoto, T., Asakawa, Y. & Noma, Y. (2002). Ferulagone: a new monoterpene ester from *Ferulago thirkeana* essential oil. *Planta Medica*, 68, 564-567.
- Baser, K.H.C., Demirci, B. & Duman, H. (2001). Composition of the essential oil of *Ferulago asparagifolia* Boiss. from Turkey. *Journal of Essential Oil Research*, 13, 134-135.

- Baser, K.H.C. & Duman, H. (1997). Composition of the essential oil of *Laserpitium petrophilum* Boiss. et Heldr. *Journal of Essential Oil Research*, 9, 707-708.
- Baser, K.H.C., Ozek, G., Ozek, T. & Duran, A. (2006). Composition of the essential oil of *Chaerophyllum macropodium* Boiss. fruits obtained by microdistillation. *Journal of Essential Oil Research*, 18, 515-517.
- Baser, K.H.C., Ozek, T., Kurkcuoglu, M.K. & Guner, A. (1993). The essential oil of *Scaligeria lazica* Boiss. *Journal of Essential Oil Research*, 5, 463-464.
- Baser, K.H.C., Ozek, T., Duman, H. & Guner, A. (1996). Essential oil of *Pimpinella aromatica* Bieb. from Turkey. *Journal of Essential Oil Research*, 8, 463-464.
- Baser, K.H.C., Ozek, T. & Tabanca, N. (1999). Essential Oil of *Pimpinella anisetum* Boiss. et Bal. *Journal of Essential Oil Research*, 11, 445-446.
- Baser, K.H.C., Ozek, T., Kurkcuoglu, M. & Aytac, Z. (2000a) Essential oil of *Seseli campestre* Besser. *Journal of Essential Oil Research*, 12, 105-107.
- Baser, K.H.C., Tabanka, N., Ozek, T., Demicri, B., Duran, A. & Duman, H. (2000b). Composition of the essential oil of *Chaerophyllum aksekiense* A. Duran et Duman, a recently described endemic from Turkey. *Flavour and Fragrance Journal*, 15, 43-44.
- Bazgir, A., Shaabani, A. & Sefidkon, F. (2005). Composition of the essential oil of *Peucedanum cervariifolium* C.A. Mey. from Iran. *Journal of Essential Oil Research*, 17, 380-381.
- Becker, N., Petric, D., Zgomba, M., Boase, C., Dahl, C., Lane, J., Kaiser, A. Mosquitoes and their control, Kluwer Academic/Plenum Publishers: New York, 2003. ISBN 0306473607.
- Bernard, C. (2001). Essential oils of three *Angelica* L. species growing in France. Part II: fruit oils. *Journal of Essential Oil Research*, 13, 260-263.
- Bernard, C. & Clair, G. (1997). Essential oils of three *Angelica* L. species growing in France. I. Root oils. *Journal of Essential Oil Research*. 9, 289-294.
- Borg-Karlson, A-K., Valterova, I. & Anders Nilson, L. (1994). Volatile compounds from flowers of six species in the family Apiaceae: Bouquets for different pollinators? *Phytochemistry*, 35, 111-119.
- Boyer, S., Serandour, J., Lemperiere, G., Raveton, M. & Ravanel, P. (2006). Do herbicide treatments reduce the sensitivity of mosquito larvae to insecticides? *Chemosphere*, 65, 721-724.
- Bulatovic, V.M., Savikin-Fodulovic, K.P., Zdunic, G.M. & Popovic, M.P. (2006). Essential oil of *Seseli peucedanoides* (MB) Kos.-Pol. *Journal of Essential Oil Research*, 18, 286-287.
- Camarda, L. & Di Stefano, V. (2003). Essential oil of leaves and fruits of *Athamanta sicula* L. (Apiaceae). *Journal of Essential Oil Research*, 15, 133-134.
- Camarda, L., Di Stefano, V. & Pitonzo, R. (2008). Chemical composition of essential oils from *Athmanta sícula*. *Chemistry of Natural Compounds*, 44, 532-533.
- Campbell, C., Gries, R. & Gries, G. (2011) Forty-two compounds in eleven essential oils elicit antennal responses from *Aedes aegypti*. *Entomologia Experimentalis et Applicata*, 138, 21-32.

- Caquet, T., Roucaute, M., Le Goff, P. & Lagadic, L. (2011). Effects of repeated field applications of two formulations of *Bacillus thuringiensis* var. *israelensis* on non-target saltmarsh invertebrates in Atlantic coastal wetlands. *Ecotoxicology and Environmental Safety*, 74, 1122-1130.
- Chakraborty, S., Singha, S. & Chandra, G. (2010). Mosquito larvicidal effect of orthophosphoric acid and lactic acid individually or their combined form on *Aedes aegypti*. *Asian Pacific Journal of Tropical Medicine*, 3, 954-956.
- Chalchat, J.C., Garry, R.P., (1997). Essential oil of angelica roots (*Angelica archangelica* L.): optimization of distillation, location in plant and chemical composition. *Journal of Essential Oil Research*, 9, 311-319.
- Chizzola, R., Novak, J. & Franz, C. (1999). Fruit oil of *Laserpitium siler* L. grown in France. *Journal of Essential Oil Research*, 11, 197-198.
- Choi, W-S., Park, B-S., Lee, Y-H., Jang, D.Y., Yoon, H.Y. & Lee, S-E. (2006). Fumigant toxicities of essential oils and monoterpenes against *Lycoriella mali* adults. *Crop Protection*, 25, 398-401.
- Choochote, W., Tuetun, B., Kanjanapothi, D., Rattanachanpichai, E., Chaithong, U., Chaiwong, P., Jitpakdi, A., Tippawangkosol, P., Riyong, D. & Pitasawat, B. (2004) Potential of crude seed extract of celery, *Apium graveolens* L., against the mosquito *Aedes aegypti* (L.) (Diptera: Culicidae). *Journal of Vector Ecology*, 29, 340-346.
- Chowdhury, N., Laskar, S. & Chandra, G. (2008). Mosquito larvicidal and antimicrobial activity of protein of *Solanum villosum* leaves. *BMC Complementary and Alternative Medicine*, 8, 62, doi: 10.1186/1472-6882-8-62. Available online <http://www.biomedcentral.com/1472-6882/8/62>.
- Conti, B., Canale, A., Bertoli, A., Gozzini, F. & Pistelli, L. (2010). Essential oil composition and larvicidal activity of six Mediterranean aromatic plants against the mosquito *Aedes albopictus* (Diptera: Culicidae). *Parasitology Research*, 107, 1455-1461.
- Daaboub, J., Ben Cheikh, R., Lamari, A., Ben Jha, I., Feriani, M., Boubaker, C. & Ben Cheikh, H. (2008). Resistance to pyrethroid insecticides in *Culex pipiens pipiens* (Diptera: Culicidae) from Tunisia. *Acta Tropica*, 107, 30-36.
- Del Pilar Corena, M., Van Den Hurk, P., Zhong, H., Brock, C., Mowery, R., Johnson, J.V. & Linser, P. J. (2006). Degradation and effects of the potential mosquito larvicides methazolamide and acetazolamide in sheepshead minnow (*Cyprinodon variegatus*). *Ecotoxicology and Environmental Safety*, 64, 369-376.
- Delaunay, P., Jeannin, C., Schaffner, F. & Marty, P. (2009). Actualites 2008 sur la presence du moustique tigre *Aedes albopictus* en France metropolitaine. *Archives de Pediatrie*, 16, Supplement 2, 66-71.
- Demetzos, C., Perdetzoglou, D., Gazouli, M., Tan, K. & Economakis, C. (2000). Chemical analysis and antimicrobial studies on three species of *Ferulago* from Greece. *Planta Medica*, 66, 560-563.
- Dogan, E., Duman, H., Tosun, A., Kurkcuoglu, M. & Baser K.H.C. (2006). Essential oil composition of the fruits of *Seseli resinosum* Freyn et Sint. and *Seseli tortuosum* L. growing in Turkey. *Journal of Essential Oil Research*, 18, 57-59.

- Dung, N.X., Cu, L.D., Moi, L.D. & Leclercq, P.A. (1996). Composition of the leaf and flower oils from *Angelica sinensis* (Oliv.) diels cultivated in Vietnam. *Journal of Essential Oil Research*, 8, 503-506.
- Eckenbach, U., Lampman, R.L., Seigler, D.S., Ebinger, J. & Novak, R.J. (1999). Mosquitocidal activity of acetylenic compounds from *Cryptotaenia Canadensis*. *Journal of Chemical Ecology*, 25, 1885-1893.
- Evergetis, E., Michaelakis, A., Kioulos, E., Koliopoulos, G. & Haroutounian, S. A. (2009). Chemical composition and larvicidal activity of essential oils from six Apiaceae family taxa against the West Nile virus vector *Culex pipiens*. *Parasitology Research*, 105, 117-124.
- Fakhari, A.R. & Sonboli, A. (2006). Essential oil composition of *Pimpinella barbata* (DC.) Boiss. from Iran. *Journal of Essential Oil Research*, 18, 679-681.
- Fraternale, D., Giamperi, L., Rici, D. & Manunta, A. (2000). Composition of the essential oil of *Peucedanum verticillare*. *Biochemical Systematics and Ecology*, 28, 143-147.
- Gu, J., Liu, M., Deng, Y., Peng, H. & Chen, X. (2011). Development of an Efficient Recombinant Mosquito Dengovirus-Mediated RNA Interference System and Its Preliminary Application in Mosquito Control. *PLoS One*, 6, doi: 10.1371/journal.pone.0021329.
- Hancock, P.A., Sinkins, S.P. & Godfray, H.C.J. (2011). Strategies for Introducing Wolbachia to Reduce Transmission of Mosquito-Borne Diseases. *PLoS Neglected Tropical Diseases*, 5, doi: 10.1371/journal.pntd.0001024.
- Hayes, S.R., Hudon, M. & Park, H.W. (2011). Isolation of novel *Bacillus* species showing high mosquitocidal activity against several mosquito species. *Journal of Invertebrate Pathology*, 107, 79-81.
- Imbahale, S.S., Fillinger, U., Githeko, A., Mukabana, W.R. & Takken, W. (2010). An exploratory survey of malaria prevalence and people's knowledge, attitudes and practices of mosquito larval source management for malaria control in western Kenya. *Acta Tropica*, 115, 248-256.
- Isman, M.B. (2000). Plant essential oils for pest and disease management. *Crop Protection*, 19, 603-608.
- Ivanic, R., Savin, K. & Robinson, F.V. (1983). Essential oil from *Pimpinella serbica* fruits. *Planta medica*, 48, 60-61.
- Jaimand, K., Ashorabadi, E.S. & Dini, M. (2006). Chemical constituents of the leaf and seed oils of *Peucedanum officinale* L. cultivated in Iran. *Journal of Essential Oil Research*, 18, 670-671.
- Ka, M.H., Chol, E.H., Chun, H.S. & Lee, K.G. (2005). Antioxidative activity of volatile extracts from *Angelica tenuissimae* roots, peppermint leaves, pine needles, and sweet flags leaves. *Journal of Agricultural and Food Chemistry*, 53, 4124-4129.
- Kanzok, S.M. & Jacobs-Lorena M. (2006). Entomopathogenic fungi as biological insecticides to control malaria. *Trends in Parasitology*, 22, 49-51.
- Kaul, P.N., Mallavarapu, G.R. & Chamoli, R.P. (1996). The essential oil composition of *Angelica glauca* roots. *Planta medica*, 62, 80-81.

- Khater, H.F. & Shalaby, A.A.S. (2008). Potential of biologically active plant oils to control mosquito larvae (*Culex pipiens*, Diptera: Culicidae) from an Egyptian locality. *Revista do Instituto de Medicina Tropical de São Paulo*, 50, 107-112.
- Kilama, W.L. (2009). Health research ethics in public health: Trials and implementation of malaria mosquito control strategies. *Acta Tropica*, 112, Supplement 1, 37-47.
- Killeen, G.F. & Smith, T.A. (2007). Exploring the contributions of bed nets, cattle, insecticides and excitorepellency to malaria control: a deterministic model of mosquito host-seeking behaviour and mortality. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 101, 867-880.
- Kim, D-H., Kim, S-I., Chang, K-S. & Ahn, Y-J. (2002). Repellent Activity of Constituents Identified in *Foeniculum vulgare* Fruit against *Aedes aegypti* (Diptera: Culicidae). *Journal of Agricultural and Food Chemistry*, 50, 6993-6996.
- Kim, M.R., El-Aty A.M.A., Kim, S-I. & Shim J.H. (2006). Determination of volatile flavor components in danggui cultivars by solvent free injection and hydrodistillation followed by gas chromatographic-mass spectrometric analysis. *Journal of Chromatography A*, 1116, 259-264.
- Kim, S-I., Park, C., Ohh, M-H., Cho, H-C. & Ahn, Y-J. (2003a). Contact and fumigant activities of aromatic plant extracts and essential oils against *Lasioderma serricornis* (Coleoptera: Anobiidae). *Journal of Stored Products Research*, 39, 11-19.
- Kim, S-I., Roh, J-Y., Kim, D-H., Lee, H-S. & Ahn, Y-J. (2003b). Insecticidal activities of aromatic plant extracts and essential oils against *Sitophilus oryzae* and *Callosobruchus chinensis*. *Journal of Stored Products*, 39, 293-303.
- Knio, K.M., Usta, J., Dagher, S., Zournajian, H. & Kreydiyyeh, S. (2008). Larvicidal activity of essential oils extracted from commonly used herbs in Lebanon against the seaside mosquito, *Ochlerotatus caspius*. *Bioresource Technology*. 99, 763-768.
- Kokkalou, E. & Stefanou, E. (1989). The volatiles of *Chaerophyllum bulbosum* wild in Greece. *Pharmaceutica Acta Helvetiae*, 64, 133-134.
- Kwon, J-H. & Ahn, Y-J. (2002). Acaricidal Activity of Butylidenephthalide Identified in *Cnidium officinale* Rhizome against *Dermatophagoides farinae* and *Dermatophagoides pteronyssinus* (Acari: Pyroglyphidae). *Journal of Agricultural and Food Chemistry*, 50, 4479-4483.
- Lamiri, A., Lhaloui, S., Benjilali, B., Berrada, M. (2001). Insecticidal effects of essential oils against Hessian fly, *Mayetiola destructor* (Say). *Field Crops Research*, 71, 9-15.
- Lavialle-Defaix, C., Apaire-Marchais, V., Legros, C., Penner, C., Mohamed, A., Licznar, P., Corbel, V. & Lapied, B. (2011). *Anopheles gambiae* mosquito isolated neurons: A new biological model for optimizing insecticide/repellent efficacy. *Journal of Neuroscience Methods*, doi:10.1016/j.jneumeth.2011.06.003.
- Lee, H-S. (2004). Acaricidal Activity of Constituents Identified in *Foeniculum vulgare* Fruit Oil against *Dermatophagoides* spp. (Acari: Pyroglyphidae). *Journal of Agricultural and Food Chemistry*, 52, 2887-2889.
- Lee, H-S. (2006). Mosquito larvicidal activity of aromatic medicinal plant oils against *Aedes aegypti* and *Culex pipiens pallens*. *Journal of the American Mosquito Control Association*, 22, 292-295.

- Letchamo, W., Korolyk, E.A. & Tkachev, A.V. (2005). Chemical screening of essential oil bearing flora of Siberia. V. Composition of the essential oil of *Chaerophyllum prescottii* DC tops from Altai region. *Journal of Essential Oil Research*, 17, 560-562.
- Lima, E.P., Paiva, M.H.S., de Araújo, A.P., da Silva, E.V.G., da Silva, U.M., de Oliveira, L.N., Santana, A.E.G., Barbosa, C.N., de Paiva Neto, C.C., Goulart, M.O.F., Wilding, G.S., Ayres, C.F.J. & de Melo Santos, M.A.V. (2011). Insecticide resistance in *Aedes aegypti* populations from Ceará, Brazil. *Parasites & Vectors*, 4, doi: 10.1186/1756-3305-4-5.
- Mamedova, S.A. & Akhmedova, E.R. (1991). Essential oil of turnip-root chervil. *Chemistry of Natural Compounds*, 27, 248-249.
- Mamedova, S.A. (1994). Essential oil of *Chaerophyllum macrospermum*. *Chemistry of Natural Compounds*, 30, 267-277.
- Manolakou, S., Pitarokili, D., Koliopoulos, G., Michaelakis, A. & Tzakou, O. (2009). Essential Oil Composition of Different Parts of Greek *Foeniculum vulgare* and Larvicidal Activity of the Stem Oil. In: "Essential Oils and Aromas: Green Extraction and Application" edit by Prof. Farid Chemat, "Har Krishan Bhalla and Sons" (Publisher Journal of Essential Oil Bearing Plants), France.
- Masoudi, S., Akhgar, M.R. & Rustaiyan, A. (2004a). Essential oils of *Peucedanum scoparium* (Boiss.) Boiss. and *Serotinocarpum insignis* Mozaffarian. from Iran. *Journal of Essential Oil Research*, 16, 117-119.
- Masoudi, S., Rustaiyan, A. & Ameri, N. (2004b). Volatile oils of *Ferulago phialocarpa* Rech. f. et H. Reidl. and *Leutea elbursensis* Mozaffarian from Iran. *Journal of Essential Oil Research*, 16, 143-144.
- Massada, Y. (1976). Analysis of essential oil by Gas Chromatography and Spectrometry. New York: Wiley.
- Melkani, A.B., Javed, M.S., Melkani, K.B., Dev. V. & Beauchamp, P.S. (2006). Terpenoid composition of the essential oil from *Pimpinella acuminata* (Edgew.) CB Clarke. *Journal of Essential Oil Research*, 18, 312-314.
- Menut, C., Mve-Mba, C.E., Lamaty, G., Amvan Zollo, P.H., Tchoumboungang, F. & Bessiere, J.M. (1995). Aromatic plants of tropical central Africa. XVIII: Essential oils of leaf and rhizome of *Peucedanum zenkeri* Engl. from Cameroon. *Journal of Essential Oil Research*, 7, 77-79.
- Miyazawa, M., Tsukamoto, T., Anzai, J. & Ishikawa, Y. (2004). Insecticidal Effect of Phthalides and Furanocoumarins from *Angelica acutiloba* against *Drosophila melanogaster*. *Journal of Agricultural and Food Chemistry*, 52, 4401-4405.
- Moleken, U., Sinnwell, V. & Kubeczka, K.H. (1998a). The essential oil composition of fruits from *Smurnium perfoliatum*. *Phytochemistry*, 47, 1079-1083.
- Moleken, U., Sinnwell, V. & Kubeczka, K.H. (1998b). Essential oil composition of *Smurnium olusatrum*. *Phytochemistry*, 49, 1709-1714.
- Momin, R.A. & Nair, M.G. (2001). Mosquitocidal, nematocidal, and antifungal compounds from *Apium graveolens* L. seeds. *Journal of Agricultural and Food Chemistry*, 49, 142-145.

- Nematollahi, F., Akhgar, M.R., Larijani, K., Rustaiyan, A. & Masoudi, S. (2005). Essential oils of *Chaerophyllum macropodium* Boiss. and *Chaerophyllum crinitum* Boiss. from Iran. *Journal of Essential Oil Research*, 17, 71-72.
- Nivinskiene, O., Butkiene, R. & Mockute, D. (2005). The chemical composition of the essential oil of *Angelica archangelica* L. roots growing wild in Lithuania. *Journal of Essential Oil Research*, 17, 373-377.
- Nykanen, I. & Nykanen, L. (1991). Composition of angelica root oils obtained by Supercritical CO₂ extraction and steam distillation. *Journal of Essential Oil Research*, 3, 229-236.
- Orozco, O.L., Lentz, D.L. (2005). Poisonous plants and their uses as insecticides in Cajamarca, Peru. *Economic Botany*, 59, 166-173.
- Pandey, S.K., Upadhyay, S. & Tripathi A.K. (2009). Insecticidal and repellent activities of thymol from the essential oil of *Trachyspermum ammi* (Linn) Sprague seeds against *Anopheles stephensi*. *Parasitology Research*, 105, 507-512.
- Papachristos, D.P. & Stamopoulos D.C. (2002). Repellent, toxic and reproduction inhibitory effects of essential oil vapours on *Acanthoscelides obtectus* (Say) (Coleoptera: Bruchidae). *Journal of Stored Products Research*, 38, 117-128.
- Patersson, R.S., Lofgren, C.S. & Boston, M.D. (1968). The sterile-male technique for control of mosquitoes: a field cage study with *Anophles quadrimaculatus*. *The Florida Entomologist*, 51, 77-82.
- Pavela, R. (2008). Larvicidal effects of various Euro-Asiatic plants against *Culex quinquefasciatus* Say larvae (Diptera: Culicidae), *Parasitology Research*, 102, 555-559.
- Pavela, R. (2009). Larvicidal effects of some Euro-Asiatic plants against *Culex quinquefasciatus* Say larvae (Diptera: Culicidae), *Parasitology Research*, 105, 887-892.
- Pedro, L.G., Da Silva, J.A., Barroso, J.G., Figueiredo, A., Cristina, D., Stanley, G., Looman, A. & Scheffer, J.J.C. (1999). Composition of the essential oil of *Chaerophyllum azoricum* Trel., an endemic species of the Azores archipelago. *Flavour And Fragrance Journal*, 14, 287-289.
- Pennetier, C., Chabi, J., Martin, T., Chandre, F., Rogier, C., Hougard, J-M. & Pages, F. (2010). New protective battle-dress impregnated against mosquito vector bites. *Parasites & Vectors*, 3, 81.
- Peterson, R.K.D., Macedo, P.A. & Davis R.S. (2006). A Human-Health Risk Assessment for West Nile Virus and Insecticides Used in Mosquito Management. *Environmental Health Perspectives*, 114, 366-372.
- Pimenov, M.G. & Leonov, M.V. (1993). The genera of the Umbelliferae, Royal Botanic Gardens Kew & Botanical Garden of Moscow State University, Kent U.K., p. 156.
- Pitasawat, B., Champakaew, D., Chochote, W., Jitpakdi, A., Chaithong, U., Kanjanapothi, D., Rattanachanpichai, E., Tippawangkosol, P., Riyong, D., Tuetun, B. & Chaiyasit, D. (2007). Aromatic plant-derived essential oil: an alternative larvicide for mosquito control. *Fitoterapia*, 78, 205-210.
- Prajapati, V., Tripathi, A.K., Aggarwal, K.K. & Khanuja, S.P.S. (2005). Insecticidal, repellent and oviposition-deterrent activity of selected essential oils against *Anopheles*

- stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*. *Bioresource technology*, 96, 1749-1757.
- Promsiri, S., Naksathit, A., Kruatrachue, M. & Thavara, U. (2006). Evaluations of larvicidal activity of medicinal plant extracts to *Aedes aegypti* (Diptera: Culicidae) and other effects on a non target fish. *Insect Science*, 13, 179-188.
- Raghavendra, K., Barik, T.K., Bhatt, R.M., Srivastava, H.C., Sreehari, U. & Dash, A.P. (2011). Evaluation of the pyrrole insecticide chlorfenapyr for the control of *Culex quinquefasciatus* Say. *Acta Tropica*, 118, 50-55.
- Rajanarendar E., Reddy, M.N., Murthy, K.R., Reddy, K.G., Raju, S., Srinivas, M., Praveen, B. & Rao, M.S. (2010). Synthesis, antimicrobial, and mosquito larvicidal activity of 1-aryl-4-methyl-3,6-bis-(5-methylisoxazol-3-yl)-2-thioxo-2,3,6,10b-tetrahydro-1H-pyrimido[5,4-c]quinolin-5-ones. *Bioorganic & Medicinal Chemistry Letters*, 20, 6052-6055.
- Ray, D. (2010). Organochlorine and Pyrethroid Insecticides. *Comprehensive Toxicology*, 13, 445-447.
- Regnault-Roger, C., Hamraul, A. (1994). Inhibition of reproduction of *Acanthoselides obtectus* Say (Coleoptera), a kidney bean (*Phaseolus vulgaris*) bruchid, by aromatic essential oils. *Crop Protection*, 13, 624-628.
- Riaz, M.A., Poupardin, R., Reynaud, S., Strode, C., Ranson, H. & David, J-P. (2009). Impact of glyphosate and benzo[a]pyrene on the tolerance of mosquito larvae to chemical insecticides. Role of detoxification genes in response to xenobiotics. *Aquatic Toxicology*, 93, 61-69.
- Rustaiyan, A., Komeilizadeh, H., Mojab, F., Khazaie, A., Masoudi, S. & Yari, M. (2001). Essential oil composition of *Peucedanum petiolare* (DC) Boiss. from Iran. *Journal of Essential Oil Research*, 13, 49-50.
- Rustaiyan, A., Monfared, A., Masoudi, S. & Ameri, N. (2002a). Essential oils of the stem and root of *Ferula galbaniflua* Boiss. et Bushe. from Iran. *Journal of Essential Oil Research*, 14, 286-287.
- Rustaiyan, A., Neekpoor, N., Rabani, M., Komeilizadeh, H., Masoudi, S. & Monfared, A. (2002b). Composition of the essential oil of *Chaerophyllum macrospermum* (Spreng.) Fisch. and C.A. Mey. from Iran. *Journal of Essential Oil Research*, 14, 216-217.
- Rustaiyan, A., Nadimi, M., Mazloomifar, H. & Massudi, S. (2005). Composition of the essential oil of *Ferula macrocolea* (Boiss.) Boiss. from Iran. *Journal of Essential Oil Research*, 17, 55-56.
- Santos, P.A.G., Figueiredo, A.C., Oliveira, M.M., Barroso, J.G., Pedro, L.G., Deans, S.G., Younous, A.K.M. & Scheffer, J.J.C. (1998). Essential oils from hairy root cultures and from fruits and roots of *Pimpinella anisum*. *Phytochemistry*, 48, 455-460.
- Sedghat, S., Rustaiyan, A., Khosravi, M., & Masoudi, S. (2003). Chemical constituents of the essential oil of *Lomatopodium khorassanicum* Mozaffarian-a species endemic to Iran. *Journal of Essential Oil Research*, 15, 416-417.
- Sefidkon, F., Khajavi, M.S. & Mirza, M. (1997). Essential oil of *Lomatopodium staurophyllum* (Rech. f.) Rech. f. *Journal of Essential Oil Research*, 9, 471-472.


- Shaalán, E., Canyon, D., Younes, M-W., Abdel-Wahad, H. & Mansour, A-H. (2005). A review of botanical phytochemicals with mosquitocidal potential. *Environment International*, 31, 1149-1166.
- Stevens, M.M., Burdett, A.S., Mudford, E.M., Helliwell, S. & Doran, G. (2011). The acute toxicity of fipronil to two non-target invertebrates associated with mosquito breeding sites in Australia. *Acta Tropica*, 117, 2 125-130.
- Sukumar, K., Perich, M-J. & Boobar, L-R. (1991). Botanical derivatives in mosquito control: a review. *Journal of the American Mosquito Control Association*, 7, 210-237.
- Tabanca, N., Demirci, B., Baser, K.H.C., Mincsovcis, E., Khan, I.A., Jacob, D.E. & Wedge, D.E. (2007). Characterization of volatile constituents of *Scaligeria tripartita* and studies on the antifungal activity against phytopathogenic fungi. *Journal of Chromatography B*, 850, 221-229.
- Tabanca, N., Demirci, B., Kirimer, N., Baser, K.H.C., Bedir, E., Khan, I.A. & Wedge, D.E. (2005). Gas chromatographic-mass spectrometric analysis of essential oils from *Pimpinella aurea*, *Pimpinella corymbosa*, *Pimpinella peregrina* and *Pimpinella puberula* gathered from Eastern and Southern Turkey. *Journal of chromatography A*, 1097, 192-198.
- Talontsi, F.M., Matasyoh, J.C., Ngoumfo, R.M. & Chepkorir, R. (2011). Mosquito larvicidal activity of alkaloids from *Zanthoxylum lemairei* against the malaria vector *Anopheles gambiae*. *Pesticide Biochemistry and Physiology*, 99, 82-85.
- Tedesco, C., Ruiz, M. & McLafferty, S. (2010). Mosquito politics: Local vector control policies and the spread of West Nile Virus in the Chicago region. *Health & Place*, 16, 1188-1195.
- Tepe, B., Akpulat, H.A., Sokmen, M., Daferera, D., Yumrutas Aydin, E., Polissiou, M. & Sokmen, A. (2006). Screening of the antioxidative properties of the essential oils of *Pimpinella anisetum* and *Pimpinella flabellifolia* from Turkey. *Food Chemistry*, 97, 719-734.
- Tilquin, M., Paris, M., Reynaud, S., Despres, L., Ravanel, P., Geremia, R.A., & Gury, J. (2008). Long Lasting Persistence of *Bacillus thuringiensis* subsp. *israelensis* (Bti) in Mosquito Natural Habitats. *PLoS ONE*, 3, e3432.
- Tirillini, B.B., Stoppini, M.A.M. & Pellegrino, R.R. (1996). Essential oil components in the epigeous and hypogeous parts of *Smyrniium perfoliatum* L. *Journal of Essential Oil Research*, 8, 611-614.
- Tirillini, B. & Tosi, B. (1992). Presence of *a*-pinene in plant callus cultures of *Smyrniium perfoliatum* L. *Journal of Essential Oil Research*, 4, 431-432.
- Tomic, A., Petrovic, S., Pavlovi, M., Tzakou, O., Couladis, M., Milenkovic, M., Vučicevic, D. & Lakušić, B. (2009). Composition and antimicrobial activity of the rhizome essential oils of two *Athamanta turbith* subspecies. *Journal of Essential Oil Research*, 21, 276-279.
- Trabousli, I, El-Haj, A.F., Tueni, S., Taoubi, M., Nader, K. & Mrad, N.A. (2005). Repellency and toxicity of aromatic plant extracts against the mosquito *Culex pipiens molestus* (Diptera: Culicidae). *Pest Management Science*, 6, 597-604.

- Tsukamoto, T., Ishikawa, Y. & Miyazawa, M. (2005). Larvicidal and Adulticidal Activity of Alkylphthalide Derivatives from Rhizome of *Cnidium officinale* against *Drosophila melanogaster*. *Journal of Agricultural and Food Chemistry*, 53, 5549-5553.
- Tuetun, B., Choochote, W., Kanjanapothi, D., Rattanachanpichai, E., Chaithong, U., Chaiwong, P., Jitpakdi, A., Tippawangkosol, P., Riyong, D. & Pitasawat, B. (2005). Repellent properties of celery, *Apium graveolens* L., compared with commercial repellents, against mosquitoes under laboratory and field conditions. *Tropical Medicine & International Health*, 10, 1190-1198.
- Tuetun, B., Choochote, W., Pongpaibul, Y., Junkum, A., Kanjanapothi, D., Chaithong, U., Jitpakdi, A., Riyong, D., Wannasan, A. & Pitasawat, B. (2009). Field evaluation of G10, a celery (*Apium graveolens*)-based topical repellent, against mosquitoes (Diptera: Culicidae) in Chiang Mai province, northern Thailand. *Parasitology Research*, 104, 3, pp 515-521
- Tunc, I., Berger, B.M., Erler, F. & Dagh, F. (2000). Onicidal activity of essential oil from five plants against two stored-product insects. *Journal of Stored Product Research*, 36, 161-168.
- Tutin, T.G., Heywood, V.H., Burges, N.A., Moore, D.M., Valentine, D.H., Walters, S.M., Webb, D.A., Ball, P.W, Chater A.O., & Ferguson I.K. (1968). *Flora Europaea* vol. 2, Cambridge University Press, pp 315-375.
- Vajs, V., Milosavljevic, S., Tesevic, V., Zivanovic, P., Jancic, R., Todorovic, B. & Slavkovska, V. (1995). *Chaerophyllum coloratum* L.: essential oils of ripe fruits and umbels. *Journal of Essential Oil Research*, 7, 529-531.
- Van Breukelen, F.R., Haemers S., Wijffels R.H. & Rinzema, A. (2011). Bioreactor and substrate selection for solid-state cultivation of the malaria mosquito control agent *Metarhizium anisopliae*. *Process Biochemistry*, 46, 751-757.
- Van den Dool, H. & Kratz, P.D.A. (1963). Generalization of the retention index system including linear temperature programmed gasliquid partition chromatography. *Journal of Chromatography A*, 11, 463-471.
- Velasco-Negueruela, A., Perez-Alonso, M.J., Perez de Paz, P.L., Pala-Paul, J. & Sanz, J. (2003). Analysis by gas chromatography-mass spectrometry of the essential oil from the aerial parts of *Pimpinella junoniae* Ceb. & Ort., gathered in La Comera, Canary Islands, Spain. *Journal of Chromatography A*, 1011, 241-244.
- Velasco-Negueruela, A., Pérez-Alonso, M.J., Pérez de Paz, P.L., Palá-Paúl, J. & Sanz, J. (2005). Analysis by gas chromatography-mass spectrometry of the essential oils from the aerial parts of *Pimpinella anagodendron* Bolle and *Pimpinella rupicola* Svent., two endemic species to the Canary islands, Spain. *Journal of Chromatography A*, 1095, 180-184.
- Verykokidou, E., Tzakou, O., Loukis, A. & Roussis, V. (1995). Chemical composition of the essential oil of *Athamanta macedonica* (L.) Sprengel subsp. *macedonica*, from Greece. *Journal of Essential Oil Research*, 7, 335-336.
- WHO. (1981). Instructions for determining the susceptibility or resistance of mosquito larvae to insecticides. Vol. WHO/VBC/81.807. 1981, Geneva: World Health Organization. p. 6.

- Yadouleton, A., Martin, T., Padonou, G., Chandre, F., Asidi, A., Djogbenou, L., Dabiré, R., Aïkpon, R., Boko, M., Glitho, I. & Akogbeto, M. (2011). Cotton pest management practices and the selection of pyrethroid resistance in *Anopheles gambiae* population in Northern Benin. *Parasites & Vectors*, 4, 60, doi: 10.1186/1756-3305-4-60.
- Zivanovic, P., Djokovic, D., Vajs, V., Slavkovska, V., Todorovic, B., & Milosavljevic, S. (1994). Essential oils flowers and fruits of *Athamanta haynaldii* Bort. Et Uchtr. (Apiaceae). *Pharmazie*, 49, 463-464.

IntechOpen

IntechOpen



Integrated Pest Management and Pest Control - Current and Future Tactics

Edited by Dr. Sonia Soloneski

ISBN 978-953-51-0050-8

Hard cover, 668 pages

Publisher InTech

Published online 24, February, 2012

Published in print edition February, 2012

Integrated Pest Management is an effective and environmentally sensitive approach that relies on a combination of common-sense practices. Its programs use current and comprehensive information on the life cycles of pests and their interactions with the environment. This information, in combination with available pest control methods, is used to manage pest damage by the most economical means and with the least possible hazard to people, property, and the environment.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Epeameinondas Evergetis, Antonios Michaelakis and Serkos A. Haroutounian (2012). Essential Oils of Umbelliferae (Apiaceae) Family Taxa as Emerging Potent Agents for Mosquito Control, Integrated Pest Management and Pest Control - Current and Future Tactics, Dr. Sonia Soloneski (Ed.), ISBN: 978-953-51-0050-8, InTech, Available from: <http://www.intechopen.com/books/integrated-pest-management-and-pest-control-current-and-future-tactics/essential-oils-of-umbelliferae-apiaceae-family-taxa-as-emerging-potent-agents-for-mosquito-control>

INTECH

open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen