

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



The Role of Landscape in Contact Zones of Sister Species of Lizards

Gabriela Cardozo¹, Sergio Naretto¹,
Marcelo Zak² and Margarita Chiaraviglio¹

¹Laboratorio de Biología del Comportamiento, Universidad Nacional de Córdoba

²Departamento de Geografía, Universidad Nacional de Córdoba
Argentina

1. Introduction

Elucidating the factors regulating the spatial distribution of ecologically similar species is a key pursuit (Dammhahn and Kappeler, 2008; Peres-Neto, 2006). Many biotic and abiotic variables might influence species distribution and determine allopatry or sympatry (Di Cola & Chiaraviglio, 2010). Moreover, species patterns are strongly associated with habitat variables at different spatial scales (Hatten & Paradzick, 2003). However, the role of landscapes in contact zones is not completely understood.

Contact zones have long been recognized as natural laboratories of evolution (Bridle et al., 2001). The geographic structure of contact zones determines dynamic evolutionary processes; however, since landscape structure influences population processes (Cardozo et al., 2007; Cardozo & Chiaraviglio, 2008) the maintenance of contact zones is likely to depend on landscape patterns.

Morphologically similar species are more likely to interact than morphologically dissimilar ones simply because a major portion of the behavioral and ecological activities of animals is associated with morphology (Losos, 1990; Pianka, 1986). Morphological similarity among coexisting animal species induces potential interactions that may lead to niche segregation (Huey, 1974; Huey & Pianka 1977). It is widely accepted that niche differentiation is often the basis for the coexistence of competitors (MacArthur & Levins, 1967; Roughgarden, 1979); however, how the coexisting species use landscape-scale resources is not clear. Interspecific competition might favour niche differentiation between competitors because it may optimise their behaviour in different ways (Law et al., 1997; Maynard Smith & Parker, 1976). Thus, niche differentiation in ecologically similar species might induce divergence of landscape-scale habitat use.

In this work, we focused on two closely related lizard species: *Tupinambis merianae* and *Tupinambis rufescens*; they are particularly interesting because they occupy the southernmost area of *Tupinambis* group distribution in South America (Peters & Donoso-Barros, 1986). *T. rufescens* would be restricted to the dry Chaco whereas *T. merianae* would occur in diverse regions (Ceï, 1993; Colli et al., 1998; Lopes & Abe, 1999) from southern Amazonia to

northern Patagonia (Carvalho et al., 2006). The contact zone of the two lizards coincides principally with the arid South American Gran Chaco.

The species have similar body size and external morphological traits, as well as overlapping macro-habitat use and general foraging mode (Castro & Galetti, 2004; Williams et al., 1993). Therefore, a potentially extensive interespecific interaction would represent a significant pressure in sympatric areas. The combination of morphological similarity, typically terrestrial habits and territoriality renders *Tupinambis* lizards ideal models for examining differential use of resources in sympatric areas based on landscape structure.

Considering that habitat loss is a serious environmental problem in many ecosystems (Ishwar et al., 2003; Luiselli & Capizzi, 1997; Mac Nally & Brown, 2001), the conservation status of landscapes in key wildlife habitats, such as contact zones, becomes strikingly relevant for species conservation. Numerous research works indicate that several species are globally threatened by habitat loss, and how changes in spatial patterns influence ecological processes has received great attention. For instance, Cardozo & Chiaraviglio (2008) found that landscape influences life history parameters and spatial distribution of reproductive individuals in snakes, leading to geographical variations in mating systems and therefore variations in reproductive potential. Furthermore, Cardozo et al. (2007) showed that landscape fragmentation affects dispersal patterns, reducing gene flow.

Investigations on landscape-scale Squamata habitats may provide essential knowledge to understand interspecific interactions and to implement measures for the conservation of herpetological communities (Filippi & Luiselli, 2006). Nevertheless, not only does habitat loss pose a threat to individual species but also landscape modifications could affect species interactions. Thus, the understanding of the associations between landscape conservation status and the distribution of sister species in contact zones could be useful to design conservation plans not only for individual species but also for ecological systems.

We examined landscape-scale habitat use in contact and allopatric zones between the two teiid lizards (*T. merianae* and *T. rufescens*) that occur in the Chaco region of central Argentina. Habitat heterogeneity is expected to increase the probability of coexistence among sister species (Tews et al., 2004). Therefore, we hypothesized divergence in landscape use in contact zones by both species, which would exploit high quality resources and take advantage of habitat heterogeneity. Within a regional context, animals that need either to maximize the availability of resources or to minimize interspecific interactions may select areas dominated by patches of a particular vegetation type (Jonshon et al., 2004).

We generated useful knowledge to guide conservation efforts including landscape-level process-oriented considerations, to contribute to avoid disruption of the evolutionary process and to ensure healthy biodiversity at all levels.

2. Methods

2.1 Study area

The study area was located in the province of Córdoba, central Argentina, which is an ideal natural scenario for the study of landscape-scale niche differentiation of *T. merianae* and *T. rufescens* because this area includes the southernmost contact zone between the species

distributions. The province of Córdoba has a central-western mountain area with a maximum elevation of 2790 m a.s.l. surrounded by vast plains of 600-900 m a.s.l. The study area lies largely within the Gran Chaco, which is the largest dry forest in South America; vegetation in the region comprises a mosaic of xerophytic forests and scrubs (Zak & Cabido, 2002). The Gran Chaco is a highly threatened wooded region, strongly affected by extensive livestock raising, extractive forestry and poorly planned agricultural expansion (Zak et al., 2004, 2008). To the east, the study area also includes the Pampas region, which was originally composed of natural grasslands but which is currently severely degraded mainly due to the advance of crop farming (Cozzani et al., 2004).

2.2 Species data

We used a database that includes approximately 700 records of the presence of *T. rufescens* and *T. meriana* in central Argentina, which were gathered during field work conducted in the framework of a major project on lizard ecology developed by our research group at the Universidad Nacional de Córdoba, Argentina. Presence records were classified according to their locality of origin (69 localities of presence of *T. meriana* and 32 localities of presence of *T. rufescens*) (Fig. 1).




Fig. 1. Localities of presence of *T. meriana* (dark green: allopatry, light green: sympatry) and *T. rufescens* (dark red: allopatry, light red: sympatry) in the province of Córdoba, central Argentina.

External polygons, considering presence of each species and the geographical characteristics of the study area were drawn. Localities were categorized as sympatric or allopatric depending on whether or not they were included within the intersection of the polygons or next to any heterospecific record of presence within a radius of 10 km (58 allopatric and 11 sympatric localities of presence of *T. merianae*, and 19 allopatric and 13 sympatric localities of presence of *T. rufescens*).

We defined the sample unit as a buffer area around the exact geographical coordinates of the locality of presence (Westphal et al., 2003). Sample units were circular plots of 2-km radius, which is equivalent to twice the area that contains the ecological range reported for other *Tupinambis* spp. (Mendoza & Noss, 2003; Winck, 2007). We intersected those areas with landscape cover features, i.e., we selected "mini-landscapes", for further characterization with landscape metrics (Westphal et al., 2003). To determine landscape availability we selected mini-landscapes at random within the distribution area delimited by the external polygon of the localities of presence for each species. We also quantified landscape availability in the contact zone considering the external polygon of the localities classified as sympatric.

2.3 Landscape analysis

The vegetation land-cover map of the province of Córdoba was created by the Multidisciplinary Institute of Plant Biology (IMBIV) of the National University of Córdoba and CONICET, Argentina (Zak, 2008). This map was obtained from the classification of Landsat 5 TM images together with phytosociological data. The researchers originally identified, described and mapped 19 land-cover types (Zak & Cabido, 2002). The classification of the Landsat imagery was based on the application of a maximum likelihood classifier using the sixth bands of the TM images and their Normalized Difference Vegetation Index (NDVI). Training sites were determined after analysis and field recognition of clusters defined by previous unsupervised classifications and the multivariate analysis of Braun-Blanquet (1950) phytosociological relevés.

We grouped the original vegetation land-cover map according to the ecological function of the land-cover types for the bioecology of the study species. Among the environmental factors that might influence behaviors in Squamata, vegetation structure would be of great importance (Blouin-Demers & Weatherhead, 2001; Chiaraviglio & Bertona, 2007; Row & Blouin-Demers, 2006) because it provides alternative thermal environments for thermoregulation (Chiaraviglio, 2006) and might affect reproductive processes and life history traits (Cardozo & Chiaraviglio, 2008; Cardozo & Chiaraviglio, 2011). Therefore, according to the complexity of the vertical structure of the land covers, we determined three major vegetation classes: forest, shrublands and low vertical structures (LVS). Forest includes lowland forests and highland forests; shrublands includes lowland scrubs and highland scrubs; and LVS vegetation includes natural grasslands, halophytes, cordgrass, palustrine vegetation, cultural vegetation, saline zones, waterlogged soils, highland grasslands and bare soils.

To obtain consistent fragmentation metrics, we refined image classification by applying a moving window using the majority analysis (Baldi et al., 2006; Cardozo et al., 2007). We assessed the configuration of the three major vegetation classes in each mini-landscape by

calculating the following fragmentation metrics: a. Class Metrics: proportional abundance in the landscape (PLAND) ; mean patch area (hectare) (AREA_MN); number of patches (NP); landscape shape index (LSI), which is a measure of the total edge of each class and increases as the patch type becomes more disaggregated; shape complexity (PARA_MN), which is calculated as mean perimeter area ratio; heterogeneity (IJI), which increases when the corresponding patch type becomes equally adjacent to all other patch types (i.e., maximally interspersed and juxtaposed to other patch types); proximity (PROX_MN), which increases as the neighborhood (defined as 2830 m to include the entire the area of the mini-landscape) is increasingly occupied by patches of the same type and as those patches become closer and more contiguous (or less fragmented) in distribution. b. Land Metrics: the total edge length in the landscape (LSI); contagion (CONTAG), which increases when all patch types are maximally aggregated and minimally interspersed (equal proportions of all pair-wise adjacencies); diversity (SHDI), which increases as the number of different patch types (i.e., patch richness, PR) increases and/or the proportional distribution of area among patch types becomes more equitable; contrast-weighted edge density (CWED) (metres per hectare), which increases as the amount of edge in the landscape increases and/or as the contrast in edges increases. We introduced the following edge contrast weights: forest-shrubs=0.25, forest-LVS=0.99, shrubs-LVS=0.75. All metrics were calculated using FragStats 3.3 (Cardozo & Chiaraviglio, 2008; McGarigal & Marks, 1995; Rutledge, 2003).

2.4 Analyses

We compared the landscape-scale habitat availability in the allopatric and sympatric localities areas of *T. merianae* and *T. rufescens* with Kruskal-Wallis test of landscape features of the random mini-landscapes. Then we compared the landscape-scale habitat use and selection between species by applying Wilcoxon test. Niche differentiation analysis was based on the comparison of landscape features of the sympatric localities between the two species. Intraspecific variations in habitat use were determined by comparing landscape features between allopatric and sympatric localities for each species by applying Wilcoxon test. We also analyzed the results in an information theoretical framework, which allowed us to examine various models including interactions among variables. We fit the generalized linear models (GLM) and employed the Akaike Information Criterion (AIC) to select the models (Westphal et al., 2003) that best identify the landscape features determining species distribution. The model with the lowest AIC was selected as the 'best' model (Mazerolle, 2006). These analyses were performed with R: A Language and Environment for Statistical Computing (2011).

3. Results

3.1 Landscape-scale habitat availability

The configuration of the available landscape presented a gradient from the distribution area of *T. merianae* to that of *T. rufescens* (SE-NW) (Table 1) of decreasing proportion (PLAND) and mean area (AREA_MN) of LVS patches, and increasing values of these metrics for the forest and shrubland areas. Along this gradient, toward the NW, LVS vegetation exhibited greater edge length because of increasing disaggregation (LSI), and increased patch shape complexity (PARA_MN). Shrublands became more interspersed (IJI), with more irregular edges (PARA_MN), than to the SE.

		Landscape availability <i>T. merianae</i> N=87	Landscape availability contact zone N=21	Landscape availability <i>T. rufescens</i> N=32	P value
<i>Class Metrics</i>					
PLAND	LVS	93.23	74.33	64.11	<0.0001
	Forest	4.26	25.64	26.09	0.0002
	Shrublands	2.51	0.03	9.80	>0.9999
AREA_MN	LVS	1347.24	849.46	684.06	<0.0001
	Forest	27.01	127.40	147.10	0.0002
	Shrublands	5.01	0.20	38.67	>0.9999
NP	LVS	1.47	3.19	4.28	>0.9999
	Forest	1.23	4.00	3.69	0.0042
	Shrublands	1.34	0.86	3.00	>0.9999
LSI	LVS	1.27	1.96	2.21	0.0004
	Forest	2.48	2.49	2.84	>0.9999
	Shrublands	3.02	1.30	2.58	>0.9999
PARA_MN	LVS	40.35	107.40	145.21	0.0005
	Forest	236.77	186.41	181.42	>0.9999
	Shrublands	262.30	635.03	382.32	0.0013
PROX_MN	LVS	21.99	28.67	54.19	>0.9999
	Forest	9.63	26.59	26.84	>0.9999
	Shrublands	16.64	0.01	21.28	>0.9999
IJI	LVS	77.77	10.71	48.06	>0.9999
	Forest	72.86	11.33	45.34	>0.9999
	Shrublands	69.73	97.25	86.62	0.0040
<i>Land Metrics</i>					
LSI		1.29	1.80	2.18	<0.0001
CWED		2.19	7.99	9.20	<0.0001
CONTAG		92.73	74.81	69.63	<0.0001
IJI		74.77	13.58	49.86	0.0005
SHDI		0.11	0.35	0.46	<0.0001

Table 1. Landscape availability in the distribution areas of *T. merianae* and *T. rufescens* in central Argentina.

The proximity (PROX_MN) among patches of the landscape cover types did not vary. Regarding mean landscape metrics, diversity (SHDI), landscape shape index (LSI) and contrast-weighted edge density (CWED) increased in the contact zone and in the distribution area of *T. rufescens*. By contrast, connectivity (CONTAG) decreased, showing more heterogeneous landscapes.

3.2 Landscape-scale habitat use and selection

Although occurring in landscapes with prevailing LVS vegetation distributed in few large patches, *T. meriana* selected landscapes with a greater proportion of forest and shrublands – about 20%– than the available average –7%– (forest : $W=6542.00$, $P<0.0001$; shrubland: $W=6166.00$, $P=0.0008$). *Tupinambis meriana* required forest patches of an average of 50 ha and shrub patches of 30 ha (Fig. 2 a).

Tupinambis rufescens also selected landscapes that are different from those available, with low proportion of LVS vegetation disaggregated in patches, and a high proportion of forest and shrubs, reaching 56%, which is slightly higher than landscape availability – 36%– (LVS: $W=1210.00$, $P<0.0223$; forest: $W=931.00$, $P=0.1240$; shrublands: $W=951.00$, $P=0.2235$). Landscapes selected by *T. rufescens* presented more forest patches than landscapes selected by *T. meriana*. *Tupinambis rufescens* required forest patches of approximately 224 ha and shrub patches of 188 ha (Fig. 2 b).




Fig. 2. Landscape use of *T. meriana* (green) and *T. rufescens* (red): (a) Percentage of landscape (PLAND); (b) Mean patch area (AREA_MN).

Species differed in the landscape-scale habitat use in terms of proportion of land cover types and mean area of the patches (LVS: PLAND, $W=946.00$, $P<0.0001$, AREA_MN, $W=955.00$, $P<0.0001$; forest, PLAND, $W=2238.5$, $P<0.0001$, AREA_MN, $W=2212.5$, $P<0.0001$; shrublands, PLAND, $W=1995.00$, $p=0.0047$, AREA_MN, $W=2010.00$, $P=0.0032$). Moreover, landscapes used by *T. rufescens* presented patches of LVS vegetation with more complex shape (PARA_MN) and more compact forest patches than landscapes used by *T. meriana*. *Tupinambis rufescens* used landscapes where LVS patches and forest patches exhibit greater proximity than landscapes used by *T. meriana*. The amount and contrast of edges (LSI and CWED) and diversity (SHDI) differed between landscapes used by both species (Table 2).

		<i>T. merianae</i> N=69	<i>T. rufescens</i> N=32	P value
<i>Class Metrics</i>				
NP	LVS	2.52	5.16	0.0001
	Forest	3.45	4.31	0.0389
	Shrublands	3.09	3.56	0.1161
LSI	LVS	1.72	2.50	0.0001
	Forest	2.68	2.68	0.6500
	Shrublands	2.96	2.73	0.6385
PARA_MN	LVS	76.87	201.44	<0.0001
	Forest	244.85	183.95	0.0132
	Shrublands	369.32	289.04	0.1980
PROX_MN	LVS	21.24	38.74	0.0003
	Forest	8.35	30.09	0.0021
	Shrublands	22.75	14.55	0.6023
IJI	LVS	58.22	56.44	0.6224
	Forest	53.35	47.75	0.6815
	Shrublands	72.92	73.85	>0.9999
<i>Land Metrics</i>				
LSI		1.77	2.31	0.0013
CWED		5.75	9.57	0.0033
CONTAG		82.60	62.56	0.0001
IJI		58.02	57.42	0.8055
SHDI		0.27	0.57	0.0002

Table 2. Landscape use of *T. merianae* and *T. rufescens* in central Argentina.

3.3 Landscape-scale habitat use and selection: Sympatry and allopatry

Tupinambis merianae both in allopatry and sympatry used similar landscapes in terms of proportions of land-cover types (LVS: $W=307.50$, $P=0.1943$; forest: $W=451.00$, $P=0.2690$; shrubs: $W=360$, $P=0.6507$). By contrast, *T. rufescens* in allopatry and sympatry used the landscape differentially. In sympatry, this species used landscapes with lower proportion of forest and shrubs, and higher proportion of LVS vegetation than in allopatry (LVS: $W=289.00$, $P=0.0042$; forest: $W=161.00$, $P=0.0399$; shrubs: $W=151.50$, $P=0.0136$). Landscape use did not differ between *T. rufescens* and *T. merianae* in sympatry, according to proportion of land cover types (LVS: $W=145.50$, $P=0.6423$; forest: $W=134.00$, $P=0.8386$; shrubs: $W=136.50$, $P=0.9483$) (Fig. 3a). Landscape-scale habitat use did not differ from landscape availability in either species (PLAND: LSV: $H=55$, $P=0.7601$; forest: $H=1.67$, $P=0.4315$; shrubs: $H=0.10$, $P=0.9401$).

Landscape-scale habitat use did not differ between *T. rufescens* and *T. merianae* in sympatry in terms of mean patch area (LVS: $W=149.50$, $P=0.4859$; forest: $W=134.00$, $P=0.8386$; shrubs: $W=136.50$, $P=0.9483$) or number of patches. *Tupinambis rufescens* in sympatry used landscapes with fewer patches of forest and shrubs than those used in allopatry (Table 3). In addition, shrubland patches were smaller and LVS vegetation patches were larger than in allopatry (LVS: $W=284.00$, $P=0.0076$; forest: $W=177.00$, $P=0.1498$; shrubs: $W=155.00$, $P=0.0197$) (Fig. 3b).




Fig. 3. Landscape use of *T. merianae* (green) and *T. rufescens* (red) in allopatry (dark) and sympatry (light). (a) Percentage of landscape (PLAND); (b) Mean patch area (AREA_MN).

Shape complexity (PARA_MN) of land cover types did not vary within or between species in sympatry (Table 3). *Tupinambis rufescens* in sympatry used landscapes with more dispersed forest patches (PROX_MN) than those used in allopatry. Landscapes used by *T. rufescens* in sympatry presented lower amount of edge (LSI) than those used in allopatry. *Tupinambis merianae* in sympatry used landscapes with shrubland patches more dispersed and with lower amount of edge than those used in allopatry.

		T. <i>merianae</i> allopatry N=58	T. <i>merianae</i> sympatry N=11	P value	T. <i>rufescens</i> allopatry N=19	T. <i>rufescens</i> sympatry N=13	P value	P value Interspecific sympatry
Class Metrics	NP							
	LVS	2.59	2.18	0.8619	5.32	4.92	0.2704	0.1541
	Forest	3.40	3.73	0.2711	5.68	2.31	0.0020	0.3034
	Shrublands	3.55	0.64	0.4358	5.00	1.46	0.0084	0.9739
LSI	LVS	1.70	1.85	0.2344	2.76	2.14	0.0654	0.6843
	Forest	2.78	2.32	0.4050	2.78	2.45	0.3893	0.7440
	Shrublands	3.24	1.52	0.0190	2.93	2.09	0.1265	0.2809
PARA_MN	LVS	80.35	58.51	0.4166	229.22	162.98	0.0926	0.3529
	Forest	248.15	233.32	0.2870	202.17	145.50	0.1761	0.1777
	Shrublands	335.32	539.34	0.0613	287.42	294.24	0.8359	0.1473
PROX_MN	LVS	20.67	24.24	0.8061	48.75	24.87	0.0988	0.2314
	Forest	10.28	1.57	0.1304	40.33	8.49	0.0581	0.0972
	Shrublands	27.30	0.0016	0.0033	17.88	3.90	0.4035	0.1258

		<i>T. merianae</i> allopatry N=58	<i>T. merianae</i> sympatry N=11	P value	<i>T. rufescens</i> allopatry N=19	<i>T. rufescens</i> sympatry N=13	P value	P value Interspecific sympatry	
Land Metrics	IJI	LVS	66.51	16.77	0.0105	55.44	60.19	0.8415	0.2207
		Forest	57.40	33.08	0.3301	45.35	56.77	0.6892	0.8065
		Shrublands	71.81	78.52	0.1722	74.83	70.19	0.6886	0.5221
	LSI	1.79	1.66	0.4460	2.64	1.83	0.0015	0.8845	
	CWED	5.65	6.27	0.3355	11.07	7.39	0.1028	0.9537	
	CONTAG	82.59	82.70	0.3614	57.21	70.39	0.1112	0.3236	
	IJI	65.71	19.58	0.0059	56.06	62.53	0.4839	0.2207	
	SHDI	0.28	0.23	0.4612	0.68	0.40	0.0148	0.2958	

Table 3. Comparison of landscape use in allopatry and sympatry of *T. merianae* and *T. rufescens* in central Argentina.

The amount of edge of the land-cover types in the landscapes did not vary between species in sympatry. *Tupinambis rufescens* in sympatry used less diverse landscapes (SHDI) than in allopatry. *Tupinambis merianae* in sympatry used less interspersed landscapes (IJI) than in allopatry because the LVS vegetation was less juxtaposed than in allopatry. Land metrics and proximity of land-cover types did not differ between species in sympatry.

3.4 Model selection

According to the lowest AIC, the presence of *T. merianae* at landscape scale is determined by the proportion and the area of the patches of the shrublands, and the number of patches of forest (residual deviance: 80.55, AIC: 86.55), and the presence of *T. rufescens* by the mean area of the patches of shrublands and the proportion of forest in the landscape (residual deviance: 81.63, AIC: 87.63).

4. Discussion

The role of the landscape for reptiles has been largely discussed (Blouin-Demers & Weatherhead, 2001; Cardozo et al., 2007; Driscoll, 2004; Luiselli & Capizzi, 1997; Mac Nally & Brown, 2001; Marchand & Litvaitis, 2004; Stow & Sunnucks, 2004). However, the novelty of our approach lies in the importance of landscape conservation to the maintenance of ecological interactions between lizard sister species. Since landscape ecology analysis is useful to gain a better understanding of environmental suitability (Fouquet et al., 2010), the present work provides useful knowledge for the conservation of these species.

Identifying the habitat characteristics that regulate the ecological processes of reptiles is imperative to determine threats to the species (Urbina-Cardona et al., 2006). Our results showed that the configuration of the available landscape presented a gradient from the distribution area of *T. merianae* to that of *T. rufescens* (SE-NW) of decreasing proportion of mean area of LVS patches and increasing proportion of forest and shrublands area. Along

this gradient, the landscape became more heterogeneous. These results show that landscape configuration is a main factor regulating the spatial distribution of ecologically similar species and has a central role in contact zones. Therefore, changes in distribution-related factors at landscape scale, such as habitat loss, might pose a threat for herpetofauna (Filippi & Luiselli, 2000).

Moreover, besides understanding how landscape patterns provide resource heterogeneity in the species distribution areas, it is vitally important to elucidate how these species use that heterogeneity i.e., species might reveal diverse responses to the spatial variations in habitat resources (Cagle, 2008; Urbina-Cardona et al., 2006). We observed that species are selective on landscape patterns; for example, although occurring in landscapes with prevailing LVS vegetation distributed in few large patches, *T. merianae* selected landscapes with a greater proportion of forest and shrubs. *Tupinambis rufescens* also selected landscapes that are different from those available, with low proportion of LVS vegetation disaggregated in patches, and a high proportion of forest and shrubs. Considering the relevance of forest and shrublands to both species, the present results might guide conservation efforts including landscape-level process-oriented considerations.

Although morphological similarity among species induces niche similarity because behavioral and ecological activities of animals are associated with morphology (Losos, 1990; Pianka, 1986) we observed that these sister species differ in several features of the landscape-scale habitat use (e.g., shape complexity, patch proximity, length of edges and diversity). Taking into account the diversity of responses of the species, conservation strategies should be species-specific (Keogh et al., 2001). Our results indicate that *T. rufescens* and *T. merianae* have complex habitat requirements. Specialized habitat requirements make species more vulnerable to extinction (CITES, 2010; Santos et al., 2009; Webb et al., 2002). Since *T. rufescens* and *T. merianae* showed differences in landscape-scale habitat requirements, planning of conservation strategies should consider such interspecific heterogeneity.

Although the results obtained showed that species differed in the landscape-scale habitat use in their distribution areas, in the contact zone where species are in sympatry, and therefore have the same landscape-scale habitat availability, interspecific interactions would represent a significant pressure on the use of the resources. When we analyzed niche differentiation in terms of landscape-scale resources, we observed that species did not differ in the use of landscape resources in contact zones. Both species used landscapes with similar proportions of land-cover types, mean patch area and number of patches; further research is needed to elucidate if the coexistence of these species could be explained by niche divergence at local scale. Furthermore, the results obtained enable us to get a better understanding of the strategies of the species in sympatric zones in terms of landscape-scale habitat use and selection. *Tupinambis merianae* both in allopatry and sympatry used similar landscapes. By contrast, *T. rufescens* in allopatry and sympatry used the landscape differentially, showing niche modification. In sympatry this species used landscapes with lower proportion of forest and shrubs, and higher proportion of LVS vegetation than in allopatry. Therefore, despite their ecological and morphological similarity, the species respond differently to spatial changes in landscape structure. Moreover, we remark the importance of landscape heterogeneity for the maintenance of species interactions in the contact zone.

5. Conclusion

Understanding the associations between landscape conservation status and distribution of sister species in contact zones might be useful to design conservation plans not only for individual species but also for ecological systems. Among the weaknesses of the regional conservation plans, poor information regarding behavioural ecology is one of the fundamental issues (The Nature Conservancy et al., 2005). To know whether species might be threatened by habitat change it is necessary to determine the relation between ecological processes and environmental patterns (Cardozo & Chiaraviglio, 2008). Our study provides useful knowledge about the important role of native forest and shrublands in allopatric and sympatric distribution areas of the lizard species. Furthermore, similar allopatry-sympatry systems might be occurring in this contact zone, which –as we mentioned above– coincides principally with the arid South American Gran Chaco. Considering that only 9% of the South American Gran Chaco is protected (The Nature Conservancy et al., 2005), we underline the need for efficient control of deforestation, protection of forest remnants and establishment of corridors. According to Beaudry et al. (2010) regional-scale conservation planning has to answer specific questions, such as the type of habitat that is needed and where it should be protected. The present work provides information that may be useful to guide conservation plans. Efforts to prevent habitat loss should involve preserving not only allopatric areas but also these critical heterogeneous sympatric areas where biological interactions might modify ecological processes of species (Brito et al., 2009; Santos et al., 2009).

6. Acknowledgments

We are grateful to the institutions that funded this study: Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Fondo para la Investigación Científica y Tecnológica (FONCyT), MinCyT Córdoba -Préstamo BID-PID No. 013/2009-, Secretaría de Ciencia y Tecnología (SeCyT) and Universidad Nacional de Córdoba, Argentina. GC is scientist of the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). SN is student of the Biological Sciences Doctorate of the Universidad Nacional de Córdoba and fellowship holder of the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET). MZ is Professor of the National University of Córdoba and senior scientist at the IMBIV. MCH is Professor and senior scientist of the National University of Córdoba.

7. References

- Baldi, G., Guerschman, J.P. & Paruelo, J.M. (2006). Characterizing fragmentation in temperate South America grasslands. *Agr. Ecosyst. Environ*, Vol.116, pp. 197–208
- Beaudry, F., Pidgeon, A.M., Radeloff, V.C., Howe, R.W., Mladenoff, D.J. & Bartelt, G.A. (2010). Modeling regional-scale habitat of forest birds when land management guidelines are needed but information is limited. *Biological Conservation*, Vol.143, pp. 1759-1769
- Blouin-Demers, G. & Weatherhead, P.J. (2001). Habitat use by black rat snakes (*Elaphe obsoleta obsoleta*) in fragmented forests. *Ecology*, vol.82, pp. 2882–2896
- Braun-Blanquet, J. (1950). *Sociologia Vegetal*. ACME, Buenos Aires
- Bridle J.R., Baird S. & Butlin, R.K. (2001). Spatial structure and habitat variation in a grasshopper hybrid zone. *Evolution*, Vol.55, pp. 1832–1843

- Brito, J.C., Acosta, A.L., Álvares, F. & Cuzin, F. (2009). Biogeography and conservation of taxa from remote regions: An application of ecological-niche based models and GIS to North-African Canids. *Biological Conservation*, Vol.142, pp. 3020-3029
- Cagle, N. (2008). Snake species distributions and temperate grasslands: a case study from the American tallgrass prairie. *Biological Conservation*, Vol.141, pp. 744-755
- Cardozo, G., Rivera, P.C., Lamfri, M., Scavuzzo, M., Gardenal, C. & Chiaraviglio, M. (2007). Effects of habitat loss on the genetic structure of the Argentine Boa Constrictor (*Boa constrictor occidentalis*) populations, In: *Biology of the Boa and Pythons*, Henderson, R., Powell, R. (Eds.), pp. (329-338), Eagle Mountain Publishing, Utah
- Cardozo, G. & Chiaraviglio, M. (2008). Landscape changes influence the reproductive behavior of a key capital breeder snake (*Boa constrictor occidentalis*) in the Gran Chaco region. *Biological Conservation*, Vol.141, (2008), pp. 3050-3058.
- Cardozo, G. & Chiaraviglio, M. (2011). Phenotypic plasticity of life history traits in relation to reproductive strategies in *Boa constrictor occidentalis*. *Evolutionary ecology*, (February 2011)
- Carvalho, R.L., Antoniazzi, M.M., Jared, C., Silva, A.M.J., Santos, A.A. & Egami, M.I. (2006). Morphological, cytochemical, and ultrastructural observations on the blood cells of the reptile *Tupinambis merianae* (Squamata). *Comp Clin Pathol*, Vol.15, pp. 169-174
- Castro, E.R. & Galetti, M. (2004). Frugivoria e dispersao de sementes pelo lagarto Teiu *Tupinambis merianae* (Reptilia: Teiidae). *Papéis Avulsos de Zoologia*, Vol.44, No.6, pp. 91-97
- Cei, J.M. (1993). *Reptiles del noroeste, nordeste y este de la Argentina*. Mus. Reg. Sci. Nat. Monografía XIV, Torino.
- Chiaraviglio, M. & Bertona, M. (2007). Reproduction and termoregulation as main factors influencing habitat choice in the Argentine Boa constrictor. In: *Biology of the boa and pythons*, Henderson R, Powell R (Eds), pp. (478-488), Eagle Mountain Publishing, Utah
- Chuvieco, E. (2002). *Teledetección Ambiental: la observación de la Tierra desde el Espacio* (first ed.) Ariel, Barcelona
- CITES- Convención sobre el Comercio Internacional de Especies Amenazadas de Fauna y Flora Silvestres (CITES). 2010. Criterios biológicos para la inclusión de especies. <http://www.cites.org/esp/res/09/09-24R14.shtml#FN0>
- Colli, G.R., Peres, A.K. & Cunha, H.J. (1998). A new species of *Tupinambis* (Squamata: Teiidae) from central Brazil, with an analysis of morphological and genetic variation in the genus. *Herpetologica*, Vol.54, pp. 477-492
- Cozzani, N.C., Sanchez, R. & Salba S. (2004). Nidificación de la loica pampeña (*Sturnella defilippii*) en la provincia de Buenos Aires. Argentina. *Hornero*, Vol.19, No.2, pp. 47-52
- Dammhahn, M. & Kappeler, P. (2008). Small-scale coexistence of two mouse lemur species (*Microcebus berthae* and *M. murinus*) within a homogeneous competitive environment. *Oecologia*, Vol.157, pp.473-483
- Di Cola, V. & Chiaraviglio, M. (2010). Establishing species' environmental requirements to understand how the southernmost species of south american pitvipers (*Bothrops*, Viperidae) are distributed: a niche based modelling approach. *Austral Ecology*, Vol. 36, pp. 90-98
- Driscoll, D.A. (2004). Extinction and outbreaks accompany fragmentation of a reptile community. *Ecol. Appl*, Vol.14, pp. 220-240

- Filippi, E. & Luiselli, L. (2000). Status of the Italian snake fauna and assessment of conservation threats. *Biological Conservation*, Vol.93, pp. 219–225
- Filippi, E. & Luiselli, L. (2006). Changes in community composition, habitats and abundance of snakes over 10 + years in a protected area in Italy: conservation implications. *Herpetol. J*, Vol.16, pp.29–36
- Fouquet, A., Ficetola, G.F., Haigh, A. & Gemmill, N. (2010). Using ecological niche modelling to infer past, present and future environmental suitability for *Leiopelma hochstetteri*, an endangered New Zealand native frog. *Biological Conservation*, Vol.143, pp. 1375–1384
- Goodman, R.M., Burton, F.J. & Echternacht, A.C. (2005). Habitat use of the endangered iguana *Cyclura lewisi* in a human-modified landscape on Grand Cayman. *Animal Conservation* (2005) Vol.8, pp. 397–405
- Guisan, A. & Zimmermann, N.E. (2000). Predictive habitat distribution models in ecology. *Ecol. Model.*, Vol.135, pp. 147–186
- Hatten, J. R. & Paradzick, C.E. (2003). A multiscaled model of southwestern willow flycatcher breeding habitat. *Journal of Wildlife Management*, Vol. 67, No. 4, (October 2003), pp. 774–815
- Huey, R.B. (1974). Ecological Character Displacement in a Lizard. *Amer. Zool*, Vol.14, No.4, pp. 1127–1136
- Huey, R.B. & Pianka, E.R. (1977). Patterns of niche overlap among broadly sympatric versus narrowly sympatric Kalahari lizards (Scincidae: *Mabuya*). *Ecology*, Vol.58, pp. 119–128
- Ishwar, N.M., Chellam, R., Kumar, A. & Noo, B.R. (2003). The response of agamid lizards to rainforest fragmentation in the Southern Western Ghats, India. *Conserv. Soc.* Vol.1, pp. 79–86
- Johnson, C.J., Boyce, M.S., Mulders, R., Gunn, A., Gau, R.J., Cluff, H.D. & Case, R.L. (2004). Quantifying patch distribution at multiple spatial scales: applications to wildlife-habitat models. *Landscape Ecology*, (2004), pp. 869–882
- Keogh, J.S., Barker, D.G. & Shine, R. (2001). Heavily exploited but poorly known: systematics and biogeography of commercially harvested pythons (*Phyton curtus* group) in Southeast Asia. *Biological Journal of Linnean Society*, Vol.79, pp. 113–129
- Law, R., Mallow, P. & Dieckman, U. (1997). On evolution under asymmetric competition. *Evol Ecol* Vol.11, pp. 485–501
- Lopes, H.R. & Abe, A.S. (1999). Biología reproductiva e comportamiento do teiú, *Tupinambis merrianae*, em cativeiro (Reptilia, Teiidae). In: *Manejo y Conservación de Fauna Silvestre en América Latina*. Fang, T.G., Montenegro, O.L., Bodmer, R.E. (Eds.), Editorial Instituto e Ecología, pp. (259–272) La Paz, Bolivia
- Losos J.B. (1990). The evolution of form and function: morphology and locomotor performance in West Indian *Anolis* lizards. *Evolution*, Vol. 44, pp. 1189–1203
- Luiselli, L. & Capizzi, D. (1997). Influences of area, isolation and habitat features on distribution of snakes in Mediterranean fragmented woodlands. *Biodivers. Conserv.* Vol.6, pp. 1339–1351
- Mac Nally, R. & Brown, G.W. (2001). Reptiles and habitat fragmentation in the box-ironbark forests of central Victoria, Australia: predictions, compositional change and faunal nestedness. *Oecologia*, Vol.128, pp. 116–125.
- MacArthur, R. & Levins, R. (1967). The limiting similarity, convergence and divergence of coexisting species. *The American Naturalist*, Vol.101, No.921, pp. 377–385

- Mc Garigal, K. & Marks, B. (1995). FRAGSTATS: spatial analysis program for quantifying landscape structure. USDA Forest Service General Technical Report PNW-GTR-351.
- Marchand, M.N. & Litvaitis, J.A. (2004). Effects of habitat features and landscape composition on the population structure of a common aquatic turtle in a region undergoing rapid development. *Conserv. Biol*, Vol.18, pp. 758–767
- Maynard Smith, J. & Parker, G.A. (1976) The logic of asymmetric contests. *Anim Behav.*, Vol.24, pp. 159-75
- Mazerolle, M. (2006) Improving data analysis in herpetology: using Akaike's information criterion (AIC) to assess the strength of biological hypotheses. *Amphib Reptil*, Vol.27, pp. 169–180
- Mendoza, F. & Noss, A. (2003). Radiotelemetría del peni (*Tupinambis rufescens*) en una zona del Chaco boliviano. In: *Manejo de fauna silvestre en Amazonía y Latinoamérica*. Selección de trabajos V Congreso Internacional. Rocío Polanco-Ochoa, ed. pp. (178-180), CITES, Fundación Natura, Bogotá
- Pearce, J. & Ferrier, S. (2000). Evaluating the predictive performance of habitat models developed using logistic regression. *Ecol. Modell*, Vol.133, pp. 225-245
- Peres-Neto P.R. (2006). Patterns in the co-occurrence of fish species in streams: the role of site suitability, morphology and phylogeny versus species interactions. *Oecologia*, Vol.140, pp. 352–360
- Peters, J.A. & Donoso-Barros, R. (1986). *Catalogue of the Neotropical Squamata. Part II. Lizards and Amphisbaenians*. Rev. ed. Smithsonian Institution Press, Washington, DC.
- Pianka E.R. (1986). *Ecology and natural history of desert lizards: analyses of the ecological niche and community structure*. Princeton University Press, Princeton, NJ.
- R: A Language and Environment for Statistical Computing. (2011). R Development Core Team. R Foundation for Statistical Computing. Vienna, Austria. ISBN 3-900051-07-0. <http://www.R-project.org>
- Roughgarden J. 1979. *Theory of population genetics and evolutionary ecology: an introduction*. Macmillan, New York.
- Row, J.R. & Blouin-Demers, G. (2006). Thermal quality influences effectiveness of thermoregulation, habitat use, and behaviour in milk snakes. *Oecologia*, Vol.148, pp. 1–11
- Rutledge, D. (2003). *Landscape Indices as Measures of the Effects of Fragmentation: can Pattern Reflect Process?* DOC Science International Series 98. Department of Conservation, Wellington
- Santos, X., Brito, J.C., Caro, J., Abril, A.J., Lorenzo, M., Sillero, N. & Pleguezuelos, J. M. (2009). Habitat suitability, threats and conservation of isolated populations of the smooth snake (*Coronella austriaca*) in the southern Iberian Peninsula. *Biological Conservation*, Vol.142, pp. 344-352
- Schaefer, J.F., Duvernell, D.D. & Kreiser, B.R. (2011). Ecological and genetic assessment of spatial structure among replicate contact zones between two topminnow species. *Evol Ecol* DOI 10.1007/s10682-011-9461-2
- Stow, A.J. & Sunnucks, P. (2004). Inbreeding avoidance in Cunningham's skinks (*Egernia cunninghami*) in natural and fragmented habitat. *Mol. Ecol.*, Vol.13, pp. 443–447
- Templeton, A.R., Robertson, R.J., Brisson, J., & Strasburg, J. (2001). Disrupting evolutionary processes: The effect of habitat fragmentation on collared lizards in the Missouri Ozarks. *PNAS*, Vol.98, No.10, (May 2001), pp. 5426–5432

- Tews, J., Brose, U., Grimm, V., Tielborger, K., Wichmann, M.C., Schwager, M. & Jeltsch, F. (2004). Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone structures. *J Biogeogr*, Vol.31, pp. 79–92
- The Nature Conservancy (2005), *Evaluación Ecorregional del Gran Chaco Americano/Gran Chaco Americano*. Fundación para el Desarrollo Sustentable del Chaco (De.S.del Chaco) y Wildlife Conservation Society Bolivia (WCS). Fundación Vida Silvestre. Buenos Aires. Argentina
- Urbina-Cardona, J.N., Olivares-Perez, M. & Reynoso, V.H. (2006). Herpetofauna diversity and microenvironment correlates across a pasture–edge–interior ecotone in tropical rainforest fragments in the Los Tuxtlas Biosphere Reserve of Veracruz, Mexico. *Biological Conservation*, Vol.132, pp. 61–75
- Webb, J., Brook, B.W. & Shine, R. (2002). What makes a species vulnerable to extinction? Comparative life-history traits of two sympatric snakes. *Ecol. Res*, Vol.17, pp. 59–67
- Westphal, M.I., Field, S.A., Tyre, A.J., Paton, D. & Possingham, H.P. (2003). Effects of Landscape Pattern on Bird Species Distribution in the Mt. Lofty Ranges, South Australia. *Landscape Ecology* Vol.18, (2003), pp. 413–426
- Williams, J.D., Donadio, O.E. & Ré, I. (1993). Notas relativas a la dieta de *Tupinambis rufescens* (Reptilia: Sauria) del noroeste argentino. *Neotropica*, Vol.39, pp. 45–51
- Winck, G.R. (2007). *Historia natural de Tupinambis merianae (Squamata, Teiidae) na estacao ecologica do Taim. Extremo sul do Brasil*. Santa Maria. RC., Brasil
- Zak, M.R. & Cabido, M. (2002). Spatial patterns of the Chaco vegetation of central Argentina: integration of remote sensing and phytosociology. *Applied Vegetation Science*, Vol.5, pp. 213–226
- Zak, M.R., Cabido, M. & Hodgson, J. (2004). Do subtropical seasonal forests in the Gran Chaco, Argentina, have a future? *Biological Conservation*, Vol.120, pp. 589–598.
- Zak, M. & Cabido, M. (2008). What Drives Accelerated Land Cover Change in Central Argentina? Synergistic Consequences of Climatic, Socioeconomic, and Technological Factors. *Environmental Management*, Vol.42, pp. 181–189

IntechOpen



Perspectives on Nature Conservation - Patterns, Pressures and Prospects

Edited by Prof. John Tiefenbacher

ISBN 978-953-51-0033-1

Hard cover, 270 pages

Publisher InTech

Published online 29, February, 2012

Published in print edition February, 2012

Perspectives on Nature Conservation demonstrates the diversity of information and viewpoints that are critical for appreciating the gaps and weaknesses in local, regional and hemispheric ecologies, and also for understanding the limitations and barriers to accomplishing critical nature conservation projects. The book is organized to emphasize the linkages between the geographic foci of conservation projects and the biological substances that we conceptualize as "nature", through original research. The reader moves through perspectives of diminishing spatial scales, from smaller to larger landscapes or larger portions of the Earth, to learn that the range of factors that promote or prevent conservation through the application of scholarship and academic concepts change with the space in question. The book reflects disciplinary diversity and a co-mingling of science and social science to promote understanding of the patterns of, pressures on and prospects for conservation.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Gabriela Cardozo, Sergio Naretto, Marcelo Zak and Margarita Chiaraviglio (2012). The Role of Landscape in Contact Zones of Sister Species of Lizards, Perspectives on Nature Conservation - Patterns, Pressures and Prospects, Prof. John Tiefenbacher (Ed.), ISBN: 978-953-51-0033-1, InTech, Available from: <http://www.intechopen.com/books/perspectives-on-nature-conservation-patterns-pressures-and-prospects/the-role-of-landscape-in-contact-zones-of-sister-species-of-lizards>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen