

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Hepatocellular Carcinoma: Epidemiology and Etiology

Davide Degli Esposti^{1,2}, Morando Soffritti^{3,*}, Antoinette Lemoine^{1,2},
Eva Tibaldi³ and Marco Manservigi³

¹AP-HP, Hôpital Paul Brousse, Service de Biochimie et Biologie
Moléculaire, Inserm; Université Paris-Sud 11;
PRES Universud-Paris; Paul Vaillant Couturier

²Laboratoire de Biochimie et Biologie Cellulaire, Faculté de Pharmacie,
Université Paris-Sud 11, Jean Baptiste Clément

³Cesare Maltoni Cancer Research Center, Ramazzini Institute,
Castello di Bentivoglio, Via Saliceto, Bentivoglio, Bologna,

^{1,2}France

³Italy

1. Introduction

Hepatocellular carcinoma (HCC) is a major public health problem, accounting for about 600,000 deaths in the world in 2004 (WHO, 2008). HCC is the sixth most common cancer worldwide with about 500,000 new cases annually, representing the third largest cause of cancer-related death (Parkin, 2005; Ferlay et al., 2010). A slight decrease in the HCC incidence has been reported in high-rate areas, such as China and Japan (McGlynn et al., 2001). However, a steadily increasing trend has been reported in historically low-rate countries, particularly the United States and some European countries, such as Italy, France, UK and Germany (IARC, 2008a; El-Seragh et al., 2007). In particular, HCC incidence rates doubled in the United States in the period 1985-2002, an earlier age of onset has been observed (with a shift towards 45-60 years old), and HCC has become the fastest growing cause of cancer-related death in men (El-Seragh et al., 2004). Interestingly, it has been reported that in the United States 15-50% of HCC patients had no established risk factors, such as viral hepatitis infections, heavy alcohol consumption or aflatoxin B1 exposure (El-Seragh et al., 2007). Moreover, approximately 10% of all HCC cases in the USA occur in patients with non-cirrhotic livers (Shaw & Shah, 2011). In Europe an analysis of mortality rates from HCC trends in the last 20 years has shown increasing rates for men in 11 countries and for women in 6 countries out of 17 whose data were considered (Bosetti et al., 2008).

The observed increase in the incidence rates of HCC has been concomitant with the obesity epidemic observed in the last 30 years in western countries. Obesity is one of the clinical

* Corresponding Author

manifestations of metabolic syndrome and, in the last decade, epidemiological and experimental studies have shown that metabolic syndrome and high fat diets are associated with an increased risk of HCC incidence/mortality (Bugianesi 2007; Starley et al., 2010; Welzel et al., 2011). However, other causes may be involved in the increased incidence of HCC and chemical-induced liver carcinogenesis appears to be a less considered etiology. In this chapter, we will review recent acquisitions in epidemiology and experimental studies on HCC and will focus on chemical risk factors and possible new mechanisms of liver carcinogenesis, in particular those concerning metabolic disruption.

2. Chemical risk agents of hepatocellular carcinoma

Most human HCC occurs following viral hepatitis (mainly HBV or HCV) infections or aflatoxin B1 exposure caused by ingestion of contaminated food (IARC 2008a). However, the epidemiological evidence shows that the human liver is susceptible to chemical-induced carcinogenesis (Blonski et al., 2010; Degli Esposti et al., 2009) and the increased incidence of HCC in patients not having established risk factors (El-Seragh et al., 2007) suggests that some underestimated or new but still not recognized risk factors exist (Blonski et al., 2010). In particular, many natural and artificial agents have been shown by experimental or epidemiological studies to induce HCC (Table 1). In this section we will review the chemical risk factors of HCC as emerging from the epidemiological and experimental data.

2.1 Human hepatocarcinogens

Various classes of chemicals are reported to induce HCC in humans: drugs or hormonal therapies (azathioprine, tamoxifen and estrogen-progesteron oral contraceptives) (IARC, 2011); radioisotopes or heavy metals (Plutonium-239, Radium-224, Thorium-232; arsenic in drinking water) (IARC, 2001; IARC, 2004b); complex mixtures of polyaromatic hydrocarbons (PAH) and combustion products (soots and tobacco smoking)(IARC, 1987; IARC, 2004a); organochlorines such as vinyl chloride monomer (VCM) or 2,3,7,8 tetrachloride-dibenzo-para-dioxin (TCDD) (IARC, 2008b, IARC, 1997); and plant derivatives (betel or *Areca catechu*) (IARC, 2004c). Recently, some psychoactive substances, like cannabinoids, have been reported to worsen liver steatosis and fibrosis, in particular in the presence of HCV infections (Hézode et al., 2008, Parfieniuk & Flisiak, 2008). However, no evidence of carcinogenicity has been shown for delta 9-tetrahydrocannabinol (the principal psychoactive ingredient in marijuana) in rats and mice (Chan et al., 1996). More research is warranted to assess the long-term carcinogenic or co-carcinogenic effects of cannabinoids, particularly in the liver, as assumption of them during cannabis smoking may result in cannabinoid exposure for a large population. Finally, recent reviews have focused on a possible underestimation of non-viral causes of HCC (Blonski et al., 2010; Degli Esposti et al., 2009). In particular, metabolic disorders (Non-Alcoholic Fatty Liver Disease (NAFLD), obesity and diabetes), hormonal drugs (oral contraception, tamoxifen), organochlorine compounds, polycyclic aromatic hydrocarbons, tobacco smoking, betel quid chewing and dietary exposures (in particular arsenic in drinking water and aflatoxin B1, a well known hepatocarcinogen) are indicated as important contributing factors for HCC (Blonski et al., 2010; Degli Esposti et al., 2009).

Agents		Human exposure	Evidence of carcinogenicity		References
Category	Type		Humans	Experimental animals	
Natural	1. Aflatoxins	food contaminant (rice, peanuts, etc.)	+	+	Wogan and Newbern 1967; Wogan et al. 1974; Yeh et al. 1985; Olsen et al. 1988; IARC 1993; Soffritti et al. 1988
	2. Alcohol	lifestyle dependent	+		Hakulinen et al. 1974; Adelstein and White 1976; Hirayama 1981; IARC 1988
	3. Hepatitis B virus	blood transfusion	+	+	Snyder et al. 1982; Buendia 1992
	4. Sterigmatocystin	food contaminant (grain, legumes)		+	Purchase and van der Watt 1970
	5. Luteoschirina	food contaminant rice		+	Uraguchi et al. 1972
	6. Cyclochlorotina	food contaminant rice		+	IARC 1976
	7. Pyrazolidinic Alkaloids	plants contaminant		+	Swoboda and Reddy 1972
	8. Cycasin	alimentary exposure		+	Laquer et al. 1963
	9. Safrole	flavouring substance		+	Long et al. 1963; Hagan et al. 1965

+ = strong evidence; (+) = limited evidence

Table 1. Agents inducing Hepatocellular Carcinoma based on experimental/epidemiological evidence (Part I)

Agents		Human exposure	Evidence of carcinogenicity		References
Category	Type		Humans	Experimental animals	
Artificial	1. Thorotrast	iatrogenic	(+)	+	Guimares et al. 1955; Commission of European Communities (CEC) 1984
	2. Radioactive colloidal gold	iatrogenic		+	Upton et al. 1956
	3. Gamma radiation	occupational or accidental			Upton et al. 1968
	4. Vinyl Chloride	occupational	(+)	+	Gokel et al. 1976; Koischwitz et al. 1981; Evans et al. 1983; Maltoni et al. 1984; Dietz et al. 1985, Pirastu et al. 1990
	5. Benzidine	occupational		+	IARC 1982
	6. 2-Acetylamino-fluorene	occupational		+	Wilson et al. 1941; Teebor and Becker 1971
	7. 4-Diethylamino-azobenzene	occupational		+	Kinosita 1936 Terayama 1967
	8. Dimethyl-nitrosamina	occupational		+	Magee and Barnes 1956
	9. Diethylnitrosamina	occupational		+	Schmal et al. 1960
	10. Steroidal oral contraceptives	iatrogenic	(+)	(+)	Klatskin 1977; Jick and Hermann 1978; IARC 1979
	11. Androgen steroids	iatrogenic	(+)		Johnson et al. 1972

+ = strong evidence; (+) = limited evidence

Table 1. Agents inducing Hepatocellular Carcinoma based on experimental/epidemiological evidence (Part II)

The diversity of chemical agents that induce liver tumors in humans may be at least partially explained by the multiplicity of molecular pathways that have been found altered both in human and animal hepatic tumors (Degli Esposti et al., 2009; Saffroy et al., 2007).

Hormonal-induced liver tumors, such those observed after estrogen-progesteron oral contraceptives and tamoxifen administration, have been in part explained by promotion of the epithelium proliferation and the generation of reactive oxygen species (ROS) caused by estrogen reactive metabolites (Russo & Russo, 2006) and ER-dependent liver responses, such as hepatocyte mitogenesis (Vickers et al., 1991). Liver carcinogenicity of alpha-particle emitters, in particular Plutonium-239, Radium-224, Thorium-232, and heavy metals such as arsenic may be explained by their ability to induce direct or indirect genotoxic damage (Lehnert et al., 1997; IARC, 2004b). The carcinogenicity of VCM has been linked to the capacity of its two active metabolites (chloroethylene oxide and chloroacetaldehyde) to react with nucleic acid bases to form adducts (IARC, 2008b) and to induce p53 mutations both in humans and in rats (Barbin et al., 1997). It is worth noting that 10-40% of HCC are characterized by global genomic instability as shown by microsatellite analyses and that the exact mechanisms behind this instability still need to be assessed (Salvucci et al., 1999, Chiappini et al., 2004).

Gene polymorphisms and gene-environment interactions may also be risk factors for HCC, particularly concerning gene coding for metabolizing enzymes (glutathione-S-transferase, epoxide hydrolase, cytochrome p4502E1) or DNA repairing enzymes (XRCC1, UDP-glucuronosyltransferase1A7)(Wong et al., 2000, Borentain, 2007). However, until now, these studies have reported contrasting results. Meta-analysis and additional studies with larger samples should be performed in order clarify the role of genetic polymorphisms in the onset of HCC (El-Seragh, 2007; White et al., 2008).

Overall, these data clearly indicate that the human liver is sensitive to chemical-induced carcinogenesis and that xenobiotic exposure could play an underestimated role in inducing hepatocellular carcinomas, alone or associated with already known etiologies.

2.2 Long-term carcinogenicity bioassays as a tool to identify potential hepatocarcinogens

Long-term carcinogenicity bioassays, mainly using rats and mice, have become a consolidated tool for identifying potentially carcinogenic chemical or physical agents (Huff, 2002; Maronpot et al., 2004; Soffritti et al., 2002). The two most extensive bioassay programs in the world are those of the American National Toxicology Program (NTP) and of the Italian Ramazzini Institute (Bucher, 2002; Huff, 2002; Soffritti et al., 2002). Although differences in the etiology exist between rodent and human HCC, there are also significant similarities in the genetic alterations leading to liver cancer in mice, rats and humans (Hoenerhoff et al., 2011; Feo et al., 2009). A recent study performed a global gene profiling of spontaneous (naturally occurring) HCC in B6C3F1 mice used in the NTP two-year bioassay (Hoenerhoff et al., 2011). The authors identified the dysregulation of genes similarly altered in human HCC, namely re-expression of fetal oncogenes, upregulation of protooncogenes, downregulation of tumor suppressor genes, and abnormal expression of cell cycle mediators, growth factors, apoptosis regulators, angiogenesis and extracellular matrix remodeling factors (Hoenerhoff et al., 2011). The use of a pathway-centered approach has lead to the identification of important targets that may be relevant to human HCC and, despite differences in etiology and pathogenesis from human HCC, the molecular readout has proved very similar, thus providing further support for applying this animal model to the study of HCC (Hoenerhoff et al., 2011). Importantly, the results from the two-year

carcinogenicity bioassays conducted by the NTP show that 188 out of 577 tested agents induced liver tumors in Fischer 344 rats or B6C3F1 mice (Table 2). Interestingly, 76% of these agents (145 out of 188) induced cancer in other sites than the liver (NTP, 2011). In particular, concerning the use of B6C3F1 mice, it has been suggested that this model might be a very sensitive system to detect chemicals that are likely to cause molecular events leading to cancer (Hoenerhoff et al., 2011). Results from the long term carcinogenicity bioassay program performed by the *Cesare Maltoni* Cancer Research Center of the Ramazzini Institute reveal that, among the studies already concluded and published, 52 agents showed clear evidence of carcinogenicity and 5 of them induced liver tumors in Sprague-Dawley rats or Swiss mice (Table 3). Interestingly, all these agents also induced cancer in other sites than the liver. These results confirm the finding that different strains of rats and mice present a different susceptibility to the development of HCC (Feo et al., 2009; Maronpot 2009). It should be noted that recent studies on genetic susceptibility and epigenetic regulation of the signaling pathways involved in hepatocarcinogenesis in rats have shown that most alterations responsible for a resistant or susceptible phenotype in rats also have a similar contribution to the prognosis of human HCC (Feo et al., 2009). In conclusion, the results of both long-term carcinogenicity studies and of genetic investigation of HCC from rodent models may provide important insights into the mechanisms of hepatocarcinogenesis, helping to identify some critical initiating events that lead to carcinogenesis, as well as progression markers and therapeutic targets (Feo et al., 2009; Hoenerhoff et al., 2011).

N.	Agents Type	Route of exposure	Evidence of carcinogenicity ^b				Carcinogenicity in other site	Human exposure
			Rats		Mice			
			M	F	M	F		
1	Acetonitrile	Inhalation	E	NE	NE	NE	No	chemical industry
2	Aldrin	Dosed-feed	E	E	CE	NE	Yes	pesticide
3	2-Aminoanthraquinone	Dosed-feed	CE	IS	CE	CE	Yes	dye industry
4	1-Amino-2,4-dibromoanthraquinone	Dosed-feed	CE	CE	CE	CE	Yes	dye industry
5	3-Amino-9-ethylcarbazole HCl	Dosed-feed	CE	CE	CE	CE	Yes	dye industry
6	11-Aminoundecanoic acid	Dosed-feed	CE	NE	E	NE	Yes	car industry; food packaging
7	Androstenedione	Gavage	E	E	CE	CE	Yes	hormonal treatment
8	Anthraquinone	Dosed-feed	SE	CE	CE	CE	Yes	dye industry
9	Aroclor 1254	Dosed-feed	E	E			Yes	insulator for current transformers
10	Benzofuran	Gavage	NE	SE	CE	CE	Yes	painting industry; food packaging

N.	Agents Type	Route of exposure	Evidence of carcinogenicity ^b				Carcino- genicity in other site	Human exposure
			Rats		Mice			
11	Benzophenone	Dosed-feed	SE	E	SE	SE	Yes	printing industry
12	Benzyl acetate	Gavage	E	NE	SE	SE	Yes	cosmetic industry
13	2-Chloro-1- methylethyl ether	Gavage			CE	CE	Yes	dye industry; pharmaceutics
14	Bromodi- chloromethane	Gavage	CE	CE	CE	CE	Yes	water contaminant
15	1,3 Butadiene	Inhalation			CE	CE	Yes	chemical industry; plastic and rubber
16	2-Butoxyethanol	Inhalation	NE	E	SE	SE	Yes	painting industry
17	Chloral hydrate	Gavage			SE		No	drugs treatment
18	Chloramben	Dosed-feed	NE	NE	E	CE	No	herbicide
19	Chlordane	Dosed-feed	NE	NE	CE	CE	No	pesticide
20	Chlordecone	Dosed-feed	CE	CE	CE	CE	No	insecticide, fungicide
21	Chlorendic acid	Dosed-feed	CE	CE	CE	NE	Yes	plastic industry
22	Chlorinated paraffins: C12 60% chlorine	Gavage	CE	CE	CE	CE	Yes	combustion products; oils
23	Chlorinated paraffins: C23 43% chlorine	Gavage	NE	E	CE	E	Yes	combustion products; oils
24	p-Chloroaniline hydrochloride	Gavage	CE	E	SE	NE	Yes	dye industry
25	Chlorobenzene	Gavage	E	NE	NE	NE	No	chemical industry; solvents
26	Chlorobenzilate	Dosed-feed	E	E	CE	CE	Yes	insecticide
27	Chlorodi- bromomethane	Gavage	NE	NE	E	SE	No	water contaminant
28	Chloroethane	Inhalation	E	E	IS	CE	Yes	chemical industry; anaesthetic
29	Chloroform	Gavage	CE	NE	CE	CE	Yes	chemical industry; anaesthetic
30	4-Chloro-m phenylenediamide	Dosed-feed	CE	NE	NE	CE	Yes	dye industry

N.	Agents Type	Route of exposure	Evidence of carcinogenicity ^b				Carcino- genicity in other site	Human exposure
			Rats		Mice			
31	4-Chloro-o phenylenediamide	Dosed-feed	CE	CE	CE	CE	Yes	dye industry
32	Chloroprene	Inhalation	CE	CE	CE	CE	Yes	chemical industry
33	5-Chloro-o-toluidine	Dosed-feed	NE	NE	CE	CE	Yes	dye industry
34	C.I. Acid red 114	Dosed- water	CE	CE			Yes	dye industry
35	C.I. Direct blue 15	Dosed- water	CE	CE			Yes	dye industry
36	C.I. Direct blue 218	Dosed-feed	SE	NE	CE	CE	Yes	dye industry
37	C.I. Disperse blue 1	Dosed-feed	CE	CE	E	NE	Yes	hair colouring
38	C.I. Disperse yellow 3	Dosed-feed	CE	NE	NE	CE	Yes	dye for clothes
39	C.I. Pigment red 3	Dosed-feed	SE	SE	SE	NE	Yes	dye industry
40	C.I. Basic red 9 monohydrochloride	Dosed-feed	CE	CE	CE	CE	Yes	dye industry; clothes, paper,leather
41	Cinnamyl anthranilate	Dosed-feed	CE	NE	CE	CE	Yes	synthetic flavour
42	C.I. solvent yellow 14	Dosed-feed	CE	CE	NE	NE	Yes	dye industry
43	Coconut oil acid diethanolamine	Topical application	NE	E	CE	CE	Yes	cosmetic industry
44	Coumarin	Gavage	SE	E	SE	CE	Yes	pharmaceutical use
45	p-Cresidine	Dosed-feed	CE	CE	CE	CE	Yes	dye industry
46	Cumene	Inhalation	CE	SE	CE	CE	Yes	chemical industry
47	Cupferron	Dosed-feed	CE	CE	CE	CE	Yes	chemical industry
48	Daminozide	Dosed-feed	NE	CE	E	NE	Yes	plants growing regulator
49	D & C red 9	Dosed-feed	CE	E	NE	NE	Yes	cosmetic industry
50	D & C yellow 11	Dosed-feed	SE	SE			Yes	cosmetic industry
51	Decabromodiphenyl oxide	Dosed-feed	Se	SE	E	NE	Yes	plastic industry
52	Decalin	Inhalation	CE	NE	NE	E	Yes	industrial solvent

N.	Agents Type	Route of exposure	Evidence of carcinogenicity ^b			Carcino- genicity in other site	Human exposure	
			Rats	Mice				
53	2,4 Diaminotoluene	Dosed-feed	CE	CE	NE	CE	Yes	dye industry
54	1,2 Dibromoethane	Gavage	CE	CE	CE	CE	Yes	gasoline additive; pesticide
55	2,3 Dibromo-1- propanol	Topical application	CE	CE	CE	CE	Yes	pesticide
56	1,4 Dichlorobenzene	Gavage	CE	NE	CE	CE	Yes	moth killer; deodorant
57	2,7 Dichlorodibenzo dioxin-	Dosed-feed	NE	NE	E	NE	Yes	pesticides contaminant
58	p,p'-Dichlorodiphenyl dichloroethylene	Dosed-feed	NE	NE	CE	CE	No	insecticide
59	2,6- Dichloro-p- phenylenediamine	Dosed-feed	NE	NE	CE	CE	No	chemical industry
60	1,2- Dichloropropane	Gavage	NE	E	SE	SE	Yes	chemical industry
61	1,3-Dichloropropene	Gavage	CE	SE	IS	CE	Yes	pesticide
62	Dicofol	Dosed-feed	NE	NE	CE	NE	No	miticide
63	Dieldrin	Dosed-feed	NE	NE	E	NE	No	insecticide
64	Diethanolamine	Topical application	NE	NE	CE	CE	Yes	chemical industry
65	Di(2-ethylhexyl) adipate	Dosed-feed	NE	NE	CE	CE	No	polivynil plastic
66	Di(2-ethylhexyl) phthalate	Dosed-feed	CE	CE	CE	CE	No	polivynil plastic
67	Di(p-ethylphenyl) dichloroethane	Dosed-feed	NE	NE	NE	E	No	insecticide
68	Diethyl phthalate	Topical application	NE	NE	E	E	No	plastic industry
69	3,4-Dihydrocoumarin	Gavage	SE	NE	NE	SE	Yes	pharmaceutical use
70	3,3'- Dimetoxybenzidine dihydrochloride	Dosed- water	CE	CE			Yes	dye industry
71	1,4 Dioxane	Dosed- water	CE	CE	CE	CE	Yes	dye, cosmetic and plastic industry

N.	Agents Type	Route of exposure	Evidence of carcinogenicity ^b			Carcino- genicity in other site		Human exposure
			Rats	Mice				
72	5,5-Diphenylhydantoin	Dosed-feed	E	NE	NE	CE	No	pharmaceutical use
73	2,5-Dithiobiurea	Dosed-feed	NE	NE	NE	E	No	film material
74	Elmiron	Gavage	NE	NE	SE	SE	Yes	pharmaceutical use
75	Ethylbenzene	Inhalation	CE	SE	SE	SE	Yes	chemical industry
76	Ethylene thiourea	Dosed-feed	CE	CE	CE	CE	Yes	rubber industry
77	Eugenol	Dosed-feed	NE	NE	E	E	No	flavouring compound
78	Fluometuron	Dosed-feed	NE	NE	E	NE	No	herbicide
79	Formamide	Gavage	NE	NE	CE	E	No	pharmaceutical use
80	Fumonisin B1	Dosed-feed	CE	NE	NE	CE	Yes	toxin
81	Furan	Gavage	CE	CE	CE	CE	Yes	polymers industry
82	Furfural	Gavage	SE	NE	CE	SE	Yes	food additive
83	Glycidol	Gavage	CE	CE	CE	CE	Yes	plastic industry
84	Goldenseal root powder	Dosed-feed	CE	CE	SE	NE	No	homeopathic use
85	HC blue 1	Dosed-feed	E	SE	CE	CE	Yes	hair colouring
86	HC red 3	Gavage	NE	NE	E	IS	No	hair colouring
87	Heptachlor	Dosed-feed	NE	E	CE	CE	Yes	insecticide
88	1,2,3,6,7,8- Hexachloro dibenzo-p-dioxin	Gavage	E	CE	CE	CE	No	pesticides contaminant
89	Hexachloroethane	Gavage	NE	NE	CE	CE	No	chemical industry; veterinary use
90	Hydrazobenzene	Dosed-feed	CE	CE	NE	CE	Yes	dye industry
91	Hydrochlorothiazide	Dosed-feed	NE	NE	E	NE	No	pharmaceutical use
92	Hydroquinone	Gavage	SE	SE	NE	SE	Yes	rubber and film industry
93	5-(Hydroxymethyl)-2- -furfural	Gavage	NE	NE	NE	SE	No	food additive

N.	Agents	Route of exposure	Evidence of carcinogenicity ^b		Carcinogenicity in other site	Human exposure		
	Type		Rats	Mice				
94	Indium phosphide	Inhalation	CE	CE	CE	Yes	electronic industry	
95	Isoeugenol	Gavage	E	NE	CE	E	Yes	flavouring compound
96	Isophorone	Gavage	SE	NE	E	NE	Yes	solvent
97	Lauric acid diethanolamine condensate	Topical application	NE	NE	NE	SE	No	pharmaceutical use
98	Leucomalachite green	Dosed-feed	E	E		SE	Yes	dye industry
99	Malachite green	Dosed-feed		E		NE	Yes	dye industry
100	2-Mercaptobenzo thiazole	Gavage	SE	SE	NE	E	Yes	rubber industry
101	Methyl carbamate	Gavage	CE	CE	NE	NE	No	textil industry
102	4,4'-Methylenebis (N,N-dimethyl) benzenamine	Dosed-feed	CE	CE	E	CE	Yes	dye industry
103	Methylene chloride	Inhalation	SE	CE	CE	CE	Yes	chemical industry
104	4,4' - Methylenedianiline dihydrochloride	Dosed-water	CE	CE	CE	CE	Yes	chemical industry
105	Methyleugenol	Gavage	CE	CE	CE	CE	Yes	flavouring industry
106	2-Methylimidazole	Dosed-feed	SE	CE	SE	SE	Yes	chemical industry
107	Methyl isobutyl ketone	Inhalation	SE	E	SE	SE	Yes	solvent
108	2-Methyl-1-nitro anthraquinone	Dosed-feed	CE	CE	CE	CE	Yes	dye industry
109	N-Methylolacrylamide	Gavage	NE	NE	CE	CE	Yes	adhesive industry
110	Methylphenidate hydrochloride	Dosed-feed	NE	NE	SE	SE	No	pharmaceutical use
111	alpha-Methylstyrene	Inhalation	SE	NE	E	CE	Yes	plastic industry
112	Michler's Ketone	Dosed-feed	CE	CE	CE	CE	Yes	dye industry

N.	Agents Type	Route of exposure	Evidence of carcinogenicity ^b				Carcino- genicity in other site	Human exposure
			Rats		Mice			
113	Mirex	Dosed-feed	CE	CE			Yes	insecticide
114	Monuron	Dosed-feed	CE	NE	NE	NE	Yes	herbicide
115	beta-Myrcene	Gavage	CE	E	CE	E	Yes	cosmetic industry
116	1,5-Naphthalene diamine	Dosed-feed	NE	CE	CE	CE	Yes	chemical industry
117	Nithiazide	Dosed-feed	NE	CE	CE	E	Yes	veterinary use
118	5-Nitroacenaphthene	Dosed-feed	CE	CE	NE	CE	Yes	dye industry
119	3-nitro-p- acetophenetide	Dosed-feed	NE	NE	CE	NE	No	pharmaceutical use
120	5-nitro-o-aniside	Dosed-feed	CE	CE	E	CE	Yes	dye industry
121	o-Nitroanisole	Dosed-feed	CE	CE	CE	SE	Yes	chemical industry
122	6-Nitrobenzimidazole	Dosed-feed	NE	NE	CE	CE	No	film industry
123	Nitrofen	Dosed-feed	IS/NE	CE/NE	CE	CE	Yes/N o	pesticide
124	Nitromethane	Inhalation	NE	CE	CE	CE	Yes	engine fuel
125	2-Nitro-p-phenyl enediamine	Dosed-feed	NE	NE	NE	CE	No	hair colouring
126	3 Nitropropionic acid	Gavage	E	NE	NE	NE	Yes	chemical industry
127	p-Nitrosodiphenyl- amine	Dosed-feed	CE	NE	CE	NE	No	rubber industry
128	o-Nitrotoluene	Dosed-feed	CE	CE	CE	CE	Yes	dye industry
129	5-Nitro-o-toluidine	Dosed-feed	NE	NE	CE	CE	Yes	dye industry
130	Oxazepam	Dosed-feed			CE	CE	Yes	pharmaceutical use
131	4,4' Oxydianiline	Dosed-feed	CE	CE	CE	CE	Yes	metallurgic industry
132	Oxymetholone	Gavage	E	CE			Yes	pharmaceutical use
133	Petachloroethane	Gavage	E	NE	CE	CE	Yes	chemical industry
134	Pentachlorophenol Dowicide EC-7	Dosed-feed			CE	CE	Yes	insecticide
135	Pentachlorophenol technical	Dosed-feed			CE	SE	Yes	insecticide

N.	Agents Type	Route of exposure	Evidence of carcinogenicity ^b			Carcino- genicity in other site	Human exposure	
			Rats	Mice				
136	Phenazopyridine hydrochloride	Dosed-feed	CE	CE	NE	CE	Yes	pharmaceutical use
137	Phenylbutazone	Gavage	E	SE	SE	NE	Yes	pharmaceutical use
138	Picloram	Dosed-feed	NE	E	NE	NE	No	herbicide
139	Piperonyl sulfoxide	Dosed-feed	NE	NE	CE	NE	No	insecticide
140	Polibrominated biphenyl mixture (Firemaster FF-1)	Dosed-feed and Gavage	CE	CE	CE	CE	No	engine fuel
141	Primidone	Dosed-feed	E	NE	CE	CE	Yes	pharmaceutical use
142	Probenecid	Gavage	NE	NE	NE	SE	No	pharmaceutical use
143	Proflavin hydrochloride	Dosed-feed	E	NE	E	E	Yes	pharmaceutical use
144	Propylene glycol mono-t-butyl ether	Inhalation	E	NE	CE	CE	Yes	solvent
145	Pulegone	Gavage	NE	SE	CE	CE	Yes	cosmetic use
146	Pyridine	Dosed- water	SE	E	CE	CE	Yes	chemical industry
147	Riddelliine	Gavage	CE	CE	CE	CE	Yes	food contaminant
148	Salicylazosulfa- pyridine	Gavage	SE	SE	CE	CE	Yes	pharmaceutical use
149	Selenium sulfide	Gavage	CE	CE	NE	CE	Yes	cosmetic use
150	Stoddard solvent (type 11C)	Inhalation	SE	NE	NE	E	Yes	solvent
151	Binary mixture PCB 126/153 (TEF evaluation)	Gavage		CE			Yes	chemical industry
152	PECDF (TEF evaluation)	Gavage		SE			Yes	chemical industry
153	PCB 118 (TEF evaluation)	Gavage		CE			Yes	chemical industry

N.	Agents Type	Route of exposure	Evidence of carcinogenicity ^b			Carcino- genicity in other site	Human exposure	
			Rats	Mice				
154	PCB mixture PCB 126/118 (TEF evaluation)	Gavage		CE		Yes	chemical industry	
155	TCDD (TEF evaluation)	Gavage		CE		Yes	chemical industry	
156	3,3',4,4' Tetrachloro azobenzene	Gavage	CE	CE	CE	CE	Yes	pesticide contaminant
157	2,3,7,8 Tetrachloro Dibenzo-p-dioxin	Gavage	CE	CE	CE	CE	Yes	herbicide contaminant
158	1,1,1,2 Tetrachloroethane	Gavage	E	NE	CE	CE	No	solvent
159	1,1,2,2 Tetrachloroethane	Gavage	E	NE	CE	CE	No	solvent
160	Tetrachloroethylene	Gavage	IS	IS	CE	CE	No	stain remover
		Inhalation	CE	SE	CE	CE	Yes	
161	Tetrachlorvinphos	Dosed-feed	NE	CE	CE	CE	Yes	insecticide
162	Tetrafluoroethylene	Inhalation	CE	CE	CE	CE	Yes	chemical industry
163	Tetrahydrofuran	Inhalation	SE	NE	NE	CE	Yes	chemical industry
164	Tetralin	Inhalation	SE	SE	NE	E	Yes	chemical industry
165	4,4'-Thiodianiline	Dosed-feed	CE	CE	CE	CE	Yes	dye industry
166	2,4-& 2,6- Toluene diisocyanate	Gavage	CE	CE	NE	CE	Yes	chemical industry
167	o-Toluidine hydrochloride	Dosed-feed	CE	CE	CE	CE	Yes	dye industry
168	Toxaphene	Dosed-feed	E	E	CE	CE	Yes	insecticide
169	Dioxin mixture (TEF evaluation)	Gavage		CE			Yes	chemical industry
170	PCB 153 (TEF evaluation)	Gavage		E			No	chemical industry
171	PCB 126 (TEF evaluation)	Gavage		CE			Yes	chemical industry

N.	Agents		Route of exposure		Evidence of carcinogenicity ^b		Carcinogenicity in other site	Human exposure
	Type		Rats	Mice				
172	Triamterene	Dosed-feed	E	NE	SE	SE	No	pharmaceutical use
173	1,1,2-Trichloroethane	Gavage	NE	NE	CE	CE	Yes	chemical industry
174	Trichloroethylene	Gavage	NE/IS	NE	CE	CE	No	solvent
175	2,4,6-Trichlorophenol	Dosed-feed	CE	NE	CE	CE	Yes	insecticide
176	1,2,3-Trichloropropane	Gavage	CE	CE	CE	CE	Yes	solvent
177	Triethanolamine	Topical application			E	SE	No	chemical industry
178	Trifluralin	Dosed-feed	NE	NE	NE	CE	Yes	pesticide
179	2,4,5-Trimethylaniline	Dosed-feed	CE	CE	E	CE	Yes	dye industry
180	tris(2,3-Dibromopropyl) phosphate	Dosed-feed	CE	CE	CE	CE	Yes	engine fuel
181	tris(2-Ethylhexyl) phosphate	Gavage	E	NE	NE	SE	Yes	engine fuel
182	Turmeric, oleoresin (curcumin)	Dosed-feed	NE	E	E	E	Yes	pharmaceutical use
183	Urethane	Dosed-water			CE	CE	Yes	chemical industry
184	Bromochloroacetic acid	Dosed-water	CE	CE	CE	CE	Yes	water disinfection byproducts
185	Dibromoacetic acid	Dosed-water	SE	SE	CE	CE	Yes	water disinfection byproducts
186	Dibromoacetonitrile	Dosed-water	CE	SE	CE	CE	Yes	water disinfection byproducts
187	2,6 -Xylidine	Dosed-feed	CE	CE			Yes	dye industry
188	Zearalenone	Dosed-feed	NE	NE	CE	CE	Yes	micotoxin

^a Data from: <http://ntp.niehs.nih.gov/>

^b CE= clear evidence; SE= some evidence; E= equivocal evidence; NE= no evidence; IS= inadequate experiment

Table 2. Chemicals industrial agents associated with tumor induction in liver based on long term carcinogenicity bioassays performed in Fischer 344 rats and B6C3F1 mice males and females by the *National Toxicology Program (NTP) USA*^a

Agents		Route of exposure	Evidence of carcinogenicity				References
N.	Type		Rats		Mice		
			M	F	M	F	
1	Vynil Chloride	Inhalation	+	+	-	-	Maltoni et al. 1984
2	Trichloroethylene	Ingestion	-	-	NS	NS	Maltoni et al. 1986
		Inhalation	-	-	+	+	
3	Benzene	Inhalation	E	(+)	NS	NS	Maltoni et al. 1989
4	Tamoxifen	Ingestion	E	E	-	-	Minardi et al. 1994
		Ingestion	+	+	NS	NS	
5	Aspartame	Ingestion	-	-	+	-	Soffritti et al. 2010

Table 3. Agents inducing Hepatocellular Carcinoma identified in the framework of the long term carcinogenicity bioassays program performed by the Cesare Maltoni Cancer Research Center of the Ramazzini Institute

3. New risk factors of hepatocellular carcinoma

The epidemic of obesity has been correlated to an increased risk of various types of cancer (Kaidar-Person et al., 2011). In the last few decades, experimental and epidemiological studies have shown a strong correlation between obesity or its related co-morbidities and HCC incidence or mortality. In this section we will review the data linking metabolic syndrome and HCC risk, as well as some recent results published in the literature suggesting novel paths to be explored in liver cancer etiology.

3.1 Metabolic syndrome and hepatocellular carcinoma: Experimental data

Although no long-term carcinogenic bioassays have been specifically performed to investigate the carcinogenic potential of a high-fat diet, some recent studies provide interesting clues, in particular concerning liver carcinogenesis. A 20-month study on C57BL/6J male mice fed with a high fat western style diet showed that treated animals developed NASH (non-alcoholic steatohepatitis) at 14 months while at 20 months primary liver dysplastic nodules were found (VanSaun et al., 2009). Similar results were found by another group which showed that a high fat diet induced NASH and HCC in C57BL/6J mice but not in A/J mice (Hill-Baskin et al., 2009). Interestingly, the switch from the high-fat to low-fat diet after 100 days of administration until the end of the experiment (400 days of life) in C57BL/6J males prevented the development of obesity by the end of study and reversed the progression of the disease (Hill-Baskin et al., 2009). Again, a study using a diethylnitrosamine-initiated hepatocarcinogenesis model on Sprague-Dawley rats showed

that animals fed with high-fat diets had an increased incidence of preneoplastic liver foci after 6 weeks compared to control animals (Wang et al., 2009).

These data may represent an early warning as to a possible liver cancer epidemic in coming decades. While obvious measures to contrast obesity have been undertaken in the last few years in many countries (WHO, 2006; Ministère de la Santé et de la Solidarité, 2006; White House Task Force on Childhood Obesity, 2010) and need to be strengthened, the impact of chemical indoor and outdoor pollution on obesity and HCC risk has been poorly studied and may play an underestimated and synergistic role. In this chapter, we will focus on chemical agents recognized as liver carcinogens and on novel mechanisms of hepatocarcinogenesis, in particular concerning metabolic and endocrine disruption.

3.2 Metabolic syndrome and hepatocellular carcinoma: The dimension

Metabolic syndrome and its characteristic manifestations, such as obesity (or central obesity), insulin resistance/type 2 diabetes, dyslipidemia or hypertension, are the most well-studied emerging risk factors of HCC (Bugianesi 2007; Starley et al., 2010; Welzel et al., 2011). In a recent US study, increase in Body Mass Index (BMI) results in a statistically significant increasing trend for HCC mortality both for women (highest calculated RR 1.68, CI 0.93-3.05, *p* for trend 0.04) and for men (highest calculated RR 4.52, CI 2.94-6.94, *p* for trend <0.001) (Calle et al., 2003). Other studies have shown that obesity is a risk factor for developing HCC, increasing the risk from 1.5 to 4 times (Møller et al., 1994; Wolk et al., 2001; Oh et al., 2005). Diabetes has been also shown to be an independent risk factor for HCC, with an increase in the risk ranging from 1.8 to 4 times in Swedish, Danish and Greek cohorts (Adami et al., 1996; Wideroff et al., 1997; Lagiou et al., 2000). Recently, metabolic syndrome has been reported to be more common in persons who developed HCC (37.1%) than in persons who did not (17.1%, *p*<0.0001) and it was significantly associated with increased risk of HCC after multiple logistic regression analyses (OR=2.13; 95%CI=1.96-2.31, *p*<0.0001) (Welzel et al., 2011). It is worth noting that a concurrent increase in the incidence and prevalence of NAFLD has been observed with the obesity epidemic (McCullough, 2004; Williams et al., 2011). NAFLD is a spectrum of liver disease ranging from simple steatosis to non-alcoholic steatohepatitis and cirrhosis (Bugianesi, 2007). The prevalence of NAFLD in the general population is estimated to range from 17 to 33% (McCullough, 2004). However, a recent prospective study on 328 persons (average age 54 years) reported that the prevalence of NAFLD was 46%, the global prevalence of NASH was 12.2% (Williams et al., 2011) and 2.7% of patients presented advanced NASH (fibrosis graded more than 2). The results of this study confirmed previous reports, as NAFLD and NASH were more frequently diagnosed in overweight/obese males, with a history of hypertension and diabetes (Williams et al., 2011); it suggests that NAFLD and its complications may be more widespread than previously reported. Importantly, some studies and several case reports have shown a direct relationship between NAFLD and HCC (Page and Harrison, 2009). In a Danish cohort study, the risk for primary liver cancer among NAFLD patients was elevated with a standardized incidence ratio of 4.4 (CI 1.2-11.4) (Sørensen et al., 2003). In a US single center case series, NAFLD accounted for 13% of the cases of HCC (Marrero et al., 2002). Moreover, at least 67 cases of HCC arising in a context of NASH have been reported and in 33% of cases HCC seems to have arisen in the absence of cirrhosis (Page and Harrison, 2009). In this context, it is of major concern that NAFLD is becoming the most common cause of liver disease in children and adolescents (Sundaram et al., 2009). A

large pediatric autopsy study found a NAFLD global prevalence of 9.6%, with 38% prevalence in obese children (Schwimmer et al., 2006). The prevalence of fatty liver increased with age from 0.7% in 2-4 year olds to 17.3% in 15-19 year olds (Schwimmer et al., 2006). Of even greater concern is the higher initial incidence of fibrosis or cirrhosis reported in pediatric cases than in adults: a study of obese children with liver steatosis and elevated aminotransferases found NASH in 88% and fibrosis in 71% of patients (Nadeau et al., 2003; Rashid & Roberts 2000).

3.3 Endocrine disruption: Linking metabolic disorder to cancer etiology in the liver?

Endocrine disruption is a term coined in the 1990s which refers to the hormone-like effects of various synthetic chemicals present in the environment (Wingspread consensus statement, 1992; Kavlock et al., 1996). Since then, most laboratory and epidemiological research conducted on endocrine disruptors (ED) has focused on reproductive system pathologies, such as decreased sperm quality, malformations of male genital tract (cryptorchidism or hypospadias), prostate and breast cancers, and on thyroid dysfunction (Soto & Sonnenschein, 2010). However, in the last decade, several ED, such as derivatives of alkylphenols (i.e. 4-nonylphenol), phthalates, polybrominated diphenyl ethers (PBDEs) or organotin compounds (i.e. tributyltin), have been shown to alter adipose tissue development and to promote fat accumulation in both adipose and liver tissues (Masumo et al., 2002; Grün et al., 2006; Grün & Blumberg 2009). Thus, some ED may also be addressed as “obesogens” and more generally considered as metabolic disruptors, suggesting that the various pathophysiological effects of ED may be due to pleiotropic effects on multiple metabolic pathways and that analyses of toxicological effects should not be limited to the endocrine system, but extended to all organs and tissues, with particular attention to adipose tissue and the liver (Grün & Blumberg, 2009; Casals-Casas & Desvergne, 2011).

Most data on the potential carcinogenic effects of ED derive from studies based on evidence of estrogen carcinogenicity. Estrogen carcinogenicity was experimentally demonstrated on the mouse mammary gland as early as the 1930s by Lacassagne (Lacassagne, 1936). In humans the most notorious case of estrogen carcinogenicity is diethylstilbestrol, a synthetic estrogen prescribed to pregnant women to prevent miscarriage in the 1940s-early 1970s. Diethylstilbestrol has been associated with an increased risk of developing clear-cell carcinoma of the vagina and breast cancer in daughters of exposed women (Herbst et al., 1971; Goodman et al., 2011; Palmer et al., 2006). Other ED, such as bisphenol A, tamoxifen or 2,3,7,8-tetrachlorodibenzodioxin, have been shown to alter mammary development and to induce precancerous and cancerous lesions in rodents (Vandenberg et al., 2008; Fenton et al., 2002; Soto & Sonnenschein, 2010). However, information on the potential carcinogenicity of ED for the liver is still elusive while the marshalling of epidemiological and experimental evidence for a potential carcinogenic effect by ED in the liver remains to be properly addressed. Interaction with nuclear receptors (NR) is one potential mechanism that may be involved in any pathological effects of ED on the liver. Indeed, many NR are signally expressed in the liver and have been reported to play an important role in liver diseases, such as steatosis and HCC (Wagner et al., 2011). Since many ED are small lipophilic compounds, their effects are thought to be mediated mostly by direct interaction with NR, modulating downstream gene expression (Casals-Casas & Desvergne, 2011). However, activation of various receptors, such as G protein-coupled membrane receptors, is also possible, as reported at least in the case of the well-established ED Bisphenol A (Chevalier et

al., 2011). The NR superfamily encompasses 48 members and ED are able to alter the signaling pathways mediated by many of them, including estrogen and androgen receptors, thyroid hormone receptor (TR), glucocorticoid receptor (GR), mineralcorticoid receptor (MR), retinoid X receptor (RXR), peroxisome proliferator-activated receptors (PPARs), liver X receptors (LXRs), farnesoid X receptor (FXR), constitutive androstane receptor (CAR), and pregnane X receptor (PXR) (Arrese & Karpen, 2010; Casals-Casas & Desvergne, 2011). The ability of ED to interact with various NR at nanomolar concentrations explains the diversity of induced metabolic perturbation and the increased effects observed particularly when fetal and neonatal exposures occur (Heindel, 2003; Soto & Sonnenschein, 2010; Casals-Casas & Desvergne, 2011). It is worth noting that in addition to sex steroid receptors, this receptor superfamily includes transcription factors playing a central role in integrating metabolic and developmental signaling pathways (Wagner et al., 2011). In particular, data from knockout mice showed that functional PPAR-alpha or FXR are essential in the regulation of hepatic lipid metabolism, as deletion or deficiency of them induces hepatic steatosis in mice. Moreover, PXR and CAR (NRs for which many xenobiotics are ligands) promote hepatic lipid storage by decreasing fatty acid beta-oxidation (Wagner et al., 2011). Importantly, FXR, CAR and PXR are also variously involved in HCC formation. FXR knockout mice suffer from chronic bile acid-induced chronic inflammation and are prone to develop HCC, while CAR and PXR are important in the liver proliferative response, although their exact role in liver tumor promotion is not clear (Yang et al., 2007; Wagner et al., 2011). These data suggest that important interactions exist between NR signaling, lipid metabolism and liver carcinogenesis. This hypothesis is supported by recent findings showing a direct implication of impaired lipid metabolism in the development of hepatocellular carcinoma in animal models. A recent study showed that mice double mutants (due to a growth hormone-activated signal transducer and activator of transcription (STAT5) and GR) developed liver steatosis that progressed to hepatocellular carcinomas (Mueller et al., 2011). Altered STAT5/GR signaling was associated with insulin resistance, high reactive oxygen species levels and increased liver and DNA injuries; that it function correctly was essential for maintenance of lipid homeostasis (Mueller et al., 2011). A second study showed that transgenic CAR $-/-$ mice fed with a NASH-inducing diet are protected from diethylnitrosamine-induced hepatocarcinogenesis (Takizawa et al., 2011). These results suggest that the nuclear receptor CAR might play an important role in promoting hepatocellular carcinomas against a background of NASH. Interestingly, human HCC have been associated with altered lipid metabolism, in particular choline and phospholipid metabolism with an increased synthesis in lysophosphatidic acid, which may provide a mitogenic and proliferative microenvironment through activation of G-protein-coupled receptors (Skill et al., 2011). Moreover, stimulation of lipid biosynthesis and perpetuating chronic hepatic metabolic disease by activation of the transactivator of stress proteins HSF1 promotes HCC development in mice (Jin et al., 2011). Finally, an interesting paper based on a microarray analysis of a hepatitis-induced HCC murine model showed that pro-inflammatory cytokines may promote HCC predominantly in males causing the loss of a gender-identifying hepatic molecular signature (Rogers et al., 2007), suggesting that HCC may be associated with liver-gender disruption in male mice and supporting the idea that endocrine disruption may have broader effects than those reported in reproductive and sexual organs.

Although the scientific data on ED and their effects on liver disease are mostly anecdotal, a few toxicological studies have shown that perinatal exposure to Bisphenol A (BPA) or organotin compounds alters not only adipogenesis in rodents, but also increases both the expression of

lipogenic genes and lipid accumulation in the liver of animals exposed (Somm et al., 2009; Grün et al., 2006). Ecotoxicological studies have shown that ED may bioaccumulate and induce altered gene expression in the liver of various animal species (Ter Veld et al., 2008). In particular, BPA, PCBs and PBDE were shown to bioaccumulate in fish livers caught in Italian seas or US lakes (Mita et al., 2011; Pérez-Fuenteaja et al., 2010) and nonylphenol (NP) was reported to induce the expression of female specific proteins in male lizard livers (Verderame et al., 2011). Although the relevance of NR stimulation to human liver cancer is still unclear, this is a well-known non-genotoxic mechanism of rodent liver cancer, in particular concerning PPAR-alpha activation (Ren et al., 2010). Interestingly, it has been shown that rodent carcinogens show higher *in vitro* potency for human NR than do non-carcinogens (Shah et al., 2011). NP was shown to activate human CAR *in vitro* and *in vivo* using transgenic mouse models (Hernandez et al., 2007). Moreover, CAR is involved in hepatic injury and in the development of HCC in a dietary model of NASH, probably by its cell proliferation promoting activity (Takizawa et al., 2011).

Available data on ED-induced hepatic alterations and on their potential action on systemic and hepatic lipid metabolism suggest that ED may have a role in the establishment and progression of some liver diseases, in particular NAFLD and HCC. However, systematic research aiming to investigate interactions between endocrine/metabolic disruption in liver disease/carcinogenesis is still lacking. We will report on some recent advances in the identification of common new subcellular targets in metabolic disease and liver carcinogenesis, with a particular focus on mitochondria and endoplasmic reticulum alterations and their cross-talks with NR signaling perturbation.

4. Molecular aspects of hepatocellular carcinoma

HCC is a cancer caused by a variety of etiologies, including HBV and HCV infection, alcohol over-consumption, aflatoxin B1 exposure and chemical agents. Thus, it is not surprising that a variety of HCC-associated molecular alterations have been detected and no universal molecular signature is definitively associated with all hepatic tumors, either in humans or in experimental animals (Degli Esposti et al., 2009; Pei et al., 2009). In particular, in humans HCC alterations are classically reported in four genetic pathways, namely p53, retinoblastoma (Rb), TGF-beta pathway and Wnt-beta-catenin pathways (Laurent-Puig, 2001; Saffroy et al., 2007). However, the carcinogenic process in the liver looks to be more complex, as a recent review on omics-based studies seems to suggest (Pei et al., 2009). In particular, comparative genomic hybridizations, high-throughput methods used to identify deletion or amplification in genomic DNA, have shown chromosomal aberrations in fourteen human chromosomes, not uniquely associated with viral infections (Pei et al., 2009). Various studies have also reported epigenetic alterations in HCC, in particular hypermethylation and subsequent silencing of some tumor suppressor genes (Pei et al., 2009). Microarray studies have been performed on both mRNA and microRNA (miRNA), but no consistent expression signatures seem to arise, due to alleged differences in technology and experimental design (Pei et al., 2009). All together, these data reinforce the view that the development of HCC, and probably all cancer types, involves multiple factors and interactions at a molecular level. While meta-analysis and integration of -omics data could prove a helpful approach for biomarker identification in cancer (Zender et al., 2006; Ludwig & Weinstein, 2005), it seems urgent we adopt a comprehensive framework

including pathophysiological and developmental aspects of the disease with consistent molecular, biochemical and biophysical interactions (Soto & Sonnenschein, 2004).

As we reported in the previous section, the case of endocrine and metabolic perturbation in liver disease and carcinogenesis offers a useful starting point for discussing newly identified or potential carcinogenic pathways and future directions to clarify the complex picture of hepatocellular carcinoma and, maybe more generally, of cancer.

4.1 New aspects of hepatocarcinogenesis: Metabolic disruption and altered cellular homeostasis

Alterations in the metabolism of cancerous tissues have been found since 1930s. Warburg (1930) described that, even in the presence of oxygen, cancer tissues had acquired an irreversible glycolytic metabolism. Increased glycolysis proves to allow the utilization of glycolytic intermediates into the various biosynthetic pathways, including nucleoside and amino acid synthesis, required by highly replicating cells (Potter, 1958, Vander Heiden et al., 2009). This feature is also described in rapidly dividing embryonic tissues, even though they are able to switch to oxidative metabolism as proliferation ceases and cells differentiate, suggesting that cancer development may be understood as altered embryonic development (Barger & Plas, 2010, Cooper, 2009, Soto & Sonnenschein, 2004). As a result of Warburg's observations, defects in mitochondrial function have been suspected as contributing to cancer development and progression (Chatterjee et al., 2011). Recently, not only mitochondria but also endoplasmic reticulum have been found to be implicated in controlling the lipid metabolism, in particular in the liver. Thus, since reprogramming of the energy metabolism has been rediscovered as a hallmark of cancer (Hanahan & Weinberg, 2011) while alteration of the lipid metabolism seems to be associated with HCC development, the role of mitochondria and endoplasmic reticulum in metabolic disruption during liver carcinogenesis forms a testable hypothesis in the context of liver carcinogenesis.

Mitochondria are key organelles both for energy (ATP) production (by oxidative phosphorylation of components of the tricarboxylic acid cycle and lipid beta-oxidation) and for integration of pro-survival/pro-death signaling in cells of every tissue and organ. These functions are essential in determining cellular and tissue homeostasis. In the last decade, a lot of research has focused on the role of mitochondria in liver diseases and mitochondrial dysfunctions have been described both in NAFLD and in HCC (Begriche et al., 2006; Chang et al., 2005; Sato, 2007; Rector et al., 2010). A recent study compared liver histology and function in the obese rodent model Otsuka Long-Evans Tokushima Fatty (OLETF) rat with its lean homolog LETO rat (Rector et al., 2010). The ultrastructure of hepatic mitochondria proved to be impaired and the total mitochondrial content decreased in OLETF rats. Moreover, mitochondrial and total fatty acid oxidation were already reduced as early as the fifth week of age in obese animals, before hepatic steatosis and insulin resistance were observed, suggesting that mitochondrial dysfunction may be a very early event in the natural history of NAFLD (Rector et al., 2010). Various mitochondrial alterations have been reported in patients with NASH. Megamitochondria with ultrastructure abnormalities have been found in NASH patients (Caldwell et al., 1999a; Sanyal et al., 2001). Severe depletion of mitochondrial DNA (mtDNA) has been also reported in patients with NASH or hepatic fibrosis (Caldwell et al., 1999b; Ducluzeau et al., 1999). mtDNA depletion may contribute to impairment of the respiratory chain, a common feature in drug-induced and primary NASH (Begriche et al., 2006). It should be noted that in primary NASH impairment of the

respiratory chain is concomitant with an increase in beta-oxidation flux (due to insulin resistance), leading to production of a high level of reactive oxygen species (ROS) (Begriche et al., 2006). Increased ROS generation by mitochondria has also been observed in genetically obese ob/ob mice (Yang et al., 2000) and in rat fed on a choline-deficient diet, a model of steatosis and NASH (Hensley et al., 2000). An increased production of ROS in fatty livers induces lipid peroxidation and subsequently reactive aldehyde formation (Begriche et al., 2006). ROS and aldehydes may further damage mitochondria, generating a vicious circle, and increase the expression of pro-inflammatory cytokines, such as TGF-beta, TNF-alpha, IL-8 or Fas ligand, worsening liver injury (Pessayre & Fromenty, 2005). Mitochondria are also involved in hepatocarcinogenesis. A decrease in mtDNA-dependent cytochromes with disturbed electron transfer has been reported in liver carcinomas, with a subsequent increase in ROS production that can induce nuclear gene mutation in carcinogenesis (Sato, 2007). Moreover, a higher frequency of somatic mutation in regulatory and coding regions of mtDNA has recently been reported in HCC compared to adjacent non-cancerous tissue (Yin et al., 2010). Interestingly, an experimental study showed that long-term administration of L-carnitine, a key molecule in fatty acid transport to mitochondria, decreases the occurrence of hepatic preneoplastic lesions in Long-Evans Cinnamon rats (Chang et al., 2005). These data reinforce the idea that mitochondrial dysfunction, and in particular its role in lipid catabolism, is involved in hepatocarcinogenesis. It is interesting to note that some chemicals with endocrine disruptive properties have been shown to target mitochondrion functionality in various tissues and organs, including the liver (Kovacic, 2010). In particular, five-month exposure to 10-50 ppm diethyl phthalate induced liver impairment, triglyceride accumulation and mitochondrial proliferation in Wistar rats (Pereira et al., 2006). Since phthalates are plasticizers present in plastics used for medical reasons, such as storage bags for blood conservation or instruments for dialysis, it is interesting to note that liver biopsies on dialyzed patients show peroxisome proliferation (Ganning et al., 1984) and that phthalate leakage from blood bags has been proposed as potentially pro-inflammatory (Rael et al., 2009). Furthermore, male mice treated perinatally with 160 or 480 mg/kg of BPA for 14 days showed an increase in cell death mediated by the mitochondrial apoptotic pathway in the testes (Wang et al., 2010). While the hepatic effects on the liver were not evaluated in this study, these results suggest that BPA may directly or indirectly target mitochondria. Other evidence of ED toxicity in mitochondria is provided by results showing that tamoxifen decreased ATP production in a model of isolated perfused rat liver (Marek et al., 2011) and that it impaired mitochondrial respiration, increased cytochrome c release, mitochondrial lipid peroxidation and mitochondrial protein nitration by stimulating mitochondrial nitric oxide synthase (Nazarewicz et al., 2007). Importantly, although epidemiological studies available in humans did not identify any increased risk of liver cancer in women who were administered tamoxifen for breast cancer, several experiments have shown that tamoxifen induces hepatocarcinomas in rats when administered at high doses (Maltoni et al., 1997; IARC, 2011). Another reported ED and human carcinogen, 2,3,7,8 TCDD, has been reported to induce cytotoxicity and mitochondrial dysfunction in isolated rat hepatocytes (Aly & Domènech, 2009) and to mediate tumor progression by activating signaling pathways similar to mtDNA depletion (Biswas et al., 2008). In this context, it would be helpful to consider the interactions between mitochondria, lipid metabolism and nuclear receptors in order to improve our understanding of liver disease and carcinogenesis. In point of fact, several members of the nuclear receptor superfamily are lipid-sensing factors that affect many aspects of lipid metabolism (Alaynick, 2008). PPARs, LXRs, interacting with their transcriptional coactivator PPARgamma Coactivator 1 alpha (PGC-1alpha) have been

shown to regulate insulin sensitivity and lipid metabolism (Alaynick, 2008). Interestingly, PGC-1 α is a known regulator of mitochondrial biogenesis and also able to modulate hepatic steatosis (Puigserver et al., 1998; Sonoda et al., 2007; Wu et al., 1999). Moreover, the mitochondrial protein ANT, a translocase that provides mitochondria with ADP allowing ATP synthesis, has recently been shown to be essential for the functioning of PGC-1 α (Kim et al., 2010) while another NR, the estrogen-related receptor α (ERR- α), important for adaptive energy metabolism (Villena & Kralli, 2008), has been shown to be an effector of PGC-1 α , regulating the expression of genes involved in oxidative phosphorylation and mitochondrial biogenesis (Schreiber et al., 2004). Thus, PGC-1 α may be a key molecule linking NR signaling to mitochondrial function and activity. Finally, it is important to note that ED and other NR ligands may also act independently on the receptor action and mitochondria can be a direct target of this mechanism, as proposed for PPAR agonists (Scatena et al., 2004). Hence, it has been suggested that characterization of reciprocal influences between mitochondria and PPAR physiology would be fundamental for a better understanding of cancer biology (Scatena et al., 2008).

Endoplasmic reticulum is an organelle responsible for protein synthesis, folding, maturation, quality control and trafficking, as well as for Ca²⁺ homeostasis. Every condition that stresses its folding ability, such as an excess of protein synthesis or alteration of energy availability, causes a physiological response called Unfolded Protein Response (UPR). UPR activation aims to increase the folding capacity of endoplasmic reticulum by inducing transcription of chaperons and by globally decreasing protein synthesis (Schroeder & Kaufman, 2005). In recent years, it has been shown that endoplasmic reticulum plays a central role in the multi-organ coordination of systemic metabolism through the integration of synthetic and catabolic pathways (Kammoun et al., 2009b, Hotamisligil, 2010). Obesity and diabetes have been shown to induce ER stress in both adipose tissue and the liver (Ozcan et al., 2004, Kammoun et al., 2009a). Puri et al., examined the role of endoplasmic reticulum stress in human NAFLD, showing UPR activation in liver biopsies from patients with NAFLD and NASH compared to subjects with the metabolic syndrome and normal liver histology (Puri et al., 2008). Moreover, free fatty acids (FFA) may be important mediators of cell dysfunction (lipotoxicity) not only through death receptors or the mitochondrial-lysosomal pathway, but also via endoplasmic reticulum stress (Alkhoury et al., 2009). Endoplasmic reticulum stress and activation of the UPR are also present in solid cancers, often characterized by hypoxia, nutrient starvation, oxidative stress and other metabolic deregulation, factors that cause endoplasmic reticulum impairment (Li et al., 2011). Depending on the duration and degree of ER stress, the UPR can provide either survival signals by activating adaptive and antiapoptotic pathways, or death signals by inducing cell death programs (Schroeder & Kaufman, 2005; Li et al., 2011). In hepatocellular carcinoma, higher accumulation of the Bip/GRP78 and nuclear localization of ATF6, characteristic of UPR activation, were found in moderately to poorly differentiated human HCC tissue samples (Shuda et al., 2003). Although direct demonstrations at a molecular level are still lacking, the cross-links between endoplasmic reticulum stress, NAFLD and HCC seem to be numerous and future research should address this potential new carcinogenic pathway. However, endoplasmic reticulum seems also to be a target of endocrine disruption. Recently, BPA has been shown, in a murine liver cell line, to increase the gene expression of various actors involved in endoplasmic reticulum stress, such as C/EBP homologous protein, caspase 12 and GRP78 (Asahi et al., 2010). This ties up interestingly with a microarray study in which activation of genetic networks involved in

endoplasmic reticulum stress was also detected in mouse testicular Sertoli cells treated with BPA at a concentration of 200 microM (Tabuchi et al., 2006). Furthermore, a number of ED, such as nonylphenol, octylphenol, bisphenol A, and butylated hydroxytoluene, have been shown to inhibit endoplasmic reticulum Ca^{2+} ATPase pumps in a low micromolar concentration (Hughes et al., 2000), suggesting that alterations in endoplasmic reticulum homeostasis may be a common action mechanism by BPA in various different organs. In this connection, some recent papers have highlighted a possible direct interaction between nuclear receptor signaling, lipid metabolism and endoplasmic reticulum stress. A protein deacetylase (SIRT1) has been shown to positively regulate the nuclear receptor PPAR alpha (Purushotham et al., 2009). In particular, hepatic-specific deletion of SIRT1 impairs PPAR alpha signaling and SIRT1 knockout mice develop hepatic steatosis, liver inflammation and endoplasmic reticulum stress (Purushotham et al., 2009). Moreover, the endoplasmic reticulum stress-induced transcription factor ATF6 has been shown to suppress insulin gene expression through the up-regulation of a transcriptional partner of nuclear receptors, SHP, in pancreatic beta-cells and in pancreatic islets of OLETF rats (Seo et al., 2008), suggesting that endoplasmic reticulum stress signaling may also act via NR signaling. As for mitochondria, while no extensive data are available, interactions between endoplasmic reticulum homeostasis and NR physiology have been shown and may play an important yet still under-explored role in the initiation and progression of liver steatosis and HCC. Perturbation of this network by ED is possible, although more research is needed to address the effects of environmental concentrations and to identify the biochemical pathways affected and any long-term pathophysiological consequences, in particular in liver carcinogenesis.

5. Conclusions

In this chapter, we have reviewed some recent acquisitions in HCC epidemiology, in particular regarding the association between obesity and metabolic syndrome and HCC incidence and mortality. We have also reviewed the evidence for liver susceptibility to chemical-induced carcinogenesis, in both rodents and humans, and have shown that long-term carcinogenicity bioassays are a useful tool for identifying potential hepatocarcinogens. In the last part of the chapter, we suggest that endocrine and metabolic disruption, a mechanism involved in the toxic effects of various chemicals, might be a plausible and testable hypothesis in the pathophysiology of NAFLD and its progression toward HCC, in particular concerning the alteration of mitochondria and endoplasmic reticulum in the liver and other tissues. We suggest that long-term carcinogenicity bioassays are a valuable approach to integrating pathological end-points, such as tumor induction, and the analysis of early chronic alteration in tissue and cellular homeostasis, such as mitochondrial and endoplasmic reticulum dysfunctions. This approach could provide important insights into chemical-induced carcinogenic mechanisms, in particular for non-genotoxic carcinogens such as most endocrine disruptors. Moreover, the comparison of experimental results with human biopsies of neoplastic and pre-neoplastic lesions, such as NASH, in well-characterized patients may help to develop specific early markers towards identifying the population at higher risk of HCC.

6. Acknowledgement

The authors thank Ralph Nisbet for his assistance in preparing the manuscript.

7. References

- Adami, H.O., Chow W.H., Nyrén, O., et al. (1996). Excess risk of primary liver cancer in patients with diabetes mellitus. *J Natl Cancer Inst*, 88, 1472-1477.
- Adelstein, A. & White G. (1976). Alcoholism and mortality. *Popul Trends*, 6, 7-13.
- Alaynick, W.A. (2008). Nuclear receptors, mitochondria and lipid metabolism. *Mitochondrion*, 8(4), 329-337.
- Alkhoury, N., Dixon, L.J., Feldstein, A.E. (2009). Lipotoxicity in nonalcoholic fatty liver disease: not all lipids are created equal. *Expert Rev Gastroenterol Hepatol*, 3, 445-451.
- Aly, H.A. & Domènech, O. (2009). Cytotoxicity and mitochondrial dysfunction of 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) in isolated rat hepatocytes. *Toxicol Lett*, 191(1), 79-87.
- Arrese, M., & Karpen, S.J. (2010). Nuclear receptors, inflammation, and liver disease: insights for cholestatic and fatty liver diseases. *Clin Pharmacol Ther*, 87, 473-478.
- Asahi, J., Kamo, H., Baba, R., et al. (2010). Bisphenol A induces endoplasmic reticulum stress-associated apoptosis in mouse non-parenchymal hepatocytes. *Life Sci*, 87(13-14), 431-438.
- Barbin, A., Froment, O., Boivin, S. et al. (1997). p53 gene mutation pattern in rat liver tumors induced by vinyl chloride. *Cancer Res*, 57, 1695-1698.
- Barger, J.F., & Plas, D.R. (2010). Balancing biosynthesis and bioenergetics: metabolic programs in oncogenesis. *Endocrine-Related Cancer*, 17, R287-R304.
- Begrache, K., Igoudjil, A., Pessayre, D., et al. (2006). Mitochondrial dysfunction in NASH: causes, consequences and possible means to prevent it. *Mitochondrion*, 6(1), 1-28.
- Biswas, G., Srinivasan, S., Anandatheerthavarada, H.K. et al. (2008). Dioxin-mediated tumor progression through activation of mitochondria-to-nucleus stress signaling. *Proc Natl Acad Sci U S A*, 105(1), 186-191.
- Blonski, W., Kotlyar, D.S., Forde, K.A. (2010). Non-viral causes of hepatocellular carcinoma. *World J Gastroenterol*, 16(29), 3603-3615.
- Borentain P, Gérolami V, Ananian P, et al. (2007). DNA-repair and carcinogen-metabolising enzymes genetic polymorphisms as an independent risk factor for hepatocellular carcinoma in Caucasian liver-transplanted patients. *Eur J Cancer*, 43(17), 2479-2486.
- Bosetti, C., Levi, F., Boffetta, P., et al. (2008). Trends in mortality from hepatocellular carcinoma in Europe, 1980-2004. *Hepatology*, 48, 137-145.
- Bucher, JR. (2002). The National Toxicology Program rodent bioassay. Design, interpretations, and scientific contributions. *Ann NY Acad Sci*, 982, 198-207.
- Buendia, M.A. (1992). Hepatitis B viruses and hepatocellular carcinoma. *Adv Cancer Res*, 59, 167-226.
- Bugianesi E (2007). Non-alcoholic steatohepatitis and cancer. *Clin Liver Dis*, 11, 191-207.
- Caldwell, S.H., Oelsner, D.H., Iezzoni, J.C., et al. (1999a). Cryptogenic cirrhosis: clinical characterization and risk factors for underlying disease. *Hepatology*, 29, 664-669.
- Caldwell, S.H., Swerdlow, R.H., Khan, E.M., et al. (1999b). Mitochondrial abnormalities in non-alcoholic steatohepatitis. *J Hepatol*, 31(3), 430-434.
- Calle, E.E., Rodriguez C., Walker-Thurmond K., et al. (2003). Overweight, obesity, and mortality from cancer in a prospectively studied cohort of U.S. adults. *N Engl J Med*, 348, 1625-1638.

- Casals-Casas, C., & Desvergne, B. (2011). Endocrine disruptors: from endocrine to metabolic disruption. *Annu Rev Physiol*, 73, 135-162.
- Chan, P.C., Sills, R.C., Braun, A.G., et al. (1996). Toxicity and carcinogenicity of delta 9-tetrahydrocannabinol in Fischer rats and B6C3F1 mice. *Fundamental Appl Toxicol*, 30, 109-117.
- Chang, B., Nishikawa, M., Nishiguchi, S., et al. (2005). L-carnitine inhibits hepatocarcinogenesis via protection of mitochondria. *Int J Cancer*, 113(5), 719-729.
- Chatterjee, A., Dasgupta, S., Sidransky, D. (2011). Mitochondrial subversion in cancer. *Cancer Prev Res (Phila)*, 4(5): 638-54.
- Chevalier, N., Bouskine, A., Fenichel, P. (2011). Bisphenol A promotes testicular seminoma cell proliferation through GPER/GPR30. *Int J Cancer* In press.
- Chiappini, F., Gross-Goupil, M., Saffroy, R., et al. (2004). Microsatellite instability mutator phenotype in hepatocellular carcinoma in non-alcoholic and non-virally infected normal livers. *Carcinogenesis*, 25(4): 541-547.
- Commission of the European Communities (CEC). The German thorostrast study. (1984). Directorate general Science, Research and Development, EUR 9504 EN.
- Cooper, M. (2009). Regenerative pathologies: stem cells, teratomas and theories of cancer. *Medicine Studies*, 1, 55-66.
- Degli Esposti, D., Pham, P., Saffroy, R., et al. (2009). Environmental risk factors in primary liver cancer: a review of the literature and perspectives for primary prevention and early detection. *Eur J Oncol*, 3, 133-150.
- Dietz, A., Langbein, G., & Permutter, W. (1985). Das Vinylchloride induzierte hepatocellulare Karzinom. *Klin Wochenschr*, 63, 325-331.
- Ducluzeau, P.H., Lachaux, A., Bouvier, R., et al. (1999). Depletion of mitochondrial DNA associated with infantile cholestasis and progressive liver fibrosis. *J Hepatol*, 30(1), 149-155.
- El-Seragh, H.B. (2004). Hepatocellular carcinoma: recent trends in the United States. *Gastroenterology*, 127 Suppl. 5: S27-S34.
- El-Seragh, H. & Rudolph, K.L. (2007). Hepatocellular Carcinoma: Epidemiology and Molecular carcinogenesis. *Gastroenterology*, 132: 2557-2576.
- Evans, D.M.D., Jones, W.W., & Kung, I.T.M. (1983). Angiosarcoma and hepatocellular carcinoma in vinyl chloride workers. *Histopathology*, 7, 377-388.
- Fenton, S.E., Hamm, J.T., Birnbaum, L., et al. (2002). Persistent abnormalities in the rat mammary gland following gestational and lactation exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD). *Toxicol Sci*, 67,63-74.
- Feo, F., Frau, M., Tomasi, M.L., et al. (2009). Genetic and epigenetic control of molecular alterations in hepatocellular carcinoma. *Exp Biol Med*, 234, 726-736.
- Ferlay, J., Shin, H.R., Bray, F., et al. (2010). GLOBOCAN 2008 v1.2, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 10. IARC Press, Lyon, France. Available from: <http://globocan.iarc.fr>
- Ganning, A.E., Brunk, U., Dallner, G. (1984). Phthalate esters and their effect on the liver. *Hepatology*, 4(3), 541-547.
- Gokel, J.M., Liebezeit, E. & Eder, M. (1976). Hemangiosarcoma and hepatocellular carcinoma of the liver following vinylchloride exposure: a report of two cases. *Virchows Arch (A)*, 372, 195-203.

- Goodman A, Schorge J, Greene MF. (2011). The long-term effects of in utero exposures--the DES story. *N Engl J Med*, 364(22), 2083-2084.
- Grün, F., & Blumberg, B. (2009). Endocrine disruptors as obesogens. *Mol Cell Endocrinol*, 304, 19-29.
- Grün, F., Watanabe, H., Zamanian, Z., et al. (2006). Endocrine disrupting organotin compounds are potent inducers of adipogenesis in vertebrates. *Mol Endocrinol*, 9, 2141-2155.
- Guimaraes, J.P., Lamerton, L.F. & Christensen, W.R. (1955). The late effects of thorostrast administration, a review and an experimental study. *Br J Cancer*, 11, 253-267.
- Hagan EC, Jenner PM, Jones WI, Fitzhugh OG, Long EL, Brouwer JG and Webb WK. (1965). Toxic properties of compound related to safrole. *Toxicol Appl Pharmacol*, 7, 18-24.
- Hanahan, D., & Weinberg R.A. (2011). Hallmark of cancer: the next generation. *Cell*, 144, 646-674.
- Hakulinen, T., Lehtimäki, L., Lehtonen, M. et al. (1974). Cancer morbidity among two male cohorts with increased alcohol consumption in Finland. *J Natl Cancer Inst*, 52, 1711-1714.
- Heindel, J.J. (2003). Endocrine disruptors and the obesity epidemic. *Toxicol Sci*, 76(2), 247-249.
- Hensley, K., Kotake, Y., Sang, H., et al. (2000). Dietary choline restriction causes complex I dysfunction and increased H₂O₂ generation in liver mitochondria. *Carcinogenesis*, 21(5), 983-989.
- Herbst, A.L., Ulfelder, H., Poskanzer, D.C. (1971). Adenocarcinoma of the vagina. Association of maternal stilbestrol therapy with tumor appearance in young women. *N Engl J Med*, 284(15), 878-881.
- Hernandez, J.P., Huang, W., Chapman, L.M., et al. (2007). The environmental estrogen, nonylphenol, activates the constitutive androstane receptor. *Toxicol Sci*, 98(2), 416-426.
- Hézode, C., Zafrani, E.S., Roudot-Thoraval, F., et al. (2008). Daily cannabis use: a novel risk factor of steatosis severity in patients with chronic hepatitis C. *Gastroenterology*, 134, 432-439.
- Hill-Baskin, A.E., Markiewski, M.M., Buchner, D.A., et al. (2009). Diet-induced hepatocellular carcinoma in genetically predisposed mice. *Hum Mol Genet* 18(16), 2975-2988.
- Hirayama T. (1981). A large-scale cohort study on the relationship between diet and selected cancers of digestive organs. In: *Gastrointestinal cancer: endogenous factor*, Bruce, W.R., Correa, P., Lipkin, M., Tannenbaum, S.R., Wilkins, T.D. (Banbury Report 7), 409-429. CSH Press, Cold Spring Harbor, NY.
- Hoenerhoff, M.J., Pandiri, A.R., Lahousse, S.A., et al. (2011). Global gene profiling of spontaneous hepatocellular carcinoma in B6CEF1 mice: similarities in the molecular landscape with human liver cancer. *Toxicol Pathol*, 39, 678-699.
- Hotamisligil, G.S. (2010). Endoplasmic reticulum stress and the inflammatory basis of metabolic disease. *Cell*, 140, 900-917.
- Huff, J. (2002). Chemicals studied and evaluated in long-term carcinogenesis bioassays by both the Ramazzini Foundation and the National Toxicology Program: in tribute to Cesare Maltoni and David Rall. *Ann NY Acad Sci*, 982, 208-230.

- Hughes, P.J., McLellan, H., Lowes, D.A., et al. (2000). Estrogenic alkylphenols induce cell death by inhibiting testis endoplasmic reticulum Ca(2+) pumps. *Biochem Biophys Res Commun*, 277(3), 568-574.
- International Agency for Research on Cancer (IARC). (1976). Monographs on evaluation of the carcinogenic risk of chemicals to man. Vol. 10. Some naturally occurring substances. IARC Press, Lyon, France.
- International Agency for Research on Cancer (IARC). (1979). Monographs on evaluation of the carcinogenic risk of chemicals to humans. Vol. 21. Sex hormones (II). IARC Press, Lyon, France.
- International Agency for Research on Cancer (IARC). (1982). Monographs on evaluation of the carcinogenic risk of chemicals to humans. Vol. 29. Some industrial chemicals and dyestuff. IARC Press, Lyon, France.
- International Agency for Research on Cancer (IARC). (1987). Monographs on the evaluation of the carcinogenic risk to humans. Suppl. 7. Overall Evaluations of Carcinogenicity: An Updating of IARC Monographs Volumes 1 to 42. IARC Press, Lyon, France.
- International Agency for Research on Cancer (IARC). (1988). Monographs on evaluation of the carcinogenic risk of chemicals to man. Vol. 44. Alcohol drinking. IARC Press, Lyon, France.
- International Agency for Research on Cancer (IARC). (1993). Evaluation of the carcinogenic risk of chemicals to man. Vol. 56. Some naturally occurring substances. Food items and constituents, heterocyclic aromatic amines and mycotoxins. IARC Press, Lyon, France.
- International Agency for Research on Cancer (IARC). (1997). Monographs on the evaluation of the carcinogenic risk of chemical to humans. Vol. 69. Polychlorinated dibenzo-para-dioxins and polychlorinated dibenzofurans. IARC Press, Lyon, France.
- International Agency for Research on Cancer (IARC). (2001). Monographs on the evaluation of the carcinogenic risk to humans. Vol. 78. Ionizing radiation part 2: some internally deposited radionucleotides. IARC Press, Lyon, France.
- International Agency for Research on Cancer (IARC). (2004a). Monographs on the evaluation of the carcinogenic risk of chemical to humans. Vol. 83. Tobacco smoke and involuntary smoking. IARC Press, Lyon, France.
- International Agency for Research on Cancer (IARC). (2004b). Monographs on the evaluation of the carcinogenic risk of chemical to humans. Vol. 84. Some drinking-water disinfectants and contaminants, including arsenic. IARC Press, Lyon, France.
- International Agency for Research on Cancer. (2004c). Monographs on the evaluation of the carcinogenic risk of chemical to humans. Vol. 85. Betel quid and areca-nut chewing and some areca-nut-derived nitrosamines. IARC Press, Lyon, France.
- International Agency for Research on Cancer. (2008a). World Cancer Report 2008. IARC Press, Lyon, France.
- International Agency for Research on Cancer. (2008b). Monographs on the evaluation of the carcinogenic risk of chemical to humans. Vol. 97. 1,3-Butadiene, ethylene oxide, and vinyl halides (vinyl fluoride, vinyl chloride, vinyl bromide). IARC Press, Lyon, France.

- International Agency for Research on Cancer (IARC). (2011). Monographs on the evaluation of the carcinogenic risk to humans. Vol. 100A. A review of human carcinogens: pharmaceuticals. IARC Press, Lyon, France.
- Jick, H., & Herman R. (1978). Oral contraceptive-induced benign liver tumors. The magnitude of the problem. *J Am Med Assoc*, 240, 828-829.
- Jin, X., Moskophidis, D., Mivechi, N.F. (2011). Heat shock transcription factor 1 is a key determinant of HCC development by regulating hepatic steatosis and metabolic syndrome. *Cell Metab*, 14(1), 91-103.
- Johnson, F.L., Lerner, K.G., and Siegal M. (1972). Association of androgenic-anabolic steroid therapy with development of hepatocellular carcinoma. *Lancet*, 2, 1273-1276.
- Kaidar-Person, O., Bar-Sela, G., Person B. (2011). The two major epidemics of the twenty-first century: obesity and cancer. *Obes Surg*, in press.
- Kammoun, H.L., Chabanon, H., Hainault, I., et al. (2009a). GRP78 expression inhibits insulin and ER stress-induced SREBP-1c activation and reduces hepatic steatosis in mice. *J Clin Invest*, 119(5), 1201-1215.
- Kammoun, H.L., Hainault, I., Ferré, P., et al. (2009b). Nutritional related liver disease: targeting the endoplasmic reticulum stress. *Curr Opin Clin Nutr Metab Care*, 12, 575-582.
- Kavlock, R.J., Daston, G.P., DeRosa, C., et al. (1996). Research needs for the risk assessment of health and environmental effects of endocrine disruptors: a report of the U.S. EPA-sponsored workshop. *Environ Health Perspect*, 104 Suppl 4, 715-740.
- Kim, E.H., Koh, E.H., Park, J.Y., et al. (2010) Adenine nucleotide translocator as a regulator of mitochondrial function: implication in the pathogenesis of metabolic syndrome. *Korean Diabetes J*, 34(3), 146-153.
- Kinosita, R. (1936). Researches on the carcinogenesis of the various chemical substances. *Gann*, 30, 423.
- Klatskin, G. (1977). Hepatic tumors. Possible relation to use of oral contraceptives. *Gastroenterology*, 73, 386-394.
- Koischwitz, V.D., Lelbach, W.K., Lackner, K. et al. (1981). Das vinylchloridinduzierte Leberangiosarkoma und hepatozelluläre Karzinom. *Fortschr Rontgenstr*, 134, 283-290.
- Kovacic, P. (2010). How dangerous are phthalate plasticizers? Integrated approach to toxicity based on metabolism, electron transfer, reactive oxygen species and cell signaling. *Med Hypotheses*, 74(4), 626-628.
- Lacassagne, A. (1936). Hormonal pathogenesis of adenocarcinoma of the breast. *Am J Cancer*, 27: 217-225.
- Lagiou, P., Kuper, H., Stuver, S.O., et al. (2000). Role of diabetes mellitus in the etiology of hepatocellular carcinoma. *J Natl Cancer Inst*, 92, 1096-1099.
- Laquer, G.L., Mickelsen, O., Whiting, M.G., et al. (1963). Carcinogenic properties of nuts from *Cycas circinalis* L. indigenous to Guam. *J Natl Cancer Inst*, 31, 919-951.
- Laurent-Puig, P., Legoux, P., Bluteau, O., et al. (2001). Genetic alterations associated with hepatocellular carcinomas define distinct pathways of hepatocarcinogenesis. *Gastroenterology*, 120, 1763-1773.

- Lehnert, B.E., Goodwin, E.H., Deshpande, A. (1997). Extracellular factor(s) following exposure to alpha particles can cause sister chromatid exchanges in normal human cells. *Cancer Res*, 57, 2164-2171.
- Li, X., Zhang, K., Li, Z. (2011). Unfolded protein response in cancer: the physician's perspective. *J Hematol Oncol*, 4 (8) 1-10.
- Long, E.L., Nelson, A.A., Fitzhugh, O.G., et al. (1963). Liver tumors produced in rats by feeding safrole. *Arch Pathol*, 75, 595-604.
- Ludwig, J.A., & Weinstein, J.N. (2005). Biomarkers in cancer staging, prognosis and treatment selection. *Nat Rev Cancer*, 11, 845-56.
- Magee, P.N., & Barnes, J.M. (1956). The production of malignant primary hepatic tumors in the rat by feeding dimethylnitrosamine. *Br J Cancer*, 10, 114-122.
- Maltoni, C., Lefemine, G., Ciliberti, A., et al. (1984). *Experimental research on vinyl chloride carcinogenesis. Archives of Research on Industrial Carcinogenesis, vol II*. Princeton Scientific Publishers, Princeton, NJ.
- Maltoni, C., Lefemine, G., & Cotti, G. (1986). *Experimental research on trichloroethylene carcinogenesis. Archives of Research on Industrial Carcinogenesis, vol V*. Princeton Scientific Publishers, Princeton, NJ.
- Maltoni, C., Ciliberti, A., Cotti, G., et al. (1989). Benzene, an experimental multipotential carcinogen: results of the long-term bioassays performed at the Bologna Institute of Oncology. *Environ Health Perspect*, 82, 109-124.
- Maltoni, C., Minardi, F., Pinto C., et al. (1997). Results of three life-span experimental carcinogenicity and anticarcinogenicity studies on tamoxifen in rats. In: Preventive strategies for Living in a chemical World, *Ann NY Acad Sci*, 837, 469-512.
- Marek, C.B., Peralta, R.M., Itinose, A.M., et al. (2011). Influence of tamoxifen on gluconeogenesis and glycolysis in the perfused rat liver. *Chem Biol Interact*, 193(1), 22-33.
- Maronpot, R.R. (2009). Biological basis of differential susceptibility to hepatocarcinogenesis among mouse strains. *J Toxicol Pathol*, 22, 11-33.
- Maronpot, R.R., Flake, G., Huff, J. (2004). Relevance of animal carcinogenesis findings to human cancer predictions and prevention. *Toxicol Pathol* 32(Suppl. 1): 40-48.
- Marrero, J.A., Fontana, R.J., Su, G.L., et al. (2002). NAFLD may be a common underlying liver disease in patients with hepatocellular carcinoma in the United States. *Hepatology*, 36, 1349-1354.
- Masumo, H., Kidani, T., Sekiya, K., et al. (2002). Bisphenol A in combination with insulin can accelerate the conversion of 3T3L1 fibroblasts to adipocytes. *J Lipid Res*, 43, 676-684.
- McCullough, A. (2004). The epidemiology and risk factors of NASH. In: *Fatty liver disease: NASH and related disorders*. Farrel GC, George J, Hall P, McCullough A Eds, 23-37, Blackwell Publishing, Oxford.
- McGlynn, K.A., Tsao, L., Hsing A.W., et al. (2001). International trends and pattern of primary liver cancer. *Int J Cancer*, 94, 290-296.
- Minardi, F., Belpoggi, F., Soffritti, M., et al. (1994). Studio sperimentale sugli effetti a lungo termine del tamoxifen sul fegato di ratto. In: *I tumori primitivi e secondari del fegato*, Natale, C., Laricchiuta, D. et al. Società Italiana di Prevenzione Diagnosi e Terapia dei Tumori, Monduzzi Editore, Bologna, Italy.

- Ministère de la Santé et de la Solidarité. (2006). *Programme National Nutrition Santé*. Retrieved from http://www.sante.gouv.fr/IMG/pdf/synthese_et_sommaire_PNNS_.pdf
- Mita, L., Bianco, M., Viggiano, E., et al. (2011). Bisphenol A content in fish caught in two different sites of the Tyrrhenian sea (Italy). *Chemosphere*, 82(3), 405-410.
- Møller, H., Møller, A., Lindvig, K., et al. (1994). Obesity and cancer risk: a Danish record-linkage study. *Eur J Cancer*, 30A, 344-350.
- Mueller KM, Kornfeld JW, Friedbichler K, et al. (2011). Impairment of hepatic growth hormone and glucocorticoid receptor signaling causes steatosis and hepatocellular carcinoma in mice. *Hepatology* In press.
- Nadeau, K., Klingensmith, G., & Sokol, R.J. (2003). Case report: steatohepatitis in a teenage girl with type 2 diabetes. *Curr Opin Pediatr*, 15(1), 127-131.
- National Toxicology Program (NTP). (2011). Chemicals associated with site-specific tumor induction in liver. Available on: <http://ntp.niehs.nih.gov/?objectid=E1D17854-123F-7908-7B168177A810AEDC>.
- Nazarewicz, R.R., Zenebe, W.J., Parihar, A., et al. (2007). Tamoxifen induces oxidative stress and mitochondrial apoptosis via stimulating mitochondrial nitric oxide synthase. *Cancer Res*, 67(3), 1282-1290.
- Oh, S.W., Yoon, Y.S., Shin, S.A. (2005). Effects of excess weight on cancer incidences depending on cancer sites and histologic findings among men: Korea National Health Insurance Corporation Study. *J Clin Oncol*, 23, 4742-4754.
- Olsen, J.H., Dragsted, L., & Autrup H. (1988). Cancer risk and occupational exposure to aflatoxins in Denmark. *Br J Cancer*, 58, 392-396.
- Ozcan, U., Cao, Q., Yilmaz, E., et al. (2004). Endoplasmic reticulum stress links obesity, insulin action, and type 2 diabetes. *Science*, 306, 457-461.
- Page, J.M. & Harrison, S.A. (2009). NASH and HCC. *Clin Liv Dis*, 13, 631-647.
- Palmer, J.R., Wise, L.A., Hatch, E.E., et al. (2006). Prenatal diethylstilbestrol exposure and risk of breast cancer. *Cancer Epidemiol Biomarkers Prev* 8, 1509-1514.
- Parfieniuk, A. & Flisiak, R. (2008). Role of cannabinoids in chronic liver diseases. *World J Gastroenterol*, 14, 6109-6114.
- Parkin DM, Bray F, Ferlay J et al. (2005). Global cancer statistics 2002. *CA Cancer J Clin*, 55, 74-108.
- Pei, Y., Zhang, T., Renault, V., et al. (2009). An overview of hepatocellular carcinoma study by omics-based methods. *Acta Biochim Biophys Sin (Shanghai)*, 41(1), 1-15.
- Pereira, C., Mapuskar, K., Rao, C.V. (2006). Chronic toxicity of diethyl phthalate in male Wistar rats--a dose-response study. *Regul Toxicol Pharmacol*, 45(2), 169-177.
- Pérez-Fuenteaja, A., Lupton, S., Clapsadl, M., et al. (2010). PCB and PBDE levels in wild common carp (*Cyprinus carpio*) from eastern Lake Eire. *Chemosphere*, 81(4): 541-547.
- Pessayre, D. & Fromenty, B (2005). NASH: a mitochondrial disease. *J Hepatol*, 42, 928-940.
- Pirastu, R., Comba, P., Reggiani, A., et al. (1990). Mortality from liver disease among Italian vinyl chloride monomer/polyvinyl chloride manufacturers. *Am J Ind Med*, 17,155-161.
- Potter, V.R. (1958). The biochemical approach to the cancer problem. *Fed Proc*, 17, 691-697.
- Puigserver, P., Wu, Z., Park, C.W., et al. (1998). A cold-inducible coactivator of nuclear receptors linked to adaptive thermogenesis. *Cell*, 92(6), 829-839.

- Purchase, I.F.H., & Van der Watt J.J. (1970). Carcinogenicity of sterigmatocystin. *Food Cosmet Toxicol*, 8, 289-295.
- Puri, P., Mirshahi, F., Cheung, O., et al. (2008). Activation and dysregulation of the unfolded protein response in nonalcoholic fatty liver disease. *Gastroenterology*, 134, 568-576.
- Purushotham, A., Schug, T.T., Xu, Q., et al. (2009). Hepatocyte-specific deletion of SIRT1 alters fatty acid metabolism and results in hepatic steatosis and inflammation. *Cell Metab*, 9(4), 327-338.
- Rael, L.T., Bar-Or, R., Ambruso, D.R., et al. (2009). Phthalate esters used as plasticizers in packed red blood cell storage bags may lead to progressive toxin exposure and the release of pro-inflammatory cytokines. *Oxid Med Cell Longev*, 2(3), 166-171.
- Rashid, M., Roberts, E.A. (2000). Nonalcoholic steatohepatitis in children. *J Pediatr Gastroenterol Nutr*, 30(1), 48-53.
- Rector, R.S., Thyfault, J.P., Uptergrove, G.M., et al. (2010). Mitochondrial dysfunction precedes insulin resistance and hepatic steatosis and contributes to the natural history of non-alcoholic fatty liver disease in an obese rodent model. *J Hepatol*, 52(5), 727-736.
- Ren, H., Aleksunes, L.M., Wood, C., et al. (2010). Characterization of peroxisome proliferator-activated receptor alpha--independent effects of PPARalpha activators in the rodent liver: di-(2-ethylhexyl) phthalate also activates the constitutive-activated receptor. *Toxicol Sci*, 113(1), 45-59.
- Rogers, A.B., Theve, E.J., Feng, Y., et al. (2007). Hepatocellular carcinoma associated with liver-gender disruption in male mice. *Cancer Res*, 67(24), 11536-11546.
- Russo, J. & Russo, I.H. (2006). The role of estrogen in the initiation of breast cancer. *J Steroid Biochem Mol Biol*; 102, 89-96.
- Saffroy, R., Pham, P., Reffas, M., et al. (2007). New perspectives and strategy research biomarkers for hepatocellular carcinoma. *Clin Chem Lab Med*, 45, 1169-1179.
- Salvucci, M., Lemoine, A., Saffroy, R., et al. (1999). Microsatellite instability in European hepatocellular carcinoma. *Oncogene*, 18(1), 181-187.
- Sanyal, A.J., Campbell-Sargent, C., Mirshahi, F., et al. (2001). Nonalcoholic steatohepatitis: association of insulin resistance and mitochondrial abnormalities. *Gastroenterology*, 120, 1183-1192.
- Sato N. (2007). Central role of mitochondria in metabolic regulation of liver pathophysiology. *J Gastroenterol Hepatol*, 22 (Suppl1), S1-6.
- Scatena, R., Bottoni, P., Giardina, B. (2008). Mitochondria, PPARs, and Cancer: Is Receptor-Independent Action of PPAR Agonists a Key? *PPAR Res*, 2008, 256251.
- Scatena, R., Bottoni, P., Martorana, G.E., et al. (2004). Mitochondrial respiratory chain dysfunction, a non-receptor-mediated effect of synthetic PPAR-ligands: biochemical and pharmacological implications. *Biochem Biophys Res Commun*, 319(3), 967-973.
- Schmal, D., Preussmann, R., & Hamperl H. (1960). Leberkrebs erzeugende Wirkung von Diathylnitrosamin nach oraler Gabe bei Ratten. *Naturwissenschaften*, 47, 89.
- Schreiber, S.N., Emter, R., Hock, M.B., et al. (2004). The estrogen-related receptor alpha (ERRalpha) functions in PPARgamma coactivator 1alpha (PGC-1alpha)-induced mitochondrial biogenesis. *Proc Natl Acad Sci U S A*, 101(17), 6472-6477.

- Schröder, M. & Kaufman, R.J. (2005). The mammalian unfolded protein response. *Annu Rev Biochem*, 74, 739-789.
- Schwimmer, J.B., Deutsch, R., Kahen, T., et al. (2006). Prevalence of fatty liver in children and adolescents. *Pediatrics*, 118(4), 1388-1393.
- Seo, H.Y., Kim, Y.D., Lee, K.M., et al. (2008). Endoplasmic reticulum stress-induced activation of activating transcription factor 6 decreases insulin gene expression via up-regulation of orphan nuclear receptor small heterodimer partner. *Endocrinology*, 149(8), 3832-3841.
- Shah, I., Houck, K., Judson, R.S., et al. (2011). Using nuclear receptor activity to stratify hepatocarcinogens. *PLoS One*, 6(2), e14584.
- Shaw, J.J. & Shah, S.A. 2011. Rising incidence and demographics of hepatocellular carcinoma in the USA: what does it mean? *Expert Rev Gastroenterol Hepatol*, 5(3), 365-370.
- Shuda, M., Kondoh, N., Imazeki, N., et al. (2003). Activation of the ATF6, XBP1 and grp78 genes in human hepatocellular carcinoma: a possible involvement of the ER stress pathway in hepatocarcinogenesis. *J Hepatol*, 38(5), 605-614.
- Skill, N.J., Scott, R.E., Wu, J., et al. (2011). Hepatocellular carcinoma associated lipid metabolism reprogramming. *J Surg Res*, 169(1), 51-56.
- Snyder, R.L., Tyler, G. & Summers J. (1982). Chronic hepatitis and hepatocellular carcinoma associated with woodchuck hepatitis virus. *Am J Pathol*, 107, 422-425.
- Soffritti, M. & McConnell EE. (1988). Liver foci formation during Aflatoxin B1 carcinogenesis in the rat. *Ann NY Acad Sci*, 534, 531-540.
- Soffritti, M., Belpoggi, F., Minardi, F., et al. (2002). Ramazzini Foundation cancer program: history and major projects, life-span carcinogenicity bioassays design, chemical studies, and results. *Ann NY Acad Sci*, 982, 26-45.
- Soffritti, M., Belpoggi, F., Manservigi, M., et al. (2010). Aspartame administered in feed, beginning prenatally through life span, induces cancers of the liver and lung in male Swiss mice. *Am J Ind Med*, 53, 1197-1206.
- Somm, E., Schwitzgebel, V.M., Toulotte, A., et al. (2009). Perinatal exposure to bisphenol a alters early adipogenesis in the rat. *Environ Health Perspect*, 117(10), 1549-1555.
- Sonoda, J., Mehl, I.R., Chong, L.W., et al. (2007). PGC-1beta controls mitochondrial metabolism to modulate circadian activity, adaptive thermogenesis, and hepatic steatosis. *Proc Natl Acad Sci U S A*, 104(12), 5223-5228.
- Sørensen, H.T., Mellekjaer, L., Jepsen, P., et al. (2003). Risk of cancer in patients hospitalized with fatty liver: a Danish cohort study. *J Clin Gastroenterol*, 36, 356-359.
- Soto, A.M., & Sonnenschein, C. (2004). The somatic mutation theory of cancer: growing problems with the paradigm? *Bioassays*, 26, 1097-1107.
- Soto, A.M., & Sonnenschein, C. (2010). Environmental causes of cancer: endocrine disruptors as carcinogens. *Nat Rev Endocrinol* 6(7), 363-370.
- Starley, B.Q., Calcagno, C.J., & Harrison, S.A. (2010). Nonalcoholic fatty liver disease and hepatocellular carcinoma: a weighty connection. *Hepatology*, 51, 1820-1832.
- Sundaram, S.S., Zeitler, P., & Nadeau, K. (2009). The metabolic syndrome and nonalcoholic fatty liver disease in children. *Curr Opin Pediatr*, 21(4), 529-535.
- Swoboda, D.J., and Reddy, J.K. (1972). Malignant tumors in rats given lasiocarpine. *Cancer Res*, 32, 908-912.


- Tabuchi, Y., Takasaki, I., Kondo, T. (2006). Identification of genetic networks involved in the cell injury accompanying endoplasmic reticulum stress induced by bisphenol A in testicular Sertoli cells. *Biochem Biophys Res Commun*, 345(3), 1044-1050.
- Takizawa, D., Kakizaki, S., Horiguchi, N., et al. (2011). Constitutive active/androstane receptor promotes hepatocarcinogenesis in a mouse model of non-alcoholic steatohepatitis. *Carcinogenesis*, 32(4), 576-583.
- Teebor, G.W., & Becker FF. (1971). Regression and persistence of hyperplastic hepatic nodules induced by N-2-Fluorenilacetamide and their relationship to hepatocarcinogenesis. *Cancer Res*, 31, 1-3.
- Ter Veld, M.G., Zawadzka, E., van den Berg, J.H., et al. (2008). Food-associated estrogenic compounds induce estrogen receptor-mediated luciferase gene expression in transgenic male mice. *Chem Biol Interact*, 174(2), 126-33.
- Terayama, H. (1967). Aminoazo carcinogenesis methods and biochemical problems. *Methods Cancer Res*, 1, 399-449.
- Upton AC. Radiation carcinogenesis. (1968). *Methods Cancer Res*, 4, 53-82.
- Upton AC., Furth J. and Burnett WT. (1956). Liver damage and hepatomas in mice produced by radioactive colloidal gold. *Cancer Res*, 16, 211-215.
- Uraguchi, K., Saito, M., Noguchi, Y., et al. (1972). Chronic toxicity and carcinogenicity in mice of the purified mycotoxins, luteoskyrin and cyclochlorotene. *Food Cosmet Toxicol*, 10, 193-207.
- Vandenberg, L.N., Maffini, M.V., Schaeberle C.M., et al. (2008). Perinatal exposure to the xenoestrogen bisphenol-A induces mammary intraductal hyperplasias in adult CD-1 mice. *Reprod Toxicol*, 26(3-4), 210-219.
- Vander Heiden, M.G., Cantley, L.C., & Thompson, C.B. (2009). Understanding the Warburg effect: the metabolic requirements of cell proliferation. *Science*, 324, 1029-1033.
- VanSaun, M.N., Lee, I.K., Washington, M.K., et al. (2009). High fat diet induced hepatic steatosis establishes a permissive microenvironment for colorectal metastases and promotes primary dysplasia in a murine model. *Am J Pathol*, 175(1), 355-364.
- Verderame, M., Prisco, M., Andreucci, P., et al. (2011). Experimentally nonylphenol-polluted diet induces the expression of silent genes VTG and Era in the liver of male lizard *Podaris sicula*. *Environ Pollut*, 159(5), 1101-1107.
- Vickers, A.E. & Lucier, G.W. (1991). Estrogen receptor, epidermal growth factor receptor and cellular ploidy in elutriated subpopulations of hepatocytes during liver tumor promotion by 17 alpha-ethinylestradiol in rats. *Carcinogenesis*, 12, 391-399.
- Villena, J.A. & Kralli, A. (2008). ERRalpha: a metabolic function for the oldest orphan. *Trends Endocrinol Metab* 19(8), 269-276.
- Wagner, M., Zollner, G., Trauner, M. (2011). Nuclear receptors in liver disease. *Hepatology*, 53(3), 1023-1034.
- Wang, Q., Zhao, X.F., Ji, Y.L., et al. (2010). Mitochondrial signaling pathway is also involved in bisphenol A induced germ cell apoptosis in testes. *Toxicol Lett*, 199(2), 129-135.
- Wang, Y., Ausman, L.M., Greenberg, A.S., et al. (2009). Nonalcoholic steatohepatitis induced by a high-fat diet promotes diethylnitrosamine-initiated early hepatocarcinogenesis in rats. *Int J Cancer*, 124(3), 540-546.
- Warburg, O. (1930). The metabolism of tumors: investigation from the Kaiser Wilhelm Institute for Biology, Berlin-Dahlem.

- Welzel, T.M., Graubard, B.I., Zeuzem, S., et al. (2011). Metabolic syndrome increases the risk of primary liver cancer in the United States: a study in the SEER-medicare database. *Hepatology*, 54(2) 463-471.
- White DL, Li D, Nurgalieva Z, El-Serag HB. (2008). Genetic variants of glutathione S-transferase as possible risk factors for hepatocellular carcinoma: a HuGE systematic review and meta-analysis. *Am J Epidemiol*, 167(4), 377-389.
- White House Task Force on Childhood Obesity. (2010). *Report to the President*. Retrieved from <http://www.letsmove.gov/white-house-task-force-childhood-obesity-report-president>
- Wideroff, L., Gridley, G., Mellekjaer, L., et al. (1997). Cancer incidence in a population-based cohort of patients hospitalized with diabetes mellitus in Denmark. *J Natl Cancer Inst*, 89, 1360-1365.
- Williams, C.D., Stengel, J., Asike, M.I., et al. (2011). Prevalence of nonalcoholic fatty liver disease and nonalcoholic steatohepatitis among a largely middle-aged population utilizing ultrasound and liver biopsy: a prospective study. *Gastroenterology*, 140, 124-131.
- Wingspread consensus statement 1992, as cited in Soto, A.M. & Sonnenschein, C. (2010).
- Wilson, R.H., De Eds F., & Cox A.J. (1941). The toxicity and carcinogenic activity of 2-acetaminofluorene. *Cancer Res* 1, 595-608.
- Wogan, G.N., & Newberne, P.N. (1967). Dose-response characteristics of aflatoxin B1 carcinogenesis in rats. *Cancer Res*, 27, 2370-2376.
- Wogan, G.N., Pagliarlunga, S. & Newberne P.N. (1974). Carcinogenic effects of low dietary levels of aflatoxin B1 in rats. *Food Cosmet Toxicol*, 12, 681-685.
- Wolk, A., Gridley, G., Svensson, M., et al. (2001). A prospective study of obesity and cancer risk (Sweden). *Cancer Causes Control*, 12, 13-21.
- Wong, N., Lai, P., Pang, E., et al. (2000). Genomic aberrations in human hepatocellular carcinomas of differing etiologies. *Clin Cancer Res*, 6, 4000-4009.
- World Health Organization. (2006). *European Charter on Counteracting Obesity*. WHO Press, retrieved from http://www.euro.who.int/__data/assets/pdf_file/0009/87462/E89567.pdf
- World Health Organization. (2008). *Global Burden of Disease: 2004 update*. WHO Press , retrieved from http://www.who.int/healthinfo/global_burden_disease/2004_report_update/en/index.html
- Wu, Z., Puigserver, P., Andersson, U., et al. (1999). Mechanisms controlling mitochondrial biogenesis and respiration through the thermogenic coactivator PGC-1. *Cell*, 98(1), 115-124.
- Yang, F., Huang, X., Yi, T et al. (2007). Spontaneous development of liver tumors in the absence of the bile acid receptor farnesoid X receptor. *Cancer Res*, 67, 863-867.
- Yang, S., Zhu, H., Li, Y., et al. (2000). Mitochondrial adaptations to obesity-related oxidant stress. *Arch Biochem Biophys*, 378(2), 259-268.
- Yeh, F.S., Mo, C.C., & Yen RC. (1985). Risk factors for hepatocellular carcinoma in Guangxi, people's Republic of China. *Natl Cancer Inst Monogr*, 69, 47-48.
- Yin, P.H., Wu, C.C., Lin, J.C., et al. (2010). Somatic mutations of mitochondrial genome in hepatocellular carcinoma. *Mitochondrion*, 10(2), 174-182.

Zender, L., Spector, M.S., Xue, W., et al. (2006). Identification and validation of oncogenes in liver cancer using an integrative oncogenomic approach. *Cell*, 125(7), 1253-1267.

IntechOpen

IntechOpen



Hepatocellular Carcinoma - Clinical Research

Edited by Dr. Joseph W.Y. Lau

ISBN 978-953-51-0112-3

Hard cover, 330 pages

Publisher InTech

Published online 02, March, 2012

Published in print edition March, 2012

This book covers the clinical aspects of hepatocellular carcinoma. This book is a compendium of papers written by experts from different parts of the world to present the most up-to-date knowledge on the clinical aspects of hepatocellular carcinoma. This book is divided into three sections: (I) Diagnosis / Differential Diagnosis; (II) Surgical Treatment; (III) Non-surgical Treatment. There are 19 chapters covering topics from novel diagnostic methods to hepatic lesions mimicking hepatocellular carcinoma, from laparoscopic liver resection to major hepatectomy without allogeneic blood transfusion, from molecular targeted therapy to transarterial radioembolization, and from local ablative therapy to regional therapy. This volume is an important contribution to the clinical management of patients with hepatocellular carcinoma. The intended readers of this book are clinicians who are interested in hepatocellular carcinoma, including hepatologists, liver surgeons, interventional and diagnostic radiologists, pathologists and epidemiologists. General surgeons, general physicians, trainees, hospital administrators, and instruments and drug manufacturers will also find this book useful as a reference.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Davide Degli Esposti, Morando Soffritti, Antoinette Lemoine, Eva Tibaldi and Marco Manservigi (2012). Hepatocellular Carcinoma: Epidemiology and Etiology, *Hepatocellular Carcinoma - Clinical Research*, Dr. Joseph W.Y. Lau (Ed.), ISBN: 978-953-51-0112-3, InTech, Available from:
<http://www.intechopen.com/books/hepatocellular-carcinoma-clinical-research/hepatocellular-carcinoma-epidemiology-and-etiology>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen