

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



# Total Growth of Tomato Hybrids Under Greenhouse Conditions

Humberto Rodriguez-Fuentes<sup>1</sup>, Juan Antonio Vidales-Contreras<sup>1</sup>,  
Alejandro Isabel Luna-Maldonado<sup>1</sup> and Juan Carlos Rodriguez-Ortiz<sup>2</sup>

<sup>1</sup>Department of Agricultural and Food Engineering, Faculty of Agriculture,  
Autonomous University of Nuevo Leon, Escobedo, Nuevo Leon,

<sup>2</sup>Faculty of Agriculture, Autonomous University of San Luis Potosi, San Luis Potosi,  
Mexico

## 1. Introduction

Often in intensive production of tomato, the fertilization is applied by the farmers without consider the suitable doses in order to cover nutritional requirements according to crop physiological stages. Thus, appropriate crop management is a strategic demand to maintain or increase tomato production. In spite of many researchers conducted experiments in this subject and data is available about physiological stage requirements for plant nutrition, only few studies have been focused to nutritional parameters. The crop growth curves and nutrient uptake for tomato may determine uptake rate for a particular nutrient eluding possible deficiencies and superfluous fertilizer consumption. The daily rates of nutrient uptake are depending on crop and wheater (Scaife and Bar-Yosef, 1995; Honorato *et al.*, 1993; Magnificent *et al.*, 1979; Miller *et al.*, 1979); however, crop requirements and opportune fertilizer applications, are little known in many of fresh consumption crops. In Mexico, vegetable production is located at desert areas in the north and middle of the country where water shortages have constraints with impact on of water demands the crops of tomato, pepper and cucumber. Thus, the surface for crop production in greenhouses has increased from 350 ha in 1997 (Steta, 1999) to about 5000 ha in 2006 (Fonseca, 2006), because the increasing demand for quality products and the risk of losses on field for crop production.

Imas (1999) found that nutrient uptake and fertilization recommendations are conditions depending of crops. For example, tomato crops under hydroponic greenhouse environments have averaged 200 t ha<sup>-1</sup> which is significantly higher than the 60-80 t ha<sup>-1</sup> yield, typically observed in an open field. In contrast to tomato crops grown in an open field, nutrient uptake in greenhouse environment can be duplicated or triplicated. In practical terms, crop growth cycle is divided according to physiological stages thence different concentrations or amounts of nutrients have to be applied according to recommendations given by Department of Agriculture. In the tomato production are considered four physiological stages: establishment-flowering, flowering-fruit set, ripening of tomato fruit (the first-crop harvest and last harvest on the crop). In each stage concentrations of nitrogen (N) and phosphorus (K) are increasing while nitrogen-phosphorous are decreasing because potassium is uptaken in large quantities during the reproductive stage of the crop (Zaidan

and Avidan, 1997). In order to determine nutrient uptake more accurately, crop growth can be scheduled by chronological periods of sampling and analysis nearest one another.

The nutrient demand is the maximum amount of nutrient that a crop needs in order to meet their metabolic growth demand and development and that is calculated to maximize production goals and domestic demand price (nutritional optimal concentration of total biomass (air and / or root) at harvest time); however this criteria is not yet thought by the farmers. An appropriate method for such calculation is to use mass balance concept in a hydroponic system. In this way, nutrimental control is more efficient in the nutritive solution and its effect can be seen rapidly in the plants (Steiner, 1961). The method before mentioned is based on the composition of plant dry matter which consists of 16 essential nutrients; however, only thirteen are directly uptaken by plants from soil. Therefore, if the amount of total dry matter production (root + aerial part) during the cycle of growth and development as well as nutrimental concentration in each physiological stage is determined, the amount of nutrients that the plant absorbs can be estimated. With this analysis it is possible to establish a program of daily/weekly fertilization for crop production. This would result in a substantial economic savings on fertilizer costs and also in a decrease of negative environmental impact for its inappropriate applications.

## 2. Tomato hybrids analysis

The field study was conducted on the farm "El Cuento" located in the town of Marin, Nuevo León, Mexico (latitude N 25° 53' and longitude W 100° 02' and 400 m above sea level), where two greenhouses were used (Figure.1), first one of these greenhouses was a tunnel type of 900 m<sup>2</sup> (75 m x 12 m and 7 m high) and second one was multi-Korean tunnel type of 1300 m<sup>2</sup> (45 m x 30 m). It was used an open hydroponics system during the tests of two indeterminate hybrid tomatoes beef type.

The seeds were sown in a mixture of peat moss and perlite (1:1 v/v) in containers of 200 cavities. The transplanting was realized at 40 days after sowing. Plants were transplanted inside of 2 gallons polythene bags (white outer and black inner, Fig.2) with a mixture of perlite and peat moss (1:1, v/v). In the Korean type greenhouse it was settled 50% of plants of the hybrids Cayman (1700) and Charleston (1700). In the tunnel type greenhouse it was established 2220 plants of the hybrid Charleston, in both cases it was used a density of 2.5 plants m<sup>-2</sup>. The hydroponic irrigation system and nutrition of the plants were conducted with emitters calibrated to 4 min L<sup>-1</sup>. It was drained of 10 to 15% of the hydroponic solution applied. The nutrimental solution used was suggested by Rodríguez *et al.* (2006).

Plant sampling and were conducted according to tomato crops and greenhouse type (Fig 3). On plant sampling, three plants, visually uniform, were removed from the hydroponic system (substrate) every 15 days after plant plantation total dry matter was determined considering root + leaves + stems + flowers and fruits (if they were in the plant) To obtain the biomass model in both production and the nutrient uptake, roots, stems, leaves, and fruits were included. The samples were analyzed in the Laboratory of Soil, Water and Plant Analysis at the Agronomy School of Autonomous University of Nuevo Leon, where plant specimens were washed with deionizer water to obtain their dried weights. In addition to this parameter, dry matter, nitrogen, phosphorus, potassium, calcium and magnesium were determined per plant sample.



Fig. 1. Tunnel-type greenhouse used for experimentation located in Marin, Nuevo Leon, Mexico.



Fig. 2. Vinyl bag used for growing tomato in a hydroponics system.





Fig. 3. Tomato plants inside of the experimental greenhouse.

To determine the dry weight at constant weight, samples were milled and then placed in a forced convection oven (Riossa model F-62, Mexico) at 70° to 80°C. Samples were screened through a 50  $\mu\text{m}$  mesh. Determination of total nitrogen was done by the Kjeldhal method (Rodríguez and Rodríguez, 2011). A wet digestion microwave (MARSX, EMC Corporation, North Carolina, USA) was used in order to analyze P, K, Ca, and Mg. Three dried samples of  $0.5 \pm 0.001\text{g}$  were placed in a digestion flask (Teflon PFA vessels of a capacity of 120 mL) and then 5 mL of  $\text{HNO}_3$  were added. The instrument was programmed with a ramp of 15 min to reach a temperature of 180 degrees Celsius and a pressure of 300 PSI and keeping these conditions for 5 minutes. Finally, to the samples were left to cool for 15 minutes.

The dry matter accumulated values in the hybrid tomato were analyzed with the DM Sigma Plot 10.0 Program and Microsoft Excel Office 2003 software and it was found that the best fitted linear regression model. To calculate the equations were considered the average values of both hybrids.

### 3. Flowering period

The flowering period occurred after 20 days after plant plantation, fruits set showed up 10 days later and until 55 days after transplanting. During the beginning of flowering there was a low accumulation of dry matter. On the fruit set and harvest time of fruit an

accumulation of intense biomass was observed (Figures. 4 and 5) until the establishment of the harvest. In last stage of crop growth the dry matter is decreasing as well as the nutrient uptake. Bugarin *et al.* (2002), coincided in their research due in same crop growth stage any nutritional deficit decreases production.



Fig. 4. Tomato plants (variety Charleston) in breaking stage.

#### 4. Accumulation of dry matter production

Accumulation of dry matter production had a linear behavior up to 96-day initial crop period. Thereafter, an erratic behavior was observed probably as a consequence of the conditions of high temperatures and low relative humidity occurred in those greenhouse designs. This condition caused a severe reduction in dry matter accumulation, low pollination of fruit and fruit set. Sampling finished after 170 days since transplanting stage; however, plant health recovery did not occurred, reducing drastically fruit production to about one third of the best period, from 2.5 to 0.8 kg m<sup>2</sup> after 125 days since transplanting. The same reduction is normal but it occurs gradually as the harvest progresses and the number of clusters.

Observed data sets were analyzed for dry matter production during 96 days after transplanting. Three models of linear fitting (Table 1 and Figures 6, 7, 8) with R<sup>2</sup> values of 0.96, 0.99 and 0.91 for hybrids: Cayman (CAI), Charleston greenhouse Korean (CHK) and Charleston in greenhouse Tunnel (CHT), respectively, were obtained. The R<sup>2</sup> value for the average of both hybrids was 0.96 (Table 1, Figs. 6 and 7) to predict possible accumulation of DM with the models obtained for each hybrid, is to believe the 170 days scheduled time to remove crop residues and begin the next crop season.





Fig. 5. Pruning of unnecessary tomato shoots.




Fig. 6. Dry matter and total accumulated average in CAI: Caiman; DM A: Dry Matter average; ADT: After day trasplanting.




Fig. 7. Dry matter and total accumulated average in CHK: Charleston Korean; DM A: Dry Matter average; ADT: After day trasplanting.






Fig. 8. Dry matter and total accumulated average in CHT: Charleston Tunnel; DM A: Dry Matter average; ADT: After day trasplanting.

ADT	DMA (g/plant)			
	CAI	CHK	CHT	Average
35	48.69	30.77	30.77	36.74
50	76.81	60.47	66.66	67.98
65	116.16	108.98	98.76	107.96
80	184.80	153.06	150.00	162.62
96	260.32	201.80	272.77	244.96
R <sup>2</sup>	0.9655	0.9949	0.9095	0.9672
Rate of Increase of MS g/day/plant	3.5017	2.8607	3.7098	3.3699

Table 1. Dry matter and total accumulated average in two hybrid tomatoes. CAI: Caiman; CHK: Charleston grown in Korean greenhouse; CHT: Charleston in tunnel; DMA: Dry matter average (g/plant). ADT: after day trasplanting.

Figure 9 shows the estimated model with the average values in the three systems evaluated, if there were some mathematical models of nutrient removal, we could estimate the dose for each nutrient to apply, once it has the most of the nutrients, it is possible calculate a fertilization program for growing tomatoes, a further advantage of this procedure is that it could adjust fertilization with irrigation, if available, a pressurized irrigation system.




Fig. 9. Linear model obtained from total dry matter considering both hybrids.

## 5. Conclusion

Although the stage of harvest of tomato is not completed successfully because of environmental factors, It was possible to estimate dry matter production, the rate of increase in dry matter was higher and similar for CAI and CHT, but was lower for CHK. The average yield of the two hybrids was considered adequate to estimate dry matter production in tomato crop in this geographical area. Once we determined the removal of nutrients such as nitrogen, phosphorus, potassium, among others, and generate models of extraction of these, it can be calculated the amount of each nutrient applied and thus may establish a program of fertilization to ensure a sustainable production system; such program may be adjusted by several cycles of this same culture.

## 6. Acknowledgment


The authors would like to thank to M Sc. Gerardo Jiménez García† his contribution to accomplish this research.

## 7. References

- Bugarin, M. R.; Galvis, A.; Sanchez, P. y García, D. (2002). *Acumulación diaria de materia seca y de potasio en la biomasa aérea total de tomate*. Terra Latinoamericana. Vol.20, No.4, pp. 401-409, ISSN 0187-5779
- De Koning, A.N.M. (1989). *Development and growth of a commercially grown tomato crop*. Acta Hort. Vol.260, pp. 267-273, ISSN 0567-7572
- Galvis, S. A. (1998). *Diagnóstico y simulación del suministro edáfico para cultivos anuales*. Tesis Doctoral en Ciencias. Colegio de Postgraduados. Montecillo, Mexico.

- Gary, C.; Jones, J.W. & Tchamitchian, M. (1998). *Crop modelling in horticulture: State of the art*. Sci. Hort. Vol.74, pp. 3-20, ISSN 03044238
- Heuvelink, E. & Marcelis, L.F.M. (1989). *Dry matter distribution in tomato and cucumber*. Acta Hort. Vol.260, pp. 149-157. ISSN 0567-7572
- Honorato, R.; Gurovich, L. y Piña R. (1993). *Ritmo de absorción de N, P y K en pepino de semilla*. Cien. Inv. Agr. Vol.20, pp. 169-172 ISSN 0718-3267
- Imas, P. (1999). *Manejo de Nutrientes por Fertirriego en Sistemas Frutihortícolas*. XXII Congreso Argentino de Horticultura. Septiembre - Octubre, 1999. Argentina.
- Magnífico, V.; Lattancio, V. & Sarli, G. (1989). *Growth and nutrient removal by broccoli raab*. J. Amer. Soc. Hort. Sci. 104 (2): 201 - 203. ISSN (print): 0394-6169
- Miller, C.H.; Mccollum, R.E. & Claimon, S. (1979). *Relationships between growth of bell peppers (Capsicum annuum L.) and nutrient accumulation during ontogeny in field environments*. Jour. Amer. Soc. Hort. Sci. Vol.104, pp. 852 - 857. Print ISSN: 0003-1062
- Ortega B., R.; Correa-Benguria, M. y Olate M., E. (2005). *Determinación de las curvas de acumulación de nutrientes en tres cultivares de liliun spp. para flor de corte*. Agrociencia Vol.40, No.1, pp. 77-88, ISSN 1405-3195
- Rodríguez, J. (1990). *La fertilización de los cultivos: Un método racional*. Facultad de Agronomía. Pontificia Universidad Católica de Chile. Santiago de Chile. 406p.
- Rodríguez F., H. y Rodríguez Absi, J. (2011). *Métodos de Análisis de Suelos y Plantas. Criterios de Interpretación*. Editorial Trillas S.A. de C. V. (Segunda edición). México. ISBN 978-607-17-0593-8.
- Rodríguez F., H.; Muñoz, S. y Alcorta G., E. (2006). *El tomate rojo. Sistema hidropónico*. Editorial Trillas S.A. de C. V. (Primera edición). México. ISBN 968-24-7606-2
- Scaife, A. & Bar-Yosef, B. (1995). *Nutrient and fertilizer management in field grown vegetables*. IPI Bulletin No. 13. International Potash Institute, Basel, Switzerland).
- Steiner, A.A. (1961). *A universal method for preparing nutrient solutions of certain desired composition*. Plant and Soil. Vo.15, pp. 134-154, (print version) ISSN: 0032-079X
- Steta, M. (1999). *Status of the greenhouse industry in Mexico*. Acta Hort. Vol.481, pp. 735-738. ISSN 0567-7572
- Sigmaplot 8.0. (2003). *Users Manual*. Exact Graphs for Exact Science. Systat Software Inc, USA.
- Zaidan, O. & Avidan, A. (1997). *Greenhouses tomatoes in soilless culture*. Ministry of Agriculture, Extension Service, Vegetables and Field Service Departments, USA.





## **Horticulture**

Edited by Prof. Alejandro Isabel Luna Maldonado

ISBN 978-953-51-0252-6

Hard cover, 172 pages

**Publisher** InTech

**Published online** 16, March, 2012

**Published in print edition** March, 2012

This book is about the novel aspects and future trends of the horticulture. The topics covered by this book are the effect of the climate and soil characteristics on the nitrogen balance, influence of fertilizers with prolongation effect, diversity in grapevine gene pools, growth and nutrient uptake for tomato plants, post-harvest quality, chemical composition and antioxidant activity, local botanical knowledge and agrobiodiversity, urban horticulture, use of the humectant agents in protected horticulture as well as post-harvest technologies of fresh horticulture produce. This book is a general reference work for students, professional horticulturalists and readers with interest in the subject.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Humberto Rodriguez-Fuentes, Juan Antonio Vidales-Contreras, Alejandro Isabel Luna-Maldonado and Juan Carlos Rodriguez-Orti (2012). Total Growth of Tomato Hybrids Under Greenhouse Conditions, Horticulture, Prof. Alejandro Isabel Luna Maldonado (Ed.), ISBN: 978-953-51-0252-6, InTech, Available from: <http://www.intechopen.com/books/horticulture/growth-and-nutrient-uptake-for-two-tomato-hybrids-under-commercial-greenhouse-conditions>

**INTECH**  
open science | open minds

#### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

#### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen