

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Phytochemicals and Their Pharmacological Aspects of *Acanthopanax koreanum*

Young Ho Kim, Jeong Ah Kim and Nguyen Xuan Nhiem
Chungnam National University,
South Korea

1. Introduction

Botanical medicines have been applied for the treatment of various human diseases with thousands of years of history all over the world. In some Asian and African countries, 80 % of population depends on traditional medicine in primary health care. On the other hand, in many developed countries, 70 % to 80 % of the population has used some forms of alternative or complementary medicine. The long tradition of using plants for medicine, supplemented by pharmaceutical research, has resulted in many plant-based Western medicines. Traditional medicine has provided Western medicine with over 40 % of all pharmaceuticals (Samuelsson & Bohlin, 2004). In the past decades, therefore, research has been focused on scientific evaluation of traditional drugs of plant origin.

Acanthopanax species (Araliaceae) are widely distributed in Asia, Malaysia, Polynesia, Europe, North Africa and the America. There are about 40 species of *Acanthopanax* to be found in over the world. *Acanthopanax* species have traditionally been used as a tonic and sedative as well as in the treatment of rheumatism, and diabetes. *A. koreanum* Nakai is an indigenous plant prevalently distributed throughout South Korea. It is deciduous shrub with upright to slightly arching stems, small, fresh green, trilobed to palmately divided leaves and several axillary as well as terminal round clusters of decorative, bluish black berries in late summer and autumn. Extensive investigation of chemical components in *A. koreanum* has been reported by many researchers in the worldwide. Several types of compounds have been isolated from this plant. Major active constituents are reported as lupanes and their glycosides, diterpenes, monoterpenes, lignans, phenylpropanoids, flavonoids from whole parts of *A. koreanum*. Of these, lupane triterpenes were reported as major components of leaves and *ent*-kauranes are main components of the roots of *A. koreanum*. They showed significant biological effects by several bioassay systems such as 1) anti-inflammatory activities: inhibit lipopolysaccharide (LPS)-stimulated TNF- α , IL-6, and IL-12 p40 productions in bone marrow-derived dendritic cells, decrease the expression of inducible nitric oxide synthase (iNOS) and cyclooxygenase-2 (COX-2) proteins, and reduce iNOS and COX-2 mRNA in a dose-dependent pattern, 2) anticancer, and 3) anti-osteoporosis by effects on the differentiation of osteoblastic MC3T3-E1 cells. The desired target of this chapter is to introduce explanations of structures and pharmacological activities of novel compounds, which have been isolated and identified from *A. koreanum* since 1985. Those studies have reported and focused on bioactivities of unambiguous

compounds from *A. koreanum*, therefore we discuss new pharmacological findings on these compounds.

The depth and breadth of research involving this plant has been organized into easily accessible and comparable information. Using Chemical Abstracts, Scifinder Scholar, and BIOSIS databases, relevant research papers were selected by based on pertinence and specificity to ethnopharmacology and phytochemistry, as well as readability. This collection was then carefully reviewed, extracted, and corroborated with available characterization data from other sources.


2. Phytochemistry and pharmacology of *A. koreanum*

2.1 Lupane aglycones

Impressic acid (**1**) was isolated for the first time from *Schefflera impressa* by (Srivastava, 1992) and it was found in the roots (Cai et al., 2004b) and the leaves (Kim et al., 2010) of *A. koreanum*. Impressic acid exhibited potently NFAT inhibitory activity with IC₅₀ value of 12.6 μM. In the studies of (Kim et al., 2010), impressic acid (**1**) and (20*R*)-3α-hydroxy-29-dimethoxylupan-23,28-dioic acid (**4**) showed significantly anti-inflammatory activities by inhibiting TNF-α, IL-6, and IL-12 p40 productions in bone marrow-derived dendritic cells with LPS-stimulated. Furthermore, impressic acid was found to inhibit TNF-α-induced NF-κB activity by inhibiting the induction of COX-2 and iNOS in HepG2 cells. Impressic acid also up-regulated the transcriptional activity of PPAR by elevating the expression of PPARγ1, PPARγ2, and SREBF-2, and by suppressing the expression of Insig-2 (Kim et al., 2011). One new norlupane, 3α,11α-dihydroxy-20,23-dioxo-30-norlupane-28-oic acid (**2**) as well as two known lupane aglycones, impressic acid (**1**), 3α,11α-dihydroxy-lup-20(29)-en-23-al-28-oic acid (**3**) were isolated and determined by (Park et al., 2010). They were evaluated for the differentiation of osteoblastic MC3T3-E1 cells. Among of them, compound **1** significantly increased osteoblastic cell growth and differentiation as assessed by MTT assay and collagen content. Compound **2** significantly increased the growth of MC3T3-E1 cells and caused a significant elevation of osteoblastic cell differentiation as assessed by the alkaline phosphatase activity (Park et al., 2010). Other compounds, 3α,11α-dihydroxy-lup-20(29)-en-23-al-28-oic acid, 3α-hydroxylup-20(29)-en-23,28-dioic acid (**5**), and 3α,11α,23-trihydroxy-lup-20(29)-en-28-oic acid (**6**) were also isolated from steamed leaves (Kim et al., 2010). However, they showed weak anti-inflammatory activity. 3α-Hydroxylup-20(29)-en-23,28-dioic acid (**5**) possessed broader antiviral activity against respiratory syncytial, influenza (H1N1), coxsackie B3, and herpes simplex virus type 1 viruses with IC₅₀ values of 6.2, 25.0, 12.5, and 18.8 μg/mL, respectively (Li et al., 2007).

2.2 Lupane-triterpene glycosides

Up to date, eighteen lupane-type triterpene glycosides have been isolated from this plant and almost of them from the leaves of *A. koreanum*. They are main saponin components of the leaves of *A. koreanum*. The first lupane triterpene glycoside, acantrifoside A (**1**) was isolated from both *A. koreanum* and *A. trifoliatus* in a year of 1998 by (Yook et al., 1998). And then, two new saponins, acankoreoside A (**10**) and acankoreoside B (**11**) were isolated from the leaves of this plant (Chang et al., 1998). Our group reported seven new lupane-type triterpene glycosides, acankoreosides I-O. Their biological activities were evaluated for



Name	Parts	R ₁	R ₂	R ₃	Reference
Impressic acid (1)	leaves roots	CH ₃	OH	CH ₂	(Cai et al., 2004b)
3 α ,11 α -Dihydroxy-20,23-dioxo-30-norlupane-28-oic acid (2)	leaves	CHO	OH	=O	(Park et al., 2010)
3 α ,11 α -Dihydroxy-lup-20(29)-en-23-al-28-oic acid (3)	leaves	CHO	OH	CH ₂	(Park et al., 2010)
(20R)-3 α -Hydroxy-29-dimethoxylupane-23,28-dioic acid (4)	steamed leaves	COOH	H	CH(OCH ₃) ₂	(Kim et al., 2010)
3 α -Hydroxylup-20(29)-en-23,28-steamed dioic acid (5)	leaves	COOH	H	CH ₂	(Kim et al., 2010)
3 α ,11 α ,23-Trihydroxy-lup-20(29)-en-28-oic acid (6)	steamed leaves	CH ₂ OH	OH	CH ₂	(Kim et al., 2010)

Fig. 1. Structures of main lupane-type triterpenes isolated from *Acanthopanax koreanum*






Fig. 2. 3-O- β -D-glucopyranosyl 3 α ,11 α -dihydroxylup-20(29)-en-28-oic acid (7) and 3 α -hydroxylup-20(29)-en-23,28-oic acid 28-O-[β -D-Glucopyranosyl-(1 \rightarrow 6)- β -D-Glucopyranosyl] ester (8) (Kim et al., 2010)

cytotoxic activities including A549 (lung), HL-60 (acute promyelocytic leukemia), MCF-7 (breast), U937 (leukemia) cancer cell lines; immune enhancement activity (INF- γ and IL-2 release in spleen cells); anti-inflammatory (inhibitory TNF- α , IL-6, and IL-12 p40 productions in bone marrow-derived dendritic cells with LPS-stimulated, and RAW 264.7). Searching for anticancer activities from natural compounds, several acankoreosides showed significantly cytotoxic activities in various cancer cell lines (A549, HL-60, MCF-7, and U937). The effects of three new lupane glycosides, acankoreosides F-H (**13**, **14**, and **21**) on the LPS-induced production of nitric oxide and prostaglandin E₂ were evaluated in RAW 264.7 macrophages. Among of them, acankoreoside F (**13**) showed the most potent inhibitory PGE₂ (59.0 %) and NO (42.0 %) production at concentration of 200.0 μ M. Furthermore, eleven lupane triterpene glycosides from *A. koreanum*, including three new compounds acankoreoside M-O (**16**, **24**, and **25**) were evaluated for Con A-induced splenolytic production of IL-2 and IFN- γ . The results indicated that acankoreosides A (**10**), D (**12**), L (**24**), and acantrifoside A (**9**)




Names	Parts	R ₁	R ₂	R ₃	Ref.
Acantrifoside A (9)	leaves	CH ₃	OH	H	(Yook et al., 1998)
Acankoreoside A (10)	leaves, roots	COOH	H	H	(Chang et al., 1998) (Cai et al., 2004b)
Acankoreoside B (11)	leaves	CH ₂ OH	OH	H	(Chang et al., 1998)
Acankoreoside D (12)	leaves	CHO	OH	H	(Chang et al., 1999)
Acankoreoside F (13)	leaves	COOH	H	OH	(Choi et al., 2008)
Acankoreoside G (14)	leaves	CHO	H	OH	(Choi et al., 2008)
Acankoreoside I (15)	leaves	CHO	OH	OH	(Nhiem et al., 2009)
Acankoreoside M (16)	leaves	COOH	OH	OH	(Nhiem et al., 2010b)
3-Epibetulinic acid 28-O-glc-glc-rha (17)	leaves, roots	CH ₃	H	H	(Cai et al., 2004b)

Fig. 3. Structures of lupane-type triterpene glycosides from *A. koreanum*



Name	Part	R	Reference
Acankoreoside C (18)	leaves	H	(Chang et al., 1999)
Acankoreoside N (19)	leaves	OH	(Nhiem et al., 2010b)




Name	Part	R ₁	R ₂	R ₃	R ₄	Reference
Acankoreoside E (20)	leaves	COOH	H	H	CHO	(Park et al., 2005)
Acankoreoside H (21)	leaves	CHO	H	H	COOH	(Choi et al., 2008)
Acankoreoside J (22)	leaves	COOH	H	=O	-	(Nhiem et al., 2010a)
Acankoreoside K (23)	leaves	COOH	H	OH	Me	(Nhiem et al., 2010a)
Acankoreoside L (24)	leaves	COOH	H	OH	CH ₂ OH	(Nhiem et al., 2010a)
Acankoreoside O (25)	leaves	COOH	OH	H	CH ₃	(Nhiem et al., 2010b)

Fig. 3. Structures of lupane-type triterpene glycosides from *A. koreanum* (continued)

significantly increased both IL-2 and IFN- γ . The structure-activity relationship of these compounds was also discussed. Moreover, lupane aglycones and lupane glycosides were assayed for LPS-stimulated pro-inflammatory cytokine production. These results suggested lupane aglycone inhibited pro-inflammatory cytokine production stronger than lupane glycosides (Kim et al., 2010). This was further confirmed by the study of (Cai et al., 2004b).

2.3 Pimarane-type diterpenes


A number of pimarane-type diterpenes have been isolated and associated with significant biological activity. There are seven pimarane-type diterpenes from *A. koreanum*. All of them were isolated from the roots. Acanthoic acid was presented in roots and leaves of this plant, and was one of compounds having potent anti-inflammatory activity. Acanthoic acid, a pimarane diterpene ((-)-pimara-9(11),15-dien-19-oic acid), was isolated for the first time from *A. koreanum* in a year of 1988 by (Kim et al., 1988b) and was proved with high content of this plant. Acanthoic acid has widely exhibited of biological activities. In study of (Kang et al., 1996), acanthoic acid has potent anti-inflammatory effects by reducing the production of proinflammatory cytokines such as IL-1 and TNF- α . It was also effective in suppressing experimental silicosis and cirrhosis. Furthermore, acanthoic acid was found to suppress TNF- α gene expression (Kang et al., 1998) and TNF- α -induced IL-8 production in a dose-dependent manner. Acanthoic acid also inhibited TNF- α -induced MAPKs activation, I κ B degradation, NF- κ B nuclear translocation, and NF- κ B/DNA binding activity (Kim et al., 2004). Furthermore, acanthoic acid significantly inhibited production of both TNF- α and tryptase in trypsin-stimulated human leukemic mast cell-1 at concentrations of 10 and 100 μ g/mL with a dose-dependent manner. Acanthoic acid inhibited ERK phosphorylation and NF- κ B activation induced by trypsin treatment without blocking of trypsin activity even though 100 μ g/mL. These results suggested that acanthoic acid may inhibit the production of inflammatory mediators through inhibition of ERK phosphorylation and NF- κ B activation pathway in human mast cells (Kang et al., 2006).




Name	Part	R	Reference
(-)-Pimara-9(11),15-dien-19-ol (26)	root barks	CH ₂ OH	(Kim et al., 1988b)
Acanthoic acid (27)	roots, leaves	COOH	(Kim et al., 1988b)
(-)-Pimara-9(11),15-dien-19-ol 19-acetate (28)	root barks	CH ₂ OAc	(Kim et al., 1988b)
(-)-Pimara-9(11),15-diene (29)	root barks	CH ₃	(Kim et al., 1988b)

Fig. 4. Structures of pimarane-type diterpenes from *A. koreanum*

The hepatoprotective effects of acanthoic acid were evaluated in a D-galactosamine/lipopolysaccharide-induced fulminant hepatic failure mouse model. The effects were likely associated with a significant decrease in serum TNF- α levels, which are correlated not only with those of alanine aminotransferase and aspartate aminotransferase but also with the reduced number of apoptotic hepatocytes in the liver as confirmed using the terminal deoxynucleotidyl transferase-mediated (dUTP) nick end-labeling method and DNA fragmentation assay (Nan et al., 2008). The protective effect of acanthoic acid was investigated in acetaminophen-induced hepatic toxicity. These results indicated that acanthoic acid protected liver tissue from oxidative stress elicits by acetaminophen-induced liver damage (Wu et al., 2010). Acanthoic acid markedly suppressed the protein expression of TNF- α , COX-2, NF- κ B and chymase as well as the mRNA expression of TNF- α and COX-2 (Kang et al., 2010).



Isopimara-9(11),15-dien-19-ol (29)	roots	(Chung & Kim, 1986)
Acanthokoreoic acid A (30)	roots	(Cai et al., 2003a)




7 β -Hydroxy-ent-pimara-8(14),15-dien-19-oic acid (31)	roots	(Cai et al., 2004a)
Sumogaside (32)	roots	(Cai et al., 2004a)

Fig. 4. Structures of pimarane-type diterpenes from *A. koreanum* (continued)

In study of (Cai et al., 2003a), a new compound, acanthokoreoside acid A (30) as well as acanthoic acid (27), (-)-pimara-9(11),15-dien-19-ol (26), and sumogaside (32) were isolated from CH₂Cl₂ fraction of *A. koreanum* roots. They were evaluated for inhibitory activity on IL-8 secretion in TNF- α -stimulated HT-29 and TNF- α secretion in trypsin-stimulated HMC-1. In the TNF- α -stimulated HT-29, acanthoic acid and sumogaside significantly inhibited IL-8 secretion at concentrations of 1, 10, and 100 μ M and at concentrations of 10 and 100 μ M, respectively.

2.4 *ent*-Kaurane-type diterpenes

ent-Kaurane, a tetracyclic diterpene, has been proven to exert various biological activities like cytotoxicity, anti-inflammation, and so on. From the roots of *A. koreanum*, (Kim et al., 1988b) and (Cai et al., 2003b) isolated six *ent*-kaurane-type diterpenes, including *ent*-kaur-16-en-19-oic acid (**33**), 16 α -hydroxy-*ent*-kauran-19-oic acid (**34**), paniculose IV (**35**), 16 α H,17-isovaleryloxy-*ent*-kauran-19-oic acid (**36**), 16 α -hydroxy-17-isovaleryloxy-*ent*-kauran-19-oic acid (**37**), and 16 α ,17-dihydroxy-*ent*-kauran-19-oic acid (**38**). (Cai et al., 2003b) evaluated five *ent*-kauranes for TNF- α secretion from HMC-1, a trypsin-stimulated human leukemic mast cell line. The results indicated that 16 α H,17-isovaleryloxy-*ent*-kauran-19-oic acid (**36**) exhibited potently an inhibitory activity with IC₅₀ value of 16.2 μ M. Furthermore, these compounds were assayed for their inhibitory effect against NFAR transcription factor and 16 α -hydroxy-17-isovaleryloxy-*ent*-kauran-19-oic acid (**37**) was found to significantly inhibit NFAT transcription factor with IC₅₀ of 6.7 μ M (Cai et al., 2004a). The authors also found that remain compounds containing a hydroxyl group at C-16 or a glycoside at C-4 showed no activity.



Name	Part	Reference
<i>ent</i> -Kaur-16-en-19-oic acid (33)	roots	(Kim et al., 1988b)
16 α -Hydroxy- <i>ent</i> -kauran-19-oic acid (34)	roots	(Cai et al., 2003b)
Paniculose IV (35)	roots	(Cai et al., 2003b)
16 α H,17-Isovaleryloxy- <i>ent</i> -kauran-19-oic acid (36)	roots	(Cai et al., 2003b)
16 α -Hydroxy-17-isovaleryloxy- <i>ent</i> -kauran-19-oic acid (37)	roots	(Cai et al., 2003b)
16 α ,17-Dihydroxy- <i>ent</i> -kauran-19-oic acid (38)	roots	(Kim et al., 1988b)

Fig. 5. Structures of *ent*-kaurane-type diterpenes from *A. koreanum*

2.5 Other class compounds

Two lignans were found from the roots of *A. koreanum*. Those were acanthoside D (**39**) (Hahn et al., 1985) and ariensin (**40**) (Kim et al., 1988a). Beside these lignans, the first




Fig. 6. Structures of compounds isolated from *A. Koreanum*

Pos.	1	2	3	4	5	6	7	8	9	10	11	12
1	35.8	35.7	35.1	32.8	32.6	35.9	36.2	33.1	36.2	33.1	35.9	34.9
2	26.8	26.9	25.5	26.0	25.9	27.1	21.9	26.2	26.9	26.2	27.1	26.8
3	74.8	73.7	75.8	72.8	72.7	75.9	81.5	71.5	75.3	73.0	75.7	72.7
4	38.4	53.5	37.7	51.8	51.7	41.2	37.9	51.9	38.5	52.8	41.1	52.6
5	49.4	44.4	48.8	44.7	44.8	43.9	50.6	45.3	49.6	45.0	43.8	43.9
6	19.2	21.8	19.3	21.6	21.0	18.3	18.4	21.7	18.6	21.8	18.3	21.0
7	36.0	35.7	35.4	23.8	34.5	35.6	35.7	34.6	35.7	34.6	35.4	35.2
8	42.7	43.2	42.2	41.6	41.5	42.8	42.7	41.8	42.8	42.8	42.7	42.4
9	55.9	56.2	55.6	50.7	50.8	56.2	55.9	51.0	56.2	51.2	55.6	55.6
10	39.8	39.6	39.1	38.2	37.2	39.6	39.7	37.4	39.9	37.6	39.6	38.7
11	69.8	70.7	70.4	21.1	21.5	69.9	69.9	20.9	69.8	21.2	69.8	69.4
12	38.4	39.4	37.7	27.7	25.8	38.4	38.3	26.0	38.3	26.0	38.3	37.8
13	37.5	37.5	37.2	28.6	38.4	37.6	37.7	38.4	37.4	38.4	37.4	37.0
14	42.9	44.1	42.6	43.0	42.7	42.8	42.9	42.9	43.0	43.0	42.9	43.0
15	29.9	30.6	29.5	30.2	30.0	30.1	30.2	30.1	30.0	30.1	30.0	29.6
16	32.8	32.7	32.0	34.7	32.8	32.8	32.9	32.2	32.3	32.4	32.2	31.9
17	56.5	57.1	56.2	56.7	56.4	56.5	56.6	57.0	56.9	57.7	56.9	56.6
18	47.5	50.0	46.6	49.1	49.5	49.4	49.4	49.8	49.5	49.7	49.4	49.1
19	49.4	52.5	48.6	43.7	47.5	47.5	47.5	47.4	47.2	47.5	47.1	46.8
20	151.0	215.0	149.7	37.6	151.1	150.8	150.9	150.9	150.4	151.0	150.4	150.1
21	31.0	29.4	30.5	24.6	30.9	31.2	31.3	30.8	30.9	30.9	30.9	30.6
22	37.5	37.7	36.8	37.2	37.3	37.4	37.4	36.9	36.8	37.2	36.7	36.4
23	29.6	210.9	28.7	178.9	179.6	71.9	29.9	181.3	29.8	179.0	71.9	209.7
24	22.7	15.1	22.2	17.8	17.7	18.3	23.0	18.1	22.9	18.0	18.3	14.6
25	18.0	17.2	17.3	16.6	16.5	17.0	16.8	16.8	16.9	16.8	17.1	16.5
26	17.4	17.8	16.2	16.6	16.5	17.7	17.6	16.7	17.7	16.6	17.7	17.4
27	14.4	15.0	14.6	14.5	14.6	14.8	14.8	14.8	14.8	14.8	14.8	14.4
28	179.2	179.4	181.5	179.4	178.6	178.8	178.9	175.0	175.0	175.9	175.0	174.6
29	110.0	30.0	110.2	107.8	109.7	110.0	110.1	110.0	110.2	110.0	110.0	109.9
30	19.2		17.9	16.4	19.2	19.5	19.6	19.4	19.5	19.3	19.5	19.2
Solv.	<i>a</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>a</i>	<i>a</i>	<i>a</i>	<i>a</i>	<i>a</i>	<i>a</i>	<i>a</i>	<i>a</i>

^a recorded in pyridine-*d*₅, ^b recorded in methanol-*d*₄.

Note: NMR data were obtained from **1**: (Srivastava, 1992); **2, 3** (Park et al., 2010); **4, 8**: (Kim et al., 2010); **5, 6, 10, 11**: (Chang et al., 1998); **7, 12**: (Chang et al., 1999)

Table 1. ¹³C-NMR data of lupane aglycone moieties

Pos.	13	14	15	16	17	18	19	20	21	22	23	24	25
1	33.2	33.1	35.9	35.5	36.5	36.1	36.7	33.3	33.1	34.0	34.2	33.9	34.5
2	26.1	26.7	27.0	26.4	26.6	21.8	19.2	26.1	26.7	26.1	26.1	26.1	26.1
3	72.8	73.0	73.8	73.8	73.5	81.3	82.4	72.7	73.0	73.4	73.3	73.6	73.3
4	51.8	52.5	53.7	53.0	40.3	37.8	38.5	51.7	52.5	52.2	52.1	52.3	52.1
5	45.4	44.0	44.5	45.4	49.9	50.5	51.0	45.6	44.0	46.6	46.9	46.2	47.4
6	21.7	20.9	22.0	22.6	19.2	18.4	22.2	21.7	21.1	22.2	22.2	22.3	22.2
7	34.5	34.1	36.2	36.1	36.3	35.4	36.4	34.6	34.1	35.1	35.5	35.3	35.5
8	41.8	41.8	41.6	43.5	43.4	42.6	43.5	41.7	41.8	42.5	43.0	42.6	43.1
9	51.0	50.6	56.3	56.7	50.2	55.8	56.3	50.6	50.2	51.9	52.0	51.7	52.0
10	37.4	36.9	38.2	40.0	38.3	39.6	40.4	37.4	36.9	38.2	38.1	38.1	38.2
11	21.0	21.0	71.1	71.1	24.2	69.7	71.2	20.9	20.8	22.0	22.4	22.0	22.2
12	27.1	27.0	38.3	39.4	26.5	38.1	39.5	26.9	26.9	28.5	29.8	28.3	26.1
13	38.3	38.3	39.5	38.2	38.2	37.3	38.3	38.2	38.2	38.6	39.5	39.4	39.1
14	42.8	42.8	44.3	44.3	43.8	42.9	44.0	43.0	43.0	43.6	44.5	44.0	44.3
15	30.2	30.1	30.8	30.8	31.6	30.0	30.9	30.0	30.0	30.9	31.2	30.9	30.9
16	32.2	32.1	32.8	32.9	32.8	32.2	32.9	32.0	32.1	32.4	33.1	32.9	34.5
17	57.0	56.9	57.9	57.9	57.9	56.8	58.0	57.0	57.3	58.0	60.1	58.0	59.4
18	50.2	50.2	50.9	50.9	50.1	49.4	51.0	48.5	48.9	50.8	49.5	49.7	50.4
19	43.2	43.2	43.7	43.7	48.2	47.1	43.8	37.3	40.6	52.8	45.7	39.3	85.4
20	156.5	156.5	155.9	155.9	151.3	150.4	156.0	50.1	42.1	215.4	76.1	39.5	36.8
21	32.7	32.7	33.5	33.6	30.6	30.8	33.6	24.6	25.0	29.3	28.8	24.38	34.5
22	36.8	36.7	37.3	37.4	37.5	36.7	37.4	37.4	37.3	37.6	37.5	38.3	37.2
23	181.2	209.8	211.2	180.6	29.5	29.8	29.8	181.8	209.9	183.4	184.2	181.8	185.2
24	18.2	14.6	15.2	18.2	23.0	23.0	23.3	18.3	14.6	18.3	18.4	18.2	18.6
25	16.8	16.4	17.2	17.5	17.8	16.8	17.3	16.8	16.3	17.2	17.4	17.2	17.4
26	16.7	16.5	18.3	18.1	17.0	17.6	18.0	16.6	16.5	16.9	17.2	17.0	17.1
27	14.9	14.8	15.1	15.1	14.9	14.7	15.2	14.7	14.9	15.2	15.6	15.1	15.5
28	175.1	175.0	176.3	176.3	176.2	174.8	176.4	175.0	174.9	176.2	177.0	176.4	176.7
29	106.1	106.1	107.7	107.7	110.8	110.1	107.6	204.6	180.0	29.9	71.2	8.9	21.9
30	64.3	64.3	65.5	65.5	19.6	19.6	65.6	7.0	10.0		19.6	110.3	27.0
Solv.	<i>a</i>	<i>a</i>	<i>b</i>	<i>b</i>	<i>b</i>	<i>a</i>	<i>b</i>	<i>a</i>	<i>a</i>	<i>b</i>	<i>b</i>	<i>b</i>	<i>b</i>

^a recorded in pyridine-*d*₅, ^b recorded in methanol-*d*₄.

Note: NMR data were obtained from **13**, **14**, **21**: (Choi et al., 2008); **15**: (Nhiem et al., 2009) (Cai et al., 2004b), **16**, **19**, **25**: (Nhiem et al., 2010b); **17**: (Cai et al., 2004b); **18**: (Chang et al., 1999), **20**: (Park et al., 2005); **21**: (Choi et al., 2008); **22**, **23**, **24**: (Nhiem et al., 2010a).

Table 1. ¹³C-NMR data of lupane aglycone moieties (continued)

phenylpropanoid, syringoside (**44**) were isolated from the roots of *A. koreanum* by (Hahn et al., 1985) and then was ariensin (**43**) (Kim et al., 1988a). In study antioxidant activity of chemical components from the leaves of this plant, (Nhiem et al., 2011) isolated one new phenylpropanoid named acanthopanic acid and one known 1,2-O-dicaffeoylcyclopenta-3-ol. These compounds showed significantly antioxidant activity by the intracellular ROS radical scavenging DCF-DA assay with IC₅₀ values of 3.8 and 2.9 μM, respectively. Until now, only rutin (**45**), a quercetin glycoside was isolated from this plant with large amount. Rutin is used in many countries as medication for blood vessel protection and are ingredients of numerous multivitamin preparations and herbal remedies. Rutin has various biological activities that are beneficial to human health such as antioxidant effect (Nhiem et al., 2011), protective effect against hepatotoxicity, and anti-inflammatory effect. On the other hand, from the leaves of *A. koreanum*, two monoterpenoids, 4S)-α-terpineol O-β-D glucopyranoside (**46**) (Nhiem et al., 2011) and betulabuside B (**47**) (Park et al., 2010) were isolated. From fruits, citric, maleic succinic, malonic, fumaric, and malic acid were isolated (Shin & Kim, 1985).

3. NMR data of lupane aglycones

Lupane triterpenes are a class of the most compounds isolated from the leaves and roots of *A. koreanum*, which were determined that this type of compounds are main chemical components of this plant.

Structure of lupanes were elucidated with ¹H-NMR, ¹³C-NMR, DEPT (distortionless enhancement by polarization transfer), COSY (¹H-¹H shift correlation spectroscopy), TOCSY (total correlation spectroscopy), HMBC (heteronuclear multiple bond correlation), HSQC (heteronuclear single quantum coherence), NOESY (nuclear overhauser enhancement spectroscopy, and ROESY (rotating frame overhauser effect spectroscopy). Proton coupling networks of sugar moieties were indicated with ¹H-NMR, COSY, HMBC and HSQC. Herein, we suggest statistical results of ¹³C-NMR data of lupane-type triterpene aglycones and their derivatives in comparison with data of references (Table 1).

Observed the isolated compounds from *A. koreanum*, we found that there are four main classes including lupane triterpenoids, pimarane diterpenoids, *ent*-kaurane diterpenoids, and lignans. Among of them, lupane triterpenes were isolated as numerous of compounds with high yield. These lupanes often contain hydroxyl group at C-3, carboxyl at C-28. In some compounds, hydroxyl, aldehydic, carboxylic groups were at C-11, C-23, and C-30, glycoside was at C-28 and rarely at C-3.

From Table 1, we summarized all ¹³C-NMR characteristics of lupane aglycones as follows:

1. When hydroxyl group at C-3, chemical shift of C-3 was about 73.0 ppm and configuration of hydroxyl group at C-3 is α orientation. When glycosidation is at C-3, chemical shift of C-3 moved to down field with δ_C of 81.0 ppm.
2. Free carboxylic group at C-28 were confirmed by chemical shift about 178.0~180.0 ppm. When sugar moiety was at C-28, chemical shift of C-28 is 174.6~176.3 ppm, decreased about 2.5-3.8 ppm.

3. When 23-methyl group was replaced with aldehydic group, chemical shifts of C-23 and C-4 moved to down field from 28.0-28.8 to 209.0-210.0, 37.5-39.5 to 54.9-56.3 ppm, respectively. When 23-methyl group was replaced with carboxylic group, chemical shifts of C-23 and C-4 changed from 28.5 to 178.0, 37.8 to 53.0 respectively, and when 23-methyl group was replaced with CH₂OH, chemical shift of C-23 had a large change from 28.0 to 71.5 ppm; chemical shift of C-4 had small change about 2.0 ppm.
4. When 30-methyl group was replaced with CH₂OH, the chemical shifts of C-20 and C-30 downshifted from 151.0 to 156.5, from 19.5 to 64.5 ppm, respectively; chemical shifts of C-19 and C-29 upshifted from 47.5 to 43.0, from 110.0 to 106.0 ppm, respectively.
5. When hydroxyl group was at C-11, chemical shift of C-11 downshifted from 21.1 to 71.0 ppm. Furthermore, configuration of hydroxyl group at this position is α .

4. Conclusion

This chapter is intended to serve as a reference tool for people in all fields of ethnopharmacology and natural products chemistry. The pharmacological studies on *A. koreanum* indicated the immense potential possibility of this plant in the treatment of conditions such as inflammation, rheumatism, diabetes, cardiovascular, and virus. However, the diverse pharmacological activities of solvent extracts and phytochemicals of *A. koreanum* have only been tested in *in vitro* assay using laboratory animals, and obtained too unclearly and ambiguously for the case of human beings to be conducted on enough. However, these gaps in the studies demand to be bridged in order to exploit medicinal potential of the entire plant of *A. koreanum*. It is still clear that *A. koreanum* is massively and widely consumed, and also continuously studied expecting clinical treatment of various diseases for the future in Korea as well as in the world. From these viewpoints, imperic acid and acanthoic acid, major components of *A. koreanum* are good candidates for further studies in clinical trials, and the development of products derived from *A. koreanum* can be an important part of our biodiversity to respect and sustain for coming generation.

5. References

- Fact sheet N° 134: Traditional medicine. World Health Organization. December 2008.
- Cai, X. F.; Shen, G.; Dat, N. T.; Kang, O. H.; Kim, J. A.; Lee, Y. M.; Lee, J. J., & Kim, Y. H. (2003a). Inhibitory effect of TNF- α and IL-8 secretion by pimarane-type diterpenoids from *Acanthopanax koreanum*. *Chemical & Pharmaceutical Bulletin*, Vol.51, No.5, pp. 605-607, ISSN 0009-2363
- Cai, X. F.; Shen, G.; Dat, N. T.; Kang, O. H.; Lee, Y. M.; Lee, J. J., & Kim, Y. H. (2003b). Inhibitory effect of kaurane type diterpenoids from *Acanthopanax koreanum* on TNF- α secretion from trypsin-stimulated HMC-1 cells. *Archives of Pharmacal Research*, Vol.26, No.9, pp. 731-734, ISSN 0253-6269
- Cai, X. F.; Lee, I. S.; Dat, N. T.; Shen, G., & Kim, Y. H. (2004a). Diterpenoids with inhibitory activity against NFAT transcription factor from *Acanthopanax koreanum*. *Phytotherapy Research*, Vol.18, No.8, pp. 677-680, ISSN 0951-418X
- Cai, X. F.; Lee, I. S.; Shen, G.; Dat, N. T.; Lee, J. J., & Kim, Y. H. (2004b). Triterpenoids from *Acanthopanax koreanum* root and their inhibitory activities on NFAT transcription. *Archives of Pharmacal Research*, Vol.27, No.8, pp. 825-828, ISSN 0253-6269


- Chang, S. Y.; Yook, C. S., & Nohara, T. (1998). Two new lupane-triterpene glycosides from leaves of *Acanthopanax koreanum*. *Chemical & Pharmaceutical Bulletin*, Vol.46, No.1, pp. 163-165, ISSN 0009-2363
- Chang, S. Y.; Yook, C. S., & Nohara, T. (1999). Lupane-triterpene glycosides from leaves of *Acanthopanax koreanum*. *Phytochemistry*, Vol.50, No.8, pp. 1369-1374, ISSN 0031-9422
- Choi, H. S.; Kim, H. J.; Nam, S. G.; Kim, I. S.; Lee, K. T.; Yook, C. S., & Lee, Y. S. (2008). Lupane glycosides from the leaves of *Acanthopanax koreanum*. *Chemical & Pharmaceutical Bulletin*, Vol.56, No.11, pp. 1613-1616, ISSN 0009-2363
- Chung, B. S. & Kim, Y. H. (1986). Studies on the constituents of *Acanthopanax koreanum*. *Saengyak Hakhoechi*, Vol.17, No.1, pp. 62-66, ISSN 0253-3073
- Hahn, D.-R.; Kim, C. J., & Kim, J. H. (1985). A study on the chemical constituents of *Acanthopanax koreanum* Nakai and its pharmacological activities. *Yakhak Hoechi* Vol.29, No.6, pp. 357-361, ISSN 0513-4234
- Kang, H. S.; Kim, Y. H.; Lee, C. S.; Lee, J. J.; Choi, I., & Pyun, K. H. (1996). Suppression of interleukin-1 and tumor necrosis factor- α . production by acanthoic acid, (-)-pimara-9(11),15-dien-19-oic acid, and its antifibrotic effects in vivo. *Cellular Immunology*, Vol.170, No.2, pp. 212-221, ISSN 0008-8749
- Kang, H. S.; Song, H. K.; Lee, J. J.; Pyun, K. H., & Choi, I. (1998). Effects of acanthoic acid on TNF- α gene expression and haptoglobin synthesis. *Mediators of Inflammation*, Vol.7, No.4, pp. 257-259, ISSN 0962-9351
- Kang, O.-H.; Choi, Y. A.; Park, H. J.; Kang, C. S.; Song, B. S.; Choi, S. C.; Nah, Y. H.; Yun, K. J.; Cai, X. F.; Kim, Y. H.; Bae, K., & Lee, Y. M. (2006). Inhibition of trypsin-induced mast cell activation by acanthoic acid. *Journal of Ethnopharmacology*, Vol.105, No.3, pp. 326-331, ISSN 0378-8741
- Kang, O. H.; Kim, D. K.; Cai, X. F.; Kim, Y. H., & Lee, Y. M. (2010). Attenuation of experimental murine colitis by acanthoic acid from *Acanthopanax koreanum*. *Archives of Pharmacal Research*, Vol.33, No.1, pp. 87-93, ISSN 0253-6269
- Kim, J. A.; Kim, D. K.; Tae, J.; Kang, O. H.; Choi, Y. A.; Choi, S. C.; Kim, T. H.; Nah, Y. H.; Choi, S. J.; Kim, Y. H.; Bae, K. H., & Lee, Y. M. (2004). Acanthoic acid inhibits IL-8 production via MAPKs and NF- κ B in a TNF- α -stimulated human intestinal epithelial cell line. *Clinica Chimica Acta*, Vol. 342, No.1-2, pp. 193-202, ISSN 0009-8981
- Kim, J. A.; Yang, S. Y.; Koo, J.-E.; Koh, Y. S., & Kim, Y. H. (2010). Lupane-type triterpenoids from the steamed leaves of *Acanthopanax koreanum* and their inhibitory effects on the LPS-stimulated pro-inflammatory cytokine production in bone marrow-derived dendritic cells. *Bioorganic & Medicinal Chemistry Letters*, Vol.20, No.22, pp. 6703-6707, ISSN 0960-894X
- Kim, J. A.; Yang, S. Y.; Song, S. B., & Kim, Y. H. (2011). Effects of impressic acid from *Acanthopanax koreanum* on NF κ B and PPAR γ activities. *Archives of Pharmacal Research*, Vol.34, No.8, pp. 1347-1351, ISSN 0253-6269
- Kim, Y. H.; Chung, B. S.; Ko, Y. S., & Han, H. J. (1988a). Studies on the chemical constituents of *Acanthopanax koreanum*. (II). *Archives of Pharmacal Research*, Vol.11, No.2, pp. 159-162, ISSN 0253-6269

- Kim, Y. H.; Chung, B. S., & Sankawa, U. (1988b). Pimaradiene diterpenes from *Acanthopanax koreanum*. *Journal of Natural Products*, Vol.51, No.6, pp. 1080-1083, ISSN 163-3864
- Li, Y.; Jiang, R.; Ooi, L. S. M.; But, P. P. H., & Ooi, V. E. C. (2007). Antiviral triterpenoids from the medicinal plant *Schefflera heptaphylla*. *Phytotherapy Research*, Vol.21, No.5, pp. 466-470, ISSN 0951-418X
- Nan, J. X.; Jin, X. J.; Lian, L. H.; Cai, X. F.; Jiang, Y. Z.; Jin, H. R., & Lee, J. J. (2008). A diterpenoid acanthoic acid from *Acanthopanax koreanum* protects against D-galactosamine/lipopolysaccharide-induced fulminant hepatic failure in mice. *Biological & Pharmaceutical Bulletin*, Vol.31, No.4, pp. 738-742, ISSN 0918-6158
- Nhiem, N. X.; Tung, N. H.; Kiem, P. V.; Minh, C. V.; Ding, Y.; Hyun, J. H.; Kang, H. K., & Kim, Y. H. (2009). Lupane triterpene glycosides from leave of *Acanthopanax koreanum* and their cytotoxic activity. *Chemical & Pharmaceutical Bulletin*, Vol.57, No.9, pp. 986-989, ISSN 0009-2363
- Nhiem, N. X.; Kiem, P. V.; Minh, C. V.; Ha, D. T.; Tai, B. H.; Yen, P. H.; Tung, N. H.; Hyun, J. H.; Kang, H. K., & Kim, Y. H. (2010a). Lupane-type triterpene glycosides from the leaves of *Acanthopanax koreanum* and their in vitro cytotoxicity. *Planta Medica*, Vol.76, No.2, pp. 189-194, ISSN 0032-0943
- Nhiem, N. X.; Kiem, P. V.; Minh, C. V.; Tai, B. H.; Tung, N. H.; Ha, D. T.; Soung, K. S.; Kim, J. H.; Ahn, J. Y.; Lee, Y. M., & Kim, Y. H. (2010b). Structure-activity relationship of lupane-triterpene glycosides from *Acanthopanax koreanum* on spleen lymphocyte IL-2 and INF- γ . *Bioorganic & Medicinal Chemistry Letters*, No.20, pp. 4927-4931, ISSN 0960-894X
- Nhiem, N. X.; Kim, K. C.; Kim, A.-D.; Hyun, J. W.; Kang, H. K.; Kiem, P. V.; Minh, C. V.; Thu, V. K.; Tai, B. H.; Kim, J. A., & Kim, Y. H. (2011). Phenylpropanoids from the leaves of *Acanthopanax koreanum* and their antioxidant activity. *Journal of Asian Natural Products Research* Vol.13, No.1, pp. 56-61, ISSN 1028-6020
- Park, S. H.; Nhiem, N. X.; Kiem, P. V.; Choi, E. M.; Kim, J. A., & Kim, Y. H. (2010). A new norlupane triterpene from the leaves of *Acanthopanax koreanum* increases the differentiation of osteoblastic MC3T3-e1 cells. *Archives of Pharmacal Research*, Vol.33, No.1, pp. 75-80, ISSN 0253-6269
- Park, S. Y.; Choi, H. S.; Yook, C. S., & Nohara, T. (2005). A new lupane glycoside from the leaves of *Acanthopanax koreanum*. *Chemical & Pharmaceutical Bulletin*, Vol.51, No.1, pp. 97-99, ISSN 0009-2363
- Samuelsson, G. & Bohlin, L. (2004). *Drugs of natural origin: A textbook of pharmacognosy*. 5 Ed.; Swedish Pharmaceutical Press, ISBN 9197431842, Stockholm, Sweden
- Shin, E. T. & Kim, C. S. (1985). Composition of fatty acid and organic acid in *Acanthopanax*. *Han'guk Sikp'um Kwahakhoechi*, Vol.17, No.5, pp. 403-405, ISSN 0367-6293
- Srivastava, S. K. (1992). A new triterpenic acid from *Schefflera impressa*. *Journal of Natural Products*, Vol.55, No.3, pp. 298-302, ISSN 0163-3864
- Wu, Y. L.; Jiang, Y. Z.; Jin, X. J.; Lan, L. H.; Piao, J. Y.; Wan, Y.; Jin, H. R.; Lee, J. J., & Nan, J. X. (2010). Acanthoic acid, a diterpene in *Acanthopanax koreanum*, protects acetaminophen-induced hepatic toxicity in mice. *Phytomedicine*, Vol.17, No.6, pp. 475-479, ISSN 0944-7113

Yook, C. S.; Kim, I. H.; Hahn, D. R.; Nohara, T., & Chang, S. Y. (1998). A lupane-triterpene glycoside from leaves of two *Acanthopanax*. *Phytochemistry*, Vol.49, No.3, pp. 839-843, ISSN 0031-9422

IntechOpen

IntechOpen



Phytochemicals - A Global Perspective of Their Role in Nutrition and Health

Edited by Dr Venketeshwer Rao

ISBN 978-953-51-0296-0

Hard cover, 538 pages

Publisher InTech

Published online 21, March, 2012

Published in print edition March, 2012

Phytochemicals are biologically active compounds present in plants used for food and medicine. A great deal of interest has been generated recently in the isolation, characterization and biological activity of these phytochemicals. This book is in response to the need for more current and global scope of phytochemicals. It contains chapters written by internationally recognized authors. The topics covered in the book range from their occurrence, chemical and physical characteristics, analytical procedures, biological activity, safety and industrial applications. The book has been planned to meet the needs of the researchers, health professionals, government regulatory agencies and industries. This book will serve as a standard reference book in this important and fast growing area of phytochemicals, human nutrition and health.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Young Ho Kim, Jeong Ah Kim and Nguyen Xuan Nhiem (2012). Phytochemicals and Their Pharmacological Aspects of *Acanthopanax koreanum*, *Phytochemicals - A Global Perspective of Their Role in Nutrition and Health*, Dr Venketeshwer Rao (Ed.), ISBN: 978-953-51-0296-0, InTech, Available from: <http://www.intechopen.com/books/phytochemicals-a-global-perspective-of-their-role-in-nutrition-and-health/phytochemicals-and-their-pharmacological-aspects-of-acanthopanax-koreanum>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen