

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Intracellular Signaling Pathways Integrating the Pore Associated with P2X7R Receptor with Other Large Pores

L.G.B. Ferreira¹, R.A.M. Reis², L.A. Alves¹ and R.X. Faria¹

¹Laboratory of Cellular Communication, Oswaldo Cruz Institute,
Oswaldo Cruz Foundation

²Laboratory of Neurochemistry, Biophysical Institute,
University Federal of Rio de Janeiro
Brazil

1. Introduction

The purinergic P2X₇ receptor (P2X7R) is a member of the family of ligand-gated ion channels composed of seven subtypes, P2X₁₋₇. These receptors possess three subunits assembled as homo- or heterotrimers to make functional receptors (Nicke et al., 1998, 2005), which has been confirmed by atomic force microscopy (Barrera et al, 2005) and by crystallography (Kawate et al, 2009). The P2X7R shares an overall membrane topology with the other members of this family of receptors; it contains two putative pore forming transmembrane segments, a large cysteine-rich ligand-binding extracellular domain, and intracellularly located N and C termini (Surprenant et al, 1996). This subtype is structurally distinguished from other members of P2XRs by its long intracellular C-terminal tail with multiple protein and lipid interaction motifs, besides a cysteine-rich 18 - amino acid segment. From a pharmacological point of view, the P2X7R requires at least a 100-fold higher ATP concentration for activation compared to other P2XRs (North, 2002). Extracellular divalent cation reduction increases agonist potency (Hibell et al, 2001; Michel et al, 1999). Moreover, extracellular cations and chloride have important effects on the channel gating (Gudipaty et al., 2001; Li et al., 2003, 2008, 2010; Riedel et al., 2007b; Virginio et al., 1997). In this context, P2X7R activation can sustainably induce a wide range of different intracellular signaling responses (Dubyak, 2007; North, 2002).

P2X7R, when activated by ATP or the potent agonist 3'-O-(4-benzoyl)benzoyl-ATP (BzATP), functions as a non-selective cation channel, permeant to small cations, such as Na⁺, K⁺, and Ca²⁺ and this activation mode is dependent on extracellular divalent cations (Ding & Sachs, 2000; Jiang, 2009; Ma et al, 2006; Sperl agh et al, 2006; Virginio et al, 1997). Upon repeated or prolonged application of agonist, the P2X7R becomes permeable to larger molecules like ethidium bromide, N-methyl-D-glucamine or neurotransmitters such as glutamate and ATP (Faria et al, 2005, 2010; Hamilton et al, 2008; Jiang et al, 2005; Marcoli et al, 2008), a process termed cell "permeabilization" or large conductance channel opening.

Up to now, several ideas have been proposed as a possible explanation to this pore opening (Alloisio et al, 2010; Coutinho-silva et al, 1997; Faria et al, 2005, 2009; Jiang et al, 2005; North,

2002; Pelegrin & Surprenant, 2006; Virginio et al, 1999; Yan et al, 2008). In a general manner, cell permeabilization can be observed in cell types transfected or natively expressing P2X7R, in contrast to *Xenopus* oocytes transfected cells (Petrou et al, 1997) or lymphocytes B cells of patients with Chronic lymphocytic leukaemia (Boldt et al, 2003; Gu et al, 2000).

Initially, some groups have proposed that the opening of the pore associated to P2X7R receptor occurs after the small channel (8 pS) allosterically changes and expands over time-dilatation (Chaumont & Khakh, 2008; Virginio et al, 1999; Yan et al, 2008). Others have reported that Pannexin-1 (panx-1), a hemichannel protein, is the large conductance channel that opens independently of the cationic P2X7R (Locovei et al, 2007; Pelegrin & Surprenant, 2006, 2007). On the other hand, some groups have demonstrated another putative explanation of the dye uptake. They have reported two possible charge selectively pathways that are activated by P2X7R: one for cationic and another for anionic dyes (Cankurtaran-Sayar et al, 2009; Schachter et al, 2008). Jiang and collaborators have proposed distinct pathways to permeate inorganic monovalent and divalent cations, organic cations (NMDG⁺) and fluorescent dyes (Jiang et al, 2005). In addition, it has been suggested at least two different conductive pathways, one for Ca²⁺ and other for monovalent ions (Alloisio et al, 2010).

In this context, some groups have published that the P2X7R permeabilization depends on intracellular factors (Donnelly-Roberts et al, 2004; Faria et al, 2005, 2010; Gu et al, 2009; Le Stunff & Raymond, 2007; Shemon et al, 2004; Zhao et al, 2007) to occur.

In this chapter, we come up with novel proposals of intracellular signaling regulation that would help us to understand about the several intriguing characteristics of the ATP-induced P2X7R.

2. Intracellular signalling associated with small conductance channel

In the P2X7R receptor gating mechanism (low conductance channel) the response to agonist challenge, allow rapid, non-selective passage of cations across the cell membrane. It is permeable to Na⁺ and K⁺ and presents high permeability to Ca²⁺ (North, 2002). The cellular response to P2X7R receptor activation after agonist exposition is generally very rapid and this reaction does not depend on the production and diffusion of second messengers within the cytosol (Burnstock, 2007; Ralevic & Burnstock, 1998). In contrast, an increase in the intracellular Ca²⁺ concentration and a consequent depolarization of the cell membrane are observed after the ionic channel opening, subsequently activating voltage-gated calcium channels. In addition, evidence indicates that the large Ca²⁺ ion concentration in the cytoplasm could activate intracellular kinases like protein kinase C (PKC), mitogen activated protein kinases (MAPKs), calcium-calmodulin-dependent protein kinase II (CaMKII) (Amstrup & Novak, 2003; Bradford & Soltoff, 2002; Heo & Han, 2006), caspases (Orinska et al, 2011), phosphoinositide 3-kinase (Pi3K) (Jaques-Silva et al, 2004) and phospholipases (Alzola et al, 1998; Perez-Andres et al, 2002; Pochet et al, 2007). Further, other functions may be mediated by P2X7R receptor activation such as IL-1 β maturation (Ferrari et al, 1997), shedding of membrane proteins (Moon et al, 2006); Src (Denlinger et al, 2001), and glycogen synthase kinase 3 (Ortega et al, 2009); membrane blebbing (Morelli et al, 2003).

These different signaling pathways may contribute to a large complexity in the response of this receptor and it raises the question about how the correct coupling and fine tuning of the signaling in response to extracellular stimuli is achieved. To date, several pieces of evidence

have been described about the small conductance P2X7R receptor and interactions with other proteins (Adinolfi et al, 2003; Antonio et al, 2011; Barbieri et al, 2008; Boumechache et al, 2009; Bradley et al, 2010; Denlinger et al, 2001; Guo et al, 2007; Lemaire et al, 2007; Liu et al, 2011; Wilson et al, 2002), mainly through its longer C-terminus region or lipids (Denlinger et al, 2001; Gonnord et al, 2009; Michel & Fonfria, 2007; Takenouchi et al, 2007; Zhao et al, 2007a, 2007b). However, there are only a few studies about the low conductance P2X7R receptor gating and the mechanism of transition to large conductance channel opening. This may be due to (i) a lack of selective agonists or antagonists only to small or large channel (North, 2002) or (ii) to P2X7R receptor polymorphisms or (iii) to artificial deletions in regions of this receptor resulting in distinct intracellular or extracellular regulation of this phenomenon. In this context, the concept that the C-terminal domain of P2X7R directly regulates a complex distinct from receptor-dependent pore activity was first introduced by El-Moatassim and Dubyak (El-Moatassim & Dubyak, 1992). They demonstrated that P2X7R receptor mediated phospholipase D (PLD) activity was dependent on GTP and independent of the large conductance channel opening. Another study found that human P2X₇ receptor currents were facilitated in response to repeated or prolonged agonist applications, via dynamic calmodulin binding (Roger et al, 2008). This Ca²⁺-dependent component is related to the uptake of large compounds seen by the pore complex. These and other papers (Alloysio et al, 2010; Boldt et al, 2003; Le Stunff et al, 2004) suggest independent intracellular signaling pathways regulating the low and large conductance channel associated with P2X7R.

3. Intracellular signalling pathways associated with large conductance channel

The intracellular regulation of the P2X7R associated large conductance channel opening is still mostly unknown. The initial suggestions for dependency of cytoplasmic factors in this event were originated from electrophysiological data in outside or inside out configurations with no large conductance channel recordings (Coutinho-Silva & Persechini, 1997; Persechini et al, 1998; Petrou et al, 1997). Posteriorly, other groups have investigated the P2X7R pore formation induced by intracellular signaling and how mutated amino acids or truncated regions affect functional availability of the receptor. In this line, Smart used truncated and single-residue-mutated P2X7R receptors in HEK-293 cells and in *Xenopus* oocytes. Truncated P2X7R at residue 581 (of 595) were not able to dye uptake, but there was dye uptake similarly to the wild receptor in those cells expressing the truncated P2X7R at position 582. In contrast, the small channel function was only suppressed in the residues 380 (Smart et al, 2003). Two alternative splices variants were identified in the human P2X7R (one lacking the first transmembrane domain and the other the entire cytoplasmic tail, but they were compared to the full-length channel). The first variant exhibited a non-functional slow conductance channel, while the second did not affect the small ion channel activity, but affected the large conductance channel and caspase activation (Cheewatrakoolpong et al, 2005). In addition, threonine 283 (Thr283) has been described as a critical residue in the ectodomain for P2X7R receptor function and it has been suggested that the intracellular leucine residue (P451L) alters downstream signalling independently of ion channel activity (Young et al, 2006). Recently, Marques-da-Silva and collaborators (Marques-da-Silva et al, 2011) demonstrated that colchicine did not inhibit ATP-evoked currents in macrophages, but it decreased ATP-induced dye uptake. Large conductance channel opening on *Xenopus*

oocytes and HEK293 cells expressing P2X7R were inhibited after colchicines treatment (Marques-da-Silva et al, 2011). Yan described that extracellular Ca^{2+} concentration is a physiological negative modulator of the P2X7R low conductance channel without affecting the large conductance channel opening (Yan et al, 2011).

There is some controversial data related to the biophysical, pharmacological or molecular tools that impair the actions of intracellular enzymes involved in the P2X7R pore formation. In one of the primary papers studying the intracellular signaling of the P2X7R large conductance channel, it was described that calmidazolium, an inhibitor of the calmodulin protein, impaired the small channel activity, but had no effect in the large conductance channel opening (Virginio et al, 1997). This result was also confirmed by other groups (Donnelly-Roberts et al, 2004; Faria et al, 2010; Lundy et al, 2004). However, Roger and coworkers (Roger et al, 2008, 2010) reported that rat P2X7R induced large organic cation permeability ionic currents were dependent on critical residues of calmodulin binding domain when recorded in patch-clamp whole cell configuration. In this sense, the intracellular Ca^{2+} concentration dependence in the P2X7R pore formation is still unclear. We have shown in 2005 that intracellular Ca^{2+} acts as a second messenger in the large conductance channel opening in peritoneal macrophages and 2BH4 cells (Faria et al, 2005). Similar data were found by other groups (Cankurtaran-Sayar et al, 2009; Roger et al, 2008, 2010; Schachter et al, 2008), but others did not observe this effect (da Cruz et al, 2006; Iglesias et al, 2008; Schachter et al, 2008; Virginio et al, 1999). We continued to investigate the Ca^{2+} participation in more detail, and we described a major Ca^{2+} dependence in the P2X7R pore formation, but we also observed Ca^{2+} independent events in the same cell types (Faria et al, 2010). This variability in the responses may be due to preponderant expression of distinct P2X7R variants or activity of different large conductance channels, as was discussed above in the text. Phospholipase C (PLC) had no effect on P2X7R large conductance channel formation in THP-1 cells (Donnelly-Roberts et al, 2004), or in mouse 2BH4 cells or peritoneal macrophages cells (Faria et al, 2010). However, P2X7R pore formation was inhibited in mouse microglial cell line by PLC (Takenouchi et al, 2005). It was also shown that MAPK is associated with P2X7R activation (Donnelly-Roberts et al, 2004; Faria et al, 2005, 2010), but in other hands this was not confirmed (da Cruz et al, 2006; Michel et al, 2006). These discrepancies may be due to species variations, distinct intracellular machinery or differences in the protocol used to investigate a specific function in a same cell type (Faria et al, 2010). Other proteins have presented less divergent responses, such as: PKC (Donnelly-Roberts et al, 2004; Faria et al, 2010; Shemon et al, 2004), Ca^{2+} -insensitive Phospholipase A PLA (Chaib et al, 2000), caspase-1 and-3 (Donnelly-Roberts et al, 2004; Faria et al, 2010), PLD (Stunff & Raymond, 2007), phosphatidylinositol 4,5-bisphosphate (PIP_2) (Zhao et al, 2007), cytoskeleton components (Marques-da-Silva et al, 2011), PI3K (Faria et al, 2010), src tyrosine phosphorylation (Iglesias et al, 2008), Peroxisome proliferator-activated receptor gamma (PPAR gamma) (Nagasawa et al, 2009a) antagonists and intracellular Ca^{2+} chelants (Faria et al, 2005, 2010).

As mentioned above, one possible drawback in relation to these results is due to the diversity of responses observed in P2X₇ species and cell types (Donnelly-Roberts et al, 2009; Michel et al, 2008). The variations may be proportional to natural P2X7R receptor polymorphisms and these may be, at least partially, functional promoting gain or loss of activity (Cheewatrakoolpong et al, 2005; Feng et al, 2006; Masin et al, 2011; Shemon et al, 2006). Another related factor to this matter is the native structural state of the P2X7R

receptor. In this line, a monoclonal antibody (Ab) to P2X7R ectodomain was used to immunoprecipitate the receptor complex in central and peripheral immune cells (Kim et al, 2001b). Using western blotting, native P2X7R in peritoneal macrophage or bone marrow cells formed bound multimeric complex with numerous bands ranging in size from 25 up to 250 kDa, in contrast to P2X7R from brain glia and/or astrocytes that formed only monomeric subunits. This result suggests differential intracellular regulation of the P2X7R pore in distinct cell types (Kim et al, 2001b). Li and coworkers discovered in parotid acinar and duct cells a cell-specific assembly and gating of the P2X7R channels, in a way that upon exposure to ATP, P2X7Rs are assembled into functional channels with rapid gating. In contrast, P2X7Rs from duct cells are preassembled and continually subject to rapid gating by ATP (Li et al, 2003). Recently, other researches have found distinct pathways of dye uptake, mediated by P2X7R receptor activation after ATP treatment, possibly through different large conductance channels. Schachter and collaborators compared P2X7R-associated cation and anionic fluorescent dyes uptake of macrophages and HEK-293 cells transfected with P2X7R receptor (Schachter et al, 2008). Transfected cells did not take up anionic dyes and did not display single channel cell-attached recordings, in contrast to the native mice peritoneal macrophages. Anionic and cationic dye effluxes induced by ATP treatment were temperature independent and dependent, respectively (Schachter et al, 2008). In addition, another study examined the process of dye uptake by transfected or natively expressed P2X7R receptor leading to the pore formation. HEK-293 cells expressing rat P2X7R was permeable to cationic but not to anionic dyes in a way that intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$) increase was not necessary to be activated (via 1). In the via 2, the pore was permeated only by lucifer yellow and it was completely dependent on $[Ca^{2+}]_i$ for activation. Also, RAW 264.7 cells presented both pathways similar to the transfected cells, but they did not require intracellular Ca^{2+} (Cankurtaran-Sayar et al, 2009).

Based on all these data from different groups suggesting that more than one pore might work simultaneously after ATP treatment, we describe the intracellular enzymes activated by the P2X7R associated large conductance channel opening compared to other large conductance channels.

4. Intracellular signaling cascades activated by these pores

Since the proteins responsible to the P2X7R large conductance channel opening are still largely unknown here we compare and discuss the intracellular signaling pathways and the possible candidates associated to the P2X7R pore formation. Among them are connexin hemichannels, pannexin-1, plasma membrane voltage dependent anion channel (pI-VDAC), maxi anion, transient receptor potential vanilloid-1 (TRPV1), transient receptor potential anquirin-1 (TRPA1), Maitotoxin-induced pore and Rising of intracellular Ca^{2+} concentration induced pore.

The connexin hemichannel, a hexameric protein composed of connexin subunits expressed in vertebrates, was the first large conductance channel studied with functional similarity to P2X7R receptor large conductance channels. An initial study used two types of J774 mouse macrophages, one sensitive and another ATP-insensitive. In the sensitive cells, connexin-43 (Cx43) gap junction mRNA and protein and P2X7R were expressed and the dye was taken up, but in the insensitive lineage there was not Cx43 expression and neither dye uptake. Therefore, they proposed that connexin 43 was the pore associated to P2X7R receptor (Beyer

& Steinberg, 1991). This concept was elegantly refuted by Alves and colleagues at least in peritoneal macrophages where they demonstrated that experimental conditions known to block hemichannels and Cx43 knockout mice maintained the P2X7R large conductance channel activity (Alves et al, 1996). Also, P2X7R and Cx43 are expressed in J774 macrophage lineage and are colocalized in the cell membrane (Fortes et al, 2004).

In relation to the intracellular signaling cascades, connexins can be modulated in the C-terminal domain by phosphorylation through PKC (Bao et al, 2007; Hawat & Baroudi, 2008), MAPK (Bao et al, 2007), S-nitrosylation with covalent binding of nitric oxide (NO) to cysteine (Cys) (De Vuyst et al, 2007; Retamal et al, 2009), protein kinase A (Liu et al, 2011), intracellular redox potential (Retamal et al, 2007) and intracellular Ca^{2+} concentration (De Vuyst et al, 2006; Schalper et al, 2008; Thimm et al, 2005). Compared to the P2X7R receptor pore, the connexin hemichannel may be dependent on intracellular Ca^{2+} , PKC and MAPK. Meanwhile, the unitary conductance value (20-250pS) for all known connexins are lower compared to the ones observed to P2X7R receptor (400pS). In addition, it has been shown that connexin hemichannel blockers have no effect on the P2X7R receptor pore formation (Faria et al, 2005) and this apparently ruled out the participation of connexins at least in cell types tested.

Maitotoxin (MTX), a marine toxin, described to increase calcium in GH4C1 rat pituitary cells (Young et al, 1995), increases intracellular Ca^{2+} concentration leading to the opening of a pore with biophysical properties similar to P2X7R large conductance channel (Schilling et al, 1999a, 1999b). The dye uptake observed after this pore opening may also be dependent on extracellular Ca^{2+} (Lundy et al, 2004; Wisnoskey et al, 2004), intracellular Ca^{2+} concentration (Wisnoskey et al, 2004), calmodulin (Donnelly-Roberts et al, 2004; Lundy et al, 2004) and PLC (Donnelly-Roberts et al, 2004). Although the maitotoxin pore may be functionally similar to the P2X7R pore, they might possess different intracellular pathways. A possible explanation to this fact may be that the same large conductance channel or different similar pores functioning in conjunction might be regulated by distinct signaling pathways

In 2009, our group described a large conductance channel stimulated by rising of intracellular Ca^{2+} concentration recorded in cell attached patches. This pore was blocked by calmodulin, Calcium-calmodulin kinase type II (CamKII), PLC, MAPK and caspase-1 and-3 antagonists and it was insensitive to PKC and P2X7R receptor antagonists (Faria et al, 2009). The intracellular signaling pathways modulating this pore and the one associated with P2X7R are distinct, but they possessed some common pathways. In addition, the pore induced by MTX presents large intracellular signalling similar to the pore described by us. Since the protein responsible for the opening of the MTX induced pore also is not identified, the pore recorded in our conditions (Faria et al, 2009) may be the same as the MTX pore.

Another large conductance channel, that may be activated by rising of intracellular Ca^{2+} concentration (Locovei et al, 2006; Ma et al, 2006), is the pannexin hemichannel, which is a hexameric protein present in vertebrates and invertebrates. However, as reported by recent papers, the extracellular or intracellular Ca^{2+} did not interfere with the pannexin activity (Ma et al, 2009; Pelegrin & Surprenant, 2007). This large conductance channel might be activated by S-nitrosylation and Src kinase (Iglesias et al, 2009; Pelegrin & Surprenant, 2006; Suadicani et al, 2009). P2X7R receptor large conductance channel was inhibited by RNAi to pannexin-1, inhibitory peptide and pannexin antagonists (Pelegrin & Surprenant, 2006). In contrast, other groups did not observe inhibition of the P2X7R large conductance channel

for pannexin-1 inhibitors or RNAi (Faria et al, 2005, 2010; Nagasawa et al, 2009b; Reyes et al, 2008; Schchater et al, 2008; Yan et al, 2008, 2011). Up to now, for some groups the pannexin-1 seems to be an important player in this phenomenon, but apparently this protein is working in conjunction with other protein(s). This information is based on the partial blockage of the P2X7R receptor induced dye uptake exhibited after pannexin-1 inhibition (Iglesias et al, 2009; Locovei et al, 2007; Pelegrin & Surprenant, 2007). Alternatively, other large conductance channel such as MTX-induced pore (Pelegrin & Surprenant, 2007), P2X2R large conductance channel (Marques-da-Silva et al, 2011) and the rising of intracellular Ca^{2+} induced pore (Faria et al, 2009) were not impaired by pannexin-1 inhibitors.

Maxi-anion channel possesses a wide nanoscopic pore suitable for nucleotide transport and an ATP-binding site in the middle of the pore lumen to facilitate the passage of the nucleotide (Sabirot & Okada, 2004). Physiologically, the same large conductance channel is operational in swelling-, ischemia-, and hypoxia-induced ATP release from neonatal rat cardiomyocytes (Dutta et al, 2004). In addition, raising the intracellular Ca^{2+} concentration (Bajnath et al, 1993; Groschner & Kukovetz, 1992; Hussy, 1992; Kawahara & Tawuka, 1991) as well as protein tyrosine dephosphorylation (Toychiev et al, 2009) can activate this pore. On the other hand, PKA (Okada et al, 1997), PKC (Kokubun et al, 1991; Saiguza & Kokubun, 1988; Vaca & Kunze, 1993), G proteins (Schwiebert et al, 1992; Sun et al, 1993) and Src kinase (Kajita et al, 1995) antagonists can inhibit it. This large conductance channel also has no protein constituents identified so far (Sabirot & Okada et al, 2009) and there are no studies comparing its effects with other pores, except to pl-VDAC (Sabirot et al, 2006).

Voltage-dependent anion channels (VDACs) were originally characterized as mitochondrial porins but other evidence began to accumulate that VDACs could also be expressed in the plasma membrane (pl-VDAC). VDAC may be activated changing the applied voltage in the presence of NADH (Zizi et al, 1994) and under apoptotic conditions (Elinder et al, 2005). In relation to the intracellular signaling pathways, there are few data up to now, but some groups have shown the involvement of this pore with lipid rafts (Ferrer, 2009; Herrera et al, 2011). Moreover, the large conductance channel and pl-VDAC may be activated by excised patch, indicating an independence of the intracellular signals to open the large conductance channel (Guibert et al, 1998; Sun et al, 1993). Relevant information is about the nucleotide-binding sites in the C-terminus of the mitochondrial VDAC, which presents the same C-terminal sequence of the pl-VDAC (Yehezkel et al, 2006). The main discrepancy in relation to both pores compared to the P2X7R pore is due to the lack of single channel recordings of the P2X7R pore in excised patches (Riedel et al, 2007). This fact may indicate that maxi anion and pl-VDAC pores are different compared to the P2X7R receptor pore, since they did not depend on cytoplasmic factors to open. But, this does not rule out the participation of this pore in this phenomenon.

The capsaicin induced receptor, transient receptor potential vanilloid 1 (TRPV1), is activated not only by capsaicin but also by heat ($>43^{\circ}\text{C}$), acid and various lipids (Moran et al, 2011). Since capsaicin and its analogues, such as resiniferatoxin (RTX), are lipophilic, it is quite possible that they pass through the cell membrane and act on the binding sites present in the intracellular surface of TRPV1. It is a Ca^{2+} permeable non-specific cation channel. It was demonstrated that activation of native or recombinant rat TRPV1 leads to time- and agonist concentration-dependent increase in the relative permeability of large cations and changes in Ca^{2+} permeability (Chung et al, 2008). TRPV1 induced small channel can be modulated by calmodulin, PKC, PKA, intracellular Ca^{2+} , PLC, G protein and PIP_2 / Src (Bhave et al, 2002;

Chuang et al, 2001; Dai et al, 2004; Moriyama et al, 2003, 2005; Sugiura et al, 2002; Tominaga et al, 2001). TRPV1 induced a large conductance channel similar to the P2X7R since it depends on the C-terminus and it is modulated by PKC phosphorylation (Chung et al, 2008). Although biophysically this pore is similar to the P2X7R associated pore, the intracellular signaling is poorly understood up to now.

TRPA1 is a nonselective cation channel that belongs to the superfamily of mammalian TRP ion channels and is unique since it possesses a large number of ankyrin repeats in its N-terminal domain (Montell, 2005). TRPA1, when activated, are permeable to small cations such as Ca^{2+} , K^+ , Na^+ ; simultaneously it depolarizes the plasma membrane and raises intracellular Ca^{2+} , which subsequently triggers a variety of physiological responses. Recently, it was described that TRPA1 activation induces dye uptake, which is blocked by selective TRPA1 antagonists. In addition, outside-out patch recordings using N-methyl-D-glucamine (NMDG^+) as the sole external cation and Na^+ as the internal cation, TRPA1




Fig. 1. Pharmacological comparison of the intracellular pathway of the pore associated with P2X₇ receptor in different cell types. A- Whole cell experiments in mice peritoneal macrophages, mice cortical astrocytes or mice mesencephalic. We applied 1mM ATP, after the incubation of the cells with 10 μ M BAPTA-AM, 1 μ M Sb203580 or 1 μ M Staurosporine for 5 minutes at 37°C. B- The graphic represents the quantification of dye uptake experiments in the cell types cited above. The values represent the mean \pm SD of three to four experiments performed on different days. * $p < 0.05$ compared with the ATP treatment of each group.

activation results in dynamic changes in permeability to NMDG⁺ (Chen et al, 2008). Other groups have reproduced this data (Banke et al, 2010, 2011), but in every cell studied the intracellular signaling was not investigated. Moreover, the fact that TRPA1 associated large conductance channel permeates large cations in outside out configuration suggests a possible independence of intracellular factors.

As we can observe above, there are diverse common intracellular signaling proteins that may be used in the activation of these large conductance pores. Moreover, these pores might be biophysically and functionally similar. Thus, when we performed assays which preincubated cells with intracellular signaling pathway blockers were stimulated with 1mM ATP, there was an activation of the large conductance channel associated to P2X7R (Figure 1), as previously shown in our published papers (Faria et al, 2005, 2010). Using whole cell configuration, we used (i) neurons to evaluate the P2X7R pore formation in cells expressing TRPV1 and TRPA1; (ii) astrocytes to study the expression of Maxi anion, pI-VDAC, Connexin 43 and Pannexin-1; and (iii) macrophages to study Maitotoxin and intracellular Ca²⁺ increase induced pores.

5. Conclusions

Finally, based on data discussed here, several issues might explain why a common gate mechanism for the P2X7R pore is not yet understood: (1) a large conductance channel is activated by different signaling pathways; (2) these signaling cascades might be related to the activation of distinct pores; (3) both [(1) and (2)] mechanisms might act together in certain cells; (4) it might exist a gate modulator that is cell-type specific, (5) P2X₇ might be part of a macromolecular protein complex or a protein-lipid complex. In summary, more studies are necessary in order to comprehend the functional mechanism of the P2X₇ receptor.

6. References

- Adinolfi E, Kim M, Young MT, Di Virgilio F, Surprenant A (2003) Tyrosine phosphorylation of HSP90 within the P2X7 receptor complex negatively regulates P2X7 receptors. *J Biol Chem* 278:37344-37351.
- Alloisio S, Di Garbo A, Barbieri R, Bozzo L, Ferroni S, Nobile M (2010) Evidence for two conductive pathways in P2X receptor: differences in modulation and selectivity. *J Neurochem*. 113, 796-806.
- Alves LA, Coutinho-Silva R, Persechini PM, Spray DC, Savino W, Campos de Carvalho AC (1996) Are there functional gap junctions or junctional hemichannels in macrophages? *Blood*. 88, 328-334.
- Alzola E, Pérez-Etxebarria A, Kabré E, Fogarty DJ, Métioui M, Chaïb N, Macarulla JM, Matute C, Dehaye JP, Marino A (1998) Activation by P2X7 agonists of two phospholipases A2 (PLA2) in ductal cells of rat submandibular gland. Coupling of the calcium-independent PLA2 with kallikrein secretion. *J Biol Chem*. 273, 30208-30217.
- Amstrup J, Novak I (2003) P2X7 receptor activates extracellular signal-regulated kinases ERK1 and ERK2 independently of Ca²⁺ influx. *Biochem J*. 374, 51-61.

- Antonio LS, Stewart AP, Xu XJ, Varanda WA, Murrell-Lagnado RD, Edwardson JM (2011) P2X4 receptors interact with both P2X2 and P2X7 receptors in the form of homotrimers. *Br J Pharmacol.* 163, 1069-1077.
- Bajnath RB, Groot JA, de Jonge HR, Kansen M, Bijman J (1993) Calcium ionophore plus excision induce a large conductance chloride channel in membrane patches of human colon carcinoma cells HT-29cl.19A. *Experientia* 1993 49, 313-316.
- Banke TG, Chaplan SR, Wickenden AD (2010) Dynamic changes in the TRPA1 selectivity filter lead to progressive but reversible pore dilation. *Am J Physiol Cell Physiol.* 298, C1457-68.
- Banke TG (2011) The dilated TRPA1 channel pore state is blocked by amiloride and analogues. *Brain Res* 1381, 21-30.
- Bao X, Lee SC, Reuss L, Altenberg GA (2007) Change in permeant size selectivity by phosphorylation of connexin 43 gap-junctional hemichannels by PKC. *Proc Natl Acad Sci U S A.* 104, 4919-4924.
- Barbieri R, Alloisio S, Ferroni S, Nobile M (2008) Differential crosstalk between P2X7 and arachidonic acid in activation of mitogen-activated protein kinases. *Neurochem Int.* 53, 255-262.
- Barrera NP, Ormond SJ, Henderson RM, Murrell-Lagnado RD, Edwardson JM. (2005) Atomic force microscopy imaging demonstrates that P2X2 receptors are trimers but that P2X6 receptor subunits do not oligomerize. *J Biol Chem.* 280, 10759-65.
- Beyer EC, Steinberg TH (1991) Evidence that the gap junction protein connexin-43 is the ATP-induced pore of mouse macrophages. *J Biol Chem.* 266, 7971-7974.
- Bhave G, Zhu W, Wang H, Brasier DJ, Oxford GS, Gereau RW 4th (2002) cAMP-dependent protein kinase regulates desensitization of the capsaicin receptor (VR1) by direct phosphorylation. *Neuron.* 2002 35, 721-731.
- Boldt W, Klapperstück M, Büttner C, Sadtler S, Schmalzing G, Markwardt F (2003) Glu496Ala polymorphism of human P2X7 receptor does not affect its electrophysiological phenotype. *Am J Physiol Cell Physiol.* 284, C749-56.
- Boumechache M, Masin M, Edwardson JM, Górecki DC, Murrell-Lagnado R (2009) Analysis of assembly and trafficking of native P2X4 and P2X7 receptor complexes in rodent immune cells. *J Biol Chem* 284, 13446-13454.
- Bradford MD, Soltoff SP (2002) P2X7 receptors activate protein kinase D and p42/p44 mitogen-activated protein kinase (MAPK) downstream of protein kinase C. *Biochem J* 366, 745-755.
- Bradley HJ, Liu X, Collins V, Owide J, Goli GR, Smith M, Surprenant A, White SJ, Jiang LH (2010) Identification of an intracellular microdomain of the P2X7 receptor that is crucial in basolateral membrane targeting in epithelial cells. *FEBS Lett.* 584, 4740-4744.
- Burnstock G (2007) Purine and pyrimidine receptors. *Cell Mol Life Sci.* 64, 1471-1483.
- Cankurtaran-Sayar S, Sayar K, Ugur M (2009) P2X7 receptor activates multiple selective dye-permeation pathways in RAW 264.7 and human embryonic kidney 293 cells. *Mol Pharmacol.* 76, 1323-1332.
- Chaib N, Kabré E, Alzola E, Pochet S, Dehaye JP (2000) Bromoenol lactone enhances the permeabilization of rat submandibular acinar cells by P2X7 agonists. *Br J Pharmacol.* 129, 703-708.

- Chaumont S, Khakh BS (2008) Patch-clamp coordinated spectroscopy shows P2X2 receptor permeability dynamics require cytosolic domain rearrangements but not Panx-1 channels. *Proc Natl Acad Sci U S A.* 105, 12063-12068.
- Cheewatrakoolpong B, Gilchrest H, Anthes JC, Greenfeder S (2005) Identification and characterization of splice variants of the human P2X7 ATP channel. *Biochem Biophys Res Commun.* 332, 17-27.
- Chen J, Zhang XF, Kort ME, Huth JR, Sun C, Miesbauer LJ, Cassar SC, Neelands T, Scott VE, Moreland RB, Reilly RM, Hajduk PJ, Kym PR, Hutchins CW, Faltynek CR (2008) Molecular determinants of species-specific activation or blockade of TRPA1 channels. *J Neurosci.* 28, 5063-5071.
- Chuang HH, Prescott ED, Kong H, Shields S, Jordt SE, Basbaum AI, Chao MV, Julius D (2001) Bradykinin and nerve growth factor release the capsaicin receptor from PtdIns(4,5)P2-mediated inhibition. *Nature* 411, 957-962.
- Chung MK, Güler AD, Caterina MJ (2008) TRPV1 shows dynamic ionic selectivity during agonist stimulation. *Nat Neurosci.* 11, 555-564.
- Coddou C, Yan C, Obsil T, Huidobro-Toro JP, Stojilkovic SS (2011) Activation and Regulation of Purinergic P2X Receptor Channels. *Pharmacol Rev* print July 7, 2011, doi: 10.1124/pr.110.003129
- Coutinho-Silva R, Persechini PM (1997) P2Z purinoceptor-associated pores induced by extracellular ATP in macrophages and J774 cells. *Am J Physiol.* 273, C1793-800.
- da Cruz CM, Ventura AL, Schachter J, Costa-Junior HM, da Silva Souza HA, Gomes FR, Coutinho-Silva R, Ojcius DM, Persechini PM (2006) Activation of ERK1/2 by extracellular nucleotides in macrophages is mediated by multiple P2 receptors independently of P2X(7)-associated pore or channel formation. *Br J Pharmacol* 147, 324-334.
- Dai Y, Moriyama T, Higashi T, Togashi K, Kobayashi K, Yamanaka H, Tominaga M, Noguchi K (2004) Proteinase-activated receptor 2-mediated potentiation of transient receptor potential vanilloid subfamily 1 activity reveals a mechanism for proteinase-induced inflammatory pain. *J Neurosci.* 24, 4293-4299.
- De Vuyst E, Decrock E, Cabooter L, Dubyak GR, Naus CC, Evans WH, Leybaert L (2006) Intracellular calcium changes trigger connexin 32 hemichannel opening. *EMBO J.* 25, 34-44.
- De Vuyst E, Decrock E, De Bock M, Yamasaki H, Naus CC, Evans WH, Leybaert L (2007) Connexin hemichannels and gap junction channels are differentially influenced by lipopolysaccharide and basic fibroblast growth factor. *Mol Biol Cell.* 18, 34-46.
- Denlinger LC, Fisette PL, Sommer JA, Watters JJ, Prabhu U, Dubyak GR, Proctor RA, Bertics PJ.
- Cutting edge: the nucleotide receptor P2X7 contains multiple protein- and lipid-interaction motifs including a potential binding site for bacterial lipopolysaccharide (2001) *J Immunol.* 167, 1871-1876.
- Denlinger LC, Sommer JA, Parker K, Gudipaty L, Fisette PL, Watters JW, Proctor RA, Dubyak GR, Bertics PJ (2003) Mutation of a dibasic amino acid motif within the C terminus of the P2X7 nucleotide receptor results in trafficking defects and impaired function. *J Immunol* 171, 1304-1311
- Ding S, Sachs F (2000) Inactivation of P2X2 purinoceptors by divalent cations. *J Physiol* 522, 199-214.

- Donnelly-Roberts DL, Namovic MT, Faltynek CR, Jarvis MF (2004) Mitogen-activated protein kinase and caspase signaling pathways are required for P2X7 receptor (P2X7R)-induced pore formation in human THP-1 cells. *J Pharmacol Exp Ther* 308, 1053-1061.
- Donnelly-Roberts DL, Namovic MT, Han P, Jarvis MF (2009) Mammalian P2X7 receptor pharmacology: comparison of recombinant mouse, rat and human P2X7 receptors. *Br J Pharmacol.* 157, 1203-1214.
- Dubyak GR (2007). Go it alone no more--P2X7 joins the society of heteromeric ATP-gated receptor channels. *Mol Pharmacol.* 72, 1402-1405.
- Dutta AK, Sabirov RZ, Uramoto H, Okada Y (2004) Role of ATP-conductive anion channel in ATP release from neonatal rat cardiomyocytes in ischaemic or hypoxic conditions. *J Physiol* 559, 799-812.
- El-Moatassim C, Dubyak GR (1992) A novel pathway for the activation of phospholipase D by P2z purinergic receptors in BAC1.2F5 macrophages. *J Biol Chem* 267, 23664-23673.
- Elinder F, Akanda N, Tofighi R, Shimizu S, Tsujimoto Y, Orrenius S, Ceccatelli S (2005) Opening of plasma membrane voltage-dependent anion channels (VDAC) precedes caspase activation in neuronal apoptosis induced by toxic stimuli. *Cell Death Differ* 12, 1134-1140.
- Faria RX, Defarias FP, Alves LA (2005) Are second messengers crucial for opening the pore associated with P2X7 receptor? *Am J Physiol Cell Physiol.* 288, C260-71.
- Faria RX, Reis RA, Casabulho CM, Alberto AV, de Farias FP, Henriques-Pons A, Alves LA (2009) Pharmacological properties of a pore induced by raising intracellular Ca²⁺. *Am J Physiol Cell Physiol.* 297, C28-42.
- Faria RX, Casabulho CM, Reis RA, Alves LA (2010) Large-conductance channel formation mediated by P2X7 receptor activation is regulated through distinct intracellular signaling pathways in peritoneal macrophages and 2BH4 cells. *Naunyn Schmiedebergs Arch Pharmacol.* 382, 73-87.
- Feng YH, Li X, Wang L, Zhou L, Gorodeski GI (2006) A truncated P2X7 receptor variant (P2X7-j) endogenously expressed in cervical cancer cells antagonizes the full-length P2X7 receptor through hetero-oligomerization. *J Biol Chem.* 281, 17228-17237.
- Ferrari D, Chiozzi P, Falzoni S, Dal Susino M, Melchiorri L, Baricordi OR, Di Virgilio F (1997) Extracellular ATP triggers IL-1 beta release by activating the purinergic P2Z receptor of human macrophages. *J Immunol* 159, 1451-1458.
- Ferrer I. (2009) Altered mitochondria, energy metabolism, voltage-dependent anion channel, and lipid rafts converge to exhaust neurons in Alzheimer's disease. *J. Bioenerg. Biomembr.* 41, 425-431.
- Fortes FS, Pecora IL, Persechini PM, Hurtado S, Costa V, Coutinho-Silva R, Braga MB, Silva-Filho FC, Bisaggio RC, De Farias FP, Scemes E, De Carvalho AC, Goldenberg RC (2004) Modulation of intercellular communication in macrophages: possible interactions between GAP junctions and P2 receptors. *J Cell Sci.* 117, 4717-4726.
- Gonnord P, Delarasse C, Auger R, Benihoud K, Prigent M, Cuif MH, Lamaze C, Kanellopoulos JM (2009) Palmitoylation of the P2X7 receptor, an ATP-gated channel, controls its expression and association with lipid rafts. *FASEB J.* 23, 795-805.

- Groschner K, Kukovetz WR (1992) Voltage-sensitive chloride channels of large conductance in the membrane of pig aortic endothelial cells. *Pflugers Arch.* 421, 209-217. Erratum in: *Pflugers Arch* 1992 Sep;421, 613.
- Gu B, Bendall LJ, Wiley JS (1998) Adenosine triphosphate-induced shedding of CD23 and L-selectin (CD62L) from lymphocytes is mediated by the same receptor but different metalloproteases. *Blood* 92, 946-951.
- Gudipaty L, Humphreys BD, Buell G, Dubyak GR (2001) Regulation of P2X(7) nucleotide receptor function in human monocytes by extracellular ions and receptor density. *Am J Physiol Cell Physiol* 280, C943-C953.
- Guibert B, Dermietzel R, Siemen D (1998) Large conductance channel in plasma membranes of astrocytic cells is functionally related to mitochondrial VDAC-channels. *Int J Biochem Cell Biol.* 30, 379-391.
- Guo C, Masin M, Qureshi OS, Murrell-Lagnado RD (2007) Evidence for functional P2X4/P2X7 heteromeric receptors. *Mol Pharmacol.* 72, 1447-1456.
- Hamilton N, Vayro S, Kirchhoff F, Verkhratsky A, Robbins J, Gorecki DC, Butt AM (2008) Mechanisms of ATP- and glutamate-mediated calcium signaling in white matter astrocytes. *Glia* 56, 734-749.
- Hawat G, Baroudi G (2008) Differential modulation of unapposed connexin 43 hemichannel electrical conductance by protein kinase C isoforms. *Pflugers Arch.* 456, 519-527.
- Heo JS, Han HJ (2006) ATP stimulates mouse embryonic stem cell proliferation via protein kinase C, phosphatidylinositol 3-kinase/Akt, and mitogen-activated protein kinase signaling pathways. *Stem Cells* 24, 2637-2648.
- Herrera JL, Diaz M, Hernández-Fernaund JR, Salido E, Alonso R, Fernández C, Morales A, Marin R (2011) Voltage-dependent anion channel as a resident protein of lipid rafts: post-transductional regulation by estrogens and involvement in neuronal preservation against Alzheimer's disease. *J Neurochem.* 116, 820-827. doi: 10.1111/j.1471-4159.2010.06987.x.
- Hibell AD, Thompson KM, Xing M, Humphrey PP, Michel AD (2001) Complexities of measuring antagonist potency at P2X(7) receptor orthologs. *J Pharmacol Exp Ther.* 296, 947-57.
- Hussy N (1992) Calcium-activated chloride channels in cultured embryonic *Xenopus* spinal neurons. *J Neurophysiol.* 68, 2042-2050.
- Iglesias R, Locovei S, Roque A, Alberto AP, Dahl G, Spray DC, Scemes E (2008) P2X7 receptor-Pannexin1 complex: pharmacology and signaling. *Am J Physiol Cell Physiol.* 295, C752-60.
- Iglesias R, Dahl G, Qiu F, Spray DC, Scemes E (2009) Pannexin 1: the molecular substrate of astrocyte "hemichannels". *J Neurosci.* 29, 7092-7097.
- Kawahara K, Takuwa N (1991) Bombesin activates large-conductance chloride channels in Swiss 3T3 fibroblasts. *Biochem Biophys Res Commun.* 177, 292-298.
- Kawate T, Michel JC, Birdsong WT, Gouaux E (2009) Crystal structure of the ATP-gated P2X(4) ion channel in the closed state. *Nature.* 460, 592-598.
- Kim M, Jiang LH, Wilson HL, North RA, Surprenant A (2001a) Proteomic and functional evidence for a P2X7 receptor signaling complex. *EMBO J* 20:6347-6358.
- Kim M, Spelta V, Sim J, North RA and Surprenant A (2001b) Differential assembly of rat purinergic P2X7 receptor in immune cells of the brain and periphery. *J Biol Chem,* 276, 23262-23267.


- Kokubun S, Saigusa A, Tamura T (1991) Blockade of Cl channels by organic and inorganic blockers in vascular smooth muscle cells. *Pflugers Arch.* 418, 204-213.
- Locovei S, Scemes E, Qiu F, Spray DC, Dahl G (2007) Pannexin1 is part of the pore forming unit of the P2X(7) receptor death complex. *FEBS Lett* 581, 483-488.
- Lundy PM, Nelson P, Mi L, Frew R, Minaker S, Vair C, Sawyer TW (2004) Pharmacological differentiation of the P2X7 receptor and the maitotoxin-activated cationic channel. *Eur J Pharmacol* 487, 17-28.
- Jacques-Silva MC, Rodnight R, Lenz G, Liao Z, Kong Q, Tran M, Kang Y, Gonzalez FA, Weisman GA, Neary JT (2004) P2X7 receptors stimulate AKT phosphorylation in astrocytes. *Br J Pharmacol* 141, 1106-1117.
- Jiang LH (2009) Inhibition of P2X(7) receptors by divalent cations: old action and new insight. *Eur Biophys J.* 38, 339-346.
- Jiang LH, Rassendren F, Mackenzie A, Zhang YH, Surprenant A, North RA (2005) N-methyl-D-glucamine and propidium dyes utilize different permeation pathways at rat P2X(7) receptors. *Am J Physiol Cell Physiol.* 289, C1295-302.
- Le Stunff H, Auger R, Kanellopoulos J, Raymond MN (2004) The Pro-451 to Leu polymorphism within the C-terminal tail of P2X7 receptor impairs cell death but not phospholipase D activation in murine thymocytes. *J Biol Chem.* 279, 16918-16926.
- Le Stunff H, Raymond MN (2007) P2X7 receptor-mediated phosphatidic acid production delays ATP-induced pore opening and cytolysis of RAW 264.7 macrophages. *Cell Signal* 19, 1909-1918.
- Lemaire I, Falzoni S, Leduc N, Zhang B, Pellegatti P, Adinolfi E, Chiozzi P, Di Virgilio F (2006) Involvement of the purinergic P2X7 receptor in the formation of multinucleated giant cells. *J Immunol.* 177, 7257-7265.
- Li Q, Luo X, Zeng W, Muallem S (2003) Cell-specific behavior of P2X7 receptors in mouse parotid acinar and duct cells. *J Biol Chem.* 278, 47554-47561.
- Li M, Chang TH, Silberberg SD, Swartz KJ (2008) Gating the pore of P2X receptor channels. *Nat Neurosci* 11, 883-887.
- Li M, Kawate T, Silberberg SD, Swartz KJ (2010) Pore-opening mechanism in trimeric P2X receptor channels. *Nat Commun* 1, 44.
- Li Q, Luo X, Muallem S (2005) Regulation of the P2X7 receptor permeability to large molecules by extracellular Cl⁻ and Na⁺. *J Biol Chem* 280, 26922-26927.
- Liu Y, Xiao Y, Li Z (2011) P2X7 receptor positively regulates MyD88-dependent NF- κ B activation. *Cytokine* 55, 229-236.
- Liu J, Ek Vitorin JF, Weintraub ST, Gu S, Shi Q, Burt JM, Jiang JX (2011) Phosphorylation of connexin 50 by protein kinase A enhances gap junction and hemichannel function. *J Biol Chem.* 286, 16914-16928.
- Locovei S, Wang J, Dahl G (2006) Activation of pannexin 1 channels by ATP through P2Y receptors and by cytoplasmic Ca²⁺. *FEBS Lett* 580, 239-244.
- Ma W, Korngreen A, Weil S, Cohen EB, Priel A, Kuzin L, Silberberg SD (2006) Pore properties and pharmacological features of the P2X receptor channel in airway ciliated cells. *J Physiol* 571, 503-517.
- Ma W, Hui H, Pelegrin P, Surprenant A (2009) Pharmacological characterization of pannexin-1 currents expressed in mammalian cells. *J Pharmacol Exp Ther.* 328, 409-418.

- Marcoli M, Cervetto C, Paluzzi P, Guarnieri S, Alloisio S, Thellung S, Nobile M, Maura G (2008) P2X7 pre-synaptic receptors in adult rat cerebrocortical nerve terminals: a role in ATP-induced glutamate release. *J Neurochem.* 105, 2330-2342.
- Marques-da-Silva C, Chaves MM, Castro NG, Coutinho-Silva R, Guimaraes MZ (2011) Colchicine inhibits cationic dye uptake induced by ATP in P2X2 and P2X7 receptor-expressing cells: implications for its therapeutic action *Br J Pharmacol.* 163, 912-926
- Masin M, Young C, Lim K, Barnes SJ, Xu XJ, Marschall V, Brutkowski W, Mooney ER, Gorecki DC, Murrell-Lagnado R (2011) Expression, assembly and function of novel C-terminal truncated variants of the mouse P2X7 receptor: Re-evaluation of P2X7 knockouts. *Br J Pharmacol.* Aug 12. doi: 10.1111/j.1476-5381.2011.01624.x.
- Michel AD, Chessell IP, Humphrey PP (1999) Ionic effects on human recombinant P2X7 receptor function. *Naunyn Schmiedebergs Arch Pharmacol.* 359, 102-109.
- Michel AD, Thompson KM, Simon J, Boyfield I, Fonfria E, Humphrey PP (2006) Species and response dependent differences in the effects of MAPK inhibitors on P2X(7) receptor function. *Br J Pharmacol* 149, 948-957
- Michel AD, Fonfria E (2007) Agonist potency at P2X7 receptors is modulated by structurally diverse lipids. *Br J Pharmacol.* 152, 523-537.
- Michel AD, Clay WC, Ng SW, Roman S, Thompson K, Condreay JP, Hall M, Holbrook J, Livermore D, Senger S (2008) Identification of regions of the P2X(7) receptor that contribute to human and rat species differences in antagonist effects. *Br J Pharmacol.* 155, 738-751.
- Montell C (2005) The TRP superfamily of cation channels. *Sci STKE* 2005(272):re3.
- Moon H, Na HY, Chong KH, Kim TJ (2006) P2X7 receptor-dependent ATP-induced shedding of CD27 in mouse lymphocytes. *Immunol Lett.* 102, 98-105.
- Moran MM, McAlexander MA, Biró T, Szallasi A (2011) Transient receptor potential channels as therapeutic targets. *Nat Rev Drug Discov.* 2011 Aug 1;10(8):601-20. doi: 10.1038/nrd3456.
- Morelli A, Chiozzi P, Chiesa A, Ferrari D, Sanz JM, Falzoni S, Pinton P, Rizzuto R, Olson MF, Di Virgilio F (2003) Extracellular ATP causes ROCK I-dependent bleb formation in P2X7-transfected HEK293 cells. *Mol Biol Cell* 14, 2655-2664.
- Moriyama T, Iida T, Kobayashi K, Higashi T, Fukuoka T, Tsumura H, Leon C, Suzuki N, Inoue K, Gachet C, Noguchi K, Tominaga M (2003) Possible involvement of P2Y2 metabotropic receptors in ATP-induced transient receptor potential vanilloid receptor 1-mediated thermal hypersensitivity. *J Neurosci.* 23, 6058-6062.
- Moriyama T, Higashi T, Togashi K, Iida T, Segi E, Sugimoto Y, Tominaga T, Narumiya S, Tominaga M (2005) Sensitization of TRPV1 by EP1 and IP reveals peripheral nociceptive mechanism of prostaglandins. *Mol Pain* 17, 1:3.
- Nagasawa K, Miyaki J, Kido Y, Higashi Y, Nishida K, Fujimoto S (2009a) Possible involvement of PPAR gamma in the regulation of basal channel opening of P2X7 receptor in cultured mouse astrocytes. *Life Sci* 84, 825-831.
- Nagasawa K, Escartin C, Swanson RA (2009b) Astrocyte cultures exhibit P2X7 receptor channel opening in the absence of exogenous ligands. *Glia.* 57, 622-633.
- Nicke A, Bäumer HG, Rettinger J, Eichele A, Lambrecht G, Mutschler E, Schmalzing G (1998) P2X1 and P2X3 receptors form stable trimers: a novel structural motif of ligand-gated ion channels. *EMBO J* 17, 3016-3028

- Nicke A, Kerschensteiner D, Soto F (2005) Biochemical and functional evidence for heteromeric assembly of P2X1 and P2X4 subunits. *J Neurochem* 92, 925–933.
- Nicke A (2008) Homotrimeric complexes are the dominant assembly state of native P2X7 subunits. *Biochem Biophys Res Commun.* 377, 803-808.
- North RA (2002) Molecular physiology of P2X receptors. *Physiol Rev* 82, 1013–1067.
- Orinska Z, Hein M, Petersen F, Bulfone-Paus S, Thon L, Adam D (2011) Retraction: Extracellular ATP induces cytokine expression and apoptosis through P2X7 receptor in murine mast cells. *J Immunol.* 186, 2683.
- Ortega F, Pérez-Sen R, Delicado EG, Miras-Portugal MT (2009) P2X7 nucleotide receptor is coupled to GSK-3 inhibition and neuroprotection in cerebellar granule neurons. *Neurotox Res.* 15(3):193-204.
- Pelegrin P, Surprenant A (2006) Pannexin-1 mediates large pore formation and interleukin-1beta release by the ATP-gated P2X7 receptor. *EMBO J* 25, 5071–5082.
- Pelegrin P, Surprenant A (2007) Pannexin-1 couples to maitotoxin- and nigericin-induced interleukin-1beta release through a dye uptake-independent pathway. *J Biol Chem.* 282, 2386-2394.
- Pérez-Andrés E, Fernández-Rodríguez M, González M, Zubiaga A, Vallejo A, García I, Matute C, Pochet S, Dehaye JP, Trueba M, Marino A, Gómez-Muñoz A (2002) Activation of phospholipase D-2 by P2X(7) agonists in rat submandibular gland acini. *J Lipid Res.* 43, 1244-1255.
- Persechini PM, Bisaggio RC, Alves-Neto JL, Coutinho-Silva R (1998) Extracellular ATP in the lymphohematopoietic system: P2Z purinoceptors off membrane permeabilization. *Braz J Med Biol Res.* 31, 25-34.
- Petrou S, Ugur M, Drummond RM, Singer JJ, Walsh JV Jr (1997) P2X7 purinoceptor expression in *Xenopus* oocytes is not sufficient to produce a pore-forming P2Z-like phenotype. *FEBS Lett.* 411, 339-345.
- Pochet S, Garcia-Marcos M, Seil M, Otto A, Marino A, Dehaye JP (2007) Contribution of two ionotropic purinergic receptors to ATP responses in submandibular gland ductal cells. *Cell Signal.* 19, 2155-2164.
- Ralevic V, Burnstock G (1998) Receptors for purines and pyrimidines. *Pharmacol Rev.* 50, 413-92
- Retamal MA, Yin S, Altenberg GA, Reuss L (2009) Modulation of Cx46 hemichannels by nitric oxide. *Am J Physiol Cell Physiol.* 296, C1356-63.
- Reyes JP, Pérez-Cornejo P, Hernández-Carballo CY, Srivastava A, Romanenko VG, Gonzalez-Begne M, Melvin JE, Arreola J (2008) Na⁺ modulates anion permeation and block of P2X7 receptors from mouse parotid glands. *J Membr Biol.* 223, 73-85.
- Riedel T, Schmalzing G, Markwardt F (2007) Influence of extracellular monovalent cations on pore and gating properties of P2X7 receptor-operated single-channel currents. *Biophys J.* 93, 846-58.
- Roger S, Pelegrin P, Surprenant A (2008) Facilitation of P2X7 receptor currents and membrane blebbing via constitutive and dynamic calmodulin binding. *J Neurosci* 28, 6393–6401.
- Roger S, Gillet L, Baroja-Mazo A, Surprenant A, Pelegrin P (2010) C-terminal calmodulin-binding motif differentially controls human and rat P2X7 receptor current facilitation. *J Biol Chem.* 285, 17514-17524.

- Saigusa A, Kokubun S (1988) Protein kinase C may regulate resting anion conductance in vascular smooth muscle cells. *Biochem Biophys Res Commun.* 155, 882-889.
- Sabirov RZ, Okada Y (2004) Wide nanoscopic pore of maxi-anion channel suits its function as an ATP-conductive pathway. *Biophys J.* 87, 1672-1685.
- Sabirov RZ, Sheiko T, Liu H, Deng D, Okada Y, Craigen WJ (2006) Genetic demonstration that the plasma membrane maxi-anion channel and voltage-dependent anion channels are unrelated proteins. *J Biol Chem* 281, 1897-1904.
- Schachter J, Motta AP, de Souza Zamorano A, da Silva-Souza HA, Guimarães MZ, Persechini PM (2008) ATP-induced P2X7-associated uptake of large molecules involves distinct mechanisms for cations and anions in macrophages. *J Cell Sci.* 121, 3261-3270.
- Schalper KA, Palacios-Prado N, Retamal MA, Shoji KF, Martínez AD, Sáez JC (2008) Connexin hemichannel composition determines the FGF-1-induced membrane permeability and free $[Ca^{2+}]_i$ responses. *Mol Biol Cell.* 19, 3501-3513.
- Schilling WP, Wasylyna T, Dubyak GR, Humphreys BD, Sinkins WG (1999a) Maitotoxin and P2Z/P2X(7) purinergic receptor stimulation activate a common cytolytic pore. *Am J Physiol.* 277, C766-76.
- Schilling WP, Sinkins WG, Estacion M (1999b) Maitotoxin activates a nonselective cation channel and a P2Z/P2X(7)-like cytolytic pore in human skin fibroblasts. *Am J Physiol.* 277, C755-65.
- Schwiebert EM, Kizer N, Gruenert DC, Stanton BA (1992) GTP-binding proteins inhibit cAMP activation of chloride channels in cystic fibrosis airway epithelial cells. *Proc Natl Acad Sci U S A.* 89, 10623-10627.
- Shemon AN, Sluyter R, Conigrave AD, Wiley JS (2004) Chelerythrine and other benzophenanthridine alkaloids block the human P2X7 receptor. *Br J Pharmacol* 142, 1015-1019.
- Shemon AN, Sluyter R, Fernando SL, Clarke AL, Dao-Ung LP, Skarratt KK, Saunders BM, Tan KS, Gu BJ, Fuller SJ, Britton WJ, Petrou S, Wiley JS (2006) A Thr357 to Ser polymorphism in homozygous and compound heterozygous subjects causes absent or reduced P2X7 function and impairs ATP-induced mycobacterial killing by macrophages. *J Biol Chem.* 281, 2079-2086.
- Smart ML, Gu B, Panchal RG, Wiley J, Cromer B, Williams DA, Petrou S (2003) P2X7 receptor cell surface expression and cytolytic pore formation are regulated by a distal C-terminal region. *J Biol Chem* 278, 8853-8860.
- Sperlágh B, Vizi ES, Wirkner K, Illes P (2006) P2X7 receptors in the nervous system. *Prog Neurobiol.* 78, 327-346.
- Suadicaní SO, Iglesias R, Spray DC, Scemes E (2009) Point mutation in the mouse P2X7 receptor affects intercellular calcium waves in astrocytes. *ASN Neuro.* 2009 Apr 14;1(1). pii: e00005. doi: 10.1042/AN20090001.
- Sugiura T, Tominaga M, Katsuya H, Mizumura K (2002) Bradykinin lowers the threshold temperature for heat activation of vanilloid receptor 1. *J Neurophysiol.* 88, 544-548.
- Sun XP, Supplisson S, Mayer E (1993) Chloride channels in myocytes from rabbit colon are regulated by a pertussis toxin-sensitive G protein. *Am J Physiol* 264, G774-85.
- Surprenant A (1996) Functional properties of native and cloned P2X receptors. *Ciba Found Symp* 198:208-219; discussion 219-222.

- Takenouchi T, Ogihara K, Sato M, Kitani H (2005) Inhibitory effects of U73122 and U73343 on Ca²⁺ influx and pore formation induced by the activation of P2X7 nucleotide receptors in mouse microglial cell line. *Biochim Biophys Acta*. 1726, 177-186.
- Takenouchi T, Sato M, Kitani H (2007) Lysophosphatidylcholine potentiates Ca²⁺ influx, pore formation and p44/42 MAP kinase phosphorylation mediated by P2X7 receptor activation in mouse microglial cells. *J Neurochem*. 102, 1518-1532.
- Thimm J, Mechler A, Lin H, Rhee S, Lal R (2005) Calcium-dependent open/closed conformations and interfacial energy maps of reconstituted hemichannels. *J Biol Chem*. 280, 10646-10654.
- Tominaga M, Wada M, Masu M (2001) Potentiation of capsaicin receptor activity by metabotropic ATP receptors as a possible mechanism for ATP-evoked pain and hyperalgesia. *Proc Natl Acad Sci U S A*. 98, 6951-6956.
- Toychiev AH, Sabirov RZ, Takahashi N, Ando-Akatsuka Y, Liu H, Shintani T, Noda M, Okada Y (2009) Activation of maxi-anion channel by protein tyrosine dephosphorylation. *Am J Physiol Cell Physiol*. 297, C990-1000.
- Vaca L, Kunze DL (1993) Depletion and refilling of intracellular Ca²⁺ stores induce oscillations of Ca²⁺ current. *Am J Physiol*. 264, H1319-22.
- Virginio C, Church D, North RA, Surprenant A (1997) Effects of divalent cations, protons and calmidazolium at the rat P2X7 receptor. *Neuropharmacology* 36, 1285-1294.
- Virginio C, MacKenzie A, Rassendren FA, North RA, Surprenant A (1999) Pore dilation of neuronal P2X receptor channels. *Nat Neurosci* 2, 315-321.
- Wilson HL, Wilson SA, Surprenant A, North RA (2002) Epithelial membrane proteins induce membrane blebbing and interact with the P2X7 receptor C terminus. *J Biol Chem* 277:34017-34023.
- Wisnoskey BJ, Estacion M, Schilling WP (2004) Maitotoxin-induced cell death cascade in bovine aortic endothelial cells: divalent cation specificity and selectivity. *Am J Physiol Cell Physiol*. 287, C345-56.
- Yan Z, Li S, Liang Z, Tomić M, Stojilkovic SS (2008) The P2X7 receptor channel pore dilates under physiological ion conditions. *J Gen Physiol*. 132, 563-573.
- Yan Z, Khadra A, Sherman A, Stojilkovic SS (2011) Calcium-dependent block of P2X7 receptor channel function is allosteric. *J Gen Physiol*. 138, 437-452.
- Yehezkel G, Hadad N, Zaid H, Sivan S, Shoshan-Barmatz V (2006) Nucleotide-binding sites in the voltage-dependent anion channel: characterization and localization. *J Biol Chem*. 281, 5938-5946.
- Young RC, McLaren M, Ramsdell JS (1995) Maitotoxin increases voltage independent chloride and sodium currents in GH4C1 rat pituitary cells. *Nat Toxins*. 3, 419-427.
- Young MT, Pelegrin P, Surprenant A (2006) Identification of Thr283 as a key determinant of P2X7 receptor function. *Br J Pharmacol* 149, 261-268.
- Zhao Q, Yang M, Ting AT, Logothetis DE (2007) PIP(2) regulates the ionic current of P2X receptors and P2X(7) receptor-mediated cell death. *Channels (Austin)* 1:46-55.
- Zizi M, Forte M, Blachly-Dyson E, Colombini M (1994) NADH regulates the gating of VDAC, the mitochondrial outer membrane channel. *J Biol Chem* 269, 1614-1626.



Patch Clamp Technique

Edited by Prof. Fatima Shad Kaneez

ISBN 978-953-51-0406-3

Hard cover, 356 pages

Publisher InTech

Published online 23, March, 2012

Published in print edition March, 2012

This book is a stimulating and interesting addition to the collected works on Patch clamp technique. Patch Clamping is an electrophysiological technique, which measures the electric current generated by a living cell, due to the movement of ions through the protein channels present in the cell membrane. The technique was developed by two German scientists, Erwin Neher and Bert Sakmann, who received the Nobel Prize in 1991 in Physiology for this innovative work. Patch clamp technique is used for measuring drug effect against a series of diseases and to find out the mechanism of diseases in animals and plants. It is also most useful in finding out the structure function activities of compounds and drugs, and most leading pharmaceutical companies used this technique for their drugs before bringing them for clinical trial. This book deals with the understanding of endogenous mechanisms of cells and their receptors as well as advantages of using this technique. It covers the basic principles and preparation types and also deals with the latest developments in the traditional patch clamp technique. Some chapters in this book take the technique to a next level of modulation and novel approach. This book will be of good value for students of physiology, neuroscience, cell biology and biophysics.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

L.G.B. Ferreira, R.A.M. Reis, L.A. Alves and R.X. Faria (2012). Intracellular Signaling Pathways Integrating the Pore Associated with P2X7R Receptor with Other Large Pores, Patch Clamp Technique, Prof. Fatima Shad Kaneez (Ed.), ISBN: 978-953-51-0406-3, InTech, Available from: <http://www.intechopen.com/books/patch-clamp-technique/intracellular-signaling-pathways-integrating-the-pore-associated-to-p2x7-receptor-with-other-large-p>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen