

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Electrophysiological Techniques for Mitochondrial Channels

Rainer Schindl¹ and Julian Weghuber^{2,*}

¹*Institute of Biophysics, Johannes Kepler University, Linz*

²*University of Applied Sciences Upper Austria, Wels
Austria*

1. Introduction

The patch-clamp technique has revolutionized studies of ion channels in a wide variety of cells. Its development in the late 1970s and early 1980s made it possible to analyze the currents of single ion channels. However, electrophysiological recordings of these proteins were limited to the plasma-membrane of living cells: the small size and the double-membrane organization of some cell organelles and the reduced stability of intracellular membranes precluded an implementation of this technique to investigate the properties of channels located in mitochondria, chloroplasts, the endoplasmic reticulum or lysosomes. For the last 25 years great effort has been made to implement the patch-clamp technique besides the planar lipid bilayer approach on these organelles. In this mini-review we focus on the developments and applications of the patch-clamp technique on mitochondrial and other cytosolic membranes and discuss the challenges of different, innovative approaches.

2. First attempts

The patch-clamp technique developed by Neher and Sakmann in 1976 as an effective refinement of the voltage-clamp-assay proved the function of single ion channels in fundamental cellular processes (Neher and Sakmann, 1976). The effectiveness of this technique to study plasma-membrane localized channels soon became of interest to characterize putative ion channels in intra-cellular membranes. Special attention was paid to mitochondria and their main function to provide the cell with energy in the form of ATP. Still in 1976, members from the lab of Alan Finkelstein inserted mitochondrial membranes into planar lipid bilayers using a Triton X-100 extract of rat liver mitochondria (Schein *et al.*, 1976). Single channels were directly detected by elevated ion permeability through the bilayer. This innovative work and further studies in the following years (Roos *et al.*, 1982) provided a good deal of information on the probable behavior of the voltage-dependent anion channel (VDAC, "mitochondrial porin"). However, the permeability properties of the intact outer mitochondrial membrane (OMM) remained hidden. In 1987, Tedeschi and colleagues managed to fill gaps in information about the electrical characterization of the OMM (Tedeschi *et al.*, 1987): they successfully introduced the patch-clamp technique on

* Corresponding Author

outer membrane vesicles isolated from mitochondria of the fungus *Neurospora crassa* and giant mitochondria isolated from cuprizone-fed mice. The copper chelator Cuprizone (biscyclohexanone oxaldihydrazone) had been shown to induce an increase of the size of mitochondria more than twofold (Flatmark *et al.*, 1980). Their results confirmed the presence of the VDAC, which facilitates the exchange of ions and molecules up to 5 kDa. Kathleen W. Kinnally, a pioneer in the field of mitochondrial electrophysiology, and co-workers continued their work on channels of the OMM. Among other findings they presented first studies on the kinetics of conductance changes induced by negative voltages and speculated on the multimerization of the high-conductance channel in the OMM (Kinnally *et al.*, 1989b). It took more than 30 years after its initial discovery, to unravel the structure of the VDAC channel by NMR spectroscopy and X-ray crystallographic techniques (Bayrhuber *et al.*, 2008; Hiller *et al.*, 2008; Ujwal *et al.*, 2008).

Apart from the progress that had been made on channels of the OMM, the importance of putative channels in the inner mitochondrial membrane (IMM) was neglected. Until the late eighties it was widely believed that the IMM was unlikely to contain any channels. The small size and the double-membrane structure of mitochondria had prohibited researchers from using the patch-clamp technique on the IMM. This prevailing opinion was finally disproved by the group of Walter Stühmer in 1987 by patch-clamp recordings of the inner mitochondrial membranes (IMM) (Sorgato *et al.*, 1987). They circumvented the restriction in size and the double-membraned organization by removing the OMM using a swelling-shrinking-sonication procedure from mitochondria isolated from cuprizone-fed mice. This led to the formation of vesicles termed mitoplasts with diameters of 3-6 μm . Measurements in whole mitoplast configuration as well as single channel recordings resulted in the identification of a voltage-dependent, anion permeable channel. The properties of this 107 pS channel, which was later found to correspond to the Inner Membrane Anion channel (IMAC) (Borecky *et al.*, 1997; Schonfeld *et al.*, 2004), were reported to be remarkably different from those of the VDAC of the OMM with a calculated mean conductance of 480 pS (Roos *et al.*, 1982).

3. Pros and cons of electrophysiological methods

In order to characterize the electrophysiological properties of single channels, two methods are commonly used: a voltage-clamp assay using planar lipid bilayers and the glass-pipette based patch-clamp method. The former one is based on the incorporation of purified membrane vesicles into the bilayer. The method allows studying channels, which are not easily accessible for the patch-clamp technique. Thus, the IMM but also other intracellular organelles including endosomes, lysosomes or the Golgi-apparatus may be analyzed. Furthermore this method is capable of detecting changes in the channel properties upon modulation by other proteins or lipids, i.e. variations in the lipid or protein composition of the bilayer. The major advantage of the patch-clamp method is the suitability to study channels in their natural environment, as the isolation process for the bilayer method may be harmful for the channel. Exemplarily, the recorded VDAC currents through single channels were 350 pS in patch-clamp recordings of whole mitochondria (Sorgato *et al.*, 1987), while a single channel conductance of 480 pS was found for the same channel after reconstitution in lipid bilayers (Roos *et al.*, 1982). The differences in the conductance might originate from a distinct membrane environment. However, the purity of the vesicle preparation is crucial due to the different fusion probability of vesicles from various membranes. This problem can be partly solved by using vesicles lacking any other

contaminating membranes. Disadvantages of the patch-clamp technique should be mentioned at this point as well: first, mitochondria and possibly other cell organelles are in close contact with membranes of the endoplasmic reticulum, forming so-called “mitochondria associated membranes” (Garcia-Perez *et al.*, 2008). It might be difficult to completely remove these membranes. Second, some membranes, e.g. mitochondrial Cristae tubules, might not be accessible for the glass pipette (Zick *et al.*, 2009). Thus, the patch clamp technique might fail to detect some important ion channel currents.

4. Mitochondrial electrophysiology in the nineties

A report of the group of T. Higuti in 1991 can be considered another milestone in mitochondrial electrophysiology: they characterized a highly K⁺ selective, low conductance channel in the IMM, using mitochondria isolated from fresh liver of rats (Inoue *et al.*, 1991). To date the activity identified as corresponding to that of the ATP-sensitive K⁺ channel (mito KATP or mtK_{ATP}) remains a matter of controversy. The molecular composition of the mito KATP has not been unraveled and some reports challenge the presence of this channel in the IMM (Brustovetsky *et al.*, 2005; Das *et al.*, 2003), while most studies support its existence (Dahlem *et al.*, 2004; Zhang *et al.*, 2001). However, the chosen patch-clamp strategy for the IMM appeared promising: To circumvent the double-membrane structure and small size of mitochondria, they removed the OMM by digitonin treatment and fused the generated giant mitoplasts using a low-pH solution containing Ca²⁺. This method does not require additional lipids necessary for the vesicle fusion process that could influence the channel in its native environment. In addition the achieved size of the fused mitoplasts of up to 15 μm allows for a robust giga-seal formation with the patch-pipette tip.

A first report on the mitochondrial multiconductance channel by Kinnally and co-workers in 1989 (Kinnally *et al.*, 1989a) opened up a new chapter in mitochondrial electrophysiology, which has not lost its relevance. The MCC, also termed mitochondrial megachannel, was later described as the mitochondrial permeability transition pore (MPT pore or MPTP), which is formed under certain pathological conditions (Lemasters *et al.*, 2009). At that time, first amphiphilic drugs were found to affect the mitochondrial permeability (Beavis and Powers, 1989; Beavis, 1989), which was regarded as a prerequisite for the study of channels in the IMM (Antonenko *et al.*, 1991; Gunter and Pfeiffer, 1990). Electrophysiological data on mitoplasts created in the lab of Mario Zoratti proposed the MCC may comprise VDAC molecules (Szabo *et al.*, 1993). Further biochemical investigations by Andrew Halestrap *et al.* and electrophysiology studies performed in the Kinnally-lab identified the adenine nucleotide translocase (ANT) as the central unit forming the MPTP (Halestrap *et al.*, 1998; Lohret *et al.*, 1996). Although these findings have been challenged by studies published only a few years ago (Baines *et al.*, 2007; Kokoszka *et al.*, 2004), the efforts of these few pioneers during the nineties accelerated the progress for a better understanding of the complex behavior of mitochondrial channels.

In contrast to the work of Inoue *et al.*, 1991 (Inoue *et al.*, 1991) on the mito KATP channel, most researchers did not work on fused mitoplasts, but continued to directly patch-clamp whole mitoplasts prepared from mammalian cell lines or tissue. These included a glioma cell line (Siemen *et al.*, 1999), an osteosarcoma cell line (Murphy *et al.*, 1998), rat liver tissue (Szabo *et al.*, 1992) or brown adipose tissue from hamsters (Borecky *et al.*, 1997). However, only mitochondria isolated from a limited number of mammalian and other eukaryotic cells

formed mitoplasts with an adequate size. Thus, most IMM channel activities were recorded after reconstitution into proteoliposomes or giant liposomes (Lohret *et al.*, 1996;Lohret *et al.*, 1997;Paliwal *et al.*, 1992). This holds especially true for studies of yeast mitochondrial channels. The small size of mitoplasts from this organism (0.3-1.5 μm), either isolated by osmotic swelling or using the French press method, makes the patch clamp technique very difficult. The advantages of the model-system yeast, e.g. fast growth, easy handling and especially the possibility of a fast gene knock-out, justify the technically challenging and laborious reconstitution process. Still, the purity of the used membrane fractions and a potential influence of externally added lipids for the generation of giant liposomes must be taken into account.

5. Mitochondrial electrophysiology over the past 10 years

We may start with an outstanding finding published in 2004 (Kirichok *et al.*, 2004). Yuriy Kirichok *et al.* from the lab of David Clapham described the properties of a channel consistent with the predicted mitochondrial Calcium uniporter (miCa). The experiments were accomplished by patch-clamp recordings of whole mitoplasts without any additional fusion or reconstitution processes. An interesting feature of the chosen COS-7 cell-line is that intact mitoplasts of 2-5 μm size could be prepared. For that purpose mitochondria were isolated by differential centrifugation and mitoplasts were formed by osmotic swelling. Interestingly, no additional steps to remove the OMM had been performed, which enhances the risk that remnants of the OMM or the tightly attached ER-membrane falsify the results. According to the authors such artifacts had been excluded as these structures appeared as black spots on the transparent vesicles. Taken together extreme care has to be taken and numerous technical problems have to be solved when performing such experiments. Very recently, the molecular identity of the miCa has been clarified: a channel protein termed MCU and a regulating protein named MICU1 have been reported to fulfill the requirements for the long searched mitochondrial Calcium uniporter (De *et al.*, 2011;Perocchi *et al.*, 2010).

Inspired by the work of these pioneers our group started electrophysiological studies on mitochondrial Magnesium transport. In 2003-2006 we had shown, using Magnesium selective fluorescent dyes, that the IMM localized protein Mrs2 is a major player for the influx of Magnesium into yeast mitochondria (Kolisek *et al.*, 2003;Weghuber *et al.*, 2006). At that time we could not exclude that Mrs2p forms a transporter rather than a high-conductance channel. To answer this question we generated mitoplasts using osmotic swelling or the French press method. However, consistent with the work performed in the Kinnally lab in the 90ies, the small size of yeast < 1 μm did not allow for effective patch-clamp recordings. Thus we decided to fuse IMM with lipid vesicles: submitochondrial particles (SMPs) (Froschauer *et al.*, 2009) were mixed with asolectin vesicles. According to the report of Criado and Keller (Criado and Keller, 1987) in 1987 we finally succeeded in generating giant vesicles (5-50 μm in size) with reconstituted IMM fractions from yeast overexpressing or having *MRS2* disrupted. The used method is characterized by dehydration-rehydration cycles and has also been successfully used by other groups working on yeast mitochondrial channels (Lohret *et al.*, 1996). During our studies the crystal structure of the bacterial homologue CorA had been reported (Lunin *et al.*, 2006;Eshaghi *et al.*, 2006;Payandeh and Pai, 2006), but the mode of ion transport remained obscure. We could finally prove that Mrs2p forms a Magnesium selective channel in the IMM with a calculated conductance of 155 pS (Schindl *et al.*, 2007). We furthermore showed that co-

expression of a homologue of Mrs2p termed Lpe10p (or Mfm1p) yielded a unique, reduced conductance in comparison to the one of Mrs2p channels (Sponder *et al.*, 2010).

6. Outlook

The progress on the electrophysiological characterization of mitochondrial channels was rather limited in the nineties with only a few labs being able to deal with the emerging technical challenges. This situation has definitely changed in the last decade: while there are not more than 100 reports on patch-clamp recordings on mitochondria indexed in pubmed in the nineties, this number has changed to more than 400 in the last decade. However, there is still a lot of work to be done. The molecular identities of most channel proteins forming the MPTP, the 107 pS channel or the mito KATP have not been defined. From that point of view the studies on the yeast Magnesium channel proteins Mrs2p and its homolog Mfm1p are of even greater relevance.

The experiences in the field of protein- and membrane reconstitution and in mitochondrial electrophysiology rendered these techniques also interesting for the characterization of other intracellular membranes. First studies by a pioneer in this field Igor Pottosin in the early nineties reported on the investigation of ion channels located in chloroplast envelope membranes (Pottosin, 1993; Heiber *et al.*, 1995) or thylakoid membranes (Pottosin and Schonknecht, 1995). Further reports described the action of a voltage-gated ion channel in the outer envelope (Hinnah *et al.*, 1997), focused on the inactivation of a 50 pS envelope anion channel necessary for import of proteins into the chloroplast (van den Wijngaard *et al.*, 1999) or characterized a fast-activating channel regulating various fluxes in the native envelope of chloroplasts (Pottosin *et al.*, 2005). The methods of choice to cope with the severe technical challenges (e.g. the maintenance of the organelle integrity) were either the reconstitution of isolated membranes into lipid vesicles – mainly based on asolectin – or measurements on whole chloroplasts. In both cases great experience as well as patience appears indispensable. That is probably the main reason for the limited number of studies in this field.

To efficiently investigate intracellular ion channels, new methods will be necessary. Assays like the one presented by Schieder and co-workers (Schieder *et al.*, 2010) using solid-matrix planar glass chips, which even enables patch-clamp recordings of lysosomes, are a step in the right direction.

7. Acknowledgements

This work was supported by the Austrian Science Foundation (FWF): project P22747 to R.S. We thank Jeffrey Nesbitt (Haverford High School, PA) for critically reading the manuscript.


8. References

- Antonenko YN, Kinnally KW, Perini S, Tedeschi H, 1991. Selective effect of inhibitors on inner mitochondrial membrane channels. *FEBS Lett* 285:89-93.
- Baines CP, Kaiser RA, Sheiko T, Craigen WJ, Molkentin JD, 2007. Voltage-dependent anion channels are dispensable for mitochondrial-dependent cell death. *Nat Cell Biol* 9:550-555.

- Bayrhuber M, Meins T, Habeck M, Becker S, Giller K, Villinger S, Vonnrhein C, Griesinger C, Zweckstetter M, Zeth K, 2008. Structure of the human voltage-dependent anion channel. *Proc Natl Acad Sci U S A* 105:15370-15375.
- Beavis AD, 1989. On the inhibition of the mitochondrial inner membrane anion uniporter by cationic amphiphiles and other drugs. *J Biol Chem* 264:1508-1515.
- Beavis AD, Powers MF, 1989. On the regulation of the mitochondrial inner membrane anion channel by magnesium and protons. *J Biol Chem* 264:17148-17155.
- Borecky J, Jezek P, Siemen D, 1997. 108-pS channel in brown fat mitochondria might be identical to the inner membrane anion channel. *J Biol Chem* 272:19282-19289.
- Brustovetsky T, Shalbuyeva N, Brustovetsky N, 2005. Lack of manifestations of diazoxide/5-hydroxydecanoate-sensitive KATP channel in rat brain nonsynaptosomal mitochondria. *J Physiol* 568:47-59.
- Criado M, Keller BU, 1987. A membrane fusion strategy for single-channel recordings of membranes usually non-accessible to patch-clamp pipette electrodes. *FEBS Lett* 224:172-176.
- Dahlem YA, Horn TF, Buntinas L, Gonoï T, Wolf G, Siemen D, 2004. The human mitochondrial KATP channel is modulated by calcium and nitric oxide: a patch-clamp approach. *Biochim Biophys Acta* 1656:46-56.
- Das M, Parker JE, Halestrap AP, 2003. Matrix volume measurements challenge the existence of diazoxide/glibencamide-sensitive KATP channels in rat mitochondria. *J Physiol* 547:893-902.
- De SD, Raffaello A, Teardo E, Szabo I, Rizzuto R, 2011. A forty-kilodalton protein of the inner membrane is the mitochondrial calcium uniporter. *Nature* 476:336-340.
- Eshaghi S, Niegowski D, Kohl A, Martinez MD, Lesley SA, Nordlund P, 2006. Crystal structure of a divalent metal ion transporter CorA at 2.9 angstrom resolution. *Science* 313:354-357.
- Flatmark T, Kryvi H, Tangeras A, 1980. Induction of megamitochondria by cuprizone(biscyclohexanone oxaldihydrazone). Evidence for an inhibition of the mitochondrial division process. *Eur J Cell Biol* 23:141-148.
- Froschauer EM, Schweyen RJ, Wiesenberger G, 2009. The yeast mitochondrial carrier proteins Mrs3p/Mrs4p mediate iron transport across the inner mitochondrial membrane. *Biochim Biophys Acta* 1788:1044-1050.
- Garcia-Perez C, Hajnoczky G, Csordas G, 2008. Physical coupling supports the local Ca²⁺ transfer between sarcoplasmic reticulum subdomains and the mitochondria in heart muscle. *J Biol Chem* 283:32771-32780.
- Gunter TE, Pfeiffer DR, 1990. Mechanisms by which mitochondria transport calcium. *Am J Physiol* 258:C755-C786.
- Halestrap AP, Kerr PM, Javadov S, Woodfield KY, 1998. Elucidating the molecular mechanism of the permeability transition pore and its role in reperfusion injury of the heart. *Biochim Biophys Acta* 1366:79-94.
- Heiber T, Steinkamp T, Hinnah S, Schwarz M, Flugge Ui, Weber A, Wagner R, 1995. Ion channels in the chloroplast envelope membrane. *Biochemistry* 34:15906-15917.
- Hiller S, Garces RG, Malia TJ, Orekhov VY, Colombini M, Wagner G, 2008. Solution structure of the integral human membrane protein VDAC-1 in detergent micelles. *Science* 321:1206-1210.

- Hinnah SC, Hill K, Wagner R, Schlicher T, Soll J, 1997. Reconstitution of a chloroplast protein import channel. *EMBO J* 16:7351-7360.
- Inoue I, Nagase H, Kishi K, Higuti T, 1991. ATP-sensitive K⁺ channel in the mitochondrial inner membrane. *Nature* 352:244-247.
- Kinnally KW, Campo ML, Tedeschi H, 1989a. Mitochondrial channel activity studied by patch-clamping mitoplasts. *J Bioenerg Biomembr* 21:497-506.
- Kinnally KW, Tedeschi H, Mannella CA, FRISCH HL, 1989b. Kinetics of voltage-induced conductance increases in the outer mitochondrial membrane. *Biophys J* 55:1205-1213.
- Kirichok Y, Krapivinsky G, Clapham DE, 2004. The mitochondrial calcium uniporter is a highly selective ion channel. *Nature* 427:360-364.
- Kokoszka JE, Waymire KG, Levy SE, Sligh JE, Cai J, Jones DP, Macgregor GR, Wallace DC, 2004. The ADP/ATP translocator is not essential for the mitochondrial permeability transition pore. *Nature* 427:461-465.
- Kolisek M, Zsurka G, Samaj J, Weghuber J, Schweyen RJ, Schweigel M, 2003. Mrs2p is an essential component of the major electrophoretic Mg²⁺ influx system in mitochondria. *EMBO J* 22:1235-1244.
- Lemasters JJ, Theruvath TP, Zhong Z, Nieminen AL, 2009. Mitochondrial calcium and the permeability transition in cell death. *Biochim Biophys Acta* 1787:1395-1401.
- Lohret TA, Jensen RE, Kinnally KW, 1997. Tim23, a protein import component of the mitochondrial inner membrane, is required for normal activity of the multiple conductance channel, MCC. *J Cell Biol* 137:377-386.
- Lohret TA, Murphy RC, Drgon T, Kinnally KW, 1996. Activity of the mitochondrial multiple conductance channel is independent of the adenine nucleotide translocator. *J Biol Chem* 271:4846-4849.
- Lunin VV, Dobrovetsky E, Khutoreskaya G, Zhang R, Joachimiak A, Doyle Da, Bochkarev A, Maguire Me, Edwards AM, Koth CM, 2006. Crystal structure of the CorA Mg²⁺ transporter. *Nature* 440:833-837.
- Murphy RC, Diwan JJ, King M, Kinnally KW, 1998. Two high conductance channels of the mitochondrial inner membrane are independent of the human mitochondrial genome. *FEBS Lett* 425:259-262.
- Neher E, Sakmann B, 1976. Single-channel currents recorded from membrane of denervated frog muscle fibres. *Nature* 260:799-802.
- Paliwal R, Costa G, Diwan JJ, 1992. Purification and patch clamp analysis of a 40-pS channel from rat liver mitochondria. *Biochemistry* 31:2223-2229.
- Payandeh J, Pai EF, 2006. Crystallization and preliminary X-ray diffraction analysis of the magnesium transporter CorA. *Acta Crystallogr Sect F Struct Biol Cryst Commun* 62:148-152.
- Perocchi F, Gohil Vm, Girgis Hs, Bao Xr, McCombs Je, Palmer Ae, Mootha VK, 2010. MICU1 encodes a mitochondrial EF hand protein required for Ca(2+) uptake. *Nature* 467:291-296.
- Pottosin II, 1993. One of the chloroplast envelope ion channels is probably related to the mitochondrial VDAC. *FEBS Lett* 330:211-214.
- Pottosin Ii, Muniz J, Shabala S, 2005. fast-activating channel controls cation fluxes across the native chloroplast envelope. *J Membr Biol* 204:145-156.

- Pottosin Ii, Schonknecht G, 1995. Patch clamp study of the voltage-dependent anion channel in the thylakoid membrane. *J Membr Biol* 148:143-156.
- Roos N, Benz R, Brdiczka D, 1982. Identification and characterization of the pore-forming protein in the outer membrane of rat liver mitochondria. *Biochim Biophys Acta* 686:204-214.
- Schein Sj, Colombini M, Finkelstein A, 1976. Reconstitution in planar lipid bilayers of a voltage-dependent anion-selective channel obtained from paramecium mitochondria. *J Membr Biol* 30:99-120.
- Schieder M, Rotzer K, Bruggemann A, Biel M, Wahl-Schott C, 2010. Planar patch clamp approach to characterize ionic currents from intact lysosomes. *Sci Signal* 3:13.
- Schindl R, Weghuber J, Romanin C, Schweyen RJ, 2007. Mrs2p forms a high conductance Mg²⁺ selective channel in mitochondria. *Biophys J* 93:3872-3883.
- Schonfeld P, Sayeed I, Bohnensack R, Siemen D, 2004. Fatty acids induce chloride permeation in rat liver mitochondria by activation of the inner membrane anion channel (IMAC). *J Bioenerg Biomembr* 36:241-248.
- Siemen D, Loupatatzis C, Borecky J, Gulbins E, Lang F, 1999. Ca²⁺-activated K channel of the BK-type in the inner mitochondrial membrane of a human glioma cell line. *Biochem Biophys Res Commun* 257:549-554.
- Sorgato MC, Keller BU, Stuhmer W, 1987. Patch-clamping of the inner mitochondrial membrane reveals a voltage-dependent ion channel. *Nature* 330:498-500.
- Sponder G, Svidova S, Schindl R, Wieser S, Schweyen RJ, Romanin C, Froschauer EM, Weghuber J, 2010. Lpe10p modulates the activity of the Mrs2p-based yeast mitochondrial Mg²⁺ channel. *FEBS J* 277:3514-3525.
- Szabo I, Bernardi P, Zoratti M, 1992. Modulation of the mitochondrial megachannel by divalent cations and protons. *J Biol Chem* 267:2940-2946.
- Szabo I, De P, V, Zoratti M, 1993. The mitochondrial permeability transition pore may comprise VDAC molecules. II. The electrophysiological properties of VDAC are compatible with those of the mitochondrial megachannel. *FEBS Lett* 330:206-210.
- Tedeschi H, Mannella CA, Bowman CL, 1987. Patch clamping the outer mitochondrial membrane. *J Membr Biol* 97:21-29.
- Ujwal R, Cascio D, Colletier JP, Faham S, Zhang J, Toro L, Ping P, Abramson J, 2008. The crystal structure of mouse VDAC1 at 2.3 Å resolution reveals mechanistic insights into metabolite gating. *Proc Natl Acad Sci U S A* 105:17742-17747.
- Van den Wijngaard PW, Dabney-Smith C, Bruce BD, Vredenberg WJ, 1999. The mechanism of inactivation of a 50-pS envelope anion channel during chloroplast protein import. *Biophys J* 77:3156-3162.
- Weghuber J, Dieterich F, Froschauer EM, Svidova S, Schweyen RJ, 2006. Mutational analysis of functional domains in Mrs2p, the mitochondrial Mg²⁺ channel protein of *Saccharomyces cerevisiae*. *FEBS J* 273:1198-1209.
- Zhang DX, Chen YF, Campbell WB, Zou AP, Gross GJ, Li PL, 2001. Characteristics and superoxide-induced activation of reconstituted myocardial mitochondrial ATP-sensitive potassium channels. *Circ Res* 89:1177-1183.
- Zick M, Rabl R, Reichert AS, 2009. Cristae formation-linking ultrastructure and function of mitochondria. *Biochim Biophys Acta* 1793:5-19.



Patch Clamp Technique

Edited by Prof. Fatima Shad Kaneez

ISBN 978-953-51-0406-3

Hard cover, 356 pages

Publisher InTech

Published online 23, March, 2012

Published in print edition March, 2012

This book is a stimulating and interesting addition to the collected works on Patch clamp technique. Patch Clamping is an electrophysiological technique, which measures the electric current generated by a living cell, due to the movement of ions through the protein channels present in the cell membrane. The technique was developed by two German scientists, Erwin Neher and Bert Sakmann, who received the Nobel Prize in 1991 in Physiology for this innovative work. Patch clamp technique is used for measuring drug effect against a series of diseases and to find out the mechanism of diseases in animals and plants. It is also most useful in finding out the structure function activities of compounds and drugs, and most leading pharmaceutical companies used this technique for their drugs before bringing them for clinical trial. This book deals with the understanding of endogenous mechanisms of cells and their receptors as well as advantages of using this technique. It covers the basic principles and preparation types and also deals with the latest developments in the traditional patch clamp technique. Some chapters in this book take the technique to a next level of modulation and novel approach. This book will be of good value for students of physiology, neuroscience, cell biology and biophysics.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Rainer Schindl and Julian Weghuber (2012). Electrophysiological Techniques for Mitochondrial Channels, Patch Clamp Technique, Prof. Fatima Shad Kaneez (Ed.), ISBN: 978-953-51-0406-3, InTech, Available from: <http://www.intechopen.com/books/patch-clamp-technique/electrophysiological-techniques-for-mitochondrial-channels>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen