

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



***Vibrio cholerae* Flagellar Synthesis and Virulence**

Anastasia R. Rugel^{1,2} and Karl E. Klose¹

¹South Texas Center for Emerging Infectious Diseases, Department of Biology,
University of Texas at San Antonio, San Antonio, Texas

²Department of Microbiology and Immunology,
University of Texas Health Science Center at San Antonio, San Antonio, Texas
USA

1. Introduction

Vibrio cholerae is a Gram-negative bacterium with a single sheathed polar flagellum (Fig. 1). *V. cholerae* causes the severe diarrheal disease cholera in humans when it colonizes the small intestine and expresses various virulence factors, including cholera toxin (CT) and toxin co-regulated pilus (TCP). *V. cholerae* is also a natural inhabitant of the marine environment, where it forms biofilms on chitinous surfaces. Motility contributes to both aspects of the *V. cholerae* lifecycle. The flagellum facilitates chemotactic-directed movement toward the preferred colonization site within the intestine (Camilli and Mekalanos 1995; Butler and Camilli 2004), and also contributes to biofilm formation within the environment (Watnick and Kolter 1999). *V. cholerae* strains defective for motility are less virulent than motile strains (Guentzel and Berry 1975; Freter and O'Brien 1981; Richardson 1991). As flagellar synthesis, motility, and chemotaxis have become better understood in *V. cholerae*, it has also become clear that motility is intimately integrated into all aspects of the lifestyle of this bacterium.




Fig. 1. *Vibrio cholerae*

2. Structure

The flagellum is a motor-driven organelle present in many bacteria. Different flagellar placement and quantity are seen in different bacteria. Monotrichous bacteria have a single

polar flagellum (e.g. *V. cholerae*), lofotrichous bacteria have multiple flagella at a single pole (e.g. *Helicobacter pylori*), amphitrichous bacteria have flagella at two poles (e.g. *Campylobacter jejuni*), and peritrichous bacteria have multiple flagella emanating from the cell in all directions (e.g. *Escherichia coli*).

The base of the bacterial flagellum is composed of a secretion system related to the Type III secretion system, which facilitates export of flagellar components from the cytoplasm to the periplasm and the exterior of the cell. The basic components of the flagellum are the basal body, which extends from the cytoplasmic membrane through the periplasm and into the outer membrane (OM), connected to the flexible hook (composed of FliE) found exterior to the cell, which in turn is connected to the flagellar filament (Kojima and Blair 2004; Terashima, Kojima et al. 2008). The motor components that drive flagellar rotation are found in the cytoplasmic membrane, and the switch components (FliG, FliM, FliN) that interact with the chemotaxis signaling system and the motor (Francis, Sosinsky et al. 1994) extend into the cytoplasm from the basal body (Fig. 2.).




Fig. 2. Flagellar Motor Complex

Unlike most other bacterial flagella, the *V. cholerae* flagellum has a sheath composed of OM that coats the entire filament (Allen and Baumann 1971; Sjoblad, Emala et al. 1983; Fuerst and Perry 1988). Sheathed flagella are found in *Vibrio spp.* and a few other Gram-negative bacteria (e.g. *H. pylori*). It is hypothesized that the sheath acts as a protective covering that shields the antigenic flagellins from recognition by the host's immune response (Yoon and Mekalanos 2008). The mechanism whereby the OM is extended to cover the filament during

V. cholerae flagellar synthesis rather than the filament protruding through the OM as in other bacterial flagella is not understood.

The bacterial flagellar filament is made up of thousands of flagellin subunits, with a cap protein (FliD) at the distal end (Ikeda, Asakura et al. 1985; Ikeda, Homma et al. 1987; Homma, DeRosier et al. 1990). The structure of the *Salmonella typhimurium* flagellin FliC has been solved by cryomicroscopy. FliC is composed of domains at its N- and C-termini that interact with each other: D0 (aa 1-45 and 456-495), D1 (aa 46-180 and 408-455), and D2 (aa 181-190 and 285-407). The D2 domains, along with the D3 domain (aa 191-284) form the antigenic variable region that are present on the filament surface, (Yonekura, Maki-Yonekura et al. 2003). Interaction of the D0 and D1 domains allows the flagellins to polymerize under the cap (FliD) protein into a hollow helical filament at the growing tip of the flagellum as they are being secreted.

In contrast to most other bacteria which have filaments composed of a single flagellin subunit, *V. cholerae* has a filament composed of 5 different flagellins, FlaABCDE. These flagellins share a high degree of homology, yet only FlaA is essential for flagellar synthesis; the other four flagellins are not required for the synthesis of the filament (Klose and Mekalanos 1998). Alignment of FlaA with the other four *V. cholerae* flagellins, as well as with *S. typhimurium* FliC, reveals that the D0 and D1 domains are well-conserved, whereas the variable regions D2 and D3 are more divergent. Interestingly, the *V. cholerae* flagellins have a much shorter region corresponding to D2 and D3 (129 aa shorter) when compared to *S. typhimurium* FliC. Because this antigenic portion of the flagellins extends out from the hollow filament core, it may be that the presence of the flagellar sheath over the *V. cholerae* filament restricts the size of the antigenic region protruding from the filament.

The basal body contains the rod structure (FlgB, FlgC, FlgF and FlgG) with L (FlgH), P (FlgI), and MS rings (FliF) localized to the OM, periplasm (peptidoglycan), and cytoplasmic membranes, respectively. In *Vibrio* spp. an additional T ring is located immediately below the P ring, which is composed of the *Vibrio*-specific components MotX, MotY, and FlgT. The C-ring, which extends into the cytoplasm from the MS ring and is made up of FliG, FliM, and FliN, is difficult to preserve during microscopy and has not been visualized in its entirety in *Vibrio* spp. (Aizawa, Dean et al. 1985; Homma, Aizawa et al. 1987; Homma, Ohnishi et al. 1987; Homma, DeRosier et al. 1990; Homma, Kutsukake et al. 1990; Ueno, Oosawa et al. 1992; Francis, Sosinsky et al. 1994; Schoenhals and Macnab 1996; Terashima, Koike et al. 2010). The chemotaxis protein CheY relays information from the chemotaxis sensory system by binding to the C ring (FliM), causing the flagellum to switch rotation from counterclockwise to clockwise.

In *S. typhimurium* and *E. coli* MotA and MotB are membrane proteins that compose the motor that utilizes H⁺ motive force to drive flagellar rotation (Lloyd, Tang et al. 1996; Zhou, Lloyd et al. 1998; Zhou, Sharp et al. 1998; Braun, Poulson et al. 1999; Blair 2003). *Vibrio* spp. contain MotA and MotB homologues, alternately referred to as PomA and PomB (Dean, Macnab et al. 1984; Stader, Matsumura et al. 1986; Blair and Berg 1990; Stolz and Berg 1991; Asai, Kojima et al. 1997; Sato and Homma 2000; Sato and Homma 2000; Yorimitsu, Asai et al. 2000; Fukuoka, Yakushi et al. 2005), but they also contain *Vibrio*-specific motor proteins MotX and MotY, localized in the T ring (McCarter 1994; McCarter 1994; Okunishi, Kawagishi et al. 1996; Okabe, Yakushi et al. 2001; Okabe, Yakushi et al. 2002; Okabe,

Yakushi et al. 2005; Koerdt, Paulick et al. 2009). The *Vibrio* MotA and MotB form a membrane complex that utilizes a Na⁺ gradient (instead of H⁺ gradient) to drive flagellar rotation. A Na⁺ gradient is required to allow MotA/MotB to associate with the flagellum (through MotX/MotY) and open the Na⁺ channel; flux of Na⁺ through the channel provides the torque to generate flagellar rotation (McCarter 1994; McCarter 1994; Yorimitsu, Kojima et al. 2004; Terashima, Fukuoka et al. 2006).

Two additional proteins control flagellar number and placement in *Vibrio* spp. FlhG contains an ATPase motif and controls flagellar number; *Vibrio* cells without *flhG* synthesize multiple polar flagella, instead of a single polar flagellum (Correa, Peng et al. 2005; Kusumoto, Kamisaka et al. 2006; Kusumoto, Shinohara et al. 2008). FlhF contains a GTP binding motif and localizes to the cell pole, thus dictating polar localization of the flagellum. *Vibrio* cells without *flhF* are largely non-flagellated; however a few cells will synthesize a flagellum at a site away from the pole (Carpenter, Hanlon et al. 1992; Zanen, Antelmann et al. 2004; Salvetti, Ghelardi et al. 2007; Green, Kahramanoglou et al. 2009; Kusumoto, Nishioka et al. 2009). FlhG interacts with FlhF, and a current model suggests that FlhG interacts with FlhF to prevent additional FlhF deposition at the pole (Kusumoto, Shinohara et al. 2008). A *V. alginolyticus* strain lacking both FlhF and FlhG is mostly lacking flagella (Kojima, Nishioka et al. 2011), but a few cells possess multiple peritrichous flagella (similar to *S. typhimurium*). An unidentified suppressor mutation can lead to virtually all *flhFG* *V. alginolyticus* cells possessing peritrichous flagella and being able to swim; the identification of this suppressor mutation should lead to greater insights into control of polar flagellar synthesis in *Vibrio* spp.

Two additional outer membrane proteins, FlgO and FlgP, contribute to flagellar stability. FlgP homologues are restricted to *Vibrio*, *Helicobacter*, and *Campylobacter* spp. *V. cholerae* FlgP is a lipoprotein that affects flagellar stability; *flgP* mutants synthesize fragile flagella and appear non-motile in motility agar, presumably due to breakage of flagella during swimming (Morris, Peng et al. 2008; Martinez, Dharmasena et al. 2009). FlgO homologues are only found in *Vibrio* spp. *V. cholerae* strains lacking *flgO* have a similar phenotype as *flgP* strains, namely they produce fragile flagella that break easily while swimming (Morris, Peng et al. 2008; Martinez, Dharmasena et al. 2009).

3. Regulation

Transcription of the *V. cholerae* flagellar genes is controlled by a four-tiered transcription hierarchy (Fig. 3.) (Prouty, Correa et al. 2001). The *V. cholerae* flagellar transcription hierarchy is similar to that which controls flagellar transcription in *Pseudomonas aeruginosa*, another bacterium with a single polar flagellum (Dasgupta, Wolfgang et al. 2003). The master regulator, FlrA, is a σ^{54} -dependent transcriptional activator. FlrA represents the sole Class I gene product, and it activates transcription of Class II flagellar genes (Klose and Mekalanos 1998). It is not clear whether environmental conditions regulate transcription of *flrA*, but *flhG* (which controls flagellar number) also negatively regulates *flrA* transcription (Correa, Peng et al. 2005).

The *P. aeruginosa* FlrA homologue, FleQ, has been shown to bind to cyclic-di-GMP (cdGMP) (Hickman and Harwood 2008). Binding of cdGMP to FleQ prevents DNA binding, resulting in the absence of flagellar synthesis and de-repression in *P. aeruginosa* of genes involved in biofilm formation normally repressed by FleQ. Interestingly, cdGMP binds to

FleQ lacking the N-terminus, indicating it binds to the transcriptional activation/DNA binding domain, which shares high homology (63% identity) with *V. cholerae* FlrA. It is not yet known whether FlrA binds to and is modulated by cdGMP. *P. aeruginosa* FleQ also binds to FleN, the homologue of FlhG (Dasgupta and Ramphal 2001). FleN binding to FleQ does not inhibit DNA binding, but downregulates FleQ-dependent transcription, resulting in reduced (single) flagellar number. As mentioned above, FlhG has a negative effect on *flrA* transcription in *V. cholerae*, but it is not known whether it also binds to FlrA and negatively affects its activity.




Fig. 3. Flagellar Transcription Regulatory Hierarchy

FlrA positively regulates Class II flagellar genes. Both FlrA and σ^{54} -containing RNA polymerase are required to activate transcription of the Class II flagellar genes (Klose and Mekalanos 1998; Klose, Novik et al. 1998; Prouty, Correa et al. 2001). The class II genes encode components of the MS ring-switch-export apparatus as well as chemotaxis and regulatory proteins. Two large flagellar operons (*fliEFGHIJ* and the *flhA* operon, which contains *flhFG*, mentioned above, as well as *fliA* (σ^{28}) and a number of chemotaxis genes), and the regulatory genes *flrBC*, are activated by FlrA. The Class II flagellar genes are predicted to encode an export apparatus-basal body intermediate; it seems likely that this structure is required to be assembled prior to progression to Class III gene expression, as is the case in *Campylobacter jejuni* and *Helicobacter pylori*, which have similar classes of flagellar genes (Hendrixson and DiRita 2003; Niehus, Gressmann et al. 2004).

The regulatory proteins FlrBC are a two-component system that controls Class III gene transcription (Prouty, Correa et al. 2001). FlrB undergoes autophosphorylation, and then activates FlrC activity by transferring a phosphate to the conserved aspartate-54 (D54) residue in the amino terminus of FlrC (FlrC-P) allowing it to activate the σ^{54} -dependent transcription of Class III genes (Correa, Lauriano et al. 2000; Correa and Klose 2005). The class III genes encode the rest of the components of the hook-basal body, as well as the flagellin FlaA and the OM proteins FlgOP. FlrC binds to enhancer sites downstream of the σ^{54} -dependent Class III promoters (Correa, Lauriano et al. 2000; Correa and Klose 2005).

Most of the Class III gene products are only required in small amounts, but the FlaA flagellin is transcribed at very high levels. One mechanism for achieving these different levels of expression is the relative binding strength of the FlrC sites, which bind FlrC strongly at the *flaA* promoter, but only weakly at other Class III promoters, e.g. the *flgK* promoter (Correa and Klose 2005).

FlrC must be phosphorylated to activate σ^{54} -dependent transcription, so presumably FlrB only phosphorylates FlrC upon assembly (not function) of the Class II export apparatus-basal body intermediate; a similar event controls expression of σ^{54} -dependent Class III genes in *C. jejuni* (Joslin and Hendrixson 2009). Detection of an intermediate that is not secretion competent may explain why the genes encoding some of the components presumably required for secretion (e.g. *fliOPQ*) are Class III (i.e. activated by FlrC) rather than Class II genes. FlrB is a soluble protein and could thus directly interact with the apparatus intermediate in the cytoplasmic membrane and phosphorylate FlrC upon assembly. Deletion of *flhF* in *V. cholerae* specifically downregulates Class III gene expression (Correa, Peng et al. 2005), suggesting that FlhF regulates FlrC-dependent transcription in addition to regulating polar flagellar placement (as discussed above). An inner membrane protein, FlrD, is also a positive regulator of class III genes. Expression of FlrD is not regulated by the flagellar transcription hierarchy, but the protein possesses a HAMP domain, so it may interact with FlrB or FlrC to influence phosphorylation and Class III transcription (Moisi, Jenul et al. 2009)

The Class II gene *fliA* encodes σ^{28} , which is required for transcription of Class IV flagellar genes (Klose and Mekalanos 1998). Similar to the checkpoint in *S. typhimurium* (Karlinsky, Tanaka et al. 2000; Chevance and Hughes 2008), the *V. cholerae* anti-sigma factor FlgM prevents σ^{28} transcriptional activity until it is secreted through a functional hook-basal body complex (Correa, Barker et al. 2004). The secretion of FlgM through the sheathed flagellum indicates that the sheath does not completely enclose the flagellum, at least at the tip. Secretion of FlgM frees σ^{28} to interact with RNA polymerase and activate Class IV flagellar genes, which encode the other four flagellins, FlaBCDE, as well as motor components (MotABX) and chemotaxis proteins (Klose and Mekalanos 1998). *V. cholerae* lacking *fliA* are non-motile and synthesize a truncated flagellum. The lack of expression of the four additional Class IV (σ^{28} -dependent) flagellins (FlaBCDE) in the *fliA* strain is likely not the reason for the truncated flagellum and lack of motility, since strains lacking *flaBCDE* are still motile and synthesize a full length flagellum, whereas a strain lacking the Class III FlaA flagellin is non-motile and aflagellate (Klose and Mekalanos 1998). Rather, the lack of expression of other Class IV genes (e.g. motor genes) likely contributes to the *fliA* phenotype. The contribution of the four Class IV flagellins to flagellar synthesis and motility is mysterious, considering that only the Class III FlaA flagellin is essential for flagellar synthesis, but perhaps the other flagellins impart subtle differences to the flagellum and thus swimming behavior that are not obvious under laboratory growth conditions.

4. Motility and virulence

V. cholerae virulence has been linked to motility. Spontaneous non-motile *V. cholerae* strains were characterized as less virulent than motile strains in several *in vivo* and *in vitro* rabbit models of cholera. Mutations that adversely affect flagellar synthesis and motility generally lead to decreased intestinal colonization in infant mice (Guentzel and Berry 1975; Montie,

Doyle-Huntzinger et al. 1982; Carsiotis, Weinstein et al. 1984; Weinstein, Carsiotis et al. 1984; Schmitt, Darnell et al. 1994; Kennedy, Rosey et al. 1997; Watnick, Lauriano et al. 2001; Syed, Beyhan et al. 2009). Non-motile live attenuated *V. cholerae* vaccine strains exhibit reduced reactogenicity (disease symptoms) in human volunteers, when compared to motile isogenic strains. (Coster, Killeen et al. 1995; Kenner, Coster et al. 1995). Using a newly-developed infant rabbit model of cholera, Rui *et al.* demonstrated that flagellin expression (whether in motile or non-motile vaccine strains) causes reactogenicity in rabbits by inducing proinflammatory cytokines in the intestine (Rui, Ritchie et al. 2010).




Fig. 4. Proposed Model of Flagellar-dependent Virulence Modulation

An inverse relationship between motility and virulence had been suggested by the observation that spontaneous hypermotile mutants express almost no CT or TCP, while spontaneous non-motile mutants express increased levels of CT and TCP (Gardel and Mekalanos 1996). Utilizing whole genome transcription profiling of *V. cholerae* strains with mutations in the key flagellar regulatory genes (*rpoN*, *flrA*, *flrC*, and *fliA*), it was observed that non-flagellated strains exhibit increased transcription of known (CT, TCP) and putative virulence factors (T6SS, hemolysins, etc)(Syed, Beyhan et al. 2009). The results suggest coordinate regulation by the flagellar regulatory hierarchy over a variety of virulence factors whose regulation was previously thought to be unlinked (Syed, Beyhan et al. 2009).

It had been known that non-motile *V. cholerae* mutants exhibited enhanced hemagglutinating activity and decreased hemolytic activity, but the identity of the respective factors was unknown (Gardel and Mekalanos 1996). The transcriptional profiling of the flagellar regulatory mutants identified the flagellar-regulated hemolysin as TLH, which is encoded adjacent to HlyA, the “El Tor” hemolysin (Syed, Beyhan et al. 2009). Also identified was the flagellar-regulated hemagglutinin, FrhA, which is a large cadherin-

containing protein that enhances binding to epithelial cells *in vitro* and intestinal colonization in both infant and adult mice. The flagellar regulatory hierarchy positively regulates *frhA* transcription and negatively regulates *tlh* transcription. Regulation of *frhA* transcription by the flagellar hierarchy is mediated through an intermediate, CdgD, a cdGMP synthase. cdGMP is an important signaling molecule that modulates complex behaviors in bacteria, most notably biofilm formation (discussed below). The results demonstrate that the flagellar hierarchy controls the transcription of non-flagellar genes that contribute to other aspects of the *V. cholerae* lifecycle besides motility (Syed, Beyhan et al. 2009).

5. Chemotaxis and virulence

Chemotaxis controls flagellar rotation in response to environmental factors, and thus is intimately tied to motility. Chemoattractants stimulate the chemotaxis machinery to cause increased clockwise (CW) rotation of the flagellum, while chemorepellants enable increased counter-clockwise (CCW) rotation (Armitage 1999; Butler and Camilli 2005). The net result of these effects on flagellar rotation is net swimming towards chemoattractants and away from chemorepellants (Falke, Bass et al. 1997; Armitage 1999). *V. cholerae* encodes three clusters of chemotaxis proteins (Heidelberg, Eisen et al. 2000), but the cluster that is embedded within the flagellar gene cluster (within the Class II *flhA* operon: *cheY3*, *cheZ*, *cheA2*, *cheB2*, and *cheW1*) appears to be the major chemotaxis machinery that controls flagellar rotation under most conditions (Camilli and Mekalanos 1995; Hyakutake, Homma et al. 2005). Methyl-accepting chemotaxis proteins (MCPs) in the cytoplasmic membrane interact with chemoattractant/repellants and the signal is transmitted through CheA to CheY via phosphorylation. Phospho-CheY then interacts with the C-ring of the flagellum, which causes a reversion from CCW to CW rotation, resulting in a change of swimming direction. CheB and CheW are involved in modulating the signal transduction pathway (Freter and O'Brien 1981; Alm and Manning 1990; Everiss, Hughes et al. 1994; Harkey, Everiss et al. 1994; Lee, Butler et al. 2001; Banerjee, Das et al. 2002; Hyakutake, Homma et al. 2005).

Interestingly, *V. cholerae* in stool exhibit a transient hyper-infectious phenotype predicted to facilitate epidemic spread of cholera, and transcription profiling revealed a transient repression of chemotaxis genes (specifically *cheW*) in these bacteria (Merrell, Butler et al. 2002). In the infant mouse model, non-chemotactic *V. cholerae* are able to outcompete chemotactic *V. cholerae* for intestinal colonization, indicating that the repression of chemotaxis in stool bacteria enhances epidemic spread (Butler and Camilli 2004; Butler, Nelson et al. 2006). Preventing phosphorylation of CheY prevents chemotactic signal transduction to the flagellum and biases it toward CCW flagellar rotation (and hence longer periods of swimming in a straight direction). The flagellum can also be biased toward CW flagellar rotation (and shorter periods of swimming in a straight direction) by the introduction of mutations into CheY that inhibit its dephosphorylation. Within the intestine, only the CCW-biased *V. cholerae* dramatically outcompete chemotactic *V. cholerae*, whereas the CW-biased bacteria are defective for intestinal colonization (Butler and Camilli 2004). Chemotactic *V. cholerae* colonize the distal end of the small intestine, whereas the CCW-biased non-chemotactic *V. cholerae* colonize the entire length of the small intestine. These results suggest that chemotaxis normally facilitates the recognition of chemoattractants within the distal small intestine or, alternatively, the recognition of chemorepellants within the proximal small intestine.

6. Biofilm formation

V. cholerae readily forms biofilms in the laboratory, and it is generally thought that *V. cholerae* predominantly exists as biofilms associated with various surfaces in the aquatic environment, including close associations with shellfish and zooplankton (Costerton, Lewandowski et al. 1995; Watnick and Kolter 1999; Faruque, Biswas et al. 2006; Yildiz and Visick 2009). Biofilm growth on chitinous surfaces induces competence in *V. cholerae*, facilitating horizontal gene transfer and rapid evolution in the marine environment (Blokesch and Schoolnik 2007). *V. cholerae* biofilms are more resistant to environmental stresses such as antibiotics, chlorine, protozoan grazing, and bacteriophage infection (Vess, Anderson et al. 1993; Faruque, Albert et al. 1998; Watnick and Kolter 1999; Matz, McDougald et al. 2005). A significant amount of study has gone into understanding *V. cholerae* biofilm formation.

Biofilm formation requires an initial phase where the bacterium associates with a solid surface, followed by attachment, formation of microcolonies, and finally the formation of the mature three-dimensional biofilm structure with characteristic pillars and water channels (Costerton, Lewandowski et al. 1995; Watnick and Kolter 1999). Formation of the mature biofilm requires the expression of the *Vibrio* exopolysaccharide (VPS), which is the polysaccharide matrix that holds the structure together (Yildiz and Schoolnik 1999; Watnick, Lauriano et al. 2001; Lauriano, Ghosh et al. 2004). *V. cholerae* expressing the VPS results in obviously wrinkled (“rugose”) colony morphology, and *V. cholerae* undergoes phase variation that leads to the rugose colony phenotype and enhanced biofilm formation (Yildiz and Schoolnik 1999; Watnick, Lauriano et al. 2001; Lim, Beyhan et al. 2007). A number of regulatory factors are involved in VPS expression and biofilm formation, and one of the driving signals behind biofilm formation is increased expression of the signaling molecule c-di-GMP (Tischler and Camilli 2004; Beyhan, Tischler et al. 2006; Beyhan, Bilecen et al. 2007; Lim, Beyhan et al. 2007; Beyhan, Odell et al. 2008; Hickman and Harwood 2008; Syed, Beyhan et al. 2009; Yildiz and Visick 2009).

In an initial screen for *V. cholerae* mutants unable to form biofilms, Watnick and Kolter identified motility as a major contributor to biofilm formation (Watnick and Kolter 1999). These results suggested that flagellar-mediated motility was important to approach and colonize a surface, and also to facilitate microcolony formation. Subsequently, it was determined that the flagellar motor itself controls VPS expression, at least in some *V. cholerae* strains, because non-flagellated mutants switch to the rugose phenotype, and this is dependent on a functional motor, suggesting that the motor acts as a sensor to induce mature biofilm formation (Lauriano, Ghosh et al. 2004). The *Vibrio* Na⁺-driven motor functioning to sense environmental conditions and drive altered gene expression is not unprecedented; the *V. parahaemolyticus* Na⁺-driven polar flagellar motor functions as a sensor to drive lateral flagellar synthesis (McCarter, Hilmen et al. 1988; Kawagishi, Imagawa et al. 1996).

In general, elevated levels of cdGMP drive *V. cholerae* toward enhanced VPS expression and down-regulate motility and virulence gene expression (Tischler and Camilli 2004; Yildiz and Visick 2009). Elevated cdGMP levels cause a decrease in Class III and IV flagellar transcription, and noticeable decreases in motility in soft agar assays (Beyhan, Tischler et al. 2006). These results suggest that activity of the Class III regulator FlrC may be responsive to elevated cdGMP levels. The effect of specific cdGMP synthases/phosphodiesterases on motility is

complicated by the presence of multiple paralogs of both types of enzymes in *V. cholerae* (Lim, Beyhan et al. 2006; Beyhan, Odell et al. 2008). Moreover, the flagellar hierarchy also regulates the expression of cdGMP modulating enzymes (mentioned above), so the effect of cdGMP on flagellar synthesis and motility is likely extremely complex, involving a large number of counteracting enzymes that regulate and are regulated by the flagellar hierarchy.

7. Conclusion

The single polar flagellum of *V. cholerae* is assembled in a stepwise fashion of components that are tightly regulated by a flagellar transcriptional hierarchy. The study of some of the unique aspects of this flagellum are likely to yield further insight into the role of flagellar synthesis, motility, and chemotaxis on the virulence and environmental persistence of this important human pathogen. One of the most unique aspects is the sheath surrounding the flagellum, which is still mysterious. The presence and function of the multiple flagellins still needs to be elucidated. Regulation of the flagellar transcriptional hierarchy is still not understood, nor how this hierarchy regulates non-flagellar genes that influence virulence and biofilm formation. Clearly much remains to be illuminated in the study of the contribution of flagellar synthesis and motility to the lifecycle of *V. cholerae*.

8. Acknowledgement

Funded by NIH AI43486

9. References

- Aizawa, S. I., G. E. Dean, et al. (1985). "Purification and characterization of the flagellar hook-basal body complex of *Salmonella typhimurium*." *J Bacteriol* 161(3): 836-849.
- Allen, R. D. and P. Baumann (1971). "Structure and arrangement of flagella in species of the genus *Beneckea* and *Photobacterium fischeri*." *J Bacteriol* 107(1): 295-302.
- Alm, R. A. and P. A. Manning (1990). "Characterization of the hlyB gene and its role in the production of the El Tor haemolysin of *Vibrio cholerae* O1." *Mol Microbiol* 4(3): 413-425.
- Armitage, J. P. (1999). "Bacterial tactic responses." *Adv Microb Physiol* 41: 229-289.
- Asai, Y., S. Kojima, et al. (1997). "Putative channel components for the fast-rotating sodium-driven flagellar motor of a marine bacterium." *J Bacteriol* 179(16): 5104-5110.
- Banerjee, R., S. Das, et al. (2002). "Involvement of in vivo induced cheY-4 gene of *Vibrio cholerae* in motility, early adherence to intestinal epithelial cells and regulation of virulence factors." *FEBS Lett* 532(1-2): 221-226.
- Beyhan, S., K. Bilecen, et al. (2007). "Regulation of rugosity and biofilm formation in *Vibrio cholerae*: comparison of VpsT and VpsR regulons and epistasis analysis of vpsT, vpsR, and hapR." *J Bacteriol* 189(2): 388-402.
- Beyhan, S., L. S. Odell, et al. (2008). "Identification and characterization of cyclic diguanylate signaling systems controlling rugosity in *Vibrio cholerae*." *J Bacteriol* 190(22): 7392-7405.
- Beyhan, S., A. D. Tischler, et al. (2006). "Transcriptome and phenotypic responses of *Vibrio cholerae* to increased cyclic di-GMP level." *J Bacteriol* 188(10): 3600-3613.

- Blair, D. F. (2003). "Flagellar movement driven by proton translocation." *FEBS Lett* 545(1): 86-95.
- Blair, D. F. and H. C. Berg (1990). "The MotA protein of *E. coli* is a proton-conducting component of the flagellar motor." *Cell* 60(3): 439-449.
- Blokesch, M. and G. K. Schoolnik (2007). "Serogroup conversion of *Vibrio cholerae* in aquatic reservoirs." *PLoS Pathog* 3(6): e81.
- Braun, T. F., S. Poulson, et al. (1999). "Function of proline residues of MotA in torque generation by the flagellar motor of *Escherichia coli*." *J Bacteriol* 181(11): 3542-3551.
- Butler, S. M. and A. Camilli (2004). "Both chemotaxis and net motility greatly influence the infectivity of *Vibrio cholerae*." *Proc Natl Acad Sci U S A* 101(14): 5018-5023.
- Butler, S. M. and A. Camilli (2005). "Going against the grain: chemotaxis and infection in *Vibrio cholerae*." *Nat Rev Microbiol* 3(8): 611-620.
- Butler, S. M., E. J. Nelson, et al. (2006). "Cholera stool bacteria repress chemotaxis to increase infectivity." *Mol Microbiol* 60(2): 417-426.
- Camilli, A. and J. J. Mekalanos (1995). "Use of recombinase gene fusions to identify *Vibrio cholerae* genes induced during infection." *Mol Microbiol* 18(4): 671-683.
- Carpenter, P. B., D. W. Hanlon, et al. (1992). "flhF, a *Bacillus subtilis* flagellar gene that encodes a putative GTP-binding protein." *Mol Microbiol* 6(18): 2705-2713.
- Carsiotis, M., D. L. Weinstein, et al. (1984). "Flagella of *Salmonella typhimurium* are a virulence factor in infected C57BL/6J mice." *Infect Immun* 46(3): 814-818.
- Chevance, F. F. and K. T. Hughes (2008). "Coordinating assembly of a bacterial macromolecular machine." *Nat Rev Microbiol* 6(6): 455-465.
- Correa, N. E., J. R. Barker, et al. (2004). "The *Vibrio cholerae* FlgM homologue is an anti-sigma28 factor that is secreted through the sheathed polar flagellum." *J Bacteriol* 186(14): 4613-4619.
- Correa, N. E. and K. E. Klose (2005). "Characterization of enhancer binding by the *Vibrio cholerae* flagellar regulatory protein FlrC." *J Bacteriol* 187(9): 3158-3170.
- Correa, N. E., C. M. Lauriano, et al. (2000). "Phosphorylation of the flagellar regulatory protein FlrC is necessary for *Vibrio cholerae* motility and enhanced colonization." *Mol Microbiol* 35(4): 743-755.
- Correa, N. E., F. Peng, et al. (2005). "Roles of the regulatory proteins FlhF and FlhG in the *Vibrio cholerae* flagellar transcription hierarchy." *J Bacteriol* 187(18): 6324-6332.
- Coster, T. S., K. P. Killeen, et al. (1995). "Safety, immunogenicity, and efficacy of live attenuated *Vibrio cholerae* O139 vaccine prototype." *Lancet* 345(8955): 949-952.
- Costerton, J. W., Z. Lewandowski, et al. (1995). "Microbial biofilms." *Annu Rev Microbiol* 49: 711-745.
- Dasgupta, N. and R. Ramphal (2001). "Interaction of the antiactivator FleN with the transcriptional activator FleQ regulates flagellar number in *Pseudomonas aeruginosa*." *J Bacteriol* 183(22): 6636-6644.
- Dasgupta, N., M. C. Wolfgang, et al. (2003). "A four-tiered transcriptional regulatory circuit controls flagellar biogenesis in *Pseudomonas aeruginosa*." *Mol Microbiol* 50(3): 809-824.
- Dean, G. E., R. M. Macnab, et al. (1984). "Gene sequence and predicted amino acid sequence of the motA protein, a membrane-associated protein required for flagellar rotation in *Escherichia coli*." *J Bacteriol* 159(3): 991-999.


- Everiss, K. D., K. J. Hughes, et al. (1994). "The accessory colonization factor and toxin-coregulated pilus gene clusters are physically linked on the *Vibrio cholerae* 0395 chromosome." *DNA Seq* 5(1): 51-55.
- Falke, J. J., R. B. Bass, et al. (1997). "The two-component signaling pathway of bacterial chemotaxis: a molecular view of signal transduction by receptors, kinases, and adaptation enzymes." *Annu Rev Cell Dev Biol* 13: 457-512.
- Faruque, S. M., M. J. Albert, et al. (1998). "Epidemiology, genetics, and ecology of toxigenic *Vibrio cholerae*." *Microbiol Mol Biol Rev* 62(4): 1301-1314.
- Faruque, S. M., K. Biswas, et al. (2006). "Transmissibility of cholera: in vivo-formed biofilms and their relationship to infectivity and persistence in the environment." *Proc Natl Acad Sci U S A* 103(16): 6350-6355.
- Francis, N. R., G. E. Sosinsky, et al. (1994). "Isolation, characterization and structure of bacterial flagellar motors containing the switch complex." *J Mol Biol* 235(4): 1261-1270.
- Freter, R. and P. C. O'Brien (1981). "Role of chemotaxis in the association of motile bacteria with intestinal mucosa: fitness and virulence of nonchemotactic *Vibrio cholerae* mutants in infant mice." *Infect Immun* 34(1): 222-233.
- Fuerst, J. A. and J. W. Perry (1988). "Demonstration of lipopolysaccharide on sheathed flagella of *Vibrio cholerae* O:1 by protein A-gold immunoelectron microscopy." *J Bacteriol* 170(4): 1488-1494.
- Fukuoka, H., T. Yakushi, et al. (2005). "Assembly of motor proteins, PomA and PomB, in the Na⁺-driven stator of the flagellar motor." *J Mol Biol* 351(4): 707-717.
- Gardel, C. L. and J. J. Mekalanos (1996). "Alterations in *Vibrio cholerae* motility phenotypes correlate with changes in virulence factor expression." *Infect Immun* 64(6): 2246-2255.
- Green, J. C., C. Kahramanoglou, et al. (2009). "Recruitment of the earliest component of the bacterial flagellum to the old cell division pole by a membrane-associated signal recognition particle family GTP-binding protein." *J Mol Biol* 391(4): 679-690.
- Guentzel, M. N. and L. J. Berry (1975). "Motility as a virulence factor for *Vibrio cholerae*." *Infect Immun* 11(5): 890-897.
- Harkey, C. W., K. D. Everiss, et al. (1994). "The *Vibrio cholerae* toxin-coregulated-pilus gene *tcpI* encodes a homolog of methyl-accepting chemotaxis proteins." *Infect Immun* 62(7): 2669-2678.
- Heidelberg, J. F., J. A. Eisen, et al. (2000). "DNA sequence of both chromosomes of the cholera pathogen *Vibrio cholerae*." *Nature* 406(6795): 477-483.
- Hendrixson, D. R. and V. J. DiRita (2003). "Transcription of sigma54-dependent but not sigma28-dependent flagellar genes in *Campylobacter jejuni* is associated with formation of the flagellar secretory apparatus." *Mol Microbiol* 50(2): 687-702.
- Hickman, J. W. and C. S. Harwood (2008). "Identification of FleQ from *Pseudomonas aeruginosa* as a c-di-GMP-responsive transcription factor." *Mol Microbiol* 69(2): 376-389.
- Homma, M., S. Aizawa, et al. (1987). "Identification of the M-ring protein of the flagellar motor of *Salmonella typhimurium*." *Proc Natl Acad Sci U S A* 84(21): 7483-7487.
- Homma, M., D. J. DeRosier, et al. (1990). "Flagellar hook and hook-associated proteins of *Salmonella typhimurium* and their relationship to other axial components of the flagellum." *J Mol Biol* 213(4): 819-832.

- Homma, M., K. Kutsukake, et al. (1990). "FlgB, FlgC, FlgF and FlgG. A family of structurally related proteins in the flagellar basal body of Salmonella typhimurium." J Mol Biol 211(2): 465-477.
- Homma, M., K. Ohnishi, et al. (1987). "Identification of flagellar hook and basal body gene products (FlaFV, FlaFVI, FlaFVII and FlaFVIII) in Salmonella typhimurium." J Bacteriol 169(8): 3617-3624.
- Hyakutake, A., M. Homma, et al. (2005). "Only one of the five CheY homologs in Vibrio cholerae directly switches flagellar rotation." J Bacteriol 187(24): 8403-8410.
- Ikeda, T., S. Asakura, et al. (1985). "'Cap' on the tip of Salmonella flagella." J Mol Biol 184(4): 735-737.
- Ikeda, T., M. Homma, et al. (1987). "Localization and stoichiometry of hook-associated proteins within Salmonella typhimurium flagella." J Bacteriol 169(3): 1168-1173.
- Jobling, M. G. and R. K. Holmes (1997). "Characterization of hapR, a positive regulator of the Vibrio cholerae HA/protease gene hap, and its identification as a functional homologue of the Vibrio harveyi luxR gene." Mol Microbiol 26(5): 1023-1034.
- Joslin, S. N. and D. R. Hendrixson (2009). "Activation of the Campylobacter jejuni FlgSR two-component system is linked to the flagellar export apparatus." J Bacteriol 191(8): 2656-2667.
- Karlinsky, J. E., S. Tanaka, et al. (2000). "Completion of the hook-basal body complex of the Salmonella typhimurium flagellum is coupled to FlgM secretion and fliC transcription." Mol Microbiol 37(5): 1220-1231.
- Kawagishi, I., M. Imagawa, et al. (1996). "The sodium-driven polar flagellar motor of marine Vibrio as the mechanosensor that regulates lateral flagellar expression." Mol Microbiol 20(4): 693-699.
- Kennedy, M. J., E. L. Rosey, et al. (1997). "Characterization of flaA- and flaB- mutants of Serpulina hyodysenteriae: both flagellin subunits, FlaA and FlaB, are necessary for full motility and intestinal colonization." FEMS Microbiol Lett 153(1): 119-128.
- Kenner, J. R., T. S. Coster, et al. (1995). "Peru-15, an improved live attenuated oral vaccine candidate for Vibrio cholerae O1." J Infect Dis 172(4): 1126-1129.
- Klose, K. E. and J. J. Mekalanos (1998). "Differential regulation of multiple flagellins in Vibrio cholerae." J Bacteriol 180(2): 303-316.
- Klose, K. E. and J. J. Mekalanos (1998). "Distinct roles of an alternative sigma factor during both free-swimming and colonizing phases of the Vibrio cholerae pathogenic cycle." Mol Microbiol 28(3): 501-520.
- Klose, K. E., V. Novik, et al. (1998). "Identification of multiple sigma⁵⁴-dependent transcriptional activators in Vibrio cholerae." J Bacteriol 180(19): 5256-5259.
- Koerdt, A., A. Paulick, et al. (2009). "MotX and MotY are required for flagellar rotation in Shewanella oneidensis MR-1." J Bacteriol 191(16): 5085-5093.
- Kojima, M., N. Nishioka, et al. (2011). "Conversion of mono-polar to peritrichous flagellation in Vibrio alginolyticus." Microbiol Immunol 55(2): 76-83.
- Kojima, S. and D. F. Blair (2004). "The bacterial flagellar motor: structure and function of a complex molecular machine." Int Rev Cytol 233: 93-134.
- Kusumoto, A., K. Kamisaka, et al. (2006). "Regulation of polar flagellar number by the flhF and flhG genes in Vibrio alginolyticus." J Biochem 139(1): 113-121.

- Kusumoto, A., N. Nishioka, et al. (2009). "Mutational analysis of the GTP-binding motif of FlhF which regulates the number and placement of the polar flagellum in *Vibrio alginolyticus*." *J Biochem* 146(5): 643-650.
- Kusumoto, A., A. Shinohara, et al. (2008). "Collaboration of FlhF and FlhG to regulate polar-flagella number and localization in *Vibrio alginolyticus*." *Microbiology* 154(Pt 5): 1390-1399.
- Lauriano, C. M., C. Ghosh, et al. (2004). "The sodium-driven flagellar motor controls exopolysaccharide expression in *Vibrio cholerae*." *J Bacteriol* 186(15): 4864-4874.
- Lee, S. H., S. M. Butler, et al. (2001). "Selection for in vivo regulators of bacterial virulence." *Proc Natl Acad Sci U S A* 98(12): 6889-6894.
- Lim, B., S. Beyhan, et al. (2006). "Cyclic-diGMP signal transduction systems in *Vibrio cholerae*: modulation of rugosity and biofilm formation." *Mol Microbiol* 60(2): 331-348.
- Lim, B., S. Beyhan, et al. (2007). "Regulation of *Vibrio* polysaccharide synthesis and virulence factor production by CdgC, a GGDEF-EAL domain protein, in *Vibrio cholerae*." *J Bacteriol* 189(3): 717-729.
- Lloyd, S. A., H. Tang, et al. (1996). "Torque generation in the flagellar motor of *Escherichia coli*: evidence of a direct role for FliG but not for FliM or FliN." *J Bacteriol* 178(1): 223-231.
- Martinez, R. M., M. N. Dharmasena, et al. (2009). "Characterization of two outer membrane proteins, FlgO and FlgP, that influence *vibrio cholerae* motility." *J Bacteriol* 191(18): 5669-5679.
- Matz, C., D. McDougald, et al. (2005). "Biofilm formation and phenotypic variation enhance predation-driven persistence of *Vibrio cholerae*." *Proc Natl Acad Sci U S A* 102(46): 16819-16824.
- McCarter, L., M. Hilmen, et al. (1988). "Flagellar dynamometer controls swarmer cell differentiation of *V. parahaemolyticus*." *Cell* 54(3): 345-351.
- McCarter, L. L. (1994). "MotX, the channel component of the sodium-type flagellar motor." *J Bacteriol* 176(19): 5988-5998.
- McCarter, L. L. (1994). "MotY, a component of the sodium-type flagellar motor." *J Bacteriol* 176(14): 4219-4225.
- Merrell, D. S., S. M. Butler, et al. (2002). "Host-induced epidemic spread of the cholera bacterium." *Nature* 417(6889): 642-645.
- Moisi, M., C. Jenul, et al. (2009). "A novel regulatory protein involved in motility of *Vibrio cholerae*." *J Bacteriol* 191(22): 7027-7038.
- Montie, T. C., D. Doyle-Huntzinger, et al. (1982). "Loss of virulence associated with absence of flagellum in an isogenic mutant of *Pseudomonas aeruginosa* in the burned-mouse model." *Infect Immun* 38(3): 1296-1298.
- Morris, D. C., F. Peng, et al. (2008). "Lipidation of an FlrC-dependent protein is required for enhanced intestinal colonization by *Vibrio cholerae*." *J Bacteriol* 190(1): 231-239.
- Niehus, E., H. Gressmann, et al. (2004). "Genome-wide analysis of transcriptional hierarchy and feedback regulation in the flagellar system of *Helicobacter pylori*." *Mol Microbiol* 52(4): 947-961.
- Okabe, M., T. Yakushi, et al. (2001). "Cloning and characterization of motX, a *Vibrio alginolyticus* sodium-driven flagellar motor gene." *J Biochem* 130(6): 879-884.

- Okabe, M., T. Yakushi, et al. (2005). "Interactions of MotX with MotY and with the PomA/PomB sodium ion channel complex of the *Vibrio alginolyticus* polar flagellum." *J Biol Chem* 280(27): 25659-25664.
- Okabe, M., T. Yakushi, et al. (2002). "MotX and MotY, specific components of the sodium-driven flagellar motor, colocalize to the outer membrane in *Vibrio alginolyticus*." *Mol Microbiol* 46(1): 125-134.
- Okunishi, I., I. Kawagishi, et al. (1996). "Cloning and characterization of motY, a gene coding for a component of the sodium-driven flagellar motor in *Vibrio alginolyticus*." *J Bacteriol* 178(8): 2409-2415.
- Prouty, M. G., N. E. Correa, et al. (2001). "The novel sigma54- and sigma28-dependent flagellar gene transcription hierarchy of *Vibrio cholerae*." *Mol Microbiol* 39(6): 1595-1609.
- Richardson, K. (1991). "Roles of motility and flagellar structure in pathogenicity of *Vibrio cholerae*: analysis of motility mutants in three animal models." *Infect Immun* 59(8): 2727-2736.
- Rui, H., J. M. Ritchie, et al. (2010). "Reactogenicity of live-attenuated *Vibrio cholerae* vaccines is dependent on flagellins." *Proc Natl Acad Sci U S A* 107(9): 4359-4364.
- Salvetti, S., E. Ghelardi, et al. (2007). "FlhF, a signal recognition particle-like GTPase, is involved in the regulation of flagellar arrangement, motility behaviour and protein secretion in *Bacillus cereus*." *Microbiology* 153(Pt 8): 2541-2552.
- Sato, K. and M. Homma (2000). "Functional reconstitution of the Na(+)-driven polar flagellar motor component of *Vibrio alginolyticus*." *J Biol Chem* 275(8): 5718-5722.
- Sato, K. and M. Homma (2000). "Multimeric structure of PomA, a component of the Na+-driven polar flagellar motor of *Vibrio alginolyticus*." *J Biol Chem* 275(26): 20223-20228.
- Schmitt, C. K., S. C. Darnell, et al. (1994). "Mutation of flgM attenuates virulence of *Salmonella typhimurium*, and mutation of fliA represses the attenuated phenotype." *J Bacteriol* 176(2): 368-377.
- Schoenhals, G. J. and R. M. Macnab (1996). "Physiological and biochemical analyses of FlgH, a lipoprotein forming the outer membrane L ring of the flagellar basal body of *Salmonella typhimurium*." *J Bacteriol* 178(14): 4200-4207.
- Sjoglad, R. D., C. W. Emala, et al. (1983). "Invited review: bacterial flagellar sheaths: structures in search of a function." *Cell Motil* 3(1): 93-103.
- Stader, J., P. Matsumura, et al. (1986). "Nucleotide sequence of the *Escherichia coli* motB gene and site-limited incorporation of its product into the cytoplasmic membrane." *J Bacteriol* 166(1): 244-252.
- Stolz, B. and H. C. Berg (1991). "Evidence for interactions between MotA and MotB, torque-generating elements of the flagellar motor of *Escherichia coli*." *J Bacteriol* 173(21): 7033-7037.
- Syed, K. A., S. Beyhan, et al. (2009). "The *Vibrio cholerae* flagellar regulatory hierarchy controls expression of virulence factors." *J Bacteriol* 191(21): 6555-6570.
- Terashima, H., H. Fukuoka, et al. (2006). "The *Vibrio* motor proteins, MotX and MotY, are associated with the basal body of Na-driven flagella and required for stator formation." *Mol Microbiol* 62(4): 1170-1180.
- Terashima, H., M. Koike, et al. (2010). "The flagellar basal body-associated protein FlgT is essential for a novel ring structure in the sodium-driven *Vibrio* motor." *J Bacteriol* 192(21): 5609-5615.

- Terashima, H., S. Kojima, et al. (2008). "Flagellar motility in bacteria structure and function of flagellar motor." *Int Rev Cell Mol Biol* 270: 39-85.
- Tischler, A. D. and A. Camilli (2004). "Cyclic diguanylate (c-di-GMP) regulates *Vibrio cholerae* biofilm formation." *Mol Microbiol* 53(3): 857-869.
- Ueno, T., K. Oosawa, et al. (1992). "M ring, S ring and proximal rod of the flagellar basal body of *Salmonella typhimurium* are composed of subunits of a single protein, FlhF." *J Mol Biol* 227(3): 672-677.
- Vess, R. W., R. L. Anderson, et al. (1993). "The colonization of solid PVC surfaces and the acquisition of resistance to germicides by water micro-organisms." *J Appl Bacteriol* 74(2): 215-221.
- Watnick, P. I. and R. Kolter (1999). "Steps in the development of a *Vibrio cholerae* El Tor biofilm." *Mol Microbiol* 34(3): 586-595.
- Watnick, P. I., C. M. Lauriano, et al. (2001). "The absence of a flagellum leads to altered colony morphology, biofilm development and virulence in *Vibrio cholerae* O139." *Mol Microbiol* 39(2): 223-235.
- Weinstein, D. L., M. Carsiotis, et al. (1984). "Flagella help *Salmonella typhimurium* survive within murine macrophages." *Infect Immun* 46(3): 819-825.
- Yildiz, F. H. and G. K. Schoolnik (1999). "*Vibrio cholerae* O1 El Tor: identification of a gene cluster required for the rugose colony type, exopolysaccharide production, chlorine resistance, and biofilm formation." *Proc Natl Acad Sci U S A* 96(7): 4028-4033.
- Yildiz, F. H. and K. L. Visick (2009). "*Vibrio* biofilms: so much the same yet so different." *Trends Microbiol* 17(3): 109-118.
- Yonekura, K., S. Maki-Yonekura, et al. (2003). "Complete atomic model of the bacterial flagellar filament by electron cryomicroscopy." *Nature* 424(6949): 643-650.
- Yoon, S. S. and J. J. Mekalanos (2008). "Decreased potency of the *Vibrio cholerae* sheathed flagellum to trigger host innate immunity." *Infect Immun* 76(3): 1282-1288.
- Yorimitsu, T., Y. Asai, et al. (2000). "Intermolecular cross-linking between the periplasmic Loop3-4 regions of PomA, a component of the Na⁺-driven flagellar motor of *Vibrio alginolyticus*." *J Biol Chem* 275(40): 31387-31391.
- Yorimitsu, T., M. Kojima, et al. (2004). "Multimeric structure of the PomA/PomB channel complex in the Na⁺-driven flagellar motor of *Vibrio alginolyticus*." *J Biochem* 135(1): 43-51.
- Zanen, G., H. Antelmann, et al. (2004). "FlhF, the third signal recognition particle-GTPase of *Bacillus subtilis*, is dispensable for protein secretion." *J Bacteriol* 186(17): 5956-5960.
- Zhou, J., S. A. Lloyd, et al. (1998). "Electrostatic interactions between rotor and stator in the bacterial flagellar motor." *Proc Natl Acad Sci U S A* 95(11): 6436-6441.
- Zhou, J., L. L. Sharp, et al. (1998). "Function of protonatable residues in the flagellar motor of *Escherichia coli*: a critical role for Asp 32 of MotB." *J Bacteriol* 180(10): 2729-2735.



Cholera

Edited by Dr. Sivakumar Gowder

ISBN 978-953-51-0415-5

Hard cover, 218 pages

Publisher InTech

Published online 28, March, 2012

Published in print edition March, 2012

Cholera, a problem in Third World countries, is a complicated diarrheal disease caused by the bacterium *Vibrio cholerae*. The latest outbreak in Haiti and surrounding areas in 2010 illustrated that cholera remains a serious threat to public health and safety. With advancements in research, cholera can be prevented and effectively treated. Irrespective of "Military" or "Monetary" power, with one's "Own Power", we can defeat this disease. The book "Cholera" is a valuable resource of power (knowledge) not only for cholera researchers but for anyone interested in promoting the health of people. Experts from different parts of the world have contributed to this important work thereby generating this power. Key features include the history of cholera, geographical distribution of the disease, mode of transmission, *Vibrio cholerae* activities, characterization of cholera toxin, cholera antagonists and preventive measures.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Anastasia R. Rugel and Karl E. Klose (2012). *Vibrio cholerae* Flagellar Synthesis and Virulence, Cholera, Dr. Sivakumar Gowder (Ed.), ISBN: 978-953-51-0415-5, InTech, Available from:
<http://www.intechopen.com/books/cholera/vibrio-cholerae-flagellar-synthesis-and-virulence>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen