

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

## 4,800

Open access books available

## 122,000

International authors and editors

## 135M

Downloads

Our authors are among the

## 154

Countries delivered to

## TOP 1%

most cited scientists

## 12.2%

Contributors from top 500 universities

**WEB OF SCIENCE™**Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)

# Peroxicretion, a Novel Tool for Engineering Membrane Trafficking

Cees M.J. Sagt

*DSM Biotechnology Center, Delft*

*The Netherlands*

## 1. Introduction

The production of proteins by recombinant micro-organisms has been possible since the late 70's. Since then enormous steps have been made to improve protein titers and to expand the range of possible proteins to be produced. The development of genetic modification tools and improved ease of use of cloning strategies have played an important role in this. The time to construct a strain and develop a cost-effective bioprocess has decreased significantly. Concurrently, sequencing data, which serve as a library for donor genes to be overexpressed, have increased exponentially. Therefore, the speed and flexibility to engineer custom-made protein factories has improved tremendously during the last decades. As in any developed technology, a certain degree of standardization is desired. In order to make this ambition more tangible the concept of cell factories is often used. This concept is based on the principle that the cell factory should be able to produce any protein to a certain desired level. Based on the concept of standardized expression vectors and expression hosts the basics of this cell factory concept were built. However, in reality this concept proved to be far too simple, as a standardized input does not result in a standardized output. This is caused by the immense complexity and dynamics of cellular systems. Even when all components are described, by sequencing the genome, the interaction, compartmentalization and dynamics of these components are largely unknown. A striking example is the difference between homologous protein expression which are produced to levels up to 50 g/l in filamentous fungi whereas heterologous proteins are produced at levels which are usually 100-1000 fold less <sup>1</sup>. So even when identical tools are used in a standardized approach the final result can vary over several orders of magnitude.

The general consensus is that this large difference is linked to cellular events and responses which are caused by the overexpression of the heterologous proteins <sup>2</sup>, therefore the scientific community, together with biotech industries, have been studying these effects for several decades. The cellular stress responses and intracellular events which are linked to the use of cells as protein production factories are very well described <sup>3-9</sup>. This profound insight in the molecular mechanism of the cellular stress reactions has facilitated the increase in expression and secretion levels of some heterologous proteins. However, levels comparable to homologous proteins have not been reached <sup>1,10</sup>. It is evident that, when even more demanding sources of biodiversity are tapped, like intracellular proteins, the engineering of a robust and versatile cell factory which is able to produce this wide range of proteins becomes a major challenge.

The production of intracellular proteins in a secreted form, to avoid complex and costly downstream processing, has hardly been touched upon. Several strategies have been developed to increase protein titers, however, these efforts have all been focused on secreted proteins. The functional diversity of intracellular proteins is very broad, while extracellular enzymes are mostly hydrolases, as shown in Figure 1. If the cell would be able to secrete these intracellular enzymes in an active form a wide range of applications could be within reach. Current strategies do not suffice but when these are combined with membrane engineering to redirect vesicle trafficking this could become a very promising approach.




Fig. 1. Distribution of enzyme functionalities in *Aspergillus niger*. The genome sequence of *Aspergillus niger* was mined for annotated functionalities based on EC classifications. The extracellular (extra) or intracellular (intra) protein distribution is based on this annotated genome. The left panel indicates relative function distribution, the right panel shows absolute numbers of enzyme function distribution.

This chapter will discuss the different strategies to improve the productivity of the cellular protein factory. Especially the possibility of engineering membrane trafficking will be addressed. This technology is not only useful for developing a versatile and robust cell factory but can be very valuable to study intracellular membrane trafficking on an academic level as well.

## 2. Conventional engineering strategies for improved protein production

The conventional strategy to improve protein production is to increase the expression levels of the gene of interest by using multiple gene copies and strong promoters<sup>11, 12</sup>. As a consequence, the cell is faced with very high levels of mRNA which should rapidly be translated into active and correctly folded proteins. To handle this increased demand for folding capacity the cell often reacts with the so called Unfolded Protein Response (UPR). This UPR is a well studied mechanism<sup>13-17</sup> which is also applied to increase folding capacity. By upregulating the transcription factors needed, or the resulting chaperones and folding enzymes, the folding capacity of the cell is enhanced<sup>18-21</sup>.

For heterologous proteins the fusion peptide concept has been proven to be successful<sup>22</sup>. This concept is used for heterologous proteins which are secreted by their native host but are overexpressed in an alternative system to increase production titers. By fusing the protein of interest to a homologous protein which is naturally secreted by the host, the desired protein is also transported outside the cell, for example, *Acremonium murorum* phenol oxidase expressed in *Aspergillus awamori*<sup>23</sup>. This carrier protein is often an N-terminal part of a very well secreted homologous protein which is fused to the protein of interest. Often a cleavage site is engineered, like Kex2 with the aim to produce only the full length protein. In reality often a mixture of unprocessed protein, partially processed protein and fully processed protein is produced, which can have consequences for downstream processing in order to meet the required product specifications. On the other hand, there are also reports which describe secretion of heterologous proteins by using only signal sequences without any additional fusion proteins like the production of llama variable heavy-chain antibody fragments (VHHs)<sup>24</sup> and the production of *Arthromyces ramosus* peroxidase<sup>25</sup> both in *Aspergillus awamori*. However, the titers obtained by these systems are in the mg/l range which is generally not sufficient for an economically feasible process.

The next generation of tools to increase protein titers are based on technology developments within bio-IT combined with the possibility of designing custom-made genes. Recently, the concept of codon adaptation was demonstrated. This concept is based on the use of favorable codons so that translation of the mRNA encoding the protein of interest is not a bottleneck. An excellent *in silico* study has recently been published<sup>26</sup> describing that translational speed and, concomitantly, ribosome density are determined by the combination of coding sequences and the tRNA pool. The first dozens of codons are translated at lower rates and create an area where ribosomes are very densely packed, especially on transcripts with a high mRNA level. This strategy could prevent ribosome jamming in later stages of the translational process and therefore be of physiological advantage of the cell. This elevation of translational speed towards the end of transcripts is often seen in fungi (which are biotechnological work-horses) and provides the cell with an effective tool against late abortions of protein synthesis which is an energy-consuming process. As suggested by the authors the fine-tuning of this ramping principle could be an additional tool to increase the efficiency of the protein production factory.

As optimizing codon usage is aimed at improving the translation of mRNA into protein, the folding capacity of the cell can become limiting, especially when high copy numbers and strong promoters are combined with codon optimization. The folding and refolding of proteins is tightly linked to membrane trafficking, a very stringent quality control system existing in the endoplasmic reticulum acts as a gatekeeper for proteins to be transported further into the secretory pathway. This quality control system has been reviewed recently<sup>2</sup>. Also the removal of proteolytic activity has been applied to improve protein production<sup>27-33</sup> this approach has shown to be very effective in specific cases. The success of the different approaches is hard to predict and highly dependent on the protein of interest being overproduced. In addition, all of the strategies above are focused to improve secretion of natural secreted proteins, most of them being hydrolases.

An overview on conventional strategies to improve the productivity of the cellular protein factory is shown in Figure 2. As shown, the modification of DNA, which includes cloning strong promoters in front of the gene of interest and constructing multiple copies of this

expression cassette is a generic strategy which can lead to improvement of protein production. Also the optimization of codon(pair) usage and mRNA stabilizing elements can be applied in a generic way. The adaptation of the folding capacity of the host cell is not a generic approach. The folding and modification of proteins is intertwined and very protein specific. Improvement of regular transport of secretory vesicles is a strategy which is less well-known, but some possible routes have been described <sup>34</sup>. In addition the overexpression of SEC4, a Rab protein associated with vesicles, doubled protein production in *Pichia pastoris* <sup>35</sup>, which is a direct proof that modification of vesicle flow can result in enhanced protein production. However all these approaches are focused on enhancing the efficiency of existing protein production and secretory processes in the host cell.




Fig. 2. Conventional improvement options for protein secretion in biological processes. Indicated in blue are the major steps in biological (eukaryotic) systems which are crucial for protein secretion, indicated in orange are improvement strategies which can be applied at the different stages of the protein secretion process.

### 3. Limitations of conventional strategies

The conventional strategies described above are focused on incremental improvement, overproducing proteins which are naturally secreted products. Therefore, the large diversity of intracellular proteins remains untapped. In addition, these strategies can be seen as generic, at least in part, but very large differences in secretion efficiency exist between different proteins; changes of a few amino acids can often have a detrimental effect on protein secretion<sup>7, 36</sup>. This is probably linked to different interactions with chaperones and folding enzymes. In the case of cutinase secretion in yeast, the bottleneck was circumvented by engineering an N-glycosylation site at the amino terminus of cutinase<sup>8</sup>. The restrictions of conventional strategies are not only illustrated by the order of magnitude of differences in protein secretion but also by the limited class of proteins which are secreted. As shown in Figure 1, in *A. niger* most of the secreted proteins are hydrolases whereas the enzyme variation of intracellular enzymes is much larger. This enormous potential of enzymes is hardly being touched upon. There are strategies to overproduce intracellular proteins inside the cell but this is always followed by a complex downstream processing step to liberate the proteins and to purify them to an acceptable level<sup>37, 38</sup>. In addition, toxic proteins or compounds cannot be produced by these methods at economical levels because they will very likely damage the host cell<sup>39-41</sup>. In order to get access to the large variety of intracellular functionalities and to be able to produce toxic compounds into a cell compartment which does not impact the viability of the host cell, an additional strategy based on membrane engineering would be very valuable.

### 4. Engineering membrane trafficking as a novel concept

Whereas the conventional approaches are focused on reaching high levels of active proteins, the engineering of membrane trafficking is aimed at redesigning the membrane trafficking in the host cell enabling a custom made flow of vesicles which can expand the possible use of the microbial cell factory. This designed vesicle flow will give access to the large diversity of intracellular enzyme activities and will be important to produce toxic compounds without damaging the host cell.

Recently a novel strategy was described to engineer membrane trafficking, this concept is called peroxicretion<sup>42</sup>. The peroxicretion concept of engineering membrane trafficking is based on two features: 1) the use of cytosolic domains of SNARE proteins (soluble NSF (N-ethylmaleimide-sensitive factor) attachment receptor) which are key for specific membrane trafficking and 2) the use of transmembrane domains, which can reposition the SNARE molecules.

Every cellular compartment of the secretory route contains a specific set of SNARE molecules. They are called v-SNARE or t-SNARE molecules (vesicle or target). SNARE molecules are transmembrane molecules which direct membrane trafficking in eukaryotic cells<sup>43</sup>. More recently, SNAREs are classified into Q and R SNAREs based on structural features. This nomenclature is more precise since certain SNARE molecules act as both v- and t-SNAREs depending on the direction of vesicle flow. Besides transmembrane domains which serve as membrane anchors also lipid modifications like palmitoylation or farnesylation<sup>44, 45</sup> occur. Combinations of transmembrane domains and palmitoylation, preventing degradation of the SNARE, are also reported<sup>46</sup>. SNARE molecules are

characterized by the so-called SNARE motif which is a conserved region of approximately 65 amino acids, this region also determines the specificity of the SNARE molecule<sup>47</sup>. This important feature enables modification of the specificity of the SNARE molecules by adapting the SNARE motifs. By fusing these SNARE motifs to transmembrane domains of other proteins it becomes possible to relocalize the SNARE motifs and modify membrane trafficking in eukaryotic systems<sup>42</sup>. Hu *et al.* showed that, by flipping SNARE molecules to the outside of cell membranes, it is possible to fuse these cells, again indicating that SNARE molecules and especially the SNARE motifs are determining the specificity of fusion of membranes<sup>48</sup>. The N-terminal domains of SNAREs are thought to act like a zipper which acts from the N-terminus towards the transmembrane domain thereby bringing the membranes in close proximity followed by the actual fusion. This model has been reviewed in an excellent paper<sup>43</sup>. Most research on SNARE molecules has been performed in the field of neuronal cells<sup>49, 50</sup>, in relation to synaptic vesicle transport. In *Saccharomyces cerevisiae* a clear set of orthologous proteins can be found as well as in *Aspergillus niger* (see Figure 5).

The concept of SNARE-pin formation has been reviewed by Jahn<sup>51, 43</sup>. In this concept, four helices are interacting to form a so called SNARE pin. The helices are named Q- and R-helices, determined by the glutamine and arginine residues which are participating. The glutamine residues are specified further as Qa, Qb, and Qc. These SNARE complexes are very efficient in mediating membrane fusion. Recently, it was shown very elegantly, by FRET studies, that one SNARE complex is sufficient to enforce membrane fusion<sup>52</sup>. Again, this indicates that modification of these SNARE complexes can have a direct effect on vesicle transport.

It has been described that N-terminal domains and SNARE motifs determine the functionality of SNARE proteins<sup>53</sup>. The functions of the transmembrane domains are not very well known, but one obvious function would be to target and anchor the SNARE in the correct membrane. By using the cytosolic domain of the SNARE and fusing this part to a transmembrane domain of a protein which is located in a different membrane it is possible to engineer membrane trafficking *in vivo*<sup>42</sup>.

To reposition the SNARE molecules a transmembrane domain is needed of a protein which is located on the vesicles which need to be modified in order to engineer membrane trafficking. The protein of which the transmembrane part is used should have the N-terminus at the cytosolic side (i.e. surface) of the membrane. The concept of this approach is depicted in Figure 3.

The transmembrane domains of proteins as tools for relocalization of SNARE molecules should be selected carefully. For the proof of principle which was aimed at transforming peroxisomes into secretory vesicles in order to secrete proteins which were located in the peroxisome, the *A. niger* ortholog of a peroxisomal membrane protein described in *Arabidopsis* named PMP22<sup>54, 55</sup> was used. The final topology of the fusion protein SNARE-membrane anchor is important. The N-terminal domains and SNARE motifs should be positioned at the cytosolic side of the vesicle in order to interact with their SNARE partner. In order to enable peroxisomal fusion with the plasma membrane, it was crucial to identify a peroxisomal membrane anchor, which could be used to decorate the peroxisome with proteins involved in membrane fusion. In order to determine whether peroxisomal membrane proteins have the correct topology we performed topology predictions at the





Fig. 3. The membrane trafficking engineering concept. The concept is based on using a cytosolic SNARE domain (green) which is located on a secretory vesicle A. This SNARE domain is fused to a transmembrane domain (grey) of a transmembrane protein originally located on vesicle B. The fusion protein (grey-green) should then be localized at vesicle B, and transforms this vesicle into a secretory vesicle B.

CBS prediction server, as shown in Figure 4 for the PMP22 ortholog of *A. niger* (An04g09130).

The peroxisomal membrane protein PMP22 has been studied in CHO cells <sup>56</sup>, in *Arabidopsis* <sup>55</sup>, and in COS cells <sup>57</sup>. All these studies give a description of the membrane topology of PMP22, PMP22 contains 4 TMDs and has the N- and C-termini placed at the cytosolic side of the peroxisome. The precise role of PMP22 is not known. PMP22 contains two peroxisomal targeting regions with almost identical basic clusters which interact with PEX19 <sup>57</sup>. The N-terminus of PMP22 is placed towards the cytosolic side and this topology makes PMP22 useful for v-SNARE fusions at the N-terminus. To determine if PMP22 localizes to the peroxisomes a GFP-PMP22 fusion was constructed and expressed in *A. niger* <sup>42</sup>. Using fluorescence microscopy the localization of GFP-PMP22 was determined. The pattern of the GFP-PMP22 fusion is similar to the GFP-SKL punctuated pattern, which indicates that the PMP22 can be used as peroxisomal membrane anchor <sup>42</sup>.





MSAKFQDEAVTSIREDTKELVHKVGNRLTGDGYLALYLRLQSQSNPLRTKMLTSGVLSSLQEILASWIA  
 HDVSKHGHYFSARVPKMALYGMFISAPLGHFLIGILQRVFAGRTSIKAKILQILASNLLVSPIQNAVYL  
 CCAVIAGARTFHQVRATVRAGFMPVMKVSWVTSPIALAFAQKFLPEHTWVPPFNIVGFVIGTYVNT  
 HTKKRLEALRKCGLPDMVG

Fig. 4. Predicted topology and sequence of the PMP22 ortholog in *A. niger* (An04g09130). The upper panel displays the predicted topology of the PMP22 ortholog by the CBS prediction servers (<http://www.cbs.dtu.dk/services/>). Four transmembrane domains are predicted, although the TMD between aa 150 and 175 does not exceed the threshold and therefore, 3 TMD are possible as well. It is clear that in both scenarios the N terminus of An04g09130 is positioned towards the outside (cytosolic side) of the vesicle. The lower panel displays the amino acid sequence of the *A. niger* ortholog of PMP22 (An04g09130) used as input for the CBS prediction server TMD algorithm.

Crucial in this engineering approach is the final orientation of the novel chimeric SNARE molecule, the SNARE domain should be positioned on the outer side (i.e. surface) of the vesicle as shown in Figure 3. The resulting vesicles are decorated with a set of alternative SNARE molecules which can enable fusion of these vesicles with other cellular compartments which contain the appropriate target SNAREs.

Peroxisomes decorated with SNARE proteins have been used as alternative vesicles for transport of intracellular proteins<sup>42</sup>. The use of peroxisomes as alternative vesicles for secretion of proteins which are normally located in the cytosol can be justified by several observations. Peroxisomes are organelles with a single membrane<sup>59</sup> and a very elegant publication shows convincing evidence that peroxisomes originate at the ER<sup>60</sup>. In addition, the identification of the small GTPase Rho1p as being localized on peroxisomes by interaction of the peroxisomal membrane protein Pex25p again indicates a link of peroxisomes with the secretory machinery<sup>61</sup>, since Rho1p is known to play a role in actin reorganization and membrane dynamics<sup>62, 63</sup>. In yeast, polarized growth is regulated by Rho1p<sup>64</sup>, and also in fungal systems the link between Rho1 and polarized growth has been confirmed<sup>65</sup> in the filamentous fungus *A. niger*, polarized growth is linked to secretion of proteins<sup>66</sup>.




Fig. 5. Overview on SNARE molecules in *Aspergillus niger* and their orthologs in *Saccharomyces cerevisiae*. This overview is based on the *A. niger* sequencing work and the supplementary material of the resulting publication<sup>58</sup>. In green, the vesicle flow from Golgi towards cell membrane is shown, dark blue indicates vesicle flow from ER towards Golgi and the orange vesicles are representing retrograde transport from Golgi towards ER. Finally, the light blue vesicles are representing vacuolar vesicle flow derived from the Golgi apparatus.

More recent papers also very clearly showed that peroxisomes originate from the endoplasmic reticulum<sup>60, 67, 68-70</sup>. The recent finding that insertion of some peroxisomal membrane proteins occurs already at the ER<sup>71</sup> clearly underpins that peroxisomes are linked to the secretory pathway in the eukaryotic cell. However, the same study mentioned that no conventional signal sequences can be found in a set of Peroxisomal Membrane Proteins (PMP) in *S. cerevisiae*. A possible mechanism could be that Pex19p, which is described as a chaperone for proper insertion of PMP into peroxisomes<sup>72-74</sup> also acts a receptor for peroxisomal membrane proteins which are inserted at the ER membrane and become part of peroxisomes at a later stage. An alternative model could be that Pex19p is involved in the budding process of (pre)peroxisomes from the ER where PMP have been already inserted. This yields an interesting model which strongly suggests that peroxisomes are derivatives of the secretory pathway. It has been reported that the transportation of peroxisomes in *Arabidopsis* by actin filaments is dependent on MYA2 (Myosin XI isoform)<sup>75</sup>, similar to the transportation of secretory vesicles. This common ground makes peroxisomes an attractive vehicle for transporting intracellular proteins towards the plasma membrane.




Fig. 6. Peroxisomes originate from the endoplasmic reticulum, taken from Tabak *et al.* 2003<sup>67</sup>. In this 3D-reconstituted SEM picture<sup>67</sup> which was processed in silico, the peroxisomes (green) are attached to the smooth ER (light blue), as a continuous membrane structure. Other peroxisomes have already been released from the smooth ER. Rough ER is indicated by dark blue and the red dots represent ribosomes on the RER.

In addition, these organelles contain an import machinery capable of importing completely folded proteins<sup>76-79</sup>, and have a controlled way of proliferation<sup>80-82</sup>. So, peroxisomes have the same origin as secretory vesicles and are thus candidates for alternative trafficking of proteins. The first application of such an engineered secretory pathway would be to produce intracellular proteins which show a much broader spectrum of enzymatic activities compared to secreted proteins which have predominantly hydrolase activity (as depicted in Figure 1). Normally, the intracellular proteins are folded in the cytosol, are not N-glycosylated, and disulphide bridges are not formed in this relatively reducing environment<sup>2</sup>. The cytosol does not contain a major machinery for the formation of disulphide bridges or for N-glycosylation as the ER does<sup>83,84</sup>. These basic differences suggest that the conventional route to secrete these intracellular proteins is not compatible with the folding environment, kinetics and modifications of these proteins.

The demonstration of engineering membrane trafficking is based on the ability of peroxisomes to import completely folded proteins. The peroxisomes containing the proteins of interest are equipped with a v-SNARE, which is normally localized at the Golgi apparatus (the study of Marelli *et al.* 2004 did not identify SNARE-like proteins on peroxisomal membranes). This ensures that peroxisomes are able to bind to t-SNAREs which are localized at the plasmamembrane. After the formation of a so called SNARE-pin the




Fig. 7. Overview of the peroxicretion concept. Panel A shows the expression of SKL tagged proteins of interest together with the expression of SNC1-PMP22 fusion protein which are localized on the peroxisomal membrane. The transmembrane anchor encoded by the PMP22 is depicted in blue, the SNC1 SNARE module is encoded by red, together they ensure the decoration of peroxisomes with the *A. niger* ortholog of the v-SNARE SNC1 at the cytosolic side. Panel B shows the translocation of the SKL tagged proteins into the decorated peroxisomes. Panel C shows the complete translocation of the SKL tagged proteins in peroxisomes, the formation of peroxisomes can be increased by overexpression of PEX11 orthologs <sup>86</sup>. Panel D shows the fusion of the peroxisomes with the plasmamembrane, using the ortholog of the t-SNARE SSO1 (purple). The complete SNAREpin structure is more complex, and is not described here for the sake of simplicity.

peroxisomes fuse with the plasma membrane thereby releasing their luminal content. The *A. niger* ortholog of v-SNARE SNC1 (An12g07570) was placed on the peroxisome using the transmembrane domain of the PMP22 ortholog (An04g09130) as a membrane anchor. The transmembrane domain of SNC1 was omitted, the fusion protein is shown in Figure 8. It has been reported that this transmembrane domain of SNC1 is important for the function of SNC1 <sup>85</sup>, but the replacement of this TMD by the PMP22 TMD did not abolish the function of SNC1 (An12g07570) in *A. niger* <sup>42</sup>. The replacement of the SNC1 TMD by the PMP22 protein does not diminish the potential of SNC1 to enforce membrane fusion because the peroxisomal content is released extracellularly. The paper of Grote *et al.* 2000 <sup>85</sup> described inhibition of SNC1 function when the TMD was replaced by geranylgeranyl anchors. Probably membrane fusion is only possible when the TMD domains of the SNARE pairs bring the membrane bilayers in close contact so that spontaneous membrane fusion occurs.

Proteins which are produced by using the peroxicretion technology are C-terminally tagged with a PTS1 signal. The most commonly used PTS1 is SKL <sup>87</sup>, however variations on this

sequence have been described as well <sup>87</sup>. Since the Pex5p (PTS1 receptor) is highly conserved between yeasts and *A. niger*, it has been shown that SKL also functions as a PTS1 signal in *A. niger* <sup>42</sup>. This raises possibilities for purifying the PTS1 tagged products on an affinity column which specifically binds PTS1 sequences. In principal nature has provided us already with a very efficient PTS1 binding protein Pex5p, which is the natural receptor for PTS1.

MSEQPYDPYIPSGANGAGAGASAAQNGDPRTREIDKKIQETVDTMRSNIFKVSEGERLD  
 SLQDKTDNLATSAQGFRRGANRVRKQMWWKDMKMRS AKFQDEAVTSIREDTKELVHK  
 VGNRLTGDGYLALYLRQLQSNPLRTKMLTSGVLSSLQEILASWIAHDVSKHGHYFSARVP  
 KMALYGMFISAPLGHFLIGILQRVFAGRTSIKAKILQILASNLLVSPIQNAVYLCCMAVIAG  
 ARTFHQVRATVRAGFMPVMKVSWVTSPIALAFAQKFLPEHTWVPPFNIVGFVIGTYVNT  
 HTKKKRLEALRKCGLPDMVG

Fig. 8. The SNC1-PMP22 fusion protein which enables peroxicretion. SNC1 ortholog sequence (no TMD) is depicted in green (based on An12g07570) the PMP22 ortholog sequence (based on An04g09130) is depicted in blue.

Mutational studies to determine whether shortening of the region between SNC1 and PMP22, thereby placing SNC1 closer to the peroxisomal membrane, could enhance peroxicretion were performed by our group. This approach was partially successful, the most efficient peroxicretion occurred when SNC1 was linked to the full length PMP22 (318 aa) or a truncated version of PMP22 (300 aa). When PMP22 was truncated further the peroxicretion decreased dramatically (< 50%). This is in contrast with a previous study <sup>88</sup>, which predicts an increase of vesicle fusion when the hinge region of the v-SNARE and the TMD domain is shortened. This discrepancy is most likely caused by the unnatural TMD of the SNC1-PMP22 fusion protein. The TMD of PMP22 was not evolutionarily selected for efficient membrane fusion. Moreover, the localization information of PMP22 is most likely positioned in this hinge region. GFP fusions with the different truncated PMP22 constructs showed a localization, which is not unambiguously peroxisomal. This indicated (partial) mis-localization of SNC1-PMP22 fusion products when PMP22 is truncated to less than 300 aa, explaining the dramatic decrease of peroxicretion.

Additional modifications to fine-tune the peroxicretion concept can be foreseen, like rebuilding the peroxisome in such a way that all components of the secretory vesicles (proteins and lipids) are present on the peroxisome, making them efficient cell organelles for peroxicretion. Subcellular proteomics could supply us with these leads. One of the most promising options is to use Sec4p a protein involved in SNARE-pin formation. Sec4p is a Rab-GTPase which plays an important role in polarized secretion by tethering secretory vesicles to the plasmamembrane <sup>89, 90</sup>. Sec4p is localized at Golgi-derived vesicles with a geranylgeranyl anchor. Sec4p GEF (Sec2p) and Sec4p GAP (Msb3p and Msb4p) are described and are controlling the activity of Sec4p <sup>91</sup>. This makes Sec4p and its partners potential leads for improving the peroxicretion technology.

In the quest for alternative secretory routes the peroxicretion concept was developed <sup>42</sup>, which shows that it is possible to redirect peroxisomes to the cell membrane where they are able to fuse and release their cargo in the extracellular medium, as shown in Figure 7. The peroxisomes can be loaded with intracellular enzymes by adding a C terminal peroxisomal

signal sequence like SKL. A combination of the peroxicretion approach with more conventional approaches to improve protein production like removal of proteases and increasing folding capacity of, in this case, the cytosol may further improve protein production. This proof of principle shows that by relocalizing the SNARE molecules to alternative compartments, it is possible to redirect vesicle trafficking in cells, which could also enable production of toxic compounds in micro organisms. This exciting possibility of constructing chimeric proteins of a specific membrane anchor with a specific SNARE functionality is an important tool to engineer membrane trafficking.

Moreover, it can also be hypothesized that due to the modified peroxisomes the complete biomass could now be used for production of enzymes, and not only the hyphal tips as is being suggested for normal secretory proteins<sup>92</sup>. A speculative idea would be to determine the components that allow the peroxisomes to utilize all of the biomass to release their content into the medium and adjust the conventional secretory vesicles in a similar way which would enable secretion of secretory proteins using the complete biomass instead of just the hyphal tips. Using this approach we could even increase fermentation yields of secretory proteins.

## 5. Designing a robust and versatile cell factory, an outlook

The cell factory of the future should be able to produce a wide range of proteins and metabolites with high productivities and yields in a robust process. It is clear that current strategies contribute in an incremental way, which is important to make the bioprocess economically feasible. In order to expand the product range and design a truly versatile cell factory more innovative approaches are needed. Until now, membrane engineering has hardly been used as an additional tool for designing cell factories. Recent developments have shown that membrane engineering can be a key for designing a truly versatile cell factory. The potential of this approach is enormous, especially in combination with conventional strategies.

In this era of synthetic biology where bio-bricks are used to engineer microbes for new functionalities the concept of membrane trafficking engineering is very well positioned. It enables a true engineering methodology for intracellular trafficking thereby unlocking a complementary approach to already applied strategies like genetic devices based on bio-bricks. In addition, the availability of -omics technologies will function as a starting platform for further fine-tuning of the membrane engineering concept. Based on this holistic approach more key components will be identified which can be used to increase the efficiency of membrane engineering and the application of engineered membranes in microbial cell factories.

This quest for further fine-tuning will be exciting and will have academic impact as well. The membrane trafficking engineering concept can serve the academic society with a new way of studying the complex vesicle flow in biological systems. By combining different components of the secretory pathway into novel chimeric proteins with new functionalities a detailed overview is within reach which can describe the molecular details of the protein secretion mechanism.

This is a major challenge and as with any technology the combined effort of industry and academia can result in significant progress when strong interaction and mutual creativity are applied and recognized.

## 6. Acknowledgments

Dr. Bert Koekman is acknowledged for editing this chapter.

## 7. References

- [1] Punt PJ, Van Biezen N, Conesa A, Albers A, Mangnus J, Van Den Hondel C. Filamentous fungi as cell factories for heterologous protein production. *Trends Biotechnol* 2002;20(5):200-6.
- [2] Brodsky JL, Skach WR. Protein folding and quality control in the endoplasmic reticulum: Recent lessons from yeast and mammalian cell systems. *Curr Opin Cell Biol*.
- [3] Al-Sheikh H, Watson AJ, Lacey GA, Punt PJ, MacKenzie DA, Jeenes DJ, Pakula T, Penttilä M, Alcocer MJC, Archer DB. Endoplasmic reticulum stress leads to the selective transcriptional downregulation of the glucoamylase gene in *Aspergillus niger*. *Mol Microbiol* 2004;53(6):1731-42.
- [4] Carvalho NDSP, Arentshorst M, Kooistra R, Stam H, Sagt CM, Van Den Hondel CAMJJ, Ram AFJ. Effects of a defective ERAD pathway on growth and heterologous protein production in *Aspergillus niger*. *Appl Microbiol Biotechnol* 2011;89(2):357-73.
- [5] Lombrana M, Moralejo FJ, Pinto R, Martín JF. Modulation of *Aspergillus awamori* thaumatin secretion by modification of bipA gene expression. *Appl Environ Microbiol* 2004;70(9):5145-52.
- [6] Punt PJ, Van Gemeren IA, Drint-Kuijvenhoven J, Hessing JGM, Van Muijlwijk-Harteveld GM, Beijersbergen A, Verrips CT, Van Den Hondel CAMJJ. Analysis of the role of the gene bipA, encoding the major endoplasmic reticulum chaperone protein in the secretion of homologous and heterologous proteins in black aspergilli. *Appl Microbiol Biotechnol* 1998;50(4):447-54.
- [7] Sagt CMJ, Müller WH, Van der Heide L, Boonstra J, Verkleij AJ, Verrips CT. Impaired cutinase secretion in *Saccharomyces cerevisiae* induces irregular endoplasmic reticulum (ER) membrane proliferation, oxidative stress, and ER-associated degradation. *Appl Environ Microbiol* 2002;68(5):2155-60.
- [8] Sagt CMJ, Kleizen B, Verwaal R, De Jong MDM, Muller WH, Smits A, Visser C, Boonstra J, Verkleij AJ, Verrips CT. Introduction of an N-glycosylation site increases secretion of heterologous proteins in yeasts. *Appl Environ Microbiol* 2000;66(11):4940-4.
- [9] Van Gemeren IA, Punt PJ, Drint-Kuyvenhoven A, Broekhuijsen MP, Van 't Hoog A, Beijersbergen A, Verrips CT, Van Den Hondel CAMJJ. The ER chaperone encoding bip A gene of black aspergilli is induced by heat shock and unfolded proteins. *Gene* 1997;198(2 JAN):43-52.
- [10] Meyer V, Wu B, Ram AFJ. *Aspergillus* as a multi-purpose cell factory: Current status and perspectives. *Biotechnol Lett* 2011;33(3):469-76.
- [11] Verdoes JC, Punt PJ, Schrickx JM, van Verseveld HW, Stouthamer AH, van den Hondel CAMJJ. Glucoamylase overexpression in *Aspergillus niger*: Molecular genetic analysis of strains containing multiple copies of the glaA gene. *Transgenic Res* 1993;2(2):84-92.

- [12] Archer DB, Jeenes DJ, Mackenzie DA. Strategies for improving heterologous protein production from filamentous fungi. *Antonie van Leeuwenhoek Int J Gen Mol Microbiol* 1994;65(3):245-50.
- [13] Saloheimo M, Valkonen M, Penttilä M. Activation mechanisms of the HAC1-mediated unfolded protein response in filamentous fungi. *Mol Microbiol* 2003;47(4):1149-61.
- [14] Mulder HJ, Nikolaev I. HacA-dependent transcriptional switch releases hacA mRNA from a translational block upon endoplasmic reticulum stress. *Eukaryotic Cell* 2009;8(4):665-75.
- [15] Mulder HJ, Nikolaev I, Madrid SM. HacA, the transcriptional activator of the unfolded protein response (UPR) in *Aspergillus niger*, binds to partly palindromic UPR elements of the consensus sequence 5'-CAN(G/A)NTGT/GCCT-3'. *Fungal Genet Biol* 2006;43(8):560-72.
- [16] Davé A, Jeenes DJ, Mackenzie DA, Archer DB. HacA-independent induction of chaperone-encoding gene *bipA* in *Aspergillus niger* strains overproducing membrane proteins. *Appl Environ Microbiol* 2006;72(1):953-5.
- [17] Mulder HJ, Saloheimo M, Penttilä M, Madrid SM. The transcription factor HacA mediates the unfolded protein response in *Aspergillus niger*, and up-regulates its own transcription. *Mol Genet Genomics* 2004;271(2):130-40.
- [18] Valkonen M, Ward M, Wang H, Penttilä M, Saloheimo M. Improvement of foreign-protein production in *Aspergillus niger var. awamori* by constitutive induction of the unfolded-protein response. *Appl Environ Microbiol* 2003;69(12):6979-86.
- [19] Inan M, Aryasomayajula D, Sinha J, Meagher MM. Enhancement of protein secretion in *Pichia pastoris* by overexpression of protein disulfide isomerase. *Biotechnol Bioeng* 2006;93(4):771-8.
- [20] Conesa A, Jeenes D, Archer DB, Van den Hondel CAMJJ, Punt PJ. Calnexin overexpression increases manganese peroxidase production in *Aspergillus niger*. *Appl Environ Microbiol* 2002;68(2):846-51.
- [21] Ngiam C, Jeenes DJ, Punt PJ, Van Den Hondel CAMJJ, Archer DB. Characterization of a foldase, protein disulfide isomerase a, in the protein secretory pathway of *Aspergillus niger*. *Appl Environ Microbiol* 2000;66(2):775-82.
- [22] Gouka RJ, Punt PJ, Van Den Hondel CAMJJ. Glucoamylase gene fusions alleviate limitations for protein production in *Aspergillus awamori* at the transcriptional and (post)translational levels. *Appl Environ Microbiol* 1997;63(2):488-97.
- [23] Gouka RJ, Van Der Heiden M, Swarthoff T, Verrrips CT. Cloning of a phenol oxidase gene from *Acremonium murorum* and its expression in *Aspergillus awamori*. *Appl Environ Microbiol* 2001;67(6):2610-6.
- [24] Joosten V, Gouka RJ, Van Den Hondel CAMJJ, Verrrips CT, Lokman BC. Expression and production of llama variable heavy-chain antibody fragments (VHHs) by *Aspergillus awamori*. *Appl Microbiol Biotechnol* 2005;66(4):384-92.
- [25] Lokman BC, Joosten V, Hovenkamp J, Gouka RJ, Verrrips CT, Van Den Hondel CAMJJ. Efficient production of *Arthromyces ramosus* peroxidase by *Aspergillus awamori*. *J Biotechnol* 2003;103(2):183-90.
- [26] Tuller T, Carmi A, Vestsigian K, Navon S, Dorfan Y, Zaborske J, Pan T, Dahan O, Furman I, Pilpel Y. An evolutionarily conserved mechanism for controlling the efficiency of protein translation. *Cell* 2010;141(2):344-54.



- [27] Nemoto T, Watanabe T, Mizogami Y, Maruyama J-, Kitamoto K. Isolation of *Aspergillus oryzae* mutants for heterologous protein production from a double proteinase gene disruptant. *Appl Microbiol Biotechnol* 2009;82(6):1105-14.
- [28] Yoon J, Kimura S, Maruyama J-, Kitamoto K. Construction of quintuple protease gene disruptant for heterologous protein production in *Aspergillus oryzae*. *Appl Microbiol Biotechnol* 2009;82(4):691-701.
- [29] Van Den Hombergh JPTW, Sollewijn Gelpke MD, Van De Vondervoort PJI, Buxton FP, Visser J. Disruption of three acid proteases in *Aspergillus niger*: Effects on protease spectrum, intracellular proteolysis, and degradation of target proteins. *Eur J Biochem* 1997;247(2):605-13.
- [30] Yoon J, Maruyama J-, Kitamoto K. Disruption of ten protease genes in the filamentous fungus *Aspergillus oryzae* highly improves production of heterologous proteins. *Appl Microbiol Biotechnol* 2011;89(3):747-59.
- [31] Braaksma M, Smilde AK, van der Werf MJ, Punt PJ. The effect of environmental conditions on extracellular protease activity in controlled fermentations of *Aspergillus niger*. *Microbiology* 2009;155(10):3430-9.
- [32] Yoon J, Kimura S, Maruyama J-, Kitamoto K. Construction of quintuple protease gene disruptant for heterologous protein production in *Aspergillus oryzae*. *Appl Microbiol Biotechnol* 2009;82(4):691-701.
- [33] Moralejo FJ, Cardoza RE, Gutierrez S, Lombraña M, Fierro F, Martíni JF. Silencing of the aspergillopepsin B (pepB) gene of *Aspergillus awamori* by antisense RNA expression or protease removal by gene disruption results in a large increase in thaumatin production. *Appl Environ Microbiol* 2002;68(7):3550-9.
- [34] Bankaitis VA, Morris AJ. Lipids and the exocytotic machinery of eukaryotic cells. *Curr Opin Cell Biol* 2003;15(4):389-95.
- [35] Liu S-, Chou W-, Sheu C-, Chang MD-. Improved secretory production of glucoamylase in *Pichia pastoris* by combination of genetic manipulations. *Biochem Biophys Res Commun* 2005;326(4):817-24.
- [36] Sagt CMJ, Müller WH, Boonstra J, Verkleij AJ, Verrips CT. Impaired secretion of a hydrophobic cutinase by *Saccharomyces cerevisiae* correlates with an increased association with immunoglobulin heavy-chain binding protein (BiP). *Appl Environ Microbiol* 1998;64(1):316-24.
- [37] Liu L, Prokopakis GJ, Asenjo JA. Optimization of enzymatic lysis of yeast. *Biotechnol Bioeng* 1988;32(9):1113-27.
- [38] Hopkins TR. Physical and chemical cell disruption for the recovery of intracellular proteins. *Bioprocess Technol* 1991;12:57-83.
- [39] Huang B, Guo J, Yi B, Yu X, Sun L, Chen W. Heterologous production of secondary metabolites as pharmaceuticals in *Saccharomyces cerevisiae*. *Biotechnol Lett* 2008;30(7):1121-37.
- [40] Engels B, Dahm P, Jennewein S. Metabolic engineering of taxadiene biosynthesis in yeast as a first step towards taxol (paclitaxel) production. *Metab Eng* 2008;10(3-4):201-6.
- [41] DeJong JM, Liu Y, Bollon AP, Long RM, Jennewein S, Williams D, Croteau RB. Genetic engineering of taxol biosynthetic genes in *Saccharomyces cerevisiae*. *Biotechnol Bioeng* 2006;93(2):212-24.

- [42] Sagt CMJ, ten Haaft PJ, Minneboo IM, Hartog MP, Damveld RA, van der Laan JM, Akeroyd M, Wenzel TJ, Luesken FA, Veenhuis M, van der Klei I, de Winde JH. Peroxicretion: A novel secretion pathway in the eukaryotic cell. *BMC Biotechnol* 2009;9.
- [43] Jahn R, Scheller RH. SNAREs - engines for membrane fusion. *Nat Rev Mol Cell Biol* 2006;7(9):631-43.
- [44] Sorek N, Bloch D, Yalovsky S. Protein lipid modifications in signaling and subcellular targeting. *Curr Opin Plant Biol* 2009;12(6):714-20.
- [45] Young SG. A thematic review series: Lipid modifications of proteins. *J Lipid Res* 2005;46(12):2529-30.
- [46] Valdez-Taubas J, Pelham H. Swf1-dependent palmitoylation of the SNARE Tlg1 prevents its ubiquitination and degradation. *EMBO J* 2005;24(14):2524-32.
- [47] Paumet F, Rahimian V, Rothman JE. The specificity of SNARE-dependent fusion encoded in the SNARE motif. *Proc Natl Acad Sci U S A* 2004;101(10):3376-80.
- [48] Hu C, Ahmed M, Melia TJ, Söllner TH, Mayer T, Rothman JE. Fusion of cells by flipped SNAREs. *Science* 2003;300(5626):1745-9.
- [49] Jahn R. A neuronal receptor for botulinum toxin. *Science* 2006;312(5773):540-1.
- [50] Hanson PI, Heuser JE, Jahn R. Neurotransmitter release - four years of SNARE complexes. *Curr Opin Neurobiol* 1997;7(3):310-5.
- [51] Jahn R. Principles of exocytosis and membrane fusion. *Ann New York Acad Sci* 2004;1014:170-8.
- [52] Van Den Bogaart G, Holt MG, Bunt G, Riedel D, Wouters FS, Jahn R. One SNARE complex is sufficient for membrane fusion. *Nat Struct Mol Biol* 2010;17(3):358-64.
- [53] Paumet F, Rahimian V, Rothman JE. The specificity of SNARE-dependent fusion encoded in the SNARE motif. *Proc Natl Acad Sci U S A* 2004;101(10):3376-80.
- [54] Murphy MA, Phillipson BA, Baker A, Mullen RT. Characterization of the targeting signal of the arabidopsis 22-kD integral peroxisomal membrane protein. *Plant Physiol* 2003;133(2):813-28.
- [55] Tugal HB, Pool M, Baker A. Arabidopsis 22-kilodalton peroxisomal membrane protein. nucleotide sequence analysis and biochemical characterization. *Plant Physiol* 1999;120(1):309-20.
- [56] Pause B, Saffrich R, Hunziker A, Ansorge W, Just WW. Targeting of the 22 kDa integral peroxisomal membrane protein. *FEBS Lett* 2000;471(1):23-8.
- [57] Brosius U, Dehmel T, Gärtner J. Two different targeting signals direct human peroxisomal membrane protein 22 to peroxisomes. *J Biol Chem* 2002;277(1):774-84.
- [58] Pel HJ, De Winde JH, Archer DB, Dyer PS, Hofmann G, Schaap PJ, Turner G, De Vries RP, Albang R, Albermann K, Andersen MR, Bendtsen JD, Benen JAE, Van Den Berg M, Breestraat S, Caddick MX, Contreras R, Cornell M, Coutinho PM, Danchin EGJ, Debets AJM, Dekker P, Van Dijck PWM, Van Dijk A, Dijkhuizen L, Driessen AJM, D'Enfert C, Geysens S, Goosen C, Groot GSP, De Groot PWJ, Guillemette T, Henrissat B, Herweijer M, Van Den Hombergh JPTW, Van Den Hondel CAMJJ, Van Der Heijden RTJM, Van Der Kaaij RM, Klis FM, Kools HJ, Kubicek CP, Van Kuyk PA, Lauber J, Lu X, Van Der Maarel MJEC, Meulenberg R, Menke H, Mortimer MA, Nielsen J, Oliver SG, Olsthoorn M, Pal K, Van Peij NNME, Ram AFJ, Rinas U, Roubos JA, Sagt CMJ, Schmoll M, Sun J, Ussery D, Varga J, Vervecken W,


- Van De Vondervoort PJJ, Wedler H, Wösten HAB, Zeng A-, Van Ooyen AJJ, Visser J, Stam H. Genome sequencing and analysis of the versatile cell factory *Aspergillus niger* CBS 513.88. *Nat Biotechnol* 2007;25(2):221-31.
- [59] Walton PA, Hill PE, Subramani S. Import of stably folded proteins into peroxisomes. *Mol Biol Cell* 1995;6(6):675-83.
- [60] Hoepfner D, Schildknecht D, Braakman I, Philippsen P, Tabak HF. Contribution of the endoplasmic reticulum to peroxisome formation. *Cell* 2005;122(1):85-95.
- [61] Marelli M, Smith JJ, Jung S, Yi E, Nesvizhskii AI, Christmas RH, Saleem RA, Tam YYC, Fagarasanu A, Goodlett DR, Aebersold R, Rachubinski RA, Aitchison JD. Quantitative mass spectrometry reveals a role for the GTPase Rho1p in actin organization on the peroxisome membrane. *J Cell Biol* 2004;167(6):1099-112.
- [62] Roumanie O, Wu H, Molk JN, Rossi G, Bloom K, Brennwald P. Rho GTPase regulation of exocytosis in yeast is independent of GTP hydrolysis and polarization of the exocyst complex. *J Cell Biol* 2005;170(4):583-94.
- [63] Macías-Sánchez K, García-Soto J, López-Ramírez A, Martínez-Cadena G. Rho1 and other GTP-binding proteins are associated with vesicles carrying glucose oxidase activity from *Fusarium oxysporum* f. sp. *lycopersici*. *Antonie Van Leeuwenhoek Int J Gen Mol Microbiol* 2011;99(3):671-80.
- [64] Kozminski KG, Alfaro G, Dighe S, Beh CT. Homologues of oxysterol-binding proteins affect Cdc42p- and Rho1p-mediated cell polarization in *Saccharomyces cerevisiae*. *Traffic* 2006;7(9):1224-42.
- [65] Martínez-Rocha AL, Roncero MIG, López-Ramírez A, Mariné M, Guarro J, Martínez-Cadena G, Di Pietro A. Rho1 has distinct functions in morphogenesis, cell wall biosynthesis and virulence of *Fusarium oxysporum*. *Cell Microbiol* 2008;10(6):1339-51.
- [66] Araujo-Bazán L, Peñalva MA, Espeso EA. Preferential localization of the endocytic internalization machinery to hyphal tips underlies polarization of the actin cytoskeleton in *Aspergillus nidulans*. *Mol Microbiol* 2008;67(4):891-905.
- [67] Tabak HF, Murk JL, Braakman I, Geuze HJ. Peroxisomes start their life in the endoplasmic reticulum. *Traffic* 2003;4(8):512-8.
- [68] Tabak HF, van der Zand A, Braakman I. Peroxisomes: Minted by the ER. *Curr Opin Cell Biol* 2008;20(4):393-400.
- [69] Tabak HF, Hoepfner D, Zand Avd, Geuze HJ, Braakman I, Huynen MA. Formation of peroxisomes: Present and past. *Biochim Biophys Acta Mol Cell Res* 2006;1763(12):1647-54.
- [70] Geuze HJ, Murk JL, Stroobants AK, Griffith JM, Kleijmeer MJ, Koster AJ, Verkleij AJ, Distel B, Tabak HF. Involvement of the endoplasmic reticulum in peroxisome formation. *Mol Biol Cell* 2003;14(7):2900-7.
- [71] Van Der Zand A, Braakman I, Tabak HF. Peroxisomal membrane proteins insert into the endoplasmic reticulum. *Mol Biol Cell* 2010;21(12):2057-65.
- [72] Girzalsky W, Hoffmann LS, Schemenewitz A, Nolte A, Kunau W-, Erdmann R. Pex19p-dependent targeting of Pex17p, a peripheral component of the peroxisomal protein import machinery. *J Biol Chem* 2006;281(28):19417-25.
- [73] Halbach A, Landgraf C, Lorenzen S, Rosenkranz K, Volkmer-Engert R, Erdmann R, Rottensteiner H. Targeting of the tail-anchored peroxisomal membrane proteins

- PEX26 and PEX15 occurs through C-terminal PEX19-binding sites. *J Cell Sci* 2006;119(12):2508-17.
- [74] Matsuzono Y, Fujiki Y. In vitro transport of membrane proteins to peroxisomes by shuttling receptor Pex19p. *J Biol Chem* 2006;281(1):36-42.
- [75] Hashimoto K, Igarashi H, Mano S, Nishimura M, Shimmen T, Yokota E. Peroxisomal localization of a myosin XI isoform in *Arabidopsis thaliana*. *Plant Cell Physiol* 2005;46(5):782-9.
- [76] Baker A, Sparkes IA. Peroxisome protein import: Some answers, more questions. *Curr Opin Plant Biol* 2005;8(6):640-7.
- [77] Van Der Klei IJ, Veenhuis M. Peroxisomes: Flexible and dynamic organelles. *Curr Opin Cell Biol* 2002;14(4):500-5.
- [78] Rucktäschel R, Girzalsky W, Erdmann R. Protein import machineries of peroxisomes. *Biochim Biophys Acta Biomembr* 2011;1808(3):892-900.
- [79] de Jonge W, Tabak HF, Braakman I. Chaperone proteins and peroxisomal protein import. *Top Curr Genet* 2006;16:149-83.
- [80] Nagotu S, Veenhuis M, Van der Klei IJ. Divide et impera: The dictum of peroxisomes. *Traffic* 2010;11(2):175-84.
- [81] Orth T, Reumann S, Zhang X, Fan J, Wenzel D, Quan S, Hu J. The PEROXIN11 protein family controls peroxisome proliferation in *Arabidopsis*. *Plant Cell* 2007;19(1):333-50.
- [82] Iwata J-, Ezaki J, Komatsu M, Yokota S, Ueno T, Tanida I, Chiba T, Tanaka K, Kominami E. Excess peroxisomes are degraded by autophagic machinery in mammals. *J Biol Chem* 2006;281(7):4035-41.
- [83] Margittai ÉV, Sitia R. Oxidative protein folding in the secretory pathway and redox signaling across compartments and cells. *Traffic* 2011;12(1):1-8.
- [84] Anelli T, Sitia R. Protein quality control in the early secretory pathway. *EMBO J* 2008;27(2):315-27.
- [85] Grote E, Baba M, Ohsumi Y, Novick PJ. Geranylgeranylated SNAREs are dominant inhibitors of membrane fusion. *J Cell Biol* 2000;151(2):453-65.
- [86] Thoms S, Erdmann R. Dynamin-related proteins and Pex11 proteins in peroxisome division and proliferation. *FEBS J* 2005;272(20):5169-81.
- [87] Kal AJ, Hettema EH, Van Den Berg M, Koerkamp MG, Van Ijlst L, Distel B, Tabak HF. In silico search for genes encoding peroxisomal proteins in *Saccharomyces cerevisiae*. *Cell Biochem Biophys* 2000;32:1-8.
- [88] McNew JA, Weber T, Engelman DM, Söllner TH, Rothman JE. The length of the flexible SNAREpin juxtamembrane region is a critical determinant of SNARE-dependent fusion. *Mol Cell* 1999;4(3):415-21.
- [89] Novick P, Medkova M, Dong G, Hutagalung A, Reinisch K, Grosshans B. Interactions between rabs, tethers, SNAREs and their regulators in exocytosis. *Biochem Soc Trans* 2006;34(5):683-6.
- [90] Lipschutz JH, Mostov KE. Exocytosis: The many masters of the exocyst. *Curr Biol* 2002;12(6):R212-214.
- [91] Gao X-, Albert S, Tcheperegine SE, Burd CG, Gallwitz D, Bi E. The GAP activity of Msb3p and Msb4p for the rab GTPase Sec4p is required for efficient exocytosis and actin organization. *J Cell Biol* 2003;162(4):635-46.

- [92] Wösten HAB, Moukha SM, Sietsma JH, Wessels JGH. Localization of growth and secretion of proteins in *Aspergillus niger*. J Gen Microbiol 1991;137(8):2017-23.

IntechOpen

IntechOpen



## **Crosstalk and Integration of Membrane Trafficking Pathways**

Edited by Dr. Roberto Weigert

ISBN 978-953-51-0515-2

Hard cover, 246 pages

**Publisher** InTech

**Published online** 11, April, 2012

**Published in print edition** April, 2012

Membrane traffic is a broad field that studies the complex exchange of membranes that occurs inside the cell. Protein, lipids and other molecules traffic among intracellular organelles, and are delivered to, or transported from the cell surface by virtue of membranous carriers generally referred as "transport intermediates". These carriers have different shapes and sizes, and their biogenesis, modality of transport, and delivery to the final destination are regulated by a multitude of very complex molecular machineries. A concept that has clearly emerged in the last decade is that each membrane pathway does not represent a close system, but is fully integrated with all the other trafficking pathways. The aim of this book is to provide a general overview of the extent of this crosstalk.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Cees M.J. Sagt (2012). Peroxcretion, a Novel Tool for Engineering Membrane Trafficking, *Crosstalk and Integration of Membrane Trafficking Pathways*, Dr. Roberto Weigert (Ed.), ISBN: 978-953-51-0515-2, InTech, Available from: <http://www.intechopen.com/books/crosstalk-and-integration-of-membrane-trafficking-pathways/peroxcretion-a-novel-tool-for-engineering-membrane-trafficking>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen