

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



## Exposure to Environmental Tobacco Smoke in Babies

Lourdes Rofes Ponce<sup>1</sup>, Ricardo Almon<sup>2</sup>, Elisa Puigdomènech<sup>3</sup>,  
Manuel A Gomez-Marcos<sup>4</sup> and Carlos Martín-Cantera<sup>5</sup>

<sup>1</sup>*Drug Dependency Unit at Sant Joan de Reus Hospital, Reus,*

<sup>2</sup>*Family Medicine Research Centre School of Health and Medical Sciences,  
Örebro University, Örebro,*

<sup>3</sup>*Lifestyles Research Group from the Primary Health Care Research Unit of Barcelona,  
IDIAP Jordi Gol, Barcelona,*

<sup>4</sup>*Department of Medicine, University of Salamanca,  
Cardiovascular Research Group (RETICS RD06/0018/0027, REDIAPP)*

*Carlos III Institute of Health of the Spanish Ministry of Health,  
SACYL, Primary Care Research Unit, La Alamedilla Health Center, Salamanca,*

<sup>5</sup>*Lifestyles Research Group from the Primary Health Care Research Unit of Barcelona,  
IDIAP Jordi Gol, Lifestyles Research Group (RETICS RD06/0018/0027, REDIAPP)*

*Carlos III Institute of Health of the Spanish Ministry of Health,  
Department of Medicine, University Autonomus of Barcelona,*

*Passeig de Sant Joan Health Center, Barcelona,*

<sup>1,3,4,5</sup>*Spain*

<sup>2</sup>*Sweden*

### 1. Introduction

#### 1.1 What is passive smoking?

Passive smoking is defined as the involuntary inhalation of tobacco smoke in closed spaces. It is also known as second hand smoke (SHS) or environmental tobacco smoke (ETS) (ENHS, 2009). ETS includes:

- *Mainstream:* Smoke exhaled by smokers
- *Sidestream:* Smoke produced during the combustion of cigarettes. Approximately 70% of ETS is made by side stream. This smoke is generated by components that escape through every pore of the filter and the paper of the cigarette. The sidestream contains small gaseous particles that can reach peripheral areas of the lung. The sidestream shows 3 more times tar and nicotine than the smoke directly inhaled by the smoker, and 5 more times carbon monoxide (CO) (US Surgeon General, 2006).

In each cigarette, 4000 chemical agents have been found, 43 of them highly toxic such as arsenic, ammonia, benzene, cadmium, hydrogen cyanide, polonium 210, butane, CO, cyanide and acetone (US Surgeon General, 1989) (see Appendix 1).

Scientific research has demonstrated that pollution caused by tobacco smoke at home can generate higher contamination rates than contamination by industrial air pollution. A series of publications have shown that children are exposed to ETS by 59.4 % - 75 % because of maternal smoking. Up to 70 % of the western children live in houses where at least one of the parents smokes. Thirty percent of these children are exposed to tobacco smoke every day (King et al., 2009).

Passive smoking is the leading cause of preventable death in infancy in industrialized countries (third cause of preventable death in adults), and one of the first leading causes of cancer in developing countries (Seong et al., 2010; Royal College of Physicians, 2010). Some authors estimate that, in Spain, between 1,228-3,237 deaths of lung cancer and cardiovascular diseases are attributable to ETS exposure at home and at work. These circumstances clearly show that exposure to passive smoking is an important public health issue (Lopez et al., 2002).

In 1950 Doll and Hill were the first to publish scientific work related to the harmful health effects of tobacco consumption (Doll & Hill, 1950). In 1974 medical research began to headline harmful effects of ETS on children's health caused by parental smoking (Harlap & Davies, 1974). Furthermore, during the last two decades, international research has found that children of smokers are more likely to be adolescent smokers, probably as a result of role models or easy access to cigarettes (Roseby et al., 2003). Nowadays numerous harmful effects of ETS have been associated with child health (Us Surgeon General, 2006). According to the International Agency for the Research on Cancer (IARC) of the World Health Organization (WHO), ETS is a type A carcinogen and no safety threshold has been established yet (IARC, 2002). Pediatric illnesses and disease related to environmental tobacco smoke are: increased risk of sudden infant death syndrome (SIDS), acute respiratory diseases, respiratory symptoms, aggravation of symptoms in patients with asthma, acute and chronic diseases in the middle ear and slowed lung growth (IARC, 2020; Samet & Sockrider, 2011).

## **2. Effects of passive smoking in children**

### **2.1 Sudden Infant Death Syndrome (SIDS)**

Sudden infant death syndrome (SIDS) can be defined as the sudden and not expected and unexplained death of an infant younger than one year, after a meticulous investigation of potential factors that might have led to infant death -including an autopsy, the examination of the death scene and the revision of the clinical history (Hunt, 2001).

According to Hannah and collaborators (Kinney & Thach, 2009) there exists no consensus about how SIDS should be defined. For instance, the definition of SIDS found in Medline reads as follows: the sudden, unexplained death of an infant younger than one year old. Some people call SIDS "crib death" because many babies who die of SIDS are found in their cribs (MedlinePlus, 2011). The National Institute of Health consensus conference issued the first standardized definition of sudden infant death in 1969 and defined it as the 'sudden death of an infant or young child, which is unexpected by history, and in which a thorough post mortem examination fails to demonstrate an adequate cause of death. The definition required thus an autopsy for all infants, who died from a similar condition as SIDS to establish a set of infants showing similar characteristics for whom vital statistics, research,

and family counseling were needed' (Hannah et al., 2009). Furthermore, the authors stated that 'although SIDS was defined as a syndrome, and thus potentially the result of more than one disease, many observers still viewed SIDS as a single entity, because of its distinctive features, which included a peak incidence at 2 to 4 months of age, male predominance, and the presence of intrathoracic petechiae. Subsequent modifications of the definitions of SIDS restricted the application of the diagnosis, SIDS, to infants under the age of 12 months, added the requirement of a death-scene investigation, or linked the death to a sleep period (i.e., the time when the majority of deaths occurred). It is still unclear whether SIDS occurs during sleep or during the many transitions between sleep and arousal that occur during the night, since such deaths are typically not witnessed'. It has to be noted that heterogeneity of the definitions of SIDS has led to contradictory results in scientific studies (Kinney & Thach, 2009).

The incidence of SIDS in industrialized countries ranges between the lower rates found in Japan (0.09 cases per 1000 infants) to the higher rates found in New Zealand (0.80 cases per 1000 infants); United States has an intermediate rate (0.57 cases per 1000 infants) (Moon et al. 2007).

Epidemiological studies have identified several risk factors for SIDS, which are mainly related to pregnancy and postnatal period. Among these risk factors, mother's tobacco consumption represents one of the most important preventable causes of SIDS. For instance, if one third of the mothers smoked during and after pregnancy, 25 % of all cases of SIDS might be attributable to this (Carrion & Pellicer, 2002).

A quantitative systematic revision has found that infant, who suffered SIDS were two times more exposed to passive smoking than those who were not (OR=2.08 and 1.94 for prenatal and postnatal maternal tobacco consumption, respectively). There is evidence showing that there is a higher postnatal exposition to ETS, when both parents smoke during pregnancy (Anderson & Cook, 1997).

A recent ecological study conducted from 1995 to 2006 has shown that for every 1% absolute increase in the prevalence of smoke-free households with children, the rates of SIDS decrease by 0.4% (Behm et al., 2011).

## **2.2 Respiratory symptoms and illness**

Several studies have shown an increased frequency of common respiratory symptoms of children (cough, sputums and wheezes) when parental smoking was present (US Surgeon General, 2006; Cook & Strachan, 1999). A meta-analysis conducted by Samet and Sockrider revealed that parental smoking was associated with respiratory symptoms (asthma, wheeze, cough, rise of sputums or dyspnea of their children (OR ranged between 1.23 and 1.35). The highest risks were found when both parents smoked (see Table 1) (US Surgeon General, 2006).

### **2.2.1 Infections of the lower respiratory tract**

During the first years of life of children, increases in respiratory problems of children are related to parental smoking habits. If both parents smoke the presence of wheezes and asthma increases by 20%. At the same time, infections of the lower respiratory tract and

cough increases by 30% and by 13%, respectively (Zmirou et al., 1990). These circumstances are directly related to the consumption of tobacco by the mother. For instance, bronchial infection with wheezes are 14% more frequent when mothers smoke more than 4 cigarettes per day, and 49% more frequent when they smoke more than 14 cigarettes per day (Neuspiel et al., 1989).

	Either parent smokes			One parent smokes			Both parents smoke		
	N	OR	95%CI	N	OR	95%CI	N	OR	95%CI
Asthma	31	1.23	(1.14-1.33)	7	1.01	(0.84-1.22)	10	1.42	(1.30-1.56)
Wheeze *	45	1.26	(1.20-1.33)	13	1.18	(1.10-1.26)	14	1.41	(1.23-1.63)
Cough	39	1.35	(1.27-1.43)	18	1.27	(1.14-1.41)	18	1.64	(1.48-1.81)
Phlegm <sup>Δ</sup>	10	1.35	(1.30-1.41)	7	1.24	(1.10-1.39)	6	1.42	(1.19-1.70)
Breathlessness <sup>Δ</sup>	6	1.31	(1.14-1.50)						

N: Number of studies

\* Excluding EC study, in which the pooled odds ratio (OR) was 1.20

<sup>Δ</sup> Data for phlegm and breathlessness restricted as several comparisons are based on fewer than five studies.

Adapted from USDHHS 2006, The Surgeon General's Report and Jonathan M Samet, Marianna Sockrider, Secondhand smoke exposure: Effects in children (Available at: [www.uptodate.com](http://www.uptodate.com), last accessed 19<sup>th</sup> October 2011)

Table 1. Effects for respiratory symptoms in children whose parents smoke according to the meta-analysis conducted by Samet and Sockrider

Bronchiolitis usually takes place among the first six months of life. Children with bronchiolitis history tend to present more bronchial spasms during early years and asthma among adolescence. This is why bronchiolitis is often considered as a risk factor that precedes repeated crisis of bronchial spasms and wheezes until the age of eight. Subsequently, passive smoking constitutes a risk factor for the appearance for asthma symptoms in infancy (Martin et al., 2003). Moreover, several studies have found that after the implementation of more restrictive tobacco law, hospitalizations of infants due to asthma decreased by 18.2% (Carriazo & Cuevo, 2010; Mackay et al., 2010).

Recently, a systematic revision including a meta-analysis has been published. The objective of this study was to explore if family tobacco consumption is related to low respiratory tract infections (LRTI) in children. This meta-analysis included more than 60 studies and showed that, if the father smokes, the OR for LRTI is 1.22 (95%CI: 1.10-1.35), if both parents smoke, the OR for LRTI is 1.62 (95%CI: 1.38-1.89) and that if any other member of the family also smokes, the OR for LRTI rises up to 1.62 (95%CI: 1.45-1.73). The strongest association of passive smoking with bronchiolitis was found when any other member of the family, besides the parents, also smoked (OR: 2.51; 95%CI: 1.96-3.21). Consequently, the authors concluded that passive smoking of cohabitants in children's households is the most important risk factor for LRTI. This risk is especially high, if the mother smokes during the postnatal period (Jones et al., 2011).

In some studies LRTI has been related to maternal tobacco consumption, but several oriental countries, as for example China and Vietnam, where maternal tobacco consumption is not

frequent, show that paternal tobacco consumption itself can explain the rise of the incidence of LRTI (Jones et al., 2011; Suzuki et al., 2009; Chen et al., 1986).

Certain studies have shown that the effect of passive smoking of babies on respiratory illnesses is critical during the first year of life, since this is the period when the baby stays more time with his or her parents, even though important effects of passive smoking on respiratory illnesses have been found during school age (Oberg et al., 2011).

### **2.2.2 Asthma**

The etiology of asthma is yet not clearly established. Some authors have related asthma to a higher frequency of respiratory infections in early life or to other inflammatory mechanisms of the pulmonary epithelium (Boulay & Boulet, 2003).

Several published scientific revisions have shown that there is a clear relationship between the prevalence and severity of asthma and the exposition to tobacco smoke in early life (US Surgeon General, 2006; Cook & Strachan, 1999).

Some authors speculate that smoking during pregnancy can alter pulmonary development in utero, which can be related to a higher risk for the development of asthma later in life (Milner et al., 1999). Other authors consider that SHS might be closely related to an allergic sensitization, however not a clear association between tobacco smoke exposure and allergy has been established (Cook & Strachan, 1999).

There is clear epidemiologic evidence for an association of exposition to SHS with the risk to suffer asthma in childhood (US Surgeon General, 2006; Goksor et al., 2007). Table 1 shows the excess risk when parents or only the mother smokes. Although there is a dose response relationship between SHS and asthma, a well defined safety threshold level without risk has not been established (Institute of Medicine, 2007). Surgeon General's Report in 2006 revealed a relationship between the prevalence of SHS and asthma, but not with the incidence of asthma (US Surgeon General, 2006). Moreover, the exposure to tobacco smoke during early years is associated with a higher prevalence of asthma in adults (Larsson et al., 2001; Svanes et al., 2004).

Finally, numerous studies have related the tobacco smoke exposure to different clinical outcomes such as an increase in a) number of visits to the emergency departments due to asthma, b) clinical symptoms, c) number of medicines and d) other clinical parameters (Carlsen & Lodrup, 2005).

### **2.2.3 Other respiratory illnesses**

Altet and collaborators found a positive association between passive smoking in infancy and a higher risk to develop pulmonary tuberculosis (Altet et al., 1996).

### **2.3 Middle ear illnesses**

Some studies have suggested an association of parental smoking with the presence of middle ear illnesses in children. (Adair-Bischoff & Sauve, 1998; Kum-Nji et al. 2006). The meta-analysis conducted by the US Center of Disease Control and Prevention found a

pooled odds ratio of 1.37 (95% CI 1.10-1.70) for recurrent middle ear otitis, 1.33 (95% CI 1.12-1.58) for middle ear effusions, and 1.20 (95% CI 0.90-1.60) for clinical referrals or operative interventions for middle ear effusions, if either parent smoked (US surgeon General, 2006).

Another meta-analysis concluded that middle ear illnesses were associated to prenatal smoking (OR:1.1), postnatal smoking (OR:1.46) and paternal smoking (OR: 1.27), the association is even higher if cohabitants, apart from the parents, smoke at home (OR: 1.35) (Royal College of Physicians, 2010).

## 2.4 Meningitis

A recently published article shows a significant positive association between passive smoking with invasive meningococcal bacteriological illness, as well as an increased carriage of *N. meningitides* and *S. pneumoniae* (Lee et al., 2010). Another recent scientific revision revealed that parental smoking (one or both parents) doubled child's risk to suffer a meningitis (OR: 2.30; 95%CI: 1.74-3.06) (Tobacco Advisory Group, 2010).

## 2.5 Dental caries

A possible effect of second hand smoke on the development of dental caries in children is not clearly established; a recently published scientific revision has failed to demonstrate a potential association of second hand smoke with caries, maybe due to the heterogeneity of the studies included (Hanioka et al., 2011).

## 3. Strategies to reduce passive smoking in babies

There are four main strategies to try to reduce exposure to passive smoking of babies: education, regulation, legislation and litigation (Davis, 1998). These strategies mainly depend on administrative measures. Nevertheless, the role of health professionals concerning the education of their patients is critical, in order to protect non-smokers from passive smoking and promote smoke-free spaces (Samet & Sockrider, 2011).

The role of pediatricians and especially pediatrics residents is of vital importance as concluded in a study published in 2008 (Hymowitz et al., 2008). The study evaluated the efficacy of a special program for training pediatric residents to address tobacco and reduce exposure to tobacco smoke. As stated by authors 'The percent of parents who smoke at sites associated with the special training condition, but not of those at sites associated with standard training, who reported that residents advised them to stop smoking, offered to help them quit, and provided quit smoking materials increased significantly from baseline to year 4'.

The above listed strategies should also focus on women in reproductive age and on pregnant women, since both conditions are the two first exposure patterns of exposure to tobacco among babies. Different types of interventions have been developed and applied to promote smoking quitting during pregnancy and in the maternal-child environment (Jane et al., 2006; Salleras et al., 2003).

The emergence of health protection laws has addressed limitations of tobacco use in public. Closed spaces for smokers have been proved to be useful measures to prevent respiratory

diseases in workers. For instance, the Spanish law 42/2010 prohibits the exposure of babies to tobacco smoke in public, closed places. However it does not regulate the exposure in private areas such as in the car or at home. Hence, a pediatric program in Primary Care should be defined and promoted, including interventions among families to reduce the exposure to SHS in children. In Spain pediatricians regularly follow up children's health status, and they would be the most appropriate professional group to promote passive smoke protection among children (Altet et al., 2007; Ortega et al., 2010).

In order to reduce infant exposure to ETS in private spaces, it is necessary to encourage parental (or baby sitters) smoking cessation. If not possible, smoke-free households could be promoted as well as the avoidance of smoking in the presence of a baby.

In Spain, the above mentioned law increases the awareness of smokers about the convenience of smoking in open-air spaces to protect others at work and public places; consequently parents, who smoke are more predisposed to adopt strategies to protect their children and other family members (see Appendix 2).

According to the last national survey of health (2006), the prevalence of adult smokers in Spain is 26.44% (21.51% in women and 31.56 in men) (see Appendix 3); in young adults the prevalence ascends up to 35 %, in spite of the fact that in the last years the number of smokers generally has decreased. Nonetheless, the number of babies exposed to tobacco smoke is still considerable (INE, 2006). Consequently, the reduction of the prevalence of smoking individuals (mainly the parents or the baby sitters) continues to be the most effective way to reduce exposure to passive smoking among infants. The following policies have contributed substantially in the reduction of the prevalence of smokers: a) increment of policies to control tobacco consumption, b) increased tobacco taxes, c) promotion of awareness about tobacco related health problems by informative advertisement, campaigns, and d) the increase of sanitary protection devices to help smokers in their cessation (Tobacco advisory Group, 2010). Programs directed to reduce the prevalence of smokers are those measures that mainly contribute to achieve smoke-free domiciles (Thomson et al., 2006).

#### 4. Theories of change

In the previous chapter, we review the strategies to reduce ETS exposure in children. However, ultimately, reduce the exposure of children depends on parents and / or caregivers. Therefore, in this section, we review the theoretical foundations and implementation, based on some articles that determines the effectiveness of interventions to reduce children's exposure to passive smoking, and how has been carried out by parents. It is a systematic review of 2008 (Priest, N.), which included 36 studies. Sixteen of 36 them expressly employed a theoretical framework in the design and/or development of the intervention.

In 1994 McIntosh developed the activities for the parent manual based on behavior modification theory. Groner 2000 employed the health belief model, and Wakefield 2002 used a harm minimization approach. Abdullah 2005 based counseling strategies on the stages of change component of Prochaska's transtheoretical model. Krieger 2005 was also guided by the transtheoretical stages of change model, as well as by social cognitive theory.

The Motivational Interviewing, was used by Emmons 2001, Curry 2003 and Chan 2005. Chan 2006a used Ajzen's theory of planned behavior in the development of their educational intervention.



Greenberg 1994, Elder 1996, Conway 2004 and Fossum 2004 employed the social learning model.

Now, in this section we will review the most important theories concerning modifications of behavioral patterns will be presented. We have divided them in three levels following the ecological perspective of the scientific revision made by Glanz and collaborators in 1997 (Glanz et al., 1997).

CONCEPT	DEFINITION
<b>Intrapersonal level</b>	Individual characteristics that influence behavior, such as knowledge, attitudes, beliefs, and personality traits
<b>Interpersonal level</b>	Interpersonal processes and primary groups, including family, friends, and peers that provide social identity, support, and role definition
<b>Community level</b>	
Institutional factors	Rules, regulations, policies, and informal structures, which may constrain or promote recommended behaviors
Community factors	Social networks and norms, or standards, which exist as formal or informal among individuals, groups, and organizations
Public policy	Local, state and federal policies and laws that regulate or support healthy actions and practices for disease prevention, early detection, control and management

Note: Table published by Glanz et al. 1997

Table 2. An ecological perspective: Levels of influence

#### 4.1 Individual or Intrapersonal level

**The Health belief model** by Janz and Becker (Janz & Becker, 1984) proposes that when someone deems it best to change a behavioral pattern, both advantages and disadvantages of this change are taken in consideration, followed by a rational decision. In addition, the perception of the individual susceptibility to the problem that requires a behavioral change and the severity of the problem are both taken into account. Recently, the concept of auto-efficiency has been added to this model. Auto-efficiency, a concept originally taken from Bandura's work (Bandura, 2011), defines an individual's confidence in undertaking a specific change in conduct.

**The Health action model** by Tones (Tones, 1991) is based on the previous model, but incorporates also the concept of self-esteem. Self-esteem is defined in this model as the individual's beliefs on his or hers qualities, and how they are perceived by others. Furthermore, this model includes a number of factors, such as previous knowledge and environmental features that can influence decision-making in the process of individual behavioral change.

**The Theory of Reasoned Action / Planned Behavior** by Ajzen (Ajzen, 1991) characterizes conduct depending on behavioral intention, subjective regulations and attitudes. This model specifies that real behavior is predicted by someone's intention to realize a

behavior. Hence, intention is the result of attitudes and subjective rules. Ajzen later modified and included the concept of "perceived behavioral control" that reflects, how easy or difficult an individual can perceive the task to follow a certain behavioral pattern.

**The Transtheoretical Model** was proposed by Prochaska y DiClemente (Prochaska & DiClemente, 1983). This model describes the phases in conduct change, and is one of the newest models describing behavioral change. It is also known as 'transtheoretical model' since it incorporates the structures of other, older models. This model considers behavioral change as a dynamic process -instead of a static process-, which takes into account the fact that, rapidity for behavior change is different in each individual and that changes occur in time following five stages. This stages are: 1) precontemplative: the individual does not think about possible behavioral changes; 2) contemplative: the individual is thinking about a change; 3) preparation: the individual enters a necessary path to be able to perform a conduct change; 4) action: the individual carries out the change in a short period of time; and 5) maintenance: the individual maintains a behavioral change over a period of time, (usually measured as keeping the change for at least six months). The transtheoretical model includes the possibility of relapses in earlier phases. This means that a maintained behavior change can also be achieved after a cyclical process of progresses and relapses. This model has been broadly used, but continues to be controversially discussed (Katz, 2001; Littlell et al., 2002; van Sluijs et al., 2004; Whitelaw et al., 2000).

**The Precaution Adoption Process Model** proposed by Weinstein (Weinstein, 1988), This model includes seven phases: not being informed or aware of a problem (unprepared), without interest in the problem, deciding about intervention, determined not to act, acting and maintenance. These phases represent qualitative models of different behavioral patterns, beliefs and experiences, and the factors influencing the transitions between the different phases can vary. Weinstein believes that this model is linear, and that interventions require being adapted to each phase of the model. It displays similarities to the transtheoretical model of behavioral change.

## 4.2 Interpersonal level

**Social learning theory or social cognitive theory** of Bandura (Bandura, 2001) postulates that people learn by observing someone else's behavior and apply it to themselves. Both, the credibility and reinforcement of the learned conduct are fundamental. This theory goes beyond individual factors influencing behavioral changes. It includes environmental and social factors. The key elements of this theory are summarize in table 3 (Cooper, 2000; Redding et al., 2000). The self-efficacy has been evaluated as an important predictor of the success in the proposed behavioral change (Cheng, 1999).

**Differential Association-Reinforcement Theory** (Glanz et al., 1997) is based on the circumstance that we can adopt new behavioral patterns by looking and imitating other persons. There are positive and negative reinforcements (price and punishment, for example). It mainly takes into account what happened after conduct change. Behavioral patterns are imitated leading to different levels of reinforcement, taking into account exposure and consequences of the conduct.

<b>Theory</b>	<b>Focus</b>	<b>Key Concepts</b>
<b>Health Belief Model</b>	Considers the pros and cons for behaviour change, as well as the perception, susceptibility and severity of the illnesses	Perceived susceptibility Perceived severity Perception of benefits from action Action barriers Auto-efficiency
<b>Health Action Model</b>	To promote people's self-esteem. To find environment support	Self-esteem Own beliefs Knowledgements Favourable environment
<b>Transtheoretical Model</b>	The intention to change varies among individuals and in time periods. The relapse is a normal phase in the process of change	Precontemplation Contemplation Preparation Action Maintenance Relapse
<b>The Precaution Adoption Process Model</b>	Changes need of certain phases and intervention should be adapted to those phases.	Without information Without interest Deciding what to do Decided to act Decided not to act Acting Maintenance
<b>Social Learning Theory or Social Cognitive Theory</b>	Considers the credibility of social models. Positive reinforcement. Helps conductal maintenance	Environment Situation Conductal capacity Expectation Hope Self-control Observational learning Reinforcements Self-efficacy Emotional aspects Reciprocal determinism
<b>Differential Association-Reinforcement Theory</b>	We learn new behavioural patterns by looking at and imitating others and by positive and negative reinforcements. The subject defines a behavioural pattern as good or justified according to his or her commitment. It takes into account what happens once a conduct change initiated.	Imitation Positive reinforcements Negative reinforcements Consequences of the behaviour
<b>Self-Regulation Theory</b>	Behaviour is defined by the interaction between subject and environment. Self-efficacy is a key concept as well as the personal standards.	Interaction environment-subject Self-efficacy Personal standards
<b>Harm reduction model</b>	Necessity to reduce health risks related to drug consumption, at both individual and community level.	Respect Acceptance Support Promotion of own's capacities (empowerment)

Table 3. Summary of the theories of change

### 4.3 Community level

**Self-Regulation Theory** (Baumeister et al., 2004). Behavior is seen as the sum of interactions between the individual and environment. Self-efficacy is the key question. Response is related to our personal values. It has been studied in the following fields: exercise, stress, alimentary behavior, cardiovascular disease and drug consumption.

**Harm reduction model** developed, among others, by Newcombe and O'Hare (Newcombe et al., 1995) is based on the principles of respect, acceptance, support and promotion of the own capacities (empowerment). This model was generated as an answer to the accumulating evidence of serious health risks related to alcohol overconsumption and tobacco use. Later on, it has been used for the understanding of behavioral pattern changes related to the consumption of drugs and to HIV infection. The model aims to diminish the negative effects of drug consumption, and does not imply quitting completely the drug consumption (O'Hare et al., 1995).

Some authors (Nigg et al., 2002), postulate that these theories can help us to elucidate, why a person is motivated to initiate a behavioral change. Other authors, in contrast, emphasize more on the reasons, why a certain conduct is being hold in time. They also suggest that for different health problems, different models can be useful, which at different stages of behavioral change can explain better these changes. Finally, multiple interventions concerning a series of health related problems (for instance, obesity and tobacco use or alcohol consumption) may need inputs from several theories at different phases or stages, to be able to explain in an improved way behavioral patterns (Martin, 2006).

## 5. Conclusions

There is enough scientific evidence to conclude that indirect tobacco smoke exposure of children (and adults) has negative health effects. Tobacco smoke exposure of children can cause: low tract respiratory infections, middle ear infections, asthma, low weight at birth and sudden death syndrome.

Subsequently, strategies to reduce passive exposure to tobacco are an urgent and priority sanitary, public health aim. Scientific evidence should enforce actions performed by health care professionals of all disciplines to undertake measures in order to eradicate this avoidable risk in children.

## 6. Key points

- Environmental tobacco smoke is made from the smoke exhaled by smokers and the smoke produced during the combustion of cigarettes.
- Passive smoking is the leading cause of preventable death in infancy in industrialized countries and one of the first leading causes of the rise of cancer in developing countries.
- Up to 70 % of the western children live in households where at least one of the parents smokes. Thirty percent of these children are exposed every day to tobacco smoke.
- Sudden infant death syndrome, acute respiratory illnesses, acute and chronic middle ear infections and meningitis are related to ETS.

- Passive smoking during pregnancy lowers baby's weight at birth, highers the risk of: prematurity, congenital malformations, spontaneous abortion and fetal and perinatal mortality.
- The rise of the policies controlling tobacco consumption, the increase in of tobacco taxes, the awareness of tobacco related health problem in advertising informative campaigns, and the increase of sanitary devices to help smokers in their cessation have contributed to reduce the number of smokers.
- Reducing the prevalence of smokers among parents and baby sitters continues to be the most effective way to prevent passive smoking among children.

Compound	Does it cause cancer?
Nicotine	
Dimethylnitrosamine	
Benzopyrene	X
Pyrene	X
Napthalene	X
Phenol	
Acetone	X
Methanol	
Vinyl	
Carbon Monoxide	X
Nitrogen oxides gases	
Tolue	
Benzene	
Arsenic	
Cadmium	X
Niquel	
Lead	
Naphthylamine	X
Toluidine	X
Polonium-210	X
Ammonia	
Dibenzocridine	X
Cianamide	
DDT	
Uretane	
Butane	

Adapted from: Departament de Salut de la Generalitat de Catalunya (<http://www.gencat.cat/salut/depsalut/pdf/gtabacp.pdf>, last accessed November 10<sup>th</sup> 2011) and Cancer Society of New Zeland Inc. and Health Promotion Services Branch Health Dept. of Western Australia

Appendix 1. Main components of tobacco smoke and its carcinogenic effect

Exposure	Does NOT avoid exposure	Avoids exposure
At home	To smoke at home when the bay is not there	Not smoking at home, in any room
	To smoke in delimited areas of the house	Prohibit family members and visitors to smoke in the house
	To smoke with the window or balcony open	To always smoke outside the house. To smoke only in the balcony or terrace keeping the door closed
	To smoke and ventilate the house after smoking	
In the car	To smoke in the car when the bay is not there	Not to smoke in the car, even though the baby is not there
	To smoke in the car when the baby is there but with the window opened	

From: Ortega G, Castellà C, Martín-Cantera C, Ballvé JL, Díaz E, Saez M, et al. Passive smoking in babies: the BIBE study (Brief Intervention in babies. Effectiveness). BMC Public Health. 2010;10:772.

#### Appendix 2. Measures to avoid exposure to tobacco smoke in babies

	Number	%
<b>Men</b>		
Total	5,794.60	31.56
16-24 years old	596.3	24.96
25-34 years old	1,580.90	40.16
35-44 years old	1,382.40	37.41
45-54 years old	1,141.30	38.83
55-64 years old	654.2	28.45
65-74 years old	312.4	18.64
≥75 years old	127	8.93
<b>Women</b>		
Total	4,101.70	21.51
16-24 years old	657	28.93
25-34 years old	1,108.40	30.16
35-44 years old	1,093.00	30.73
45-54 years old	845.6	28.47
55-64 years old	286.5	11.77
65-74 years old	83.9	3.86
≥75 years old	27.3	1.37

From: Spanish Ministry of Health. National Health survey, 2006

#### Appendix 3. Tobacco consumption in Spain. Daily number of smokers (in thousands) and percentage aged ≥ 16 according to sex and age group

## 7. Abbreviations

CI: Confidence Interval

CO: Carbon monoxide

ETS: Environmental Tobacco Smoke

IARC: International Agency for the Research on Cancer

LRTI: Low respiratory tract infections

OR: Odds Ratio

SHS: Second Hand Smoke

SIDS: Sudden Infant Death Syndrome

WHO: World Health Organization

## 8. References

- Abdullah ASM, Lam TH, Mak YW, Loke AY. A randomized control trial of a smoking cessation intervention parents of young children: a randomised controlled trial. *Addiction* 2005; 100(11): 1731-40.
- Adair-Bischoff CE, Sauve RS. Environmental tobacco smoke and middle ear disease in preschool-age children. *Arch Pediatr Adolesc Med* 1998 Feb;152(2):127-33.
- Altet MN, Alcaide J, Plans P, Taberner JL, Salto E, Folguera LI, et al. Passive smoking and risk of pulmonary tuberculosis in children immediately following infection. A case-control study. *Tuber Lung Dis* 1996 Dec;77(6):537-44.
- Altet N, Pascual MT, Grupo de Trabajo sobre Tabaquismo en la Infancia. Tabaquismo en la infancia y adolescencia. Papel del pediatra en su prevención y control. *An Esp Pediatr* 2007;52:168-77.
- Ajzen I. The theory of planned behavior. *Organizational Behavior and Human Decision Processes* 1991;50:179-211.
- Anderson HR, Cook DG. Passive smoking and sudden infant death syndrome: review of the epidemiological evidence. *Thorax* 1997 Nov;52(11):1003-9.
- Bandura A. Social cognitive theory: an agentic perspective. *Annu Rev Psychol* 2001;52:1-26.
- Baumeister F, Vohs K. *Handbook of Self-Regulation. Research, Theory, and Applications*. Edited by Roy F Baumeister and Kathleen D Vohs 2004.
- Behm I, Kabir Z, Connolly GN, Alpert HR. Increasing prevalence of smoke-free homes and decreasing rates of sudden infant death syndrome in the United States: an ecological association study. *Tob Control* 2011 Apr 7.
- Boulay ME, Boulet LP. The relationships between atopy, rhinitis and asthma: pathophysiological considerations. *Curr Opin Allergy Clin Immunol* 2003 Feb;3(1):51-5.
- Carlsen KH, Lodrup Carlsen KC. Parental smoking and childhood asthma: clinical implications. *Treat Respir Med* 2005;4(5):337-46.
- Carriazo N, Cuervo J. Universalizar los espacios sin humo parece disminuir los ingresos hospitalarios en niños. *Evid Pediatría* 2010;6:84.
- Carrion F, Pellicer C. El tabaquismo pasivo en la infancia. Nuevas evidencias. *Prevencion del Tabaquismo* 2002;4(1):20-5.
- Chan SS, Lam TH, Salili F, Leung GM, Wong DC, Botelho RJ, et al. A randomized controlled trial of an individualized motivational intervention on smoking cessation for parents of sick children: a pilot study. *Applied Nursing Research* 2005;18(3):178-81.


- Chan SS, Lam TH. Preventing exposure to second-hand smoke. *Seminars in Oncology Nursing* 2003;19(4): 284-90
- Chen Y, Li W, Yu S. Influence of passive smoking on admissions for respiratory illness in early childhood. *Br Med J (Clin Res Ed)* 1986 Aug 2;293(6542):303-6.
- Cheng TL, DeWitt TG, Savageau JA, O'Connor KG. Determinants of counseling in primary care pediatric practice: physician attitudes about time, money, and health issues. *Arch Pediatr Adolesc Med* 1999 Jun;153(6):629-35.
- Conway TL, Woodruff SI, Edwards CC, Hovell MF, Klein J. Intervention to reduce environmental tobacco smoke exposure in Latino children: null effects on hair biomarkers and parents reports. *Tobacco Control* 2004; 13(1): 90-2.
- Cook DG, Strachan DP. Health effects of passive smoking-10: Summary of effects of parental smoking on the respiratory health of children and implications for research. *Thorax* 1999 Apr;54(4):357-66.
- Cooper MD. Towards a model of safety culture. *Safety Science* 2000;36(2):111-36.
- Curry SJ, Lundman EJ, Graham E, Stout J, Grothaus L, Lozano P. Pediatric-based smoking cessation intervention for low-income women: a randomized trial. *Archives of Pediatrics and Adolescent Medicine* 2003;157(3):295-302
- Davis RM. Exposure to environmental tobacco smoke: identifying and protecting those at risk. *JAMA* 1998 Dec 9;280(22):1947-9.
- Doll R, Hill AB. Smoking and carcinoma of the lung; preliminary report. *Br Med J* 1950 Sep 30;2(4682):739-48.
- Elder JP, Perry CL, Stone EJ, Johnson CC, Yang M, Edmundson EW, et al. Tobacco use measurement, prediction, and intervention in elementary schools in four states: the CATCH Study. *Preventive Medicine* 1996;25(4):486-94.
- Emmons KM, Wong M, Hammond SK, Velicer WF, Fava JL, Monroe AD et al. Intervention and policy issues related to children's exposure to environmental tobacco smoke. *Preventive Medicine* 2001;32:321-31.
- Environment and Health Information System (ENHIS). Exposure of children to second-hand tobacco smoke. fact sheet 3 4 December 2009; (Available in <http://www.euro.who.int>).
- Fossum B, Arborelius E, Bremberg S. Evaluation of a counseling method for the prevention of child exposure to tobacco smoke: an example of client-centered communication. *Preventive Medicine* 2004;38(3): 295-301.
- Glanz K, Rimer BK. Theory at a Glance. A Guide for Health Promotion Practice. National Cancer Institute, National Institutes of Health, U S 1997;Washington, DC., Department of Health and Human Services. NIH Pub. 97-3896.
- Goksor E, Amark M, Alm B, Gustafsson PM, Wennergren G. The impact of pre- and post-natal smoke exposure on future asthma and bronchial hyper-responsiveness. *Acta Paediatr* 2007 Jul;96(7):1030-5.
- Greenber RA, Strecher VJ, Bauman KE, Boat BW, et al. Evaluation of a home-based intervention program to reduce infant passive smoking and lower respiratory illness. *Journal of Behavioral medicine* 1994;17(3):273-90.
- Groner JA, Ahijevych K, Grossman LK, Rich LN. The impact of a brief intervention on maternal smoking behavior. *Pediatrics* 2000;105(1 Pt 3): 267-71.
- Hanioka T, Ojima M, Tanaka K, Yamamoto M. Does secondhand smoke affect the development of dental caries in children? A systematic review. *Int J Environ Res Public Health* 2011 May;8(5):1503-19.



- Harlap S, Davies AM. Infant admissions to hospital and maternal smoking. *Lancet* 1974 Mar 30;1(7857):529-32.
- Hymowitz et al. The pediatric residency training on tobacco project: four-year parent outcome findings. *Prev. Med.* 2008, Aug;47(2):221-4. Epub 2008 Jun 30.
- Hunt CE. Sudden infant death syndrome and other causes of infant mortality: diagnosis, mechanisms, and risk for recurrence in siblings. *Am J Respir Crit Care Med* 2001 Aug 1;164(3):346-57.
- IARC. Tobacco Smoke and Involuntary Smoking. Vol. 28. Lyon. IARC Monographs 2002.
- Institute of Medicine. Exposure to environmental tobacco smoke. In: *Clearing the Air: Asthma and Indoor Air Exposures* National Academy Press, Washington DC 2000;278.
- Instituto Nacional de Estadística. Encuesta Nacional de Salud de España. Ministerio de Sanidad y Consumo 2006.
- Jane M, Martinez C, Altet N. Guia clínica per promoure l'abandonament del consum de tabac durant l'embaràs. Departament de Salut 2006.
- Janz NK, Becker MH. The Health Belief Model: a decade later. *Health Educ Q* 1984;11(1):1-47.
- Jones LL, Hashim A, McKeever T, Cook DG, Britton J, Leonardi-Bee J. Parental and household smoking and the increased risk of bronchitis, bronchiolitis and other lower respiratory infections in infancy: systematic review and meta-analysis. *Respir Res* 2011;12:5.
- Katz DL. Behavior modification in primary care: the pressure system model. *Prev Med* 2001 Jan;32(1):66-72.
- Kinney HC, Thach BT. The sudden infant death syndrome. *N Engl J Med* 2009 Aug 20;361(8):795-805.
- King K, Martynenko M, Bergman MH, Liu YH, Winickoff JP, Weitzman M. Family composition and children's exposure to adult smokers in their homes. *Pediatrics* 2009 Apr;123(4):e559-e564.
- Krieger JK, Takaro TK, Allen C, Song L, Weaver M, Chai S, et al. The Seattle-King County healthy homes project: implementation of a comprehensive approach to improving indoor environmental quality for low-income children with asthma. *Environmental Health perspectives* 2002;110 Suppl 2: 311-22
- Kum-Nji P, Meloy L, Herrod HG. Environmental tobacco smoke exposure: prevalence and mechanisms of causation of infections in children. *Pediatrics* 2006 May;117(5):1745-54.
- Larsson ML, Frisk M, Hallstrom J, Kiviloog J, Lundback B. Environmental tobacco smoke exposure during childhood is associated with increased prevalence of asthma in adults. *Chest* 2001 Sep;120(3):711-7.
- Lee CC, Middaugh NA, Howie SRC, Ezzati M. Association of Secondhand Smoke Exposure with Pediatric Invasive Bacterial Disease and Bacterial Carriage: A Systematic Review and Meta-analysis. *PLoS Med* 2010 Dec 7;7(12):e1000374.
- Littell JH, Girvin H. Stages of change. A critique. *Behav Modif* 2002 Apr;26(2):223-73.
- Lopez MJ, Perez-Rios M, Schiaffino A, Nebot M, Montes A, Ariza C, et al. Mortality attributable to passive smoking in Spain, 2002. *Tob Control* 2007 Dec;16(6):373-7.
- Mackay D, Haw S, Ayres JG, Fischbacher C, Pell JP. Smoke-free legislation and hospitalizations for childhood asthma. *N Engl J Med* 2010 Sep 16;363(12):1139-45.

- Martin C. La factibilidad del consejo preventivo sobre accidentes de tráfico en atención primaria. Tesis Doctoral Universidad Autonoma de Barcelona 2006.
- Martin C, Roig L, Ferrer S, Jane C, Vilella E. Los niños y su exposición al tabaquismo pasivo. 3er Congreso del CNPT. Zaragoza. 2003. Ref Type: Personal Communication.
- McIntosh NA, Clark NM, Howatt WF. Reducing tobacco smoke in the environment of child with asthma: a cotinine-assisted, minimal-contact intervention. *Journal of Asthma* 1994;31(6): 453-62.
- MedlinePlus. Available at: <http://www.nlm.nih.gov/medlineplus/>
- Milner AD, Marsh MJ, Ingram DM, Fox GF, Susiva C. Effects of smoking in pregnancy on neonatal lung function. *Arch Dis Child Fetal Neonatal Ed* 1999 Jan;80(1):F8-14.
- Moon RY, Horne RS, Hauck FR. Sudden infant death syndrome. *Lancet* 2007 Nov 3;370(9598):1578-87.
- Neuspiel DR, Rush D, Butler NR, Golding J, Bijur PE, Kurzon M. Parental smoking and post-infancy wheezing in children: a prospective cohort study. *Am J Public Health* 1989 Feb;79(2):168-71.
- Newcombe R, O'Hare PA, Matthews A, Buning EC. The reduction of drug-related harm. A conceptual framework for theory, practice and research 1992; London; Routledge.
- Nigg CR, Allegrante JP, Ory M. Theory-comparison and multiple-behavior research: common themes advancing health behavior research. *Health Educ Res* 2002 Oct 1;17(5):670-9.
- Oberg M, Jaakkola MS, Woodward A, Peruga A, Pruss-Ustun A. Worldwide burden of disease from exposure to second-hand smoke: a retrospective analysis of data from 192 countries. *Lancet* 2011 Jan 8;377(9760):139-46.
- O'Hare PA, Newcombe A, Matthews EC, Buning ED. La reducció de los daños relacionados con las drogas. Barcelona, Grup Igia 1995;17-39.
- Ortega G, Castella C, Martin-Cantera C, Ballve JL, Diaz E, Saez M, et al. Passive smoking in babies: the BIBE study (Brief Intervention in babies. Effectiveness). *BMC Public Health* 2010;10:772.
- Priest N, Roseby R, Waters E. et al. Family and carer smoking control programmes for reducing children's exposure to environmental tobacco smoke (Review). *The Cochrane Database Syst Rev*. 2008, Oct 8; (4):CD001746.
- Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: toward an integrative model of change. *J Consult Clin Psychol* 1983 Jun;51(3):390-5.
- Redding C, Rossi J, Rossi S, Velicer W, Prochaska JO. Health Behavior Models. *The International Electronic Journal of Health Education* 2000;3(Special Issue):180-93.
- Roseby R, Waters E, Polnay A, Campbell R, Webster P, Spencer N. Family and carer smoking control programmes for reducing children's exposure to environmental tobacco smoke. *Cochrane Database Syst Rev* 2003;(3):CD001746.
- Royal College of Physicians. Passive smoking and children. A report by the Tobacco Advisory Group London: RCP, 2010.
- Salleras L, et al. Guia per a la prevenció i el control del tabaquisme des de l'àmbit pediàtric. Departament de Sanitat i seguretat social Generalitat de Catalunya 2003.
- Samet J, Sockrider M. Secondhand smoke exposure: Effects in children. UpToDate 2011.
- Samet J, Sockrider M. Control of secondhand smoke exposure. UpToDate 2011; <http://www.uptodate.com/>.

- Seong MW, Moon JS, Hwang JH, Ryu HJ, Kang SJ, Kong SY, et al. Preschool children and their mothers are more exposed to paternal smoking at home than school children and their mothers. *Clin Chim Acta* 2010 Jan;411(1-2):72-6.
- Suzuki M, Thiem VD, Yanai H, Matsubayashi T, Yoshida LM, Tho LH, et al. Association of environmental tobacco smoking exposure with an increased risk of hospital admissions for pneumonia in children under 5 years of age in Vietnam. *Thorax* 2009 Jun;64(6):484-9.
- Svanes C, Omenaas E, Jarvis D, Chinn S, Gulsvik A, Burney P. Parental smoking in childhood and adult obstructive lung disease: results from the European Community Respiratory Health Survey. *Thorax* 2004 Apr;59(4):295-302.
- Thomson G, Wilson N, Howden-Chapman P. Population level policy options for increasing the prevalence of smokefree homes. *J Epidemiol Community Health* 2006 Apr;60(4):298-304.
- Tobacco Advisory Group. Passive smoking and children. Royal College of Physicians 2010.
- Tones K. Health promotion, self empowerment and the concept of control. Leeds Polytechnic, Leeds 1991.
- US Surgeon General. The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General. U.S. Dept. of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion. Office on Smoking and Health, Atlanta, Georgia 2006.
- US Surgeon General. Reducing the health consequences of smoking: 25 years of progress. US Department of Health and Human Services 1989;DHHS publication No (CDC) 89-8411.
- van Sluijs EMF, van Poppel MNM, van Mechelen W. Stage-based lifestyle interventions in primary care - Are they effective? *American Journal of Preventive Medicine* 2004;26(4):330-43.
- Wakefield M, Banham D, McCaul K, Martin J, Ruffin R, Badcock N, et al. Effect of feedback regarding urinary cotinine and brief tailored advice on home smoking restrictions among low-income parents of children with asthma: a controlled trial. *Preventive Medicine* 2002;34(1):58-65.
- Weinstein ND. The precaution adoption process. *Health Psychol* 1988;7(4):355-86.
- Whitelaw S, Baldwin S, Bunton R, Flynn D. The status of evidence and outcomes in Stages of Change research. *Health Educ Res* 2000 Dec;15(6):707-18.
- Zmirou D, Blatier JF, Andre E, Ferley JP, Balducci F, Rossum F, et al. [Passive smoking respiratory risk. A quantitative synthesis of the literature]. *Rev Mal Respir* 1990;7(4):361-71.



## **Primary Care at a Glance - Hot Topics and New Insights**

Edited by Dr. Oreste Capelli

ISBN 978-953-51-0539-8

Hard cover, 446 pages

**Publisher** InTech

**Published online** 27, April, 2012

**Published in print edition** April, 2012

"Both among scientists and clinical practitioners, some find it easier to rely upon trivial explanations, while others never stop looking for answers". With these surprising words, Augusto Murri, an Italian master in clinical medicine, reminds us that medical practice should be a continuous journey towards knowledge and the quality of care. The book brings together contributions by over 50 authors from many countries, all around the world, from Europe to Africa, from Asia to Australia, from North to South America. Different cultures are presented together, from those with advanced technologies to those of intangible spirituality, but they are all connected by five professional attributes, that in the 1978 the Institute of Medicine (IOM)<sup>1</sup> stated as essentials of practicing good Primary Care: accessibility, comprehensiveness, coordination, continuity and accountability. The content of the book is organized according to these 5 attributes, to give the reader an international overview of hot topics and new insights in Primary Care, all around the world.

### **How to reference**

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Lourdes Rofes Ponce, Ricardo Almon, Elisa Puigdomènech, Manuel A Gomez-Marcos and Carlos Martín-Cantera (2012). Exposure to Environmental Tobacco Smoke in Babies, Primary Care at a Glance - Hot Topics and New Insights, Dr. Oreste Capelli (Ed.), ISBN: 978-953-51-0539-8, InTech, Available from: <http://www.intechopen.com/books/primary-care-at-a-glance-hot-topics-and-new-insights/exposure-to-environmental-tobacco-smoke-in-babies>

**INTECH**  
open science | open minds

### **InTech Europe**

University Campus STeP Ri  
Slavka Krautzeka 83/A  
51000 Rijeka, Croatia  
Phone: +385 (51) 770 447  
Fax: +385 (51) 686 166  
[www.intechopen.com](http://www.intechopen.com)

### **InTech China**

Unit 405, Office Block, Hotel Equatorial Shanghai  
No.65, Yan An Road (West), Shanghai, 200040, China  
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元  
Phone: +86-21-62489820  
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen