

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Distinctive Immunological Functions of HLA-G

Giada Amodio and Silvia Gregori

*San Raffaele Telethon Institute for Gene Therapy (HSR-TIGET),
Division of Regenerative Medicine, Stem Cells and Gene Therapy,
San Raffaele Scientific Institute, Milan,
Italy*

1. Introduction

HLA-G is a non-classical HLA class Ib molecule belonging to the Major Histocompatibility Complex (MHC) located on the short arm of chromosome 6. The tissue-restricted distribution of HLA-G, the low polymorphism in the coding region, the fact that HLA-G primary transcript is alternatively spliced in seven isoforms, and the inhibitory action on immune cells, constitute four hallmarks of HLA-G, which distinguish it from other HLA class I molecules (Carosella et al., 1999). In healthy conditions, a basal level of HLA-G gene transcription is observed in most cells and tissues. However, translation into HLA-G protein is restricted to trophoblasts at the fetal-maternal interface (Carosella et al., 2003), and in adults, to thymic epithelial, cornea, mesenchymal stem cells (MSCs), nail matrix, pancreatic- β cells, erythroid and endothelial precursors. HLA-G can be also neo-expressed in pathological conditions including malignant transformation, viral infections, inflammatory and autoimmune diseases, and allogeneic transplantation (Carosella, 2011).

HLA-G locus is very low polymorphic and, due to its structure, HLA-G molecule can recognize and present only a restricted peptide repertoire compared to classical HLA class I molecules (Clements et al., 2005). These peculiarities render HLA-G exclusively oriented towards immune inhibition and tolerance. In the late ninety, Rouas-Freiss et al. showed that HLA-G expressing trophoblasts were protected from maternal NK cell-mediated cytotoxicity, indicating for the first time HLA-G as a key molecule in fetal-maternal tolerance (Rouas-Freiss et al., 1997). From this first observation, several groups have worked to define the mode of action of HLA-G and in which settings it is involved in promoting tolerance. It is now generally accepted that HLA-G is a unique molecule with several immuno-modulatory properties: it plays an important role not only in fetal-maternal tolerance, but also in modulating immune responses and promoting and maintaining peripheral tolerance in healthy and pathological conditions. HLA-G can indeed act on both the innate and the adaptive branches of immunity and regulate short- and long-term immune-responses.

2. HLA-G molecule

The alternative splicing of HLA-G primary transcript results in seven different isoforms, four of which are membrane bound (HLA-G1 to -G4) and three are soluble (HLA-G5 to -G7) forms (Fujii et al., 1994; Ishitani and Geraghty, 1992) (Fig. 1A).




Fig. 1. HLA-G isoforms, monomers and dimers

A. The alternative splicing of a unique primary transcript yields 7 protein isoforms: truncated isoforms are generated by excision of one or two exons encoding globular (α) domains, whereas translation of intron 4 (i4) or intron 2 (i2) yields soluble isoforms that lack the transmembrane domain. **B.** HLA-G molecules can form homodimers through the generation of Cys42-Cys42 disulfide bonds.

In addition, a soluble HLA-G1 isoform (shed HLA-G1) can be generated by the membrane HLA-G1 proteolytic cleavage that is dependent on metalloproteinase activity (Park et al., 2004), and is regulated by Nitric Oxide (NO) concentration (Diaz-Lagares et al., 2009) and Tumor Necrosis Factor (TNF)- α /NF κ B pathway activation (Zidi et al., 2006). Soluble and membrane-bound HLA-G isoforms have similar functions. HLA-G1, HLA-G5, and shed HLA-G1 are the most described isoforms in healthy tissues (Paul et al., 2000) and their structure is similar to that of classical HLA class I molecules: a heavy chain of three globular domains non-covalently associated with β 2-microglobulin (β 2m) and a nona-peptide (Clements et al., 2005). The other HLA-G isoforms contain the α 1 domain but lack one or two of the other globular domains, and are not associated with β 2m and the peptide. The

presence of the $\alpha 3$ domain is important to HLA-G functions since it represents a binding site for HLA-G receptors (Clements et al., 2005; Clements et al., 2007).

HLA-G possesses two unique cysteine residues, in position 42 of the $\alpha 1$ domain and in position 147 of the $\alpha 2$ domain. Through these residues, HLA-G may dimerize by intramolecular disulfide bonds (Boyson et al., 2002). Membrane-bound or soluble HLA-G dimers were detected both *in vitro* and *in vivo* (Apps et al., 2007; Boyson et al., 2002; Gonen-Gross et al., 2005) (Fig. 1B). Dimerization of HLA-G is one of its key features since dimers bind to HLA-G receptors with higher affinity and slower dissociation rates compared to monomers (Shiroishi et al., 2006a) and, as a consequence, dimers, but not monomers, carry most of the HLA-G functions (Gonen-Gross et al., 2003; Li et al., 2009).

HLA-G acts through three inhibitory receptors: immunoglobulin-like transcript (ILT)2 (CD85j/LILRB1), ILT4 (CD85d/LILRB2), the killer cell immunoglobulin-like receptor (KIR)2DL4 (CD158d), and the co-receptor CD8 (Colonna et al., 1997; Colonna et al., 1998; Rajagopalan and Long, 1999) (Fig. 2).




Fig. 2. HLA-G receptors

Inhibitory receptors known to bind HLA-G. Basic structural organization and expression patterns are shown. The HLA-G structural configuration that these receptors are known to bind are indicated as '+' (reported binding) and '-' (reported absence of binding or minor binding).

ILT2 is expressed by B cells, some T cells (both CD4⁺ and CD8⁺), a sub-population of NK cells, and all monocytes/Dendritic Cells (DCs) (Colonna et al., 1997), whereas ILT4 is only expressed by mono/macrophages and DCs (Colonna et al., 1998). KIR2DL4 is expressed by some CD8⁺ T and NK cells (Goodridge et al., 2003).

Even if ILT2 and ILT4 recognize other HLA Class I molecules, HLA-G is their ligand of highest affinity and they bind HLA-G dimers even more strongly (Shiroishi et al., 2006a; Shiroishi et al., 2003). ILT2 and ILT4 can recognize different HLA-G structures: ILT2 binds only $\beta 2m$ -associated HLA-G1/G5 isoforms, whereas ILT4 also recognizes their $\beta 2m$ -free counterparts (Gonen-Gross et al., 2005; Shiroishi et al., 2006b). Finally, significantly higher expression of ILT2 was necessary for efficient HLA-G tetramer binding, suggesting that this

interaction may have relatively lower affinity compared to that of ILT4 (Allan et al., 1999). Interestingly, HLA-G can directly influence the expression rate of its receptors both at mRNA level and at the protein level (LeMaoult et al., 2005). Thus, HLA-G is unique in its ability to be expressed in different isoforms and to act through inhibitory receptors.

3. Mechanisms underlying HLA-G expression

Even if theoretically any tissue might express HLA-G, its physiological expression is restricted to few tissues such as trophoblasts, thymic epithelium, cornea, MSC and pancreatic β -cells (Carosella, 2011). Nonetheless, in pathological conditions, HLA-G expression can be induced and/or up-regulated. So far the reason why HLA-G can be expressed in some, but not in all tissues has not been fully elucidated. However, several factors such as immune-modulatory cytokines and hormones, were described to potentially affect transcriptional mechanisms responsible for HLA-G expression (Moreau et al., 2009). In contrast to classical HLA class I molecules, HLA-G expression is not influenced by the transcription factor NF- κ B (Gobin et al., 1998) or by classical pro-inflammatory cytokines such as Interferon (IFN)- γ (Gobin et al., 1999), since responsive elements for IFN- γ are missed in HLA-G promoter (Gobin et al., 1999; Steimle et al., 1995). In contrast, HLA-G transcriptional rate is increased by the presence of a number of anti-inflammatory cytokines including IFN- β (Lefebvre et al., 2001), Interleukin (IL)-4, IL-5, IL-6 (Deschaseaux et al., 2011), and IL-10 (Moreau et al., 1999) and can be modulated by Granulocyte Macrophage Colony-Stimulating Factor (GM-CSF) (Onno et al., 2000; Yang et al., 1996), Transforming Growth Factor (TGF)- β , or Granulocyte Colony-Stimulating Factor (G-CSF).

In addition to cytokines, hormones like glucocorticoids (dexamethasone) and progesterone were shown to increase the secretion of both HLA-G5 and HLA-G6 by trophoblasts (Akhter et al., 2011; Moreau et al., 2001; Yie et al., 2006a; Yie et al., 2006b). HLA-G expression can be also influenced by Adenosine Triphosphate (ATP) and by the tryptophan catalyzing enzyme indoleamine 2,3-dioxygenase (IDO). ATP acts as a potent inhibitor of HLA-G1 and HLA-G5 production from LPS-activated PBMCs *via* down-regulation of IL-10 (Rizzo et al., 2009). IDO, known to be involved in the generation of tolerogenic microenvironments by tryptophan depletion and in the generation of T regulatory (Treg) cells (Chung et al., 2009; Munn et al., 2002), was shown to differentially modulate HLA-G expression in monocytes and myeloid DCs. IDO blocked HLA-G cell-surface expression on monocytes (Gonzalez-Hernandez et al., 2005), but it induced HLA-G expression and shedding in myeloid DCs (Lopez et al., 2006). Overall, a number of immune mediators can influence HLA-G expression. It has to be taken in account that these molecules can be up-regulated in conditions, such as inflammation, in which HLA-G expression is needed to control or dampen the immune response.

In addition to soluble factors, it has been recently reported that “transient” HLA-G expression can be acquired *via* trogocytosis, a cell-to-cell contact-dependent uptake of membranes and associated molecules from one cell by another (reviewed in (Davis, 2007; LeMaoult et al., 2007a). Trogocytosis of HLA-G was shown for activated CD4⁺ and CD8⁺ T cells (LeMaoult et al., 2007b), activated NK cells (Caumartin et al., 2007), and monocytes (HoWangYin et al., 2011). For CD4⁺ T cells, the most important consequence of HLA-G acquisition, is that the newly-expressing cells became transiently tolerogenic (LeMaoult et al., 2007a), as discussed below. Thus, trogocytosis generates HLA-G positive cells, which

display, but do not express HLA-G. The ability of different cells to transiently acquire HLA-G expression broadens the immuno-modulatory activity of HLA-G.

4. HLA-G immunological function

The functions of HLA-G are exclusively oriented towards immune inhibition and tolerance. HLA-G exerts its inhibitory functions on several immune cells through direct binding to the inhibitory receptors ILT2, ILT4, and KIR2DL4 (Shiroishi et al., 2006b).

4.1 Short-term immuno-modulation mediated by HLA-G

HLA-G:ILT2 interaction modulates NK cell activity (Rouas-Freiss et al., 1997). HLA-G-expressing target cells are indeed resistant to the lysis mediated by NK cells (Riteau et al., 2001). Although the mechanisms underlying this effect are not completely elucidated, it has been recently defined that HLA-G:ILT2 interaction disrupts the immunological synapse (IS), a supramolecular structure responsible for both T and NK cell activation and function (Favier et al., 2010), leading to inhibition of NK cytotoxicity. HLA-G binds to KIR2DL4 on NK cells (Shiroishi et al., 2006b). However, it still remains unclear which are the effects mediated by this interaction. Ligation of KIR2DL4 activates cytokine production but not cytotoxicity (Rajagopalan et al., 2001), and it has been recently proposed that activation of KIR2DL4 by soluble HLA-G activates a pro-inflammatory/pro-angiogenic responses, consistent with a role in promoting vascularization during early pregnancy (Rajagopalan et al., 2006). HLA-G can also bind CD8, which is expressed by a subpopulation of NK cells leading to Fas ligand expression, and induction of apoptosis (Contini et al., 2003).

The interaction between HLA-G and CD8⁺ T cells impairs the antigen-specific cytotoxic T lymphocyte (CTLs) activity (Le Gal et al., 1999). HLA-G has a dose-dependent effect on the generation of allo-CTL responses, while it seems to not affect pre-existing allo-CTLs (Kapasi et al., 2000). Furthermore, during later stages of T cell activation, HLA-G:CD8 interaction promotes TCR-independent apoptosis of CD8⁺ T cells through the same mechanisms described for NK cells (Fournel et al., 2000).

During the early phase of CD4⁺ T cell activation, HLA-G:ILT2 interaction induces a cell cycle blockade at the G1 phase (Bahri et al., 2006), and inhibits allo-reactive T-cell proliferation (Naji et al., 2007a). Interestingly, it has been proposed that HLA-G isoforms have different impact on the cytokine secretion profile of CD4⁺ T cells. HLA-G1 was indeed shown to promote T helper (h)2 polarization of naïve T cells (Kanai et al., 2001b; Kapasi et al., 2000) whereas HLA-G5 induces tumor necrosis factor (TNF)- α , IFN- γ , and IL-10 (Kanai et al., 2001a). Morandi et al. recently described that HLA-G5 has an additional effect on CD4⁺ T cells since it down-regulates expression and function of CCR2, CXCR3, and CXCR4 on different subsets of activated CD4⁺ T cells, impairing their migratory capability (Morandi et al., 2010).

The immunological functions of HLA-G on B cells are poorly described. It has been shown that in patients who underwent renal transplantation, serum levels of soluble HLA-G are negatively associated with anti-HLA antibodies and with graft rejection, suggesting that soluble HLA-G may possibly act to inhibit the immune humoral responses against HLA (Qiu et al., 2006). Further investigation is needed to better highlight the role of HLA-G in modulating B cell responses.

Myeloid antigen presenting cells (APCs) express both ILT2 and ILT4, which render them targets of HLA-G-mediated modulation. In the presence of HLA-G, myeloid APCs fail to stimulate allo-proliferative T cell responses *in vitro* (LeMaoult et al., 2004) and reduce NK-mediated cytotoxicity (Gros et al., 2008). Moreover, HLA-G inhibits the up-regulation of HLA Class II, CD80 and CD86 on myeloid DCs in response to LPS or allo-activation signals *in vitro* (Ristich et al., 2007) (Gros et al., 2008). Thus, HLA-G inhibits the functions and differentiation of myeloid APCs, leading to improper T lymphocyte activation and to impair NK cytotoxic activity. However, HLA-G does not block myeloid APCs functions, but induces them to differentiate into tolerogenic cells. Several reports indeed show that HLA-G treatment up-regulates the expression of cytokines, chemokines, and chemokine receptors by myeloid APCs (Apps et al., 2007; Gros et al., 2008; Li et al., 2009; Liang et al., 2008). The first evidence that HLA-G promotes the induction of tolerogenic APCs comes from studies conducted by the group of Horuzsko demonstrating that HLA-G treatment of monocyte-derived DCs that highly express ILT2 and ILT4 results in cells with tolerogenic-like phenotype and the potential to induce T-cell anergy (Ristich et al., 2005). Results obtained in a murine model support this notion by showing that HLA-G inhibited maturation of DCs (Horuzsko et al., 2001) and that in ILT4 transgenic mice, the HLA-G:ILT4 interaction impaired DC maturation *in vivo*, leading to delayed skin allograft rejection (Liang et al., 2002). Thereby, HLA-G by interacting with ILT receptors present on myeloid APCs induces their differentiation into regulatory cells.

4.2 Long-term tolerogenic properties of HLA-G

Myeloid APCs are not only the cells more sensitive to HLA-G-mediated modulation, but they also commonly express HLA-G. Myeloid APCs may express all HLA-G isoforms (Le Friec et al., 2004), but cell-surface HLA-G1 and secreted HLA-G5 have been described the most. In pathological contexts including transplantation, cancer, viral infections, and inflammatory diseases, the expression of HLA-G on myeloid cells can be up-regulated (Carosella et al., 2008). Recently, in liver-transplanted patients, the presence of HLA-G-expressing myeloid DCs has been correlated with tolerance and graft acceptance (Castellaneta et al., 2011). Similarly, in cancer (reviewed in (Amiot et al., 2011) as well as in viral infections (reviewed in (Fainardi et al., 2011), myeloid APCs expressing HLA-G were reported, and often correlated with poor clinical outcomes. Myeloid APCs not only express membrane-bound HLA-G1 but they may also secrete or shed HLA-G molecules, contributing to the generation of a tolerogenic microenvironment. Such a microenvironment may alter the functions not only of lymphocytes, but also of the HLA-G-expressing myeloid APCs themselves, in a tolerogenic feedback loop. HLA-G can indeed directly promote its expression and the expression of its receptors on myeloid APCs (LeMaoult et al., 2005). Thus, myeloid APCs may be viewed as suppressor cells capable of inhibiting other immune effectors, and also of generating regulatory cells including Treg cells.

Treg cells are critical players for preservation of immune homeostasis and for establishment and maintenance of peripheral tolerance. Treg cells belonging to different T cell subsets, including CD4⁺ and CD8⁺, NKT, and $\gamma\delta$ T cells. The best characterized Treg populations are the forkhead box P3 (FOXP3)-expressing, CD4⁺CD25⁺ Tregs (FOXP3⁺ Tregs) (Sakaguchi et al., 2010), and the CD4⁺ IL-10-producing Tr1 cells (Akdis et al., 2004; Barrat et al., 2002; Groux et al., 1997; Roncarolo et al., 2006). FOXP3⁺ Tregs and Tr1 cells

are distinct populations of Treg cells that play a non-redundant role in maintaining tolerance, and are distinguished from one another by their distinct phenotype and cytokine profiling. FOXP3⁺ Treg cells are characterized by the constitutively-high expression of CD25 and the transcription factor FOXP3 (Sakaguchi, 2005), and their development and function is strictly dependent on FOXP3 (Bacchetta et al., 2007; Gavin et al., 2007). Tr1 cells can be induced in the periphery upon chronic antigen (Ag) stimulation in the presence of IL-10 (Roncarolo et al., 2006), and are currently identified by their unique cytokine profile consisting of high levels of IL-10, TGF- β , and low levels of IL-2 and IFN- γ and the absence of IL-4 after stimulation (Bacchetta et al., 1994; Groux et al., 1997). Once activated through their specific TCR, Tr1 cells secrete IL-10 and TGF- β that directly inhibit effector T cell proliferation and expression of HLA class II and co-stimulatory molecules on APCs, which indirectly suppress effector T-cell activation. More recently, we demonstrated that Tr1 cells specifically kill myeloid cells *via* a Granzyme B-dependent mechanism (Magnani et al., 2011).

HLA-G-expressing APCs were shown to be capable of priming naïve T cells to become Treg cells. Indeed, APC lines over-expressing membrane-bound HLA-G1 induced the differentiation of CD4⁺ and CD8⁺ T cells able to inhibit allogeneic responses (LeMaoult et al., 2004). Resulting HLA-G-induced Treg cells included CD4^{low} and CD8^{low} T cells that suppress *via* soluble factors (Naji et al., 2007a; Naji et al., 2007b). These results are in agreement with those obtained in patients who received a combined liver-kidney transplant, in which high plasma levels of HLA-G5 correlated with an increased percentage of suppressor T cells (Le Rond et al., 2004; Naji et al., 2007b). They are also in line with results showing that high HLA-G5 plasma levels in the peripheral blood of stem cell-transplanted patients are associated with the expansion in peripheral blood of CD4⁺CD25⁺CD152⁺ T cells with suppressive activity (Le Maux et al., 2008).

We recently identified and characterized a new subset of human DCs that arises in the presence of IL-10 and endogenously expresses cell-surface HLA-G (Gregori et al., 2010). DC-10 are characterized by the outstanding ability to produce IL-10 and by the expression of high levels of membrane-bound HLA-G1 and other tolerogenic signaling molecules such as ILT2, ILT3, and ILT4 (Fig. 3).

DC-10 are potent inducers of adaptive allo-specific Tr1 cells (Gregori et al., 2010). Furthermore, allergen-specific Tr1 cells can be generated *in vitro* by stimulating human T cells with autologous tolerogenic DC-10 pulsed with allergen (Pacciani et al., 2010). Interestingly, the expression of membrane-bound HLA-G1 and that of its receptors are up-regulated by IL-10 on both DC-10 and T cells, and the expression of high levels of membrane-bound HLA-G1, ILT4, and IL-10 by DC-10 is critical to the generation of Tr1 cells by DC-10 (Rossetti et al., 2011).

5. HLA-G expressing regulatory cells

In addition to myeloid cells that can constitutively or transiently express HLA-G, T cells or other immune cells can express or secrete HLA-G.

A subpopulation of CD4⁺HLA-G1⁺ and CD8⁺HLA-G1⁺ T cells has been described (Feger et al., 2007). They represent a small subset of peripheral blood cells in healthy individuals, but




Fig. 3. A unique subset of human tolerogenic DC: DC-10

A. DC-10 are present *in vivo* and characterized by the expression of the surface markers CD14, CD16, CD83. They also express high levels of HLA-G1 and its receptor ILT4. **B.** DC-10 can be differentiated *in vitro* starting from the CD14⁺ fraction of PBMCs cultured for 7 days in the presence of GM-CSF, IL-4 and IL-10.

are increased at sites of inflammation, such as in the central nervous system of patients with neuro-inflammatory disorders and in muscle tissue in patients with idiopathic myositis (Feger et al., 2007; Huang et al., 2009a; Wiendl et al., 2005). CD4⁺ HLA-G⁺ T cells represent a population of naturally occurring Treg cells distinct from nTreg cells, since they lack CD25 and FOXP3 expression. HLA-G⁺ Treg cells showed reduced proliferation in response to allogeneic (mDCs) and polyclonal stimulation (α CD3/CD28), and have a cytokine profiling different from Tr1 cells. CD4⁺ HLA-G⁺ and CD8⁺ HLA-G⁺ Treg cells inhibit the proliferation of autologous HLA-G⁻ T cells through HLA-G1/sHLA-G1 (Feger et al., 2007) and soluble factors, including IL-10 (Huang et al., 2009b). A population of induced HLA-G⁺ T cells has been also described, these allo-specific CD4⁺ and CD8⁺ T cells were shown to express HLA-G1 and to secrete HLA-G5 (Le Rond et al., 2004; Lila et al., 2001). Moreover, we demonstrated that up-regulation of HLA-G on CD4⁺ T cells is critically required for DC-10 mediated induction of adoptive Tr1 cells (Gregori et al., 2010).

Adult bone marrow MSCs can also express HLA-G and have been included in the list of regulatory cells. MSCs are multipotent cells that play an important role in tissue regeneration and also have strong immuno-modulatory properties (reviewed in (Uccelli et al., 2008)). MSCs express low levels of HLA Class I molecules, do not express HLA class II or co-stimulatory molecules, and therefore do not induce T-cell activation. It has been demonstrated that MSCs constitutively express HLA-G5 (Selmani et al., 2009). Even though HLA-G expression decrease during MSCs expansion *in vitro*, recent studies clearly identified it as a key contributor to MSCs immune-tolerogenic functions (Nasef et al., 2007; Rizzo et al.,

2008; Selmani et al., 2008). In addition to a direct suppressive function through HLA-G5, MSCs can induce *via* a cell-to-cell contact mechanism CD4⁺ CD25^{high} Foxp3⁺ Treg cells (Selmani et al., 2008).

6. HLA-G and its polymorphisms

Very few polymorphisms have been identified in the coding region of the HLA-G locus. The heavy chain encoding region exhibits 33 Single Nucleotide Polymorphisms (SNPs) but only 13 amino acid variations are observed. This peculiarity is important in determining the biological function of HLA-G, since a reduced variability leads to a limited peptide repertoire and presentation capability (Clements et al., 2005), and can influence the polymerization rate. Moreover, the 5' Up-stream Regulatory Region (5' URR), containing the promoter, and the 3' untranslated region (UTR) exhibit numerous nucleotide variations that may influence HLA-G expression (Donadi et al., 2011; Larsen and Hviid, 2009) and consequently its tissue distribution in healthy and pathological conditions (Fig. 4).




Fig. 4. HLA-G locus and its polymorphisms

Schematic representation of HLA-G locus. The major polymorphisms observed at the 5'URR and at 3'UTR regions are indicated. URR: Up-stream regulatory region; UTR: Untranslated region.

6.1 5' up-stream regulatory region (URR)

Because of the presence of regulatory elements, polymorphisms at the 5' URR might have a relevant impact in the regulation of HLA-G expression (Fig. 4). The region up-stream from the transcriptional start site of HLA-G contains 27 different polymorphisms but only 13 of them are within or very close to known transcription factor binding sites or regulatory elements (Hviid et al., 1999; Ober et al., 2003; Tan et al., 2005). Interestingly, all the 27 polymorphisms are in strong linkage disequilibrium (LD) and define 13 unique promoter haplotypes (Ober et al., 2003; Tan et al., 2005). Among the 5' URR polymorphisms, only the -725C/G/T single nucleotide polymorphism (SNP) has been associated with HLA-G expression. Ober et al. demonstrated that the presence of G in position -725 significantly increases HLA-G transcription rate in JEG-3 cell line (Ober et al., 2006) and, when it is present in both parents, it is significantly associated with fetal loss rate (Ober et al., 2003).

An additional interesting SNP located in 5' URR is the -201 A/G SNP since it resides into the enhancer A region (Hviid et al., 1999). However, it still remains to be clarified if this polymorphism has any impact on the expression of certain HLA-G alleles.

6.2 3'untranslated region (UTR)

The exon 7 of the HLA-G locus is always absent in the mature messenger(m)RNA. Moreover, due to the presence of a stop codon in the exon 6 of HLA-G, the exon 8 is not translated. Thus, exon 8 is considered the 3' UTR of the mature RNA. This region contains several regulatory elements (Kuersten and Goodwin, 2003), including polyadenylation signals, AU-rich elements (Alvarez et al., 2009), and several polymorphic sites that may potentially influence mRNA stability, turnover, mobility and splicing pattern. 3'UTR polymorphisms that can influence HLA-G expression are: i) the insertion (INS) or deletion (DEL) of a 14bp fragment (14bp INS/DEL) that has been associated with mRNA stability; ii) the SNP at position +3142, which may be a target for certain microRNAs (miRNAs), small RNA sequences which once bound to a complementary mRNA lead to its translational repression or degradation and gene silencing (Bartel, 2009); iii) additional six SNPs landed in putative binding sites for miRNAs (Castelli et al., 2009) (Fig.4).

The first identified 3'UTR HLA-G polymorphism is the 14bp INS/DEL and has been studied the most. The insertion of the 14bp fragment, results in the formation of a cryptic breakpoint in the mRNA that loose the first 92bp of exon 8 (Hviid et al., 2003). Rousseau et al. demonstrated that HLA-G transcript with the 92bp deleted seems to be more stable than the complete mRNA fragment generated by the 14bp DEL (Rousseau et al., 2003), suggesting that 14bp INS might be associated with high levels of HLA-G expression. Nonetheless, several groups have demonstrated that 14 bp INS/INS genotype is associated with lower serum and plasma level of sHLA-G1/HLA-G5 compared to those observed in 14bp INS/DEL and 14bp DEL/DEL genotypes (Chen et al., 2008; Hviid et al., 2004; Hviid et al., 2006), an observation that constitutes the "14bp paradox". Consistently, conflicting results have been obtained in different studies concerning the association of the 14bp HLA-G genotypes with autoimmune diseases, pathological pregnancy, recurrent spontaneous abortions, and pre-eclampsia. Notably, the presence of 14bp DEL has been found to be predictive for the incidence of GvHD after unrelated (La Nasa et al., 2007) and HLA-identical sibling (Sizzano et al., in press) Human Stem Cell Transplantation (HSCT) for beta-thalassemia.

Several SNPs at 3'UTR of HLA-G locus have been identified (Castelli et al., 2009). Among them, the +3142 C/G SNP has been proposed to be critically involved in HLA-G regulation since it is associated with asthma (Tan et al., 2005). The G variant of a G/C SNP at position +3142 of the 3'UTR has been hypothesized to increase the affinity of the resulting mRNA for miR-148a, miR-148b and miR-152 (Veit and Chies, 2009). Among these miRNAs, only miR-152 has been demonstrated responsible for HLA-G post-transcriptional regulation. Indeed, over-expression of miR-152 in JEG-3 cell lines resulted in decreasing HLA-G expression and increased susceptibility to NK cell-mediated cytotoxicity (Zhu et al., 2010).

In silico analysis of the 3'UTR of the HLA-G locus revealed the presence of additional six SNPs that are landed in putative binding sites for miRNAs (Castelli et al., 2009). Interestingly the 14bp INS/DEL and the 7 SNPs arrange in different combinations to generate eight distinct haplotypes of 3'UTR HLA-G region (Castelli et al., 2010) (Table 1).

Therefore, it has been hypothesized that the expression of HLA-G might be influenced by a combination of specific polymorphisms. Our group is currently investigating the impact of the HLA-G 3'UTR haplotypes on HLA-G expression. Preliminary results indicate that different 3'UTR HLA-G haplotypes are associated with variable levels of HLA-G1 expression on myeloid APCs.

Haplotype	14-bp	+3003	+3010	+3027	+3035	+3142
UTR-1	Del	T	G	C	C	C
UTR-2	Ins	T	C	C	C	G
UTR-3	Del	T	C	C	C	G
UTR-4	Del	C	G	C	C	C
UTR-5	Ins	T	C	C	T	G
UTR-6	Del	T	G	C	C	C
UTR-7	Ins	T	C	A	T	G
UTR-8	Ins	T	G	C	C	G

Table 1. 3'UTR HLA-G haplotypes

7. Conclusions and perspectives

Since the discovery of HLA-G, research has established its role in modulating immune responses and inducing tolerance. While a great deal of progress has been made in understanding the mechanisms underlying immune modulation by HLA-G, several questions remain to be clarified. First, HLA-G can promote the differentiation of regulatory cells and HLA-G-expressing cells act as regulatory cells, however additional studies are required to better define their role in promoting tolerance. Second, although it is generally accepted that 5' URR and 3' UTR are involved in HLA-G expression, further investigation are needed to elucidate the molecular mechanisms involved in HLA-G regulation. Third, 3'UTR HLA-G haplotypes have been identified, future studies are warranted to define whether they can influence the transcriptional rate of HLA-G isoforms and their functions. Answering these questions will not only bring us closer to understanding how HLA-G function, but also how to exploit or modulate its suppressive activity for targeted therapy against a wide variety of diseases.

8. Acknowledgement

This work was supported by Telethon Foundation, the Associazione Italiana per la Ricerca sul Cancro (AIRC) IG 8978, and the Italian Ministry of Health.

9. References

Akdis, M., Verhagen, J., Taylor, A., Karamloo, F., Karagiannidis, C., Cramer, R., Thunberg, S., Deniz, G., Valenta, R., Fiebig, H., *et al.* (2004). Immune responses in healthy and

- allergic individuals are characterized by a fine balance between allergen-specific T regulatory 1 and T helper 2 cells. *The Journal of experimental medicine* 199, 1567-1575.
- Akhter, A., Das, V., Naik, S., Faridi, R.M., Pandey, A., and Agrawal, S. (2011). Upregulation of HLA-G in JEG-3 cells by dexamethasone and hydrocortisone. *Archives of gynecology and obstetrics*.
- Allan, D.S., Colonna, M., Lanier, L.L., Churakova, T.D., Abrams, J.S., Ellis, S.A., McMichael, A.J., and Braud, V.M. (1999). Tetrameric complexes of human histocompatibility leukocyte antigen (HLA)-G bind to peripheral blood myelomonocytic cells. *The Journal of experimental medicine* 189, 1149-1156.
- Alvarez, M., Piedade, J., Balseiro, S., Ribas, G., and Regateiro, F. (2009). HLA-G 3'-UTR SNP and 14-bp deletion polymorphisms in Portuguese and Guinea-Bissau populations. *Int J Immunogenet* 36, 361-366.
- Amiot, L., Ferrone, S., Grosse-Wilde, H., and Seliger, B. (2011). Biology of HLA-G in cancer: a candidate molecule for therapeutic intervention? *Cell Mol Life Sci* 68, 417-431.
- Apps, R., Gardner, L., Sharkey, A.M., Holmes, N., and Moffett, A. (2007). A homodimeric complex of HLA-G on normal trophoblast cells modulates antigen-presenting cells via LILRB1. *Eur J Immunol* 37, 1924-1937.
- Bacchetta, R., Bigler, M., Touraine, J.L., Parkman, R., Tovo, P.A., Abrams, J., de Waal Malefyt, R., de Vries, J.E., and Roncarolo, M.G. (1994). High levels of interleukin 10 production in vivo are associated with tolerance in SCID patients transplanted with HLA mismatched hematopoietic stem cells. *The Journal of experimental medicine* 179, 493-502.
- Bacchetta, R., Gambineri, E., and Roncarolo, M.G. (2007). Role of regulatory T cells and FOXP3 in human diseases. *J Allergy Clin Immunol* 120, 227-235; quiz 236-227.
- Bahri, R., Hirsch, F., Josse, A., Rouas-Freiss, N., Bidere, N., Vasquez, A., Carosella, E.D., Charpentier, B., and Durrbach, A. (2006). Soluble HLA-G inhibits cell cycle progression in human alloreactive T lymphocytes. *J Immunol* 176, 1331-1339.
- Barrat, F.J., Cua, D.J., Boonstra, A., Richards, D.F., Crain, C., Savelkoul, H.F., de Waal-Malefyt, R., Coffman, R.L., Hawrylowicz, C.M., and O'Garra, A. (2002). In vitro generation of interleukin 10-producing regulatory CD4(+) T cells is induced by immunosuppressive drugs and inhibited by T helper type 1 (Th1)- and Th2-inducing cytokines. *The Journal of experimental medicine* 195, 603-616.
- Bartel, D.P. (2009). MicroRNAs: target recognition and regulatory functions. *Cell* 136, 215-233.
- Boyson, J.E., Erskine, R., Whitman, M.C., Chiu, M., Lau, J.M., Koopman, L.A., Valter, M.M., Angelisova, P., Horejsi, V., and Strominger, J.L. (2002). Disulfide bond-mediated dimerization of HLA-G on the cell surface. *Proc Natl Acad Sci U S A* 99, 16180-16185.
- Carosella, E.D. (2011). The tolerogenic molecule HLA-G. *Immunol Lett*.
- Carosella, E.D., Favier, B., Rouas-Freiss, N., Moreau, P., and Lemaoult, J. (2008). Beyond the increasing complexity of the immunomodulatory HLA-G molecule. *Blood* 111, 4862-4870.
- Carosella, E.D., Moreau, P., Le Maoult, J., Le Discorde, M., Dausset, J., and Rouas-Freiss, N. (2003). HLA-G molecules: from maternal-fetal tolerance to tissue acceptance. *Adv Immunol* 81, 199-252.

- Carosella, E.D., Rouas-Freiss, N., Paul, P., and Dausset, J. (1999). HLA-G: a tolerance molecule from the major histocompatibility complex. *Immunol Today* 20, 60-62.
- Castellaneta, A., Mazariegos, G.V., Nayyar, N., Zeevi, A., and Thomson, A.W. (2011). HLA-G level on monocytoïd dendritic cells correlates with regulatory T-cell Foxp3 expression in liver transplant tolerance. *Transplantation* 91, 1132-1140.
- Castelli, E.C., Mendes-Junior, C.T., Deghaide, N.H., de Albuquerque, R.S., Muniz, Y.C., Simoes, R.T., Carosella, E.D., Moreau, P., and Donadi, E.A. (2010). The genetic structure of 3'untranslated region of the HLA-G gene: polymorphisms and haplotypes. *Genes Immun* 11, 134-141.
- Castelli, E.C., Moreau, P., Oya e Chiromatzo, A., Mendes-Junior, C.T., Veiga-Castelli, L.C., Yaghi, L., Giuliatti, S., Carosella, E.D., and Donadi, E.A. (2009). In silico analysis of microRNAs targeting the HLA-G 3' untranslated region alleles and haplotypes. *Human immunology* 70, 1020-1025.
- Caumartin, J., Favier, B., Daouya, M., Guillard, C., Moreau, P., Carosella, E.D., and LeMaoult, J. (2007). Trogocytosis-based generation of suppressive NK cells. *EMBO J* 26, 1423-1433.
- Chen, X.Y., Yan, W.H., Lin, A., Xu, H.H., Zhang, J.G., and Wang, X.X. (2008). The 14 bp deletion polymorphisms in HLA-G gene play an important role in the expression of soluble HLA-G in plasma. *Tissue Antigens* 72, 335-341.
- Chung, D.J., Rossi, M., Romano, E., Ghith, J., Yuan, J., Munn, D.H., and Young, J.W. (2009). Indoleamine 2,3-dioxygenase-expressing mature human monocyte-derived dendritic cells expand potent autologous regulatory T cells. *Blood* 114, 555-563.
- Clements, C.S., Kjer-Nielsen, L., Kostenko, L., Hoare, H.L., Dunstone, M.A., Moses, E., Freed, K., Brooks, A.G., Rossjohn, J., and McCluskey, J. (2005). Crystal structure of HLA-G: a nonclassical MHC class I molecule expressed at the fetal-maternal interface. *Proc Natl Acad Sci U S A* 102, 3360-3365.
- Clements, C.S., Kjer-Nielsen, L., McCluskey, J., and Rossjohn, J. (2007). Structural studies on HLA-G: implications for ligand and receptor binding. *Human immunology* 68, 220-226.
- Colonna, M., Navarro, F., Bellon, T., Llano, M., Garcia, P., Samaridis, J., Angman, L., Cella, M., and Lopez-Botet, M. (1997). A common inhibitory receptor for major histocompatibility complex class I molecules on human lymphoid and myelomonocytic cells. *The Journal of experimental medicine* 186, 1809-1818.
- Colonna, M., Samaridis, J., Cella, M., Angman, L., Allen, R.L., O'Callaghan, C.A., Dunbar, R., Ogg, G.S., Cerundolo, V., and Rolink, A. (1998). Human myelomonocytic cells express an inhibitory receptor for classical and nonclassical MHC class I molecules. *J Immunol* 160, 3096-3100.
- Contini, P., Ghio, M., Poggi, A., Filaci, G., Indiveri, F., Ferrone, S., and Puppo, F. (2003). Soluble HLA-A,-B,-C and -G molecules induce apoptosis in T and NK CD8+ cells and inhibit cytotoxic T cell activity through CD8 ligation. *Eur J Immunol* 33, 125-134.
- Davis, D.M. (2007). Intercellular transfer of cell-surface proteins is common and can affect many stages of an immune response. *Nat Rev Immunol* 7, 238-243.
- Deschaseaux, F., Delgado, D., Pistoia, V., Giuliani, M., Morandi, F., and Durrbach, A. (2011). HLA-G in organ transplantation: towards clinical applications. *Cell Mol Life Sci* 68, 397-404.

- Diaz-Lagares, A., Alegre, E., LeMaoult, J., Carosella, E.D., and Gonzalez, A. (2009). Nitric oxide produces HLA-G nitration and induces metalloprotease-dependent shedding creating a tolerogenic milieu. *Immunology* 126, 436-445.
- Donadi, E.A., Castelli, E.C., Arnaiz-Villena, A., Roger, M., Rey, D., and Moreau, P. (2011). Implications of the polymorphism of HLA-G on its function, regulation, evolution and disease association. *Cell Mol Life Sci* 68, 369-395.
- Fainardi, E., Castellazzi, M., Stignani, M., Morandi, F., Sana, G., Gonzalez, R., Pistoia, V., Baricordi, O.R., Sokal, E., and Pena, J. (2011). Emerging topics and new perspectives on HLA-G. *Cell Mol Life Sci* 68, 433-451.
- Favier, B., Lemaoult, J., Lesport, E., and Carosella, E.D. (2010). ILT2/HLA-G interaction impairs NK-cell functions through the inhibition of the late but not the early events of the NK-cell activating synapse. *FASEB J* 24, 689-699.
- Feger, U., Tolosa, E., Huang, Y.H., Waschbisch, A., Biedermann, T., Melms, A., and Wiendl, H. (2007). HLA-G expression defines a novel regulatory T-cell subset present in human peripheral blood and sites of inflammation. *Blood* 110, 568-577.
- Fournel, S., Aguerre-Girr, M., Huc, X., Lenfant, F., Alam, A., Toubert, A., Bensussan, A., and Le Bouteiller, P. (2000). Cutting edge: soluble HLA-G1 triggers CD95/CD95 ligand-mediated apoptosis in activated CD8⁺ cells by interacting with CD8. *J Immunol* 164, 6100-6104.
- Fujii, T., Ishitani, A., and Geraghty, D.E. (1994). A soluble form of the HLA-G antigen is encoded by a messenger ribonucleic acid containing intron 4. *J Immunol* 153, 5516-5524.
- Gavin, M.A., Rasmussen, J.P., Fontenot, J.D., Vasta, V., Manganiello, V.C., Beavo, J.A., and Rudensky, A.Y. (2007). Foxp3-dependent programme of regulatory T-cell differentiation. *Nature* 445, 771-775.
- Gobin, S.J., Keijsers, V., van Zutphen, M., and van den Elsen, P.J. (1998). The role of enhancer A in the locus-specific transactivation of classical and nonclassical HLA class I genes by nuclear factor kappa B. *J Immunol* 161, 2276-2283.
- Gobin, S.J., van Zutphen, M., Woltman, A.M., and van den Elsen, P.J. (1999). Transactivation of classical and nonclassical HLA class I genes through the IFN-stimulated response element. *J Immunol* 163, 1428-1434.
- Gonen-Gross, T., Achdout, H., Arnon, T.I., Gazit, R., Stern, N., Horejsi, V., Goldman-Wohl, D., Yagel, S., and Mandelboim, O. (2005). The CD85J/leukocyte inhibitory receptor-1 distinguishes between conformed and beta 2-microglobulin-free HLA-G molecules. *J Immunol* 175, 4866-4874.
- Gonen-Gross, T., Achdout, H., Gazit, R., Hanna, J., Mizrahi, S., Markel, G., Goldman-Wohl, D., Yagel, S., Horejsi, V., Levy, O., *et al.* (2003). Complexes of HLA-G protein on the cell surface are important for leukocyte Ig-like receptor-1 function. *J Immunol* 171, 1343-1351.
- Gonzalez-Hernandez, A., LeMaoult, J., Lopez, A., Alegre, E., Caumartin, J., Le Rond, S., Daouya, M., Moreau, P., and Carosella, E.D. (2005). Linking two immunosuppressive molecules: indoleamine 2,3 dioxygenase can modify HLA-G cell-surface expression. *Biol Reprod* 73, 571-578.
- Goodridge, J.P., Witt, C.S., Christiansen, F.T., and Warren, H.S. (2003). KIR2DL4 (CD158d) genotype influences expression and function in NK cells. *J Immunol* 171, 1768-1774.

- Gregori, S., Tomasoni, D., Pacciani, V., Scirpoli, M., Battaglia, M., Magnani, C.F., Hauben, E., and Roncarolo, M.G. (2010). Differentiation of type 1 T regulatory cells (Tr1) by tolerogenic DC-10 requires the IL-10-dependent ILT4/HLA-G pathway. *Blood* 116, 935-944.
- Gros, F., Cabillic, F., Toutirais, O., Maux, A.L., Sebti, Y., and Amiot, L. (2008). Soluble HLA-G molecules impair natural killer/dendritic cell crosstalk via inhibition of dendritic cells. *Eur J Immunol* 38, 742-749.
- Groux, H., O'Garra, A., Bigler, M., Rouleau, M., Antonenko, S., de Vries, J.E., and Roncarolo, M.G. (1997). A CD4⁺ T-cell subset inhibits antigen-specific T-cell responses and prevents colitis. *Nature* 389, 737-742.
- Horuzsko, A., Lenfant, F., Munn, D.H., and Mellor, A.L. (2001). Maturation of antigen-presenting cells is compromised in HLA-G transgenic mice. *Int Immunol* 13, 385-394.
- HoWangYin, K.Y., Alegre, E., Daouya, M., Favier, B., Carosella, E.D., and LeMaoult, J. (2011). Different functional outcomes of intercellular membrane transfers to monocytes and T cells. *Cell Mol Life Sci* 67, 1133-1145.
- Huang, Y.H., Zozulya, A.L., Weidenfeller, C., Metz, I., Buck, D., Toyka, K.V., Bruck, W., and Wiendl, H. (2009a). Specific central nervous system recruitment of HLA-G(+) regulatory T cells in multiple sclerosis. *Ann Neurol* 66, 171-183.
- Huang, Y.H., Zozulya, A.L., Weidenfeller, C., Schwab, N., and Wiendl, H. (2009b). T cell suppression by naturally occurring HLA-G-expressing regulatory CD4⁺ T cells is IL-10-dependent and reversible. *Journal of leukocyte biology*.
- Hviid, T.V., Hylenius, S., Rorbye, C., and Nielsen, L.G. (2003). HLA-G allelic variants are associated with differences in the HLA-G mRNA isoform profile and HLA-G mRNA levels. *Immunogenetics* 55, 63-79.
- Hviid, T.V., Rizzo, R., Christiansen, O.B., Melchiorri, L., Lindhard, A., and Baricordi, O.R. (2004). HLA-G and IL-10 in serum in relation to HLA-G genotype and polymorphisms. *Immunogenetics* 56, 135-141.
- Hviid, T.V., Rizzo, R., Melchiorri, L., Stignani, M., and Baricordi, O.R. (2006). Polymorphism in the 5' upstream regulatory and 3' untranslated regions of the HLA-G gene in relation to soluble HLA-G and IL-10 expression. *Human immunology* 67, 53-62.
- Hviid, T.V., Sorensen, S., and Morling, N. (1999). Polymorphism in the regulatory region located more than 1.1 kilobases 5' to the start site of transcription, the promoter region, and exon 1 of the HLA-G gene. *Human immunology* 60, 1237-1244.
- Ishitani, A., and Geraghty, D.E. (1992). Alternative splicing of HLA-G transcripts yields proteins with primary structures resembling both class I and class II antigens. *Proc Natl Acad Sci U S A* 89, 3947-3951.
- Kanai, T., Fujii, T., Kozuma, S., Yamashita, T., Miki, A., Kikuchi, A., and Taketani, Y. (2001a). Soluble HLA-G influences the release of cytokines from allogeneic peripheral blood mononuclear cells in culture. *Mol Hum Reprod* 7, 195-200.
- Kanai, T., Fujii, T., Unno, N., Yamashita, T., Hyodo, H., Miki, A., Hamai, Y., Kozuma, S., and Taketani, Y. (2001b). Human leukocyte antigen-G-expressing cells differently modulate the release of cytokines from mononuclear cells present in the decidua versus peripheral blood. *Am J Reprod Immunol* 45, 94-99.

- Kapasi, K., Albert, S.E., Yie, S., Zavazava, N., and Librach, C.L. (2000). HLA-G has a concentration-dependent effect on the generation of an allo-CTL response. *Immunology* 101, 191-200.
- Kuersten, S., and Goodwin, E.B. (2003). The power of the 3' UTR: translational control and development. *Nat Rev Genet* 4, 626-637.
- La Nasa, G., Littera, R., Locatelli, F., Lai, S., Alba, F., Caocci, G., Lisini, D., Nesci, S., Vacca, A., Piras, E., *et al.* (2007). The human leucocyte antigen-G 14-basepair polymorphism correlates with graft-versus-host disease in unrelated bone marrow transplantation for thalassaemia. *Br J Haematol* 139, 284-288.
- Larsen, M.H., and Hviid, T.V. (2009). Human leukocyte antigen-G polymorphism in relation to expression, function, and disease. *Human immunology* 70, 1026-1034.
- Le Friec, G., Gros, F., Sebti, Y., Guilloux, V., Pangault, C., Fauchet, R., and Amiot, L. (2004). Capacity of myeloid and plasmacytoid dendritic cells especially at mature stage to express and secrete HLA-G molecules. *Journal of leukocyte biology* 76, 1125-1133.
- Le Gal, F.A., Riteau, B., Sedlik, C., Khalil-Daher, I., Menier, C., Dausset, J., Guillet, J.G., Carosella, E.D., and Rouas-Freiss, N. (1999). HLA-G-mediated inhibition of antigen-specific cytotoxic T lymphocytes. *Int Immunol* 11, 1351-1356.
- Le Maux, A., Noel, G., Birebent, B., Grosset, J.M., Vu, N., De Guibert, S., Bernard, M., Semana, G., and Amiot, L. (2008). Soluble human leucocyte antigen-G molecules in peripheral blood haematopoietic stem cell transplantation: a specific role to prevent acute graft-versus-host disease and a link with regulatory T cells. *Clin Exp Immunol* 152, 50-56.
- Le Rond, S., Le Maoult, J., Creput, C., Menier, C., Deschamps, M., Le Friec, G., Amiot, L., Durrbach, A., Dausset, J., Carosella, E.D., *et al.* (2004). Alloreactive CD4+ and CD8+ T cells express the immunotolerant HLA-G molecule in mixed lymphocyte reactions: in vivo implications in transplanted patients. *Eur J Immunol* 34, 649-660.
- Lefebvre, S., Berrih-Aknin, S., Adrian, F., Moreau, P., Poeta, S., Gourand, L., Dausset, J., Carosella, E.D., and Paul, P. (2001). A specific interferon (IFN)-stimulated response element of the distal HLA-G promoter binds IFN-regulatory factor 1 and mediates enhancement of this nonclassical class I gene by IFN-beta. *J Biol Chem* 276, 6133-6139.
- LeMaoult, J., Caumartin, J., and Carosella, E.D. (2007a). Exchanges of membrane patches (trogonocytosis) split theoretical and actual functions of immune cells. *Human immunology* 68, 240-243.
- LeMaoult, J., Caumartin, J., Daouya, M., Favier, B., Le Rond, S., Gonzalez, A., and Carosella, E.D. (2007b). Immune regulation by pretenders: cell-to-cell transfers of HLA-G make effector T cells act as regulatory cells. *Blood* 109, 2040-2048.
- LeMaoult, J., Krawice-Radanne, I., Dausset, J., and Carosella, E.D. (2004). HLA-G1-expressing antigen-presenting cells induce immunosuppressive CD4+ T cells. *Proc Natl Acad Sci U S A* 101, 7064-7069.
- LeMaoult, J., Zafaranloo, K., Le Danff, C., and Carosella, E.D. (2005). HLA-G up-regulates ILT2, ILT3, ILT4, and KIR2DL4 in antigen presenting cells, NK cells, and T cells. *FASEB J* 19, 662-664.
- Li, C., Houser, B.L., Nicotra, M.L., and Strominger, J.L. (2009). HLA-G homodimer-induced cytokine secretion through HLA-G receptors on human decidual macrophages and natural killer cells. *Proc Natl Acad Sci U S A* 106, 5767-5772.


- Liang, S., Baibakov, B., and Horuzsko, A. (2002). HLA-G inhibits the functions of murine dendritic cells via the PIR-B immune inhibitory receptor. *Eur J Immunol* 32, 2418-2426.
- Liang, S., Ristich, V., Arase, H., Dausset, J., Carosella, E.D., and Horuzsko, A. (2008). Modulation of dendritic cell differentiation by HLA-G and ILT4 requires the IL-6--STAT3 signaling pathway. *Proc Natl Acad Sci U S A* 105, 8357-8362.
- Lila, N., Rouas-Freiss, N., Dausset, J., Carpentier, A., and Carosella, E.D. (2001). Soluble HLA-G protein secreted by allo-specific CD4+ T cells suppresses the allo-proliferative response: a CD4+ T cell regulatory mechanism. *Proc Natl Acad Sci U S A* 98, 12150-12155.
- Lopez, A.S., Alegre, E., LeMaoult, J., Carosella, E., and Gonzalez, A. (2006). Regulatory role of tryptophan degradation pathway in HLA-G expression by human monocyte-derived dendritic cells. *Mol Immunol* 43, 2151-2160.
- Magnani, C.F., Alberigo, G., Bacchetta, R., Serafini, G., Andreani, M., Roncarolo, M.G., and Gregori, S. (2011). Killing of myeloid APCs via HLA class I, CD2 and CD226 defines a novel mechanism of suppression by human Tr1 cells. *Eur J Immunol* 41, 1652-1662.
- Morandi, F., Ferretti, E., Bocca, P., Prigione, I., Raffaghello, L., and Pistoia, V. (2010). A novel mechanism of soluble HLA-G mediated immune modulation: downregulation of T cell chemokine receptor expression and impairment of chemotaxis. *PLoS One* 5, e11763.
- Moreau, P., Adrian-Cabestre, F., Menier, C., Guiard, V., Gourand, L., Dausset, J., Carosella, E.D., and Paul, P. (1999). IL-10 selectively induces HLA-G expression in human trophoblasts and monocytes. *Int Immunol* 11, 803-811.
- Moreau, P., Faure, O., Lefebvre, S., Ibrahim, E.C., O'Brien, M., Gourand, L., Dausset, J., Carosella, E.D., and Paul, P. (2001). Glucocorticoid hormones upregulate levels of HLA-G transcripts in trophoblasts. *Transplant Proc* 33, 2277-2280.
- Moreau, P., Flajollet, S., and Carosella, E.D. (2009). Non-classical transcriptional regulation of HLA-G: an update. *J Cell Mol Med* 13, 2973-2989.
- Munn, D.H., Sharma, M.D., Lee, J.R., Jhaver, K.G., Johnson, T.S., Keskin, D.B., Marshall, B., Chandler, P., Antonia, S.J., Burgess, R., *et al.* (2002). Potential regulatory function of human dendritic cells expressing indoleamine 2,3-dioxygenase. *Science* 297, 1867-1870.
- Naji, A., Durrbach, A., Carosella, E.D., and Rouas-Freiss, N. (2007a). Soluble HLA-G and HLA-G1 expressing antigen-presenting cells inhibit T-cell alloproliferation through ILT-2/ILT-4/FasL-mediated pathways. *Human immunology* 68, 233-239.
- Naji, A., Le Rond, S., Durrbach, A., Krawice-Radanne, I., Creput, C., Daouya, M., Caumartin, J., LeMaoult, J., Carosella, E.D., and Rouas-Freiss, N. (2007b). CD3+CD4^{low} and CD3+CD8^{low} are induced by HLA-G: novel human peripheral blood suppressor T-cell subsets involved in transplant acceptance. *Blood* 110, 3936-3948.
- Nasef, A., Mathieu, N., Chapel, A., Frick, J., Francois, S., Mazurier, C., Boutarfa, A., Bouchet, S., Gorin, N.C., Thierry, D., *et al.* (2007). Immunosuppressive effects of mesenchymal stem cells: involvement of HLA-G. *Transplantation* 84, 231-237.

- Ober, C., Aldrich, C.L., Chervoneva, I., Billstrand, C., Rahimov, F., Gray, H.L., and Hyslop, T. (2003). Variation in the HLA-G promoter region influences miscarriage rates. *Am J Hum Genet* 72, 1425-1435.
- Ober, C., Billstrand, C., Kuldaneck, S., and Tan, Z. (2006). The miscarriage-associated HLA-G -725G allele influences transcription rates in JEG-3 cells. *Hum Reprod* 21, 1743-1748.
- Onno, M., Le Friec, G., Pangault, C., Amiot, L., Guilloux, V., Drenou, B., Caulet-Maugendre, S., Andre, P., and Fauchet, R. (2000). Modulation of HLA-G antigens expression in myelomonocytic cells. *Human immunology* 61, 1086-1094.
- Pacciani, V., Gregori, S., Chini, L., Corrente, S., Chianca, M., Moschese, V., Rossi, P., Roncarolo, M.G., and Angelini, F. (2010). Induction of anergic allergen-specific suppressor T cells using tolerogenic dendritic cells derived from children with allergies to house dust mites. *J Allergy Clin Immunol* 125, 727-736.
- Park, G.M., Lee, S., Park, B., Kim, E., Shin, J., Cho, K., and Ahn, K. (2004). Soluble HLA-G generated by proteolytic shedding inhibits NK-mediated cell lysis. *Biochem Biophys Res Commun* 313, 606-611.
- Paul, P., Cabestre, F.A., Ibrahim, E.C., Lefebvre, S., Khalil-Daher, I., Vazeux, G., Quiles, R.M., Bermond, F., Dausset, J., and Carosella, E.D. (2000). Identification of HLA-G7 as a new splice variant of the HLA-G mRNA and expression of soluble HLA-G5, -G6, and -G7 transcripts in human transfected cells. *Human immunology* 61, 1138-1149.
- Qiu, J., Terasaki, P.I., Miller, J., Mizutani, K., Cai, J., and Carosella, E.D. (2006). Soluble HLA-G expression and renal graft acceptance. *Am J Transplant* 6, 2152-2156.
- Rajagopalan, S., Bryceson, Y.T., Kuppusamy, S.P., Geraghty, D.E., van der Meer, A., Joosten, I., and Long, E.O. (2006). Activation of NK cells by an endocytosed receptor for soluble HLA-G. *PLoS Biol* 4, e9.
- Rajagopalan, S., Fu, J., and Long, E.O. (2001). Cutting edge: induction of IFN-gamma production but not cytotoxicity by the killer cell Ig-like receptor KIR2DL4 (CD158d) in resting NK cells. *J Immunol* 167, 1877-1881.
- Rajagopalan, S., and Long, E.O. (1999). A human histocompatibility leukocyte antigen (HLA)-G-specific receptor expressed on all natural killer cells. *The Journal of experimental medicine* 189, 1093-1100.
- Ristich, V., Liang, S., Zhang, W., Wu, J., and Horuzsko, A. (2005). Tolerization of dendritic cells by HLA-G. *Eur J Immunol* 35, 1133-1142.
- Ristich, V., Zhang, W., Liang, S., and Horuzsko, A. (2007). Mechanisms of prolongation of allograft survival by HLA-G/ILT4-modified dendritic cells. *Human immunology* 68, 264-271.
- Riteau, B., Menier, C., Khalil-Daher, I., Martinuzzi, S., Pla, M., Dausset, J., Carosella, E.D., and Rouas-Freiss, N. (2001). HLA-G1 co-expression boosts the HLA class I-mediated NK lysis inhibition. *Int Immunol* 13, 193-201.
- Rizzo, R., Campioni, D., Stignani, M., Melchiorri, L., Bagnara, G.P., Bonsi, L., Alviano, F., Lanzoni, G., Moretti, S., Cuneo, A., *et al.* (2008). A functional role for soluble HLA-G antigens in immune modulation mediated by mesenchymal stromal cells. *Cytherapy* 10, 364-375.

- Rizzo, R., Ferrari, D., Melchiorri, L., Stignani, M., Gulinelli, S., Baricordi, O.R., and Di Virgilio, F. (2009). Extracellular ATP acting at the P2X7 receptor inhibits secretion of soluble HLA-G from human monocytes. *J Immunol* 183, 4302-4311.
- Roncarolo, M.G., Gregori, S., Battaglia, M., Bacchetta, R., Fleischhauer, K., and Levings, M.K. (2006). Interleukin-10-secreting type 1 regulatory T cells in rodents and humans. *Immunological reviews* 212, 28-50.
- Rossetti, M., Gregori, S., and Roncarolo, M.G. (2011). Granulocyte-colony stimulating factor drives the in vitro differentiation of human dendritic cells that induce anergy in naive T cells. *Eur J Immunol* 40, 3097-3106.
- Rouas-Freiss, N., Goncalves, R.M., Menier, C., Dausset, J., and Carosella, E.D. (1997). Direct evidence to support the role of HLA-G in protecting the fetus from maternal uterine natural killer cytotoxicity. *Proc Natl Acad Sci U S A* 94, 11520-11525.
- Rousseau, P., Le Discorde, M., Mouillot, G., Marcou, C., Carosella, E.D., and Moreau, P. (2003). The 14 bp deletion-insertion polymorphism in the 3' UT region of the HLA-G gene influences HLA-G mRNA stability. *Human immunology* 64, 1005-1010.
- Sakaguchi, S. (2005). Naturally arising Foxp3-expressing CD25+CD4+ regulatory T cells in immunological tolerance to self and non-self. *Nat Immunol* 6, 345-352.
- Sakaguchi, S., Miyara, M., Costantino, C.M., and Hafler, D.A. (2010). FOXP3+ regulatory T cells in the human immune system. *Nat Rev Immunol* 10, 490-500.
- Selmani, Z., Naji, A., Gaiffe, E., Obert, L., Tiberghien, P., Rouas-Freiss, N., Carosella, E.D., and Deschaseaux, F. (2009). HLA-G is a crucial immunosuppressive molecule secreted by adult human mesenchymal stem cells. *Transplantation* 87, S62-66.
- Selmani, Z., Naji, A., Zidi, I., Favier, B., Gaiffe, E., Obert, L., Borg, C., Saas, P., Tiberghien, P., Rouas-Freiss, N., *et al.* (2008). Human leukocyte antigen-G5 secretion by human mesenchymal stem cells is required to suppress T lymphocyte and natural killer function and to induce CD4+CD25highFOXP3+ regulatory T cells. *Stem Cells* 26, 212-222.
- Shiroishi, M., Kuroki, K., Ose, T., Rasubala, L., Shiratori, I., Arase, H., Tsumoto, K., Kumagai, I., Kohda, D., and Maenaka, K. (2006a). Efficient leukocyte Ig-like receptor signaling and crystal structure of disulfide-linked HLA-G dimer. *J Biol Chem* 281, 10439-10447.
- Shiroishi, M., Kuroki, K., Rasubala, L., Tsumoto, K., Kumagai, I., Kurimoto, E., Kato, K., Kohda, D., and Maenaka, K. (2006b). Structural basis for recognition of the nonclassical MHC molecule HLA-G by the leukocyte Ig-like receptor B2 (LILRB2/LIR2/ILT4/CD85d). *Proc Natl Acad Sci U S A* 103, 16412-16417.
- Shiroishi, M., Tsumoto, K., Amano, K., Shirakihara, Y., Colonna, M., Braud, V.M., Allan, D.S., Makadzange, A., Rowland-Jones, S., Willcox, B., *et al.* (2003). Human inhibitory receptors Ig-like transcript 2 (ILT2) and ILT4 compete with CD8 for MHC class I binding and bind preferentially to HLA-G. *Proc Natl Acad Sci U S A* 100, 8856-8861.
- Steimle, V., Durand, B., Barras, E., Zufferey, M., Hadam, M.R., Mach, B., and Reith, W. (1995). A novel DNA-binding regulatory factor is mutated in primary MHC class II deficiency (bare lymphocyte syndrome). *Genes Dev* 9, 1021-1032.
- Tan, Z., Shon, A.M., and Ober, C. (2005). Evidence of balancing selection at the HLA-G promoter region. *Hum Mol Genet* 14, 3619-3628.

- Uccelli, A., Moretta, L., and Pistoia, V. (2008). Mesenchymal stem cells in health and disease. *Nat Rev Immunol* 8, 726-736.
- Veit, T.D., and Chies, J.A. (2009). Tolerance versus immune response -- microRNAs as important elements in the regulation of the HLA-G gene expression. *Transpl Immunol* 20, 229-231.
- Wiendl, H., Feger, U., Mittelbronn, M., Jack, C., Schreiner, B., Stadelmann, C., Antel, J., Brueck, W., Meyermann, R., Bar-Or, A., *et al.* (2005). Expression of the immunetolerogenic major histocompatibility molecule HLA-G in multiple sclerosis: implications for CNS immunity. *Brain* 128, 2689-2704.
- Yang, Y., Chu, W., Geraghty, D.E., and Hunt, J.S. (1996). Expression of HLA-G in human mononuclear phagocytes and selective induction by IFN-gamma. *J Immunol* 156, 4224-4231.
- Yie, S.M., Li, L.H., Li, G.M., Xiao, R., and Librach, C.L. (2006a). Progesterone enhances HLA-G gene expression in JEG-3 choriocarcinoma cells and human cytotrophoblasts in vitro. *Hum Reprod* 21, 46-51.
- Yie, S.M., Xiao, R., and Librach, C.L. (2006b). Progesterone regulates HLA-G gene expression through a novel progesterone response element. *Hum Reprod* 21, 2538-2544.
- Zhu, X.M., Han, T., Wang, X.H., Li, Y.H., Yang, H.G., Luo, Y.N., Yin, G.W., and Yao, Y.Q. (2010). Overexpression of miR-152 leads to reduced expression of human leukocyte antigen-G and increased natural killer cell mediated cytotoxicity in JEG-3 cells. *Am J Obstet Gynecol* 202, 592 e591-597.
- Zidi, I., Guillard, C., Marcou, C., Krawice-Radanne, I., Sangrouber, D., Rouas-Freiss, N., Carosella, E.D., and Moreau, P. (2006). Increase in HLA-G1 proteolytic shedding by tumor cells: a regulatory pathway controlled by NF-kappaB inducers. *Cell Mol Life Sci* 63, 2669-2681.

IntechOpen



Histocompatibility

Edited by Dr. Bahaa Abdel-Salam

ISBN 978-953-51-0589-3

Hard cover, 188 pages

Publisher InTech

Published online 02, May, 2012

Published in print edition May, 2012

This book presents some recent researches related to histocompatibility for scientists interested in this field. It includes 10 chapters, in different topics, prepared by Sundararajulu Panneerchelvam and Mohd Nor Norazmi; Giada Amodio and Silvia Gregori; Adema Ribic; Bahaa K. A. Abdel-Salam; Kai-Fu Tang; Roberto Biassoni, Irene Vanni and Elisabetta Ugolotti; Wei-Cheng Yang, Lien-Siang Chou and Jer-Ming Hu; Shatrah Othman and Rohana Yusof; Masahiro Hirayama, Eiichi Azuma and Yoshihiro Komada; Gustav Roder, Linda Geironson, Elna Follin, Camilla Thuring and Kajsa Paulsson.

How to reference

In order to correctly reference this scholarly work, feel free to copy and paste the following:

Giada Amodio and Silvia Gregori (2012). Distinctive Immunological Functions of HLA-G, Histocompatibility, Dr. Bahaa Abdel-Salam (Ed.), ISBN: 978-953-51-0589-3, InTech, Available from:
<http://www.intechopen.com/books/histocompatibility/distinctive-immunological-functions-of-hla-g>

INTECH
open science | open minds

InTech Europe

University Campus STeP Ri
Slavka Krautzeka 83/A
51000 Rijeka, Croatia
Phone: +385 (51) 770 447
Fax: +385 (51) 686 166
www.intechopen.com

InTech China

Unit 405, Office Block, Hotel Equatorial Shanghai
No.65, Yan An Road (West), Shanghai, 200040, China
中国上海市延安西路65号上海国际贵都大饭店办公楼405单元
Phone: +86-21-62489820
Fax: +86-21-62489821

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen