

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Speciation in Brazilian Atlantic Forest Mosquitoes: A Mini-Review of the *Anopheles cruzii* Species Complex

Luísa D.P. Rona¹, Carlos J. Carvalho-Pinto² and Alexandre A. Peixoto³

¹Universidade Federal do Rio de Janeiro / Polo de Xerém, Duque de Caxias - RJ,

²Departamento de Microbiologia e Parasitologia, CCB, Universidade Federal de Santa Catarina, Florianópolis - SC,

³Laboratório de Biologia Molecular de Insetos, Instituto Oswaldo Cruz, FIOCRUZ, Rio de Janeiro, Brazil

1. Introduction

Anopheles (Kerteszia) cruzii s.l. (Diptera: Culicidae) has long been known as the primary vector of human and simian malaria parasites in southern and southeastern Brazil (Deane *et al.*, 1970; 1971; Rachou, 1958). Between 1930 and 1960, *An. cruzii* together with *Anopheles (Kerteszia) bellator* and *Anopheles (Kerteszia) homunculus* were considered the main vectors of malaria once endemic in southern Brazil. Vector control has reduced or even interrupted malaria transmission in some areas, but *An. cruzii* is still responsible for several oligosymptomatic malaria cases in southern and southeastern Brazil. This mosquito is also a vector of simian malaria in Rio de Janeiro and São Paulo States (Deane *et al.*, 1970). Studies on seasonal and vertical distribution of *An. cruzii* demonstrated high vertical mobility from ground level to tree tops and this behavior could be responsible for human infection by simian *Plasmodium* species (Deane *et al.*, 1984; Marrelli *et al.*, 2007; Ueno *et al.*, 2007).

The distribution of this mosquito follows the coast of the Brazilian Atlantic forest (Consoli & Lourenço-de-Oliveira, 1994; Zavortink, 1973), which provides an excellent environment for *An. cruzii*, since it is an ecosystem abundant in bromeliads, the larval habitat for this anopheline (Pittendrigh, 1949; Rachou, 1958; Veloso *et al.*, 1956). The adults are found in a variety of habitats, from sea level in coastal areas to the mountains. Females are strongly anthropophilic and blood-feed preferably during the evening (Aragão, 1964; Corrêa *et al.*, 1961; Veloso *et al.*, 1956), perhaps biting more than one host to complete egg maturation, which is epidemiologically relevant for malaria transmission (Bona & Navarro-Silva, 2006; Wilkerson & Peyton, 1991). However, notwithstanding its importance as a malaria vector, there are not many population genetic studies of *An. cruzii* (e.g. Calado *et al.*, 2006; Carvalho-Pinto & Lourenço-de-Oliveira, 2004; Malafrente *et al.*, 2007; Ramirez & Dessen, 2000a,b; see also below).

The possibility that *An. cruzii* could represent more than one species was first suggested by morphological differences observed among populations from the states of Santa Catarina

and Rio de Janeiro (Zavortink, 1973). Later it was revealed that southern and southeastern Brazilian populations of *An. cruzii* are polymorphic for chromosomal inversions (Ramirez *et al.*, 1994; Ramirez & Dessen, 1994). The authors found evidence for the occurrence of genetically distinct *An. cruzii* populations with three different sets of inversions on the X chromosome, defined as forms A, B and C. In populations where two forms are sympatric no heterozygotes were detected, suggesting the absence or limited gene flow between the two groups (Ramirez *et al.*, 1994; Ramirez & Dessen, 1994, 2000a,b).

The possibility that *An. cruzii* may represent a complex of cryptic species was also supported by isoenzymatic profiles from 10 distinct *loci* of several *An. cruzii* populations. This analysis indicated two genetically isolated groups, one from northeastern Brazil (Itaparica Island - Bahia State) and the other from southeastern and southern Brazil (Nova Iguaçu - Rio de Janeiro State, Cananéia - São Paulo State and Florianópolis - Santa Catarina State) (Carvalho-Pinto & Lourenço-de-Oliveira, 2004).

These papers, which proposed that *An. cruzii* is a species complex, led to further studies using molecular markers to investigate the genetic differentiation among populations of this malaria vector. For example, Malafronte *et al* (2007) found some differences between ITS2 sequences comparing a number of southern and southeastern *An. cruzii* populations from Brazil. Similar results were observed by Calado *et al* (2006), using PCR-RAPD and PCR-RFLP of the ITS2 region.

We used a number of single-copy genes to investigate the molecular differentiation and gene flow among the putative sibling species of this complex (Rona *et al.*, 2009, 2010a,b). The results and the main conclusions of these analyses are discussed in more detail below.

2. Molecular markers and the genetic differentiation among Brazilian populations of *An. cruzii* s.l.

The *timeless* gene is a *locus* involved in the control of circadian activity rhythms in *Drosophila* (reviewed in Hardin 2005). It also controls mating rhythms (Sakai & Ishida, 2001) and its orthologues in mosquitoes are potentially involved in maintaining temporal reproductive isolation between closely related species. Rona *et al* (2009) isolated a fragment of the *timeless* gene in *An. cruzii* and used it to assess the genetic differentiation among six populations of this malaria vector within its geographic distribution range in Brazil: Florianópolis - Santa Catarina State, Cananéia and Juquitiba - São Paulo State, Itatiaia - Rio de Janeiro State, Santa Teresa - Espírito Santo State and Itaparica Island - Bahia State (Figure 1).

Very strong evidence was obtained for the existence of a different species in Itaparica, a finding that supports the isoenzyme study mentioned above (Carvalho-Pinto & Lourenço-de-Oliveira, 2004). Extremely high F_{ST} values and an elevated number of fixed differences (Table 1) were observed between this northeastern population and the other five studied localities. In addition, the data also suggest that some populations from southern and southeastern regions might also constitute different incipient species. Moderately high F_{ST} values were found when comparing Itatiaia with Florianópolis, Cananéia, Juquitiba and Santa Teresa, suggesting perhaps that this population is in a process of differentiation and incipient speciation (Table 1).




Fig. 1. Localities of the six Brazilian *An. cruzii* populations studied in Rona *et al.* (2009). Values in table are approximated distances between localities in km. (Source: IBGE and Google Maps). All mosquitoes used in this study were females captured at the following localities along the Brazilian Atlantic forest: Florianópolis, Santa Catarina State (SC) (27°31'S / 48°30'W), Cananéia and Jujutiba, São Paulo State (SP) (25°01'S / 47°55'W and 23°57'S / 47°03'W), Itatiaia, Rio de Janeiro State (RJ) (22°27'S / 44°36'W), Santa Teresa, Espírito Santo State (ES) (19°56'S / 40°35'W) and Itaparica Island (Jaguaripe), Bahia State (BA) (13°05'S / 38°48'W) (Modified from Rona *et al.*, 2009).

	Florianópolis	Cananéia	Juquitiba	Itatiaia	Santa Teresa	Itaparica (Bahia)
Florianópolis	-	0.055	0.087	0.145	0.158	0.835
Cananéia	00	-	0.108	0.225	0.215	0.851
Juquitiba	00	00	-	0.203	0.069	0.840
Itatiaia	00	00	00	-	0.184	0.876
Santa Teresa	00	00	00	00	-	0.862
Itaparica (Bahia)	27	29	30	30	32	-

Table 1. Genetic differentiation between *An. cruzii* populations using the *timeless* gene. The pair-wise estimates of population differentiation (F_{ST}) are shown in the upper right matrix and the numbers of fixed differences between each pair of populations are shown in the lower left matrix of the table. In all cases the F_{ST} values were significant (significance evaluated by 1000 random permutations). The sequences were aligned using ClustalX software (Thompson *et al.*, 1997) and population genetics analysis was carried out using DNASP4.0 (Rozas *et al.*, 2003) and PROSEQ V 2.91 (Filatov & Charlesworth, 1999) (Modified from Rona *et al.*, 2009).

These results were supported by a Neighbor-joining tree (Figure 2). The *An. cruzii* sequences from Itaparica (Bahia) were clearly separated in an isolated branch indicating that this northeastern population has diverged significantly from the other populations, in agreement with the isoenzyme analysis (Carvalho-Pinto & Lourenço-de-Oliveira, 2004). In addition, although no clear separation between the *timeless* sequences from Florianópolis, Cananéia, Juquitiba and Santa Teresa was observed, the sequences from Itatiaia do not appear at a random, showing some clustering. Therefore, a process of incipient speciation seems to be occurring between Itatiaia and the other studied southern and southeastern populations.

To investigate in more detail the genetic differentiation between the southern/southeastern and northeastern siblings of *An. cruzii*, a *multilocus* analysis was carried out comparing Itaparica to Florianópolis (Rona *et al.*, 2010a). The aim of this study was to determine if there is still gene flow between the two sibling species and to estimate their divergence time. This analysis was implemented using six *loci*, three circadian clock genes (*timeless*, *Clock* and *cycle*) and three encoding ribosomal proteins (*Rp49*, *RpS29* and *RpS2*). As mentioned above, circadian clock genes (Hardin, 2005), such as *timeless*, *Clock* and *cycle*, are putatively involved in the control of mating rhythms and therefore are potentially important in maintaining temporal reproductive isolation between closely related species (Sakai & Ishida, 2001, Tauber *et al.*, 2003). The analysis revealed very high F_{ST} values (ranging from 0.58 to 0.89) and fixed differences between these two cryptic species in all six *loci*, irrespective of their

function. The divergence time and the migration rate parameters were estimated for all combined *loci*. Figure 3 shows the posterior probability distributions for each of the three parameters estimated using the IM program. The results suggested that the two species have not exchanged migrants since their separation and that they possibly diverged between 1.1 and 3.6 million years ago (Rona *et al.*, 2010a). In fact, the divergence time between the southern and northeastern species fall within the Pleistocene, a period of intense climatic changes (Cantolla, 2003; Ravelo *et al.*, 2004).




Fig. 2. Neighbor-joining tree using *timeless* nucleotide sequences of the *Anopheles cruzii* populations carried out using MEGA 4.0 (Tamura *et al.*, 2007) with Kimura 2-parameters distance. Numbers on the nodes represent the percentage bootstrap values based on 1000 replications. Flo: Florianópolis population; Can: Cananéia; Juq: Juquitiba; Ita: Itatiaia; San: Santa Teresa; Bahia: Itaparica Island population. (Source: Rona *et al.*, 2009).






Fig. 3. Posterior probability distributions for each of the three demographic parameters estimated using IM: divergence time between Florianópolis and Itaparica, and migration rates in both directions. The estimated mutation rate, based on *Drosophila*, was used to convert the divergence time parameter t to the number of years since population splitting. Four IM simulations using different seed numbers were plotted for each parameter estimate. All curves are shown including the range of the priors. The IM program is an implementation of the Isolation with Migration model and is based on the MCMC (Markov Chain Monte Carlo) simulations of genealogies (Hey & Nielsen, 2004). Initial IM runs were performed in order to establish appropriate upper limits for the priors of each demographic parameter mentioned above. These preliminary simulations generated marginal

distributions that facilitated the choice of parameter values used in the final IM analyses. The convergence was assessed through multiple long runs (four independent MCMC runs with different seed numbers were carried out with at least 30,000,000 recorded steps after a burn-in of 100,000 steps) and by monitoring the ESS values, the update acceptance rates and the trend lines. The Infinite Sites model (Kimura, 1969) was chosen as the mutation model in the IM simulations because the two species are closely related and all genes are nuclear. The optimal recombination-filtered block was extracted from each gene alignment using the IM_{GC} program, which also removes haplotypes that represent likely recombinant sequences (Woerner *et al.*, 2007). See Rona *et al.* (2010a) for more details. (Modified from Rona *et al.*, 2010a).




Fig. 4. Haplotype network of *cpr* sequences. Each color represents one population of *An. cruzii*. Each circle represents a different haplotype with size proportional to its relative frequency. The number of sequences of each haplotype is given in brackets. The small white circles represent missing intermediates and the lines connecting the haplotypes represent one mutational step between two observed haplotypes. Each individual of Itatiaia population is discriminated next to the respective haplotype. The haplotype network was estimated using TCS1.21 (Clement *et al.*, 2000). (Modified from Rona *et al.*, 2010b).

As mentioned above, the analysis of the molecular polymorphism and genetic differentiation of the *timeless* gene among *An. cruzii* populations from southern and southeastern Brazil suggested that the population from Itatiaia (Rio de Janeiro State) is in a process of differentiation and incipient speciation (Rona *et al.*, 2009). To analyze the divergence between these populations, a fragment of the *cpr* gene, a *locus* involved in metabolic insecticide resistance and odorant clearance in insects, was used. High F_{ST} values, some fixed differences and few shared polymorphisms were found between Itatiaia and the other populations (Florianópolis, Cananéia, Juquitiba and Santa Teresa). Moreover, an haplotype network constructed using the *cpr* sequences shows that Itatiaia is clearly separated in an isolated group (Figure 4) suggesting that this population represents a different species in the *An. cruzii* complex (Rona *et al.*, 2010b).

In addition, a more detailed analysis of the Itatiaia *cpr* sequences revealed that this sample might enclose two different sets of individuals. Based on the number of uninterrupted AG repeats found in the intron included in the studied fragment, the Itatiaia population can be divided in two groups: one called Itatiaia A (04 to 06 AG repeats) and the second, called Itatiaia B (03 AG repeats) (Figure 5). In fact, the separation between the two groups is also

Ita02a	AGAGAGAGAG
Ita02b	AGAGAGAGAGAG
Ita03a	AGAGAGAGAGAG
Ita03b	AGAGAGAGAG
Ita04a	AGAGAGAG
Ita04b	AGAGAGAGAG
Ita05a/b	AGAGAG
Ita06a/b	AGAGAG
Ita07a/b	AGAGAG
Ita08a	AGAGAGAGAG
Ita08b	AGAGAGAGAGAG
Ita09a/b	AGAGAG
Ita10a/b	AGAGAGAGAG
Ita11a	AGAGAGAGAG
Ita11b	AGAGAGAGAGAG
Ita12a	AGAGAG
Ita12b	AGAGAGAGAGAG

Fig. 5. Schematic representation of the AG repeat variable region in the DNA sequences of the *cpr* gene fragment from the Itatiaia population. The sequences of homozygote individuals were grouped and are represented as a/b. The haplotypes with exactly three AG repeats are in red. According to this classification the individuals Ita2, Ita3, Ita4, Ita8, Ita10 and Ita11 belong to Itatiaia A (genotype “4-6/4-6”), the mosquitoes Ita5, Ita6, Ita7 and Ita9 belong to Itatiaia B (genotype “3/3”) and individual Ita12 is the only “hybrid” between the two groups (genotype “3/4-6”). Inspection of the data shows that the Itatiaia sample is not in Hardy-Weinberg equilibrium suggesting the possibility that two sympatric sibling species might exist in this locality. (Modified from Rona *et al.*, 2010b).

evident in the haplotype network of *cpr* sequences shown in Figure 4. Besides, the F_{ST} value (considering gaps as single mutations) between Itatiaia A and B is quite large (0.67) and highly significant ($P < 0.001$) despite the small sample sizes. To confirm, with another *locus*, the hypothesis that the Itatiaia population might include two incipient sympatric sibling species, the *timeless* data (Rona *et al.*, 2009) from the same sample were reanalyzed. As for the *cpr* data, the *timeless* sequences were divided into Itatiaia A and Itatiaia B. The *timeless* gene also suggests that the sequences might belong to two different sibling species with a highly significant F_{ST} value (0.34; $P < 0.001$) (Rona *et al.*, 2010b).

Further work is clearly needed in this locality and an analysis of a number of other molecular markers might allow a more precise estimate of the differentiation and gene flow between the two putative Itatiaia siblings and between this and other localities in southern Brazil. It will be also important to extend our analyses to a number of other populations along the distribution area of *An. cruzii* as this might provide a more complete representation of the evolutionary history of this species complex. These studies are currently under way.

3. Conclusion

In this chapter we reviewed some of our results on *An. cruzii* with emphasis on how the molecular data is providing insights on the evolution of this complex of cryptic species, an example of speciation in Brazilian Atlantic Forest Mosquitoes. Our results and previously published data from other groups suggest that this complex is formed by a number of siblings or incipient species with different levels of genetic divergence and gene flow.

Population genetic studies using molecular markers often revealed complexes of cryptic sibling species in *Anopheles* mosquitoes with wide geographical distributions (Krzywinski & Besansky, 2003). This is the case of the *An. cruzii* complex, an excellent model for studying ecological vicariance and endemic regions in the Brazilian Atlantic Forest due to its broad geographic range, from southern to northeastern Brazil, and its dependence on forested areas as larval habitat. The appearance of ecological barriers caused by climatic changes as in glaciation periods is a possible explanation for the genetic structure found in this species complex. *An. cruzii* is a forest obligate mosquito and these cooling periods are known to cause forest fragmentation (Cantolla, 2003; Ravelo *et al.*, 2004), which probably affected the distribution of intraspecific lineages and might have split a single ancestral species into isolated groups.

The genetic pattern exhibited by the *An. cruzii* complex is compatible with a historical scenario of populations isolated during the Pleistocene ecological changes (Carnaval *et al.*, 2009). The subdivision of the Brazilian Atlantic Forest has been recognized as a cause of endemism, for example, in bats (Martins *et al.*, 2009) and pit vipers (Grazziotin *et al.*, 2006) and climatic changes have been proposed to explain the differentiation among many forest-obligate species (Carnaval *et al.*, 2009; Marroig *et al.*, 2004; Pedro *et al.*, 2008).

Understanding the forces that shaped the Brazilian Atlantic Forest diversity is essential to explain the biodiversity of this important and endangered ecosystem and might help the conservation programs selecting the endemic areas that should be considered conservation priorities.

4. Acknowledgment

The authors would like to thank Dr Rosely Malafronte (Instituto de Medicina Tropical de São Paulo), Dr Monique Motta (FIOCRUZ - Rio de Janeiro) and Claudiney dos Santos (Unidade de Medicina Tropical - Universidade Federal do Espírito Santo) for providing most of the *An. cruzii* samples used in our work. The authors are also indebted to Dr André Pitaluga for helping prepare Figure 1. Our work is supported by grants from the Howard Hughes Medical Institute, FIOCRUZ, Faperj and CNPq.

5. References

- Aragão, M.B. (1964). Geographic distribution and abundance of *Anopheles* species (*Kerteszia*) (Diptera, Culicidae). *Rev Bras Malariol Doencas Trop.* Vol.16, pp. 73-109.
- Bona, A.C. & Navarro-Silva, M.A. (2006). *Anopheles cruzii* parity in dense rain forest in Southern Brazil. *Rev Saude Publica.* Vol.40, pp. 1118-1123.
- Calado, D.C.; Navarro-Silva, M.A. & Sallum, M.A.M. (2006). PCR-RAPD and PCR-RFLP polymorphism detected in *Anopheles cruzii* (Diptera, Culicidae). *Rev Bras Entomol.* Vol.50, pp. 423-430.
- Cantolla, A.U. (2003). *Earth's Climate History*. Servicio Central de Publicaciones del Gobierno Vasco; [<http://homepage.mac.com/uriarte/historia.html>]
- Carnaval, A.C.; Hickerson, M.J.; Haddad, C.F.; Rodrigues, M.T. & Moritz, C. (2009). Stability predicts genetic diversity in the Brazilian Atlantic forest hotspot. *Science.* Vol.323, pp. 785-789.
- Carvalho-Pinto, C.J. & Lourenço-de-Oliveira, R. (2004). Isoenzymatic analysis of four *Anopheles (Kerteszia) cruzii* (Diptera: Culicidae) populations of Brazil. *Mem Inst Oswaldo Cruz.* Vol.99, pp. 471-475.
- Clement, M.; Posada, D. & Crandall, K.A. (2000). TCS: a computer program to estimate gene genealogies. *Mol Ecol.* Vol.9, pp. 1657-1659.
- Consoli, R.A.G.B. & Lourenço-de-Oliveira, R. (1994). *Principais mosquitos de importância sanitária no Brasil*. Rio de Janeiro: Ed. Fiocruz.
- Corrêa, R.R.; Forattini, O.P.; Guarita, O.F. & Rabello, E.X. (1961). Observations on the flight of *Anopheles (Kerteszia) cruzii* and of *A. (K.) bellator*, vectors of malaria (Diptera, Culicidae). *Arq Hig Saude Publica.* Vol.26, pp. 333-342.
- Deane, L.M.; Ferreira-Neto, J.A.; Deane, S.P. & Silveira, I.P. (1970). *Anopheles (Kerteszia) cruzii*, a natural vector of the monkey malaria parasites, *Plasmodium simium* and *Plasmodium brasilianum*. *Trans R Soc Trop Med Hyg.* Vol.64, pp. 647.
- Deane, L.M.; Deane, M.P.; Ferreira-Neto, J.A. & Almeida, F.B. (1971). On the transmission of simian malaria in Brazil. *Rev Inst Med Trop.* Vol.13, pp. 311-319.
- Deane, L.M.; Ferreira-Neto, J.A. & Lima, M.M. (1984). The vertical dispersion of *Anopheles (Kerteszia) cruzii* in a forest in southern Brazil suggests that human cases of simian origin be expect. *Mem Inst Oswaldo Cruz.* Vol.79, pp. 461-463.
- Filatov, D.A. & Charlesworth, D. (1999) DNA polymorphism, haplotype structure and balancing selection in the *Leavenworthia* PgiC locus. *Genetics.* Vol.153, pp. 1423-1434.
- Grazziotin, F.G.; Monzel, M.; Echeverrigaray, S. & Bonatto, S. (2006). Phylogeography of the *Bothrops jararaca* complex (Serpentes: Viperidae): past fragmentation and island colonization in the Brazilian Atlantic Forest. *Mol Ecol.* Vol.15, pp. 3969-82.

- Hardin, P.E. (2005). The Circadian Timekeeping System of *Drosophila*. *Curr Biol.* Vol.15, pp. 714-722.
- Hey, J. & Nielsen, R. (2004). Multilocus methods for estimating population sizes, migration rates and divergence time, with applications to the divergence of *Drosophila pseudoobscura* and *D. persimilis*. *Genetics*. Vol.167, pp. 747-760.
- Kimura, M. (1969). The number of heterozygous nucleotide sites maintained in a finite population due to steady flux of mutations. *Genetics*. Vol.61, pp. 893-903.
- Krzywinski, J. & Besansky, N.J. (2003). Molecular systematics of *Anopheles*: from subgenera to subpopulations. *Annu. Rev. Entomol.* Vol.48, pp. 111-39.
- Malafrente, R.S.; Marrelli, M.T.; Ramirez, C.C.; Nassar, M.N. & Marinotti, O. (2007). Intraspecific variation of second internal transcribed spacer of nuclear ribosomal DNA among populations of *Anopheles (Kerteszia) cruzii* (Diptera: Culicidae). *J Med Entomol.* Vol.44, pp. 538-542.
- Marrelli, M.T.; Malafrente, R.S.; Sallum, M.A. & Natal, D. (2007). *Kerteszia* subgenus of *Anopheles* associated with the Brazilian Atlantic rainforest: current knowledge and future challenges. *Malar J.* Vol.6, pp. 127-134.
- Marroig, G.; Cropp, S.; Cheverud, J.M. (2004). Systematics and evolution of the *Jacchus* group of marmosets (*Platyrrhini*). *Am J Phys Anthropol.* Vol.123, pp. 11-22.
- Martins, F.M.; Templeton, A.R.; Pavan, A.C.O.; Kohlbach, B.C. & Morgante, J.S. (2009). Phylogeography of the common vampire bat (*Desmodus rotundus*): Marked population structure, Neotropical Pleistocene vicariance and incongruence between nuclear and mtDNA markers. *BMC Evol Biol.* Vol.9, pp. 294.
- Nei, M. & Kumar, S. (2000). *Molecular Evolution and Phylogenetics*. New York: Oxford University Press.
- Pedro, P.M.; Sallum, M.A. & Butlin, R.K. (2008). Forest-obligate *Sabethes* mosquitoes suggest palaeoecological perturbations. *Heredity*. Vol.101, pp. 186-195.
- Pittendrigh, C.S. (1949). The ectopic specialization of *Anopheles homunculus*, and its relation to competition with *An. bellator*. *Evolution*. Vol.4, pp. 64-78.
- Rachou, R.G. (1958). Anofelinos do Brasil: Comportamento das espécies vetoras de malária. *Rev Bras Malariol Doencas Trop.* Vol.10, pp. 145-181.
- Ramirez, C.C. & Dessem, E.M. (1994). Cytogenetics analysis of a natural population of *Anopheles cruzii*. *Rev Bras Genet.* Vol.17, pp. 41-46.
- Ramirez, C.C.; Dessem, E.M. & Otto, P.A. (1994). Inversion polymorphism in a natural population of *Anopheles cruzii*. *Caryologia*. Vol.47, pp. 121-130.
- Ramirez, C.C. & Dessen, E.M. (2000a). Chromosomal evidence for sibling species of the malaria vector *Anopheles cruzii*. *Genome*. Vol.43, pp. 143-151.
- Ramirez, C.C. & Dessen, E.M. (2000b). Chromosome differentiated populations of *Anopheles cruzii*: evidence for a third sibling species. *Genetica*. Vol.108, pp. 73-80.
- Ravelo, A.C.; Andreasen, D.H.; Lyle, M.; Olivarez, A. & Wara, M.W. (2004). Regional climate shifts caused by gradual global cooling in the Pliocene epoch. *Nature*. Vol.429, pp. 263-267.
- Rona, L.D.; Carvalho-Pinto, C.J.; Gentile, C.; Grisard, E.C. & Peixoto, A.A. (2009). Assessing the molecular divergence between *Anopheles (Kerteszia) cruzii* populations from Brazil using the *timeless* gene: Further evidence of a species complex. *Malar J.* Vol.8, pp. 60.

- Rona, L.D.; Carvalho-Pinto, C.J.; Mazzoni, C.J. & Peixoto, A.A. (2010a). Estimation of divergence time between two sibling species of the *Anopheles (Kerteszia) cruzii* complex using a multilocus approach. *BMC Evolutionary Biology*. Vol.10, pp. 91.
- Rona, L.D.; Carvalho-Pinto, C.J.; Peixoto, A.A. (2010b). Molecular evidence for the occurrence of a new sibling species within the *Anopheles (Kerteszia) cruzii* complex in south-east Brazil. *Malaria Journal*. Vol.9, pp. 33.
- Rozas, J.; Sánchez-DelBarrio, J.C.; Messeguer, X. & Rozas, R. (2003). DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics*. Vol.19, pp. 2496-2497.
- Sakai, T. & Ishida, N. (2001). Circadian rhythms of female mating activity governed by clock genes in *Drosophila*. *Proc Natl Acad Sci*. Vol.98, pp. 9221-9225.
- Tamura, K.; Dudley, J.; Nei, M. & Kumar, S. (2007). MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol*. Vol.24, pp. 1596-1599.
- Tauber, E.; Roe, H.; Costa, R.; Hennessy, J.M. & Kyriacou, C.P. (2003). Temporal mating isolation driven by a behavioral gene in *Drosophila*. *Curr Biol*. Vol.13, pp. 140-145.
- Thompson, J.D.; Gibson, T.J.; Plewniak, F.; Jeanmougin, F. & Higgins, D.G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res*. Vol.25, pp. 4876-4882.
- Ueno, H.M.; Forattini, O.P. & Kakitani, I. (2007). Vertical and seasonal distribution of *Anopheles (Kerteszia)* in Ilha Comprida, Southeastern Brazil. *Rev Saude Publica*. Vol.41, pp. 269-275.
- Veloso, H.P.; De Moura, J.V. & Klein, R.M. (1956). Ecological limitation of *Anopheles* of the Subgenus *Kerteszia* in the coastal region of Southern Brazil. *Mem Inst Oswaldo Cruz*. Vol.54, pp. 517-548.
- Wilkerson, R.C. & Peyton, E.L. (1991). The Brazilian malaria vector *Anopheles (Kerteszia) cruzii*: Life stages and biology (Diptera: Culicidae). *Mosq System*. Vol.23, pp. 110-122.
- Woerner, A.E.; Cox, M.P. & Hammer, M.F. (2007). Recombination-filtered genomic datasets by information maximization. *Bioinformatics*. Vol.23, pp. 1851-1853.
- Zavortink, T.J. (1973). A review of the subgenus *Kerteszia* of *Anopheles*. *Cont Am Entomol Inst*. Vol.9, pp. 1-54.

© 2012 The Author(s). Licensee IntechOpen. This is an open access article distributed under the terms of the [Creative Commons Attribution 3.0 License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

IntechOpen

IntechOpen