

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Clinical and Genetic Heterogeneity of Autism

---

Yu Wang and Nanbert Zhong

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/48700>

---

## 1. Introduction

Autism (MIM 209850) comprises a heterogeneous group of disorders with a complex genetic etiology, characterized by impairments in reciprocal social communication and presence of restricted, repetitive and stereotyped patterns of behavior [1]. With an early onset prior to age 3 and prevalence as high as 0.9–2.6% [2,3], autism occurs predominantly in males, with a ratio of male: female of 4 to 1. It is one of the leading causes of childhood disability and inflicts serious suffering and burden for the family and society [4].

Diagnosis of autism is based on expert observation and assessment of behavior and cognition, not etiology or pathogenic mechanism. This is further emphasized by the current trend in the DSM-V, in which the category of Asperger syndrome is removed and the diagnostic criteria for autism are modified under the new heading of autism spectrum disorder (ASD). The change in diagnostic criteria is not based on known similarities or differences in causation between these clinically defined categories, but rather on the consensus of opinions of expert clinicians. For autism, several diagnostic instruments are available. Two are commonly used in autism research: the Autism Diagnostic Interview-Revised (ADI-R) that is a semi-structured parent interview [5], and the Autism Diagnostic Observation Schedule (ADOS) uses observation and interaction with the child(ren) [6]. The Childhood Autism Rating Scale (CARS) is used widely in clinical environments to assess severity of autism based on observation of children [7]. The M-CHAT was developed in the late 1990s as a first-stage screening tool for ASD in toddlers' age 18 to 24 months, with a sensitivity of 0.87 and a specificity of 0.99 in American children [8, 9].

## 2. Clinical heterogeneity of ASD

Autistic conditions are a spectrum of disorders, rather than a distinct clinical disorder, which means that the symptoms can be present in a variety of combinations with a range of severity. The disease has variable cognitive manifestations, ranging from a non-verbal child with mental retardation to a high-functioning college student with above average IQ with

inadequate social skills [10]. Clinical heterogeneity of autism showed three major categories: idiopathic autism, autistic spectrum disorder (ASD), and syndromic autistics that usually resulted from an identified syndrome with known genetic etiology. Traditionally, ASD includes autism, Asperger syndrome, where language appears normal, Rett syndrome and pervasive developmental disorder not otherwise specified (PDD-NOS), in which children meet some but not all criteria for autism. Rett syndrome (RTT), occurring almost exclusively in females, is characterized by developmental arrest between 5 and 18 months of age, followed by regression of acquired skills, loss of speech, stereotypic movements (classically of the hands), microcephaly, seizures, and intellectual difficulties. These disorders share deficits in social communication and show variability in language and repetitive behavior domains [1]. Autistic individuals may have symptoms that are independent of the diagnosis. Mental retardation is present in approximately 75% of cases of autism, seizures in 15 to 30% of cases, attention deficit hyperactivity disorder (ADHD) in 59-75% of cases, schizophrenia (SZ) in 5% of cases, obsessive-compulsive disorder (OCD) in about 60% of cases and electroencephalographic abnormalities in 20 to 50% of cases [11]. In addition, approximately 15 to 37% of cases of autism have a comorbid medical condition such as epilepsy, sensory abnormalities, motor abnormalities, sleep disturbances, and gastrointestinal symptoms. Five to 14% of cases had a known genetic disorder or chromosomal anomaly. The 4 most common conditions associated with autistic phenotypes are fragile X syndrome, tuberous sclerosis, 15q duplications, and untreated phenylketonuria. Other conditions associated with autistic phenotypes include Angelman syndrome, Cowden disease, Smith-Lemli-Opitz syndrome, cortical dysplasia-focal epilepsy (CDFE) syndrome, Neurofibromatosis, and X-linked mental retardation.

### **3. Autism is a complex genetic disorder**

It is widely held that autism is largely genetic in origin; several dozen autism susceptibility genes have been identified in the past decade, collectively accounting for about 20% of autistic cases. There is strong evidence from twin and family studies for the importance of complex genetic factors in the development of autism [12, 13]. Family studies have shown that a recurrence rate of autism in siblings of affected proband is as high as 8–10% [12, 14]. Thus, the recurrence risk in siblings is roughly 100 times higher than that found in the general population. The substantial degree of familial clustering in ASD could reflect shared environmental factors, but twin studies strongly point to genetics. Several epidemiological studies among sex-matched twins have clearly demonstrated significant differences of concordance rates in the monozygotic (MZ) and dizygotic (DZ) twins. The largest of these studies [15] found that 60% of the MZ pairs were concordant for autism compared with none of the DZ pairs, suggesting a heritability estimate of >90% assuming a multifactorial threshold model. This is what is observed in every twin study in autism, and is overall consistent with heritability estimates of about 70–80% [15, 16]. One exception is a very recent study with a large sample of twins, which, despite showing a concordance of about 0.6 for MZ twins and 0.25 for DZ twins, comes to the conclusion that shared environment plays a larger role than genetic factors [17]. However, the question of how a shared environment

would have a more major role than genetics is not clear. Moreover, studies in families show that first-degree relatives of an autistic proband have a markedly increased risk for autism relative to the population, consistent with a strong familial or genetic effect observed in twins [18]. This is not to dispute the role of the environment but to emphasize that genes play an important role. Similar to other common diseases with genetic contributions, autism was thought to fit a model in which multiple variants, each with small to moderate effect sizes, interact with each other and perhaps in some cases, environmental factors, to lead to autism; a situation referred to as complex genetics [13].

#### 4. Genetic heterogeneity of autism

Although autism is highly heritable, the identification of candidate genes has been hindered by the heterogeneity of the disease. Autism genetics is highly complex, involving many genes/loci and different genetic variations, including translocation, deletion, single nucleotide polymorphism (SNP) and copy number variation (CNV) [13, 19, 20]. The most obvious general conclusion from all of the published genetic studies is the extraordinary etiological heterogeneity of autism. No specific gene accounts for the majority of autism; rather, even the most common genetic forms account for not more than 1–2% of cases [21]. Further, these genes, including those mentioned earlier, represent a diversity of molecular mechanisms that include cell adhesion, neurotransmission, synaptic structure, RNA processing/splicing, and activity-dependent protein translation. Genetic heterogeneity of autistic cases has been documented by identification of single gene mutations and genomic variations including CNV. The mutant genes identified from autistic patients are: *FMR1*, *MECP2*, *CNTNAP2*, *PTEN*, *DHCR7*, *CACNA1C*, *UBE3A*, *TSC2*, *NF1*, *ARX*, *NLGN3*, *NLGN4*, *NRXN1*, *FOXP1*, *FOXP2*, *GRIK2*, and *SHANK3* (Table 1). Genomic variation including copy number deletion or duplication at loci of 1q21.2, 1q42.2, 2q31.1, 3p25.3, 7q11.23, 7q22.1, 7q36.3, 11q13.3, 12q14.2, 15q11-13, 16p11.2, 16q13.3, 17q11.2, 17q12, 17q21.32, 22q13.33, or Xp22.11 may also associate with autism.

#### 5. Genotype/phenotype correlation in ASD

The presence of genetic and phenotypic heterogeneity in autism with a number of underlying pathogenic mechanisms is highlighted in this current review. There are at least three phenotypic presentations with distinct genetic underpinnings: (1) autism with syndromic phenotype characterized by rare, single-gene defects (Table 2); (2) broad autistic phenotypes caused by genetic variations in single or multiple genes, each of these variations being common and distributed continually in the general population but resulting in variant clinical phenotypes when it reaches a certain threshold through complex gene-gene and gene-environment interactions; and (3) severe and specific phenotype caused by 'de-novo' mutations in the patient or transmitted through asymptomatic carriers of such mutations (Table 3) [48, 49]. Understanding the neurobiological processes by which genotypes lead to phenotypes, along with the advances in developmental neuroscience and neuronal networks at the cellular and molecular level, are paving the way for translational research

involving targeted interventions of affected molecular pathways and early intervention programs that promote normal brain responses to stimuli and alter the developmental trajectory [50]. Recent genetic results have improved our knowledge of the genetic basis of autism. Nevertheless, identification of phenotypic markers remains challenging due to phenotypic and genotypic heterogeneity.

Gene	Genetic alteration	Location	Reference
<i>FMR1</i>	The number of CGG in <i>FMR1</i> alleles is classified as intermediate mutation (45 to 55), premutation (55 to 200), or full mutation (>200)	5'untranslated region	22-24
<i>MECP2</i>	T158M, T158A	Missense mutation	25
<i>CNTNAP2</i>	3709delG	Exon 22	26
	G731S, I869T R1119H, D1129H, I1253T, T1278I	Exon 14, 17 Exon 20, 21, 23, 24	27
	H275A	Exon 6	28
	CNV (microdeletion)	Promoter	29
<i>PTEN</i>	Deletion	Exon 2	30
<i>CACNA1C</i>	G406R	Missense mutation	31
<i>UBE3A</i>	D15S122	5' end of <i>UBE3A</i>	32, 33
<i>TSC2</i>	SNP	Intron 4, 9; exon 40	34
<i>NF1</i>	SNP	Intron 27	35
<i>NLGN3</i>	R451C	Missense mutation	36, 37
<i>NLGN4</i>	1186insT	Frameshift mutation	37
<i>NRXN1</i>	<i>De novo</i> 320-kb deletion	Promoter and initial coding exons	38, 39
	Missense structural variant	Neurexin1 $\beta$ signal peptide region	40
<i>FOXP1</i>	<i>De novo</i> intragenic deletion	Exons 4-14	41
<i>FOXP2</i>	Del CAA;	Exon 5	42, 43
	Frequency of the TT allele	Intron 15	
<i>GRIK2</i>	SNP	M867I	44
<i>SHANK3</i>	<i>De novo</i> Q321R	Stop codon	45
	1-bp insertion	Exon 11	46
	<i>De novo</i> 7.9-Mb deletion	22q13.2-qter	47

**Table 1.** Genetic alteration identified from autism

Gene/loci	Chromosome	Phenotype (human/mouse)	Mechanism involved	Risk of autism	Reference
<i>CNTNAP2</i>	7q35-q36.1	Recessive EPI syndrome, ASD, ADHD, TS, OCD	Chromosomal rearrangements and large deletions, disruption of the transcription factor <i>FOXP2</i> , SNP	Not conclusive	51-54
<i>CHD7</i>	8q12.1	CHARGE	Mutations/deletions of gene <i>CHD7</i> , Chromatin remodeling; disruption of the transcription factor <i>FOXP2</i> ; SNP;	15–50%	55, 56
<i>TSC1</i>	9q34.13	Tuberous Sclerosis type I.	Mutation in gene <i>TSC1</i> and subsequent hyperactivation of the downstream mTOR pathway, resulting in increased cell growth and proliferation.	Not conclusive	57
<i>PTEN</i>	10q23.31	Cowden disease.	Mutation of gene <i>PTEN</i>	Not conclusive	30
<i>DHCR7</i>	11q13.4	Smith-Lemli-Opitz syndrome	Mutations of gene <i>DHCR</i> , leading to a deficiency of cholesterol synthesis and an accumulation of 7-dehydrocholesterol	15–50% 3%	58-60 61, 62
<i>CACNA1C</i>	12p13.33	Timothy syndrome.	Missense mutations in the calcium channel gene <i>CACNA1H</i>	Not conclusive	63
<i>UBE3A</i>	15q11.2	Angelman syndrome	Maternal deletion, paternal UPD, deletions and epimutations at IC, mutations of <i>UBE3A</i> , Lack of expression of maternally expressed gene <i>UBE3A</i>	Not conclusive	32, 33
<i>TSC2</i>	16p13.3	Tuberous Sclerosis type II	Mutation in gene <i>TSC2</i> and subsequent hyperactivation of the downstream mTOR pathway, resulting in increased cell growth and proliferation.	Not conclusive	57
<i>NF1</i>	17q11.2	Neurofibromatosis	Polymorphisms within the intron-27, including the (AAAT)(n) and two (CA)n	Not conclusive	35
<i>DMD</i>	Xp21.2	Duchenne muscular dystrophy	Mutations of <i>DMD</i> gene resulting in absence of dystrophin protein	Not conclusive	64
<i>ARX</i>	Xp21.3	LIS, XLID, EPI, ASD	Naturally occurring mutations. Nonsense mutations, polyalanine tract expansions and missense mutations	Not conclusive	65
<i>FMR1</i>	Xq27.3	Fragile X syndrome	CGG repeat expansion and DNA methylation of <i>FMR1</i> gene, reduced <i>FMR1</i> expression	60–67% in males, 23% in female	66
<i>MECP2</i>	Xq28	Rett syndrome	Mutations in <i>MECP2</i> and <i>CDKL5</i>	Overlap in symptoms Infancy	67, 68

Abbreviations: LIS, lissencephaly; XLID, X-linked intellectual disability; EPI, epilepsy; OCD, obsessive compulsive disorder; TS, Tourette syndrome; ADHD, attention deficit hyperactivity disorder.

**Table 2.** Autism plus syndromic ASD caused by rare, single-gene disorders

Gene	Chromosome	Phenotype (human/mouse)	Mechanism involved in ASD	Reference
<i>NRXN1</i>	2p16.3	ASD, ID, SCZ, Language delay	De novo 320-kb deletion that removes the promoter and initial coding exons of the <i>NRXN1</i> gene, resulting in deletion of neurexin 1a	39
			Missense structural variants in the neurexin 1b signal peptide region	40
			CNV	69, 70
			Translocations and intragenic rearrangements in or near <i>NRXN1</i> gene	71, 72
<i>FOXP1</i>	3p13	ID, ASD, SLI	<i>De novo</i> intragenic deletion encompassing exons 4-14 of <i>FOXP1</i> , <i>de novo</i> nonsense mutation (c.1573C>T) in the conserved fork head DNA-binding domain	73
<i>GRIK2</i>	6q16.3	ASD, Recessive ID	SNP1 and SNP2 of gene <i>GRIK2</i> were associated with autism	74
<i>FOXP2</i>	7q31.1	ASD, SLI	Directly bind intron 1 of the <i>CNTNAP2</i> gene and regulate its expression	74
	11p15.5	Beckwith-Wiedemann syndrome	Overexpression of paternally expressed <i>IGF2</i> , due to a gain of DNA methylation at paternal allele of <i>IC1</i> and suppression of maternally expressed suppressing factor <i>CDKN1C</i>	75
	15q11-q13	Prader-Willi syndrome	Paternal deletions, maternal UPD at15q11-13, deletions and epimutations of <i>IC</i> , translocations disrupting <i>SNRPN</i>	76, 77
		Maternal duplication of 15q11-13 region	Maternal duplications of 15q11-13 region	78
<i>SHANK3</i>	22q13.33	ASD	Mutation at an intronic donor splice site, one missense mutation in the coding region	79
<i>NLGN4X</i>	Xp22.32-p22.31	ASD, ID, TS, ADHD	Frameshift mutation (1186insT)	37
<i>NLGN3</i>	Xq13.1	ASD	R451C mutation within the esterase domain of neuroligin 3	36, 37


Abbreviations: ID, intellectual disability; SCZ, schizophrenia; TS, Tourette syndrome; SLI, speech and language impairment; ADHD, attention deficit hyperactivity disorder

**Table 3.** Severe and specific phenotype with rare variants of genes

## 6. Copy number variation (CNV): A paradigm shift in autism

The strong genetic contribution shown in family studies and the association of cytogenetic changes, but apparent lack of common risk factors in autism, led to a hypothesis that rare sub-microscopic unbalanced changes in the form of CNVs likely contribute to the autism

phenotype. With the development of microarrays capable of scanning the genome at sub-microscopic resolution, there is accumulating evidence that multiple CNVs contribute to the genetic vulnerability to autism [80]. *de novo* CNV has been identified in up to 7–10% of sporadic autism [81, 82], but are less frequent in multiplex families, in which CNV accounts only for about 2% of families screened [80, 83]. This could possibly suggest different genetic liabilities in simplex and multiplex autism. Recurrent CNVs at 15q11-13 (1-3% of autism patients), 16p11 (1% of autism patients), and 22q11-13 have been confirmed in multiple studies [80, 83-86]. This hypothesis also has been proven largely successful in identifying autism-susceptibility candidate genes, including gains and losses at *SHANK2* [87], *SHANK3* [88], *NRXN1* [13], *NLGN3* and *NLGN4* [37], and *PTCHD1* [89, 90]. Neurexins and neuroligins are synaptic cell-adhesion molecules (CAMs) that connect pre- and postsynaptic neurons at synapses, mediate trans-synaptic signaling, and shape neural network properties by specifying synaptic functions. The Shank family of proteins provides scaffolding for signaling molecules in the postsynaptic density of glutamatergic synapses. Genes encoding CAMs play crucial roles in modulating or fine-tuning synaptic formation and synaptic specification. Localization and interacting proteins at the synapse is shown in Figure 1.



**Figure 1.** Localization of cell-adhesion molecules and their interacting proteins at the synapse. Proteins associated with ASD are underlined.

It is apparent that many different loci, each with a presumably unique yet subtle contribution to neurodevelopment, underlie the phenotype of autism. These observations have resulted in a paradigm shift away from the previously held “common disease-common variant” hypothesis to a “common disease-rare variant” model for the genetic architecture of autism. The central tenet of this model suggests a role for multiple, rare, highly penetrant, genetic risk factors for ASD, many of which are in the form of CNV. To make sense of the contribution of CNVs to autism, a “threshold” model has been proposed [80]. The model posits that different CNVs exhibit different penetrance depending on the dosage sensitivity and function (relative to autism) of the gene(s) they affect. Some CNVs have a large impact



on autism susceptibility and these are typically *de novo* in origin, cause more severe autistic symptoms, are more prevalent among sporadic forms of autism, and are less influenced by other factors like gender and parent of origin. Other CNVs have moderate or mild effects that probably require other genetic (or non-genetic) factors to take the phenotype across the autistic threshold.

## 7. Epigenetics plays an important role in autism

In addition to structural genetic factors that play causative roles for autism, environmental factors also play an important role in autism by influencing fetal or early postnatal brain development, directly or *via* epigenetic modifications. Epigenetic modifications include cytosine methylation, post-translational modification of histones, small interfering RNA and genomic imprinting. Involvement of epigenetic factors in autism is demonstrated by the central role of epigenetic regulatory mechanisms in the pathogenesis of Rett syndrome and fragile X syndrome (FXS), both are the monogenic disorders resulted from single gene defects and commonly associated with autism [38-40]. FXS is a result of a triplet expansion of CGG repeats at the 5' untranslated region of *FMR1* gene, which encodes the FMRP (fragile X mental retardation protein). FMRP is proposed to act as a translation regulator of specific mRNAs in the brain and involved in synaptic development and maturation, through its nucleo-cytoplasmic shuttle activity as an RNA-binding protein. It has been shown that FMRP uses its arginine-glycine-glycine (RGG) box domain to bind a subset of mRNA targets that form a G-quadruplex structure. FMRP has also been shown to undergo the post-translational modifications of arginine methylation and phosphorylation [91, 92]. Our recent study demonstrated that alteration of methylation patterns at loci of *Neurex1* and *ENO2* are associated with autism [Wang and Zhong, manuscript in preparation].

Genomic imprinting is the classic example of regulation of gene expression *via* epigenetic modifications, such as hypemethylation, that leads to parent of origin-specific gene expression. In addition, a growing number of genes that are not imprinted are regulated by DNA methylation, including Reelin (*RELN*) [41, 93-96], which has been considered as a candidate for autism. Several of the linkage peaks overlap or are in close proximity to regions that are subject to genomic imprinting on chromosomes 15q11-13, 7q21-31.31, 7q32.3-36.3 and possibly 4q21-31, 11p11.2-13 and 13q12.3, with the loci on chromosomes 15q and 7q demonstrating the most compelling evidence for a combination of genetic and epigenetic factors that confer risks for autism [97-101]. Genes in the imprinted cluster on chromosome 15q11-13 include *MKRN3*, *ZNF127AS*, *MAGE12*, *NDN*, *ATP10A*, *GABRA5*, *GABRB3*, and *GABRG3* [102, 103]. Genes in the imprinted cluster on chromosome 7q21.3 include *SGCE*, *PEG10*, *PPP1R9A*, *DLX5*, *CALCR*, *ASB4*, *PON1*, *PON2*, and *PON3* [104, 105].

Research has recently focused on the connections between the immune system and the early development of brain, including its possible role in the development of autism [106]. Immune aberrations consistent with a deregulated immune response may target neuronal

development and differentiation [107, 108]. Our study has suggested that a close contact with natural rubber latex (NRL) could trigger an immunoreaction to *Hevea brasiliensis* (Hev-b) proteins in NRL and resulted in autism [109]. This led us to a hypothesis that immune reactions triggered by environmental factors could damage synapse formation and neuronal connections, which would result in missing normal structure or function of synaptic proteins that are encoded by genes *NLGNs*, *NRXN1*, *CNTNAPs*, *SHANKs*, or in *deregulation of gene expression of FMR1, PTEN, FOXP*s, and *GRIK2*.

## 8. Converging molecular pathways of autism

Autism is a heterogeneous disorder with a fundamental question of whether autism represents an etiologically heterogeneous disorder in which a myriad of genetic or environmental risk factors perturb common underlying molecular pathways in the brain [110]. Two recent studies have suggested there could be convergence at the level of molecular mechanisms in autism. The first study on molecular convergence in autism identified protein interactors of known autism or autism-associated genes [111]. This interactome revealed several novel interactions, including between two autism candidate genes, *SHANK3* and *TSC1*. The biological pathways identified in this study include synapse, cytoskeleton and GTPase signaling, demonstrating a remarkable overlap with those identified by the gene expression. The second, an analysis of gene expression in postmortem autism brain, provides strong evidence for a shared set of molecular alterations in a majority of cases of autism. This included disruption of the normal gene expression pattern that differentiates frontal and temporal lobes and two groups of genes deregulated in autistic brains: one related to neuronal function, and the other related to immune/inflammatory responses [111]. Genes associated with neuronal function were enriched in metabolic signal pathways, providing evidence that these changes were causal, rather than the consequence of the disease [112]. In contrast, the immune/inflammatory changes did not show a strong genetic signal, indicating a non-genetic etiology for this process and implicating environmental or epigenetic factors instead. These results provide strong evidence for converging molecular abnormalities in autism, and implicating transcriptional and splicing deregulation as underlying mechanisms of neuronal dysfunction in this disorder.

## 9. In summary

Autism is a heterogeneous set of brain developmental disorders with complex genetics, involving interactions between genetic, epigenetic and environmental factors. The heterogeneous genetics involves many genes/loci and different genetic variations in autism, such as deletion, translocation, SNP and CNV. Recent studies have also suggested there could be convergence at the level of molecular mechanisms in autism. Although the genetic basis is well documented, considering phenotypic and genotypic heterogeneity, correspondences between genotype and phenotype have yet to be well established.

## Author details

Yu Wang<sup>1</sup>, Nanbert Zhong<sup>1,2,3,\*</sup>

<sup>1</sup>Shanghai Children's Hospital Affiliated to Shanghai Jiaotong University, Shanghai, China

<sup>2</sup>Peking University Center of Medical Genetics, Beijing, China

<sup>3</sup>New York State Institute for Basic Research in Developmental Disabilities, Staten Island, New York, USA

## Acknowledgement

This work was supported in part by the "973" program (2012CB517905) granted by the Chinese Ministry of Science and Technology, the Shanghai Municipal Department of Science and Technology (2009JC1412600), and the New York State Office of People with Developmental Disabilities (OPWDD).

## 10. References

- [1] Geschwind DH (2009) Advances in autism. *Annu Rev Med.* 60: 367–380.
- [2] Kogan MD, Blumberg SJ, Schieve LA (2007) Prevalence of parent-reported diagnosis of autism spectrum disorder among children in the US. *Pediatrics.* 124: 1395–1403.
- [3] Kim YS, Leventhal L, Koh YJ (2011) Prevalence of autism spectrum disorders in a total population sample. *Am. J. Psychiatry.* 168: 904–912.
- [4] Ganz ML (2006) The Costs of Autism In Moldin, SO and Rubenstein, JLR (eds), *Understanding Autism: from Basic Neuroscience to Treatment.* CRC Press, Boca Raton, FL, pp. 476–498.
- [5] Lord C, Pickles A, McLennan J (1997) Diagnosing autism: analyses of data from the Autism Diagnostic Interview. *Autism Dev Disord.* 27: 501-517.
- [6] Lord C, Risi S, Lambrecht L (2000) The autism diagnostic observation schedule-generic: a standard measure of social and communication deficits associated with the spectrum of autism. *Autism Dev Disord.* 30: 205-223.
- [7] Schopler E, Reichler R, Renner BR (1991) *The childhood autism rating scale.* Los Angeles: Western Psychological Services; 1988, *Psychol Monogr.* 117: 313-357.
- [8] Robins D, Fein D, Barton M, Green J (2001) The Modified Checklist for Autism in Toddlers: an initial study investigating the early detection of autism and pervasive developmental disorders. *Autism Dev Disord.* 31: 131-151.
- [9] Pinto MJ, Levy S (2004) Early diagnosis of autism spectrum disorders. *Curr Treat Options Neurol.* 6: 391-400.
- [10] Gillberg C and Coleman M (2000) *The biology of autistic syndromes*, 3<sup>rd</sup> ed. Mac Keith, London. 22p.
- [11] Fombonne E (2001) Is there an epidemic of autism? *Pediatrics.* 107: 411–412.

---

\* Corresponding Author

- [12] Szatmari P, Jones MB, Zwaigenbaum L (1998) Genetics of autism: overview and new directions. *J Autism and Dev Disord.* 28: 351–368.
- [13] Abrahams BS, Geschwind DH (2008) Advances in autism genetics: on the threshold of a new neurobiology. *Nat Rev Genet.* 9: 341–355.
- [14] Zwaigenbaum L, Bryson S, Roberts W (2005) Behavioral markers of autism in the first year of life. *Intern J. Dev Neurosci.* 23: 143–152.
- [15] Bailey A, Le Couteur A, Gottesman I (1995) Autism as a strongly genetic disorder: Evidence from a British twin study. *Psychological Medicine.* 25: 63–77.
- [16] Rosenberg RE, Law JK, Yenokyan G (2009) Characteristics and concordance of autism spectrum disorders among 277 twin pairs. *Arch Pediatr Adolesc Med.* 163: 907–914.
- [17] Hallmayer J, Cleveland S, Torres A (2011) Genetic heritability and shared environmental factors among twin pairs with autism. *Arch Gen Psychiatry.* 68: 1095–1102.
- [18] Bolton P, Macdonald H, Pickles A (1994) A case-control family history study of autism. *Child Psychol Psychiatry.* 35: 877–900.
- [19] Glessner JT, Wang K, Cai G (2009) Autism genome-wide copy number variation reveals ubiquitin and neuronal genes. *Nature.* 459: 569–573.
- [20] Wang K, Zhang H, Ma D (2009) Common genetic variants on 5p14.1 associate with autism spectrum disorders. *Nature.* 459: 528–533.
- [21] Bucan M, Abrahams BS, Wang K (2009) Genome-wide analyses of exonic copy number variants in a family-based study point to novel autism susceptibility genes. *PLoS Genet.* 5: e1000536.
- [22] Maddalena A, Richards CS, McGinniss MJ (2001) Technical standards and guidelines for Fragile X: The first of a series of disease-specific supplements to the Standards and Guidelines for Clinical Genetics Laboratories of the American College of Medical Genetics. Quality assurance subcommittee of the laboratory practice committee. *Genet Med.* 3: 200–205.
- [23] Pfeiffer BE, Huber KM (2009) The state of synapses in fragile X syndrome. *Neuroscientist.* 15: 549–567.
- [24] Tan H, Li H, Jin P (2009) RNA-mediated pathogenesis in fragile X-associated disorders. *Neurosci Lett.* 466: 103–108.
- [25] Goffin D, Allen M, Zhang L (2011) Rett syndrome mutation MeCP2 T158A disrupts DNA binding, protein stability and ERP responses. *Nat Neurosci.* 15: 274–283.
- [26] Strauss KA, Puffenberger EG, Huentelman MJ (2006) Recessive symptomatic focal epilepsy and mutant contactin-associated protein-like 2. *N Engl J Med.* 354: 1370–1377.
- [27] Bakkaloglu B, O'Roak BJ, Louvi A (2008) Molecular cytogenetic analysis and resequencing of contactin associated protein-like 2 in autism spectrum disorders. *Hum Genet.* 82: 165–173.
- [28] O'Roak BJ, Deriziotis P, Lee C (2011) Exome sequencing in sporadic autism spectrum disorders identifies severe de novo mutations. *Nat Genet.* 46: 585–589.
- [29] Nord AS, Roeb W, Dickel DE (2011) Reduced transcript expression of genes affected by inherited and de novo CNVs in autism. *Eur J Hum Genet.* 19: 727–731.

- [30] Conti S, Condò M, Posar A (2011) Phosphatase and Tensin Homolog (PTEN) Gene Mutations and Autism: Literature review and a case report of a patient with Cowden Syndrome, Autistic Disorder and Epilepsy. *J. Child Neurol.* 29: 123-126.
- [31] Splawski I, Yoo DS, Stotz SC (2006) CACNA1H mutations in autism spectrum disorders. *J. Biol Chem.* 281: 22085-22091.
- [32] Guffanti G, Strik Lievers L, Bonati MT (2011) Role of UBE3A and ATP10A genes in autism susceptibility region 15q11-q13 in an Italian population: a positive replication for UBE3A *Psychiatry Res.* 185: 33-38.
- [33] Nurmi EL, Bradford Y, Chen Y (2001) Linkage disequilibrium at the Angelman syndrome gene UBE3A in autism families. *Genomics.* 77: 105-113.
- [34] FJ Serajee, R Nabi, H Zhong (2003) Association of INPP1, PIK3CG, and TSC2 gene variants with autistic disorder: Implications for phosphatidylinositol. *J Med Genet.* 40: 119-123.
- [35] Marui T, Hashimoto O, Nanba E (2004) Association between the Neurofibromatosis-1 (NF1) locus and autism in the Japanese population. *Am J Med Genet B Neuropsychiatr Genet.* 131B: 43-47.
- [36] Jamain S, Quach H, Betancur C (2003) Mutations of the X-linked genes encoding neuroligins NLGN3 and NLGN4 are associated with autism. *Nat Genet.* 34: 27-29.
- [37] Comoletti D, De Jaco A, Jennings LL (2004) The Arg451 Cys- neuroligin-3 mutation associated with autism reveals a defect in protein processing. *J Neurosci.* 24: 4889-4893.
- [38] Friedman JM, Baross A, Delaney AD (2006) Oligonucleotide microarray analysis of genomic imbalance in children with mental retardation. *Am J Hum Genet.* 79: 500-513.
- [39] Zahir FR, Baross A, Delaney AD (2008) A patient with vertebral, cognitive and behavioural abnormalities and a de novo deletion of N RXN1a. *Med Genet.* 45: 239-243.
- [40] Feng J, Schroer R, Yan J (2006) High frequency of neurexin 1 signal peptide structural variants in patients with autism. *Neurosci Lett.* 409: 10-13.
- [41] Hamdan FF, Daoud H, Rochefort D (2010) De novo mutations in FOXP1 in cases with intellectual disability, autism, and language impairment. *Am J Hum Genet.* 87: 671-678.
- [42] Li H, Yamagata T, Mori M (2005) Absence of causative mutations and presence of autism-related allele in FOXP2 in Japanese autistic patients. *Brain Dev.* 27: 207-210.
- [43] Mukamel Z, Konopka G, Wexler E (2011) Regulation of MET by FOXP2, genes implicated in higher cognitive dysfunction and autism risk. *J Neurosci.* 31: 11437-11442.
- [44] Jamain S, Betancur C, Quach H (2002) Linkage and association of the glutamate receptor 6 gene with autism. *Mol Psychiatry.* 7: 302-310.
- [45] Durand CM, Perroy J, Loll F (2012) SHANK3 mutations identified in autism lead to modification of dendritic spine morphology via an actin-dependent mechanism. *Mol Psychiatry.* 17: 71-84.
- [46] Kolevzon A, Cai G, Soorya L (2011) Analysis of a purported SHANK3 mutation in a boy with autism: clinical impact of rare variant research in neurodevelopmental disabilities. *Brain Res.* 1380: 98-105.
- [47] Chen CP, Lin SP, Chern SR (2010) A de novo 7.9 Mb deletion in 22q13.2→qter in a boy with autistic features, epilepsy, developmental delay, atopic dermatitis and abnormal immunological findings. *Eur J Med Genet.* 53: 329-332.

- [48] Chiocchetti A, Klauck SM (2011) Genetic analyses for identifying molecular mechanisms in autism spectrum disorders. *Encephale*. 37: 68-74.
- [49] Bonnet-Brilhault F. (2011) Genotype/phenotype correlation in autism: genetic models and phenotypic characterization. *Encephale*. 37: 68-74.
- [50] Eapen V (2011) Genetic basis of autism: is there a way forward? *Curr Opin Psychiatry*. 24: 226-236.
- [51] Vernes SC, Newbury DF, Abrahams BS (2008) A functional genetic link between distinct developmental language disorders. *N Engl J Med*. 359: 2337–2345.
- [52] Newbury DF, Paracchini S, Scerri TS (2011) Investigation of dyslexia and SLI risk variants in reading- and language-impaired subjects. *Behav Genet*. 41: 90–104.
- [53] Poot M, Beyer V, Schwaab I (2010) Disruption of CNTNAP2 and additional structural genome changes in a boy with speech delay and autism spectrum disorder. *Neurogenetics*. 11: 81–89.
- [54] Sehested LT, Møller RS, Bache I (2010) Deletion of 7q34-q36.2 in two siblings with mental retardation, language delay, primary amenorrhea, dysmorphic features. *Am J Med Genet*. 152A: 3115–3119.
- [55] Teramitsu I, Kudo LC, London SE (2004) Parallel FoxP1 and FoxP2 expression in songbird and human brain predicts functional interaction. *Neurosci*. 24: 3152–3163.
- [56] Panaitof SC, Abrahams BS, Dong H (2010) Language-related Cntnap2 gene is differentially expressed in sexually dimorphic song nuclei essential for vocal learning in songbirds. *Comp. Neurol*. 518: 1995–2018.
- [57] Shoubridge C, Tan MH, Fullston T (2010) Mutations in the nuclear localization sequence of the Aristaless related homeobox; sequestration of mutant ARX with IPO13 disrupts normal subcellular distribution of the transcription factor and retards cell division. *Pathogenetics*. 3: 1.
- [58] Hartshorne TS, Grialou TL, Parker KR (2005) Autistic-like behavior in CHARGE syndrome. *Am J Med Genet A*. 133A: 257-261.
- [59] Johansson M, Rastam M, Billstedt E (2006) Autism spectrum disorders and underlying brain pathology in CHARGE association. *Dev Med Child Neurol*. 48: 40-50.
- [60] Smith IM, Nichols SL, Issekutz K (2005) Behavioral profiles and symptoms of autism in CHARGE syndrome: preliminary Canadian epidemiological data. *Am J Med Genet A*. 133A: 248-256.
- [61] Skuse DH, James RS, Bishop DV (1997) Evidence from Turner's syndrome of an imprinted X-linked locus affecting cognitive function. *Nature*. 387: 705-708.
- [62] Bianconi SE, Conley SK, Keil MF (2011) Adrenal function in Smith-Lemli-Opitz syndrome. *Am J Med Genet A*. 155A: 2732-2738.
- [63] Depil K, Beyl S, Stary-Weinzinger A (2011) Timothy mutation disrupts the link between activation and inactivation in Ca(V)1.2 protein. *J Biol Chem*. 286: 31557-31564.
- [64] Klymiuk N, Thirion C, Burkhardt K (2011) 238 tailored pig model of Duchenne muscular dystrophy. *Reprod Fertil Dev*. 24: 231.
- [65] Valerio N, Romina M, Paolo C (2009) Recent advances in neurobiology of Tuberous Sclerosis Complex. *Brain Dev*. 31: 104-113.

- [66] Bianconi SE, Conley SK, Keil MF (2011) Adrenal function in Smith-Lemli-Opitz syndrome. *Am J Med Genet A. J.* 155A: 2732-2738.
- [67] Coutinho AM, Oliveira G, Katz C (2007) MECP2 coding sequence and 3'UTR variation in 172 unrelated autistic patients. *Am J Med Genet B Neuropsychiatr Genet.* 144B: 475-483.
- [68] Shibayama A, Cook EH, Feng J (2004) MECP2 structural and 3'-UTR variants in schizophrenia, autism and other psychiatric diseases: a possible association with autism. *Am J Med Genet B Neuropsychiatr Genet.* 128B: 50-53.
- [69] Glessner JT, Wang K, Cai G (2009) Autism genome-wide copy number variation reveals ubiquitin and neuronal genes. *Nature.* 459: 569-573.
- [70] Szatmari P, Paterson AD, Zwaigenbaum L (2007) Mapping autism risk loci using genetic linkage and chromosomal rearrangements. *Nat Genet.* 39: 319-328.
- [71] Kim HG, Kishikawa S, Higgins AW (2008) Disruption of neurexin 1 associated with autism spectrum disorder. *Am J Hum Genet.* 82: 199-207.
- [72] Wisniewiecka KB, Nesteruk M, Peters SU (2010) Intragenic rearrangements in NRXN1 in three families with autism spectrum disorder, developmental delay, and speech delay. *Am J Med Genet B Neuropsychiatr Genet.* 153B: 983-993.
- [73] Hamdan FF, Daoud H, Rochefort D (2010) De novo mutations in FOXP1 in cases with intellectual disability, autism, and language impairment. *Am J Hum Genet.* 87: 671-678.
- [74] Casey JP, Magalhaes T, Conroy JM (2011) Regan RA novel approach of homozygous haplotype sharing identifies candidate genes in autism spectrum disorder. *Hum Genet.* 131: 565-579.
- [75] Kent L, Bowdin S, Kirby GA (2008) Beckwith Weidemann syndrome: a behavioral phenotype-genotype study. *Am J Med Genet B Neuropsychiatr Genet.* 147B: 1295-1297.
- [76] Descheemaeker MJ, Govers V, Vermeulen PJ (2006) Pervasive developmental disorders in Prader-Willi syndrome: the Leuven experience in 59 subjects and controls. *Am J Med Genet A.* 140: 1136-1142.
- [77] Veltman MW, Thompson RJ, Roberts SE (2004) Prader-Willi Syndrome—a study comparing deletion and uniparental disomy cases with reference to autism spectrum disorders. *Eur Child Adolesc Psychiatry.* 13: 42-50.
- [78] Hogart A, Wu D, Lasalle JM (2010) The comorbidity of autism with the genomic disorders of chromosome 15q11.2-q13. *Neurobiol Dis.* 38: 181-191.
- [79] Gauthier J, Champagne N, Lafrenière RG (2010). De novo mutations in the gene encoding the synaptic scaffolding protein SHANK3 in patients ascertained for schizophrenia. *Proc Natl Acad Sci.* 107: 7863-7868.
- [80] Cook EH, Scherer SW (2008) Copy number variations associated with neuropsychiatric conditions. *Nature.* 16: 919-923.
- [81] Sebat J, Lakshmi B, Malhotra D (2007) Strong association of de novo copy number mutations with autism. *Science.* 316: 445-449.
- [82] Marshall CR, Noor A, Vincent JB (2008) Structural variation of chromosomes in autism spectrum disorder. *Am J Hum Genet.* 82: 477-488.
- [83] Morrow EM, Yoo SY, Flavell SW (2008) Identifying autism loci and genes by tracing recent shared ancestry. *Science.* 321: 218-223.

- [84] Szatmari P, Paterson AD, Zwaigenbaum L (2007) Mapping autism risk loci using genetic linkage and chromosomal rearrangements. *Nat Genet.* 39: 319-328.
- [85] Weiss LA, Shen Y, Korn JM (2008) Association between microdeletion and microduplication at 16p11.2 and autism. *N Engl J Med.* 358: 667-675.
- [86] Kumar RA, KaraMohamed S, Sudi J (2008) Recurrent 16p11.2 microdeletions in autism. *Hum Mol Genet.* 17: 628-638.
- [87] Berkel S, Marshall CR, Weiss B (2010) Mutations in the SHANK2 synaptic scaffolding gene in autism spectrum disorder and mental retardation. *Nature Genetics.* 42: 489-491.
- [88] Durand CM, Betancur C, Boeckers TM (2007) Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. *Nature Genetics.* 39: 25-27.
- [89] Pinto D, Pagnamenta AT, Klei L (2010) Functional impact of global rare copy number variation in autism spectrum disorder. *Nature.* 466: 368-372.
- [90] Noor A, Whibley A, Marshall CR (2010) Disruption at the PTCHD1 locus on Xp22.11 in autism spectrum disorder and intellectual disability. *Sci Transl Med.* 2: 49ra68.
- [91] Auerbach BD, Osterweil EK, Bear MF (2011) Mutations causing syndromic autism define an axis of synaptic pathophysiology. *Nature.* 480: 63-68.
- [92] Evans TL, Blice-Baum AC, Mihailescu MR (2012) Analysis of the Fragile X mental retardation protein isoforms 1, 2 and 3 interactions with the G-quadruplex forming semaphorin 3F mRNA. *Mol Biosyst.* 8: 642-649.
- [93] Noh JS, Sharma RP, Veldic M (2005) DNA methyltransferase1 regulates reelin mRNA expression in mouse primary cortical cultures. *Proc Natl Acad Sci USA.* 102: 1749-1754.
- [94] Grayson DR, Chen Y, Costa E (2006) The human reelin gene: Transcription factors (t), repressors (2) and the methylation switch(t/2) in schizophrenia. *Pharmacol. Ther.* 111: 272-286.
- [95] Sato N, Fukushima N, Chang R (2006) Differential and epigenetic gene expression profiling identifies frequent disruption of the RELN pathway in pancreatic cancers. *Gastroenterology.* 30: 548-565.
- [96] Serajee FJ, Zhong H, Mahbulul AH (2006) Association of Reelin gene polymorphisms with autism. *Genomics.* 87: 75-83.
- [97] Numachi Y, Yoshida S, Yamashita M (2004) Psychostimulant alters expression of DNA methyltransferase mRNA in the rat brain. *Ann. NY Acad Sci.* 1025: 102-109.
- [98] Huang CH, Chen CH. (2006) Absence of association of a polymorphic GGC repeat at the 5' untranslated region of the reelin gene with schizophrenia. *Psychiatry Res.* 142: 89-92.
- [99] Skaar DA, Shao Y, Haines JL (2005) Analysis of the RELN gene as a genetic risk factor for autism. *Mol. Psychiatry.* 10: 563-571.
- [100] Li J, Nguyen L, Gleason C (2004) Lack of evidence for an association between WNT2 and RELN polymorphisms and autism. *Am J Med Genet B Neuropsychiatr. Genet.* 126: 51-57.
- [101] Bonora E, Beyer KS, Lamb JA (2003) Analysis of reelin as a candidate gene for autism. *Mol. Psychiatry.* 8: 885-892.



- [102] Lee S, Walker CL, Karten B (2005) Essential role for the Prader-Willi syndrome protein *neccdin* in axonal outgrowth. *Hum Mol Genet.* 14: 627–637.
- [103] Kashiwagi A, Meguro M, Hoshiya H (2003) Predominant maternal expression of the mouse *Atp10c* in hippocampus and olfactory bulb. *Hum Genet.* 48: 194–198.
- [104] Draganov DI, Teiber JF, Speelman A (2005) Human paraoxonases (PON1, PON2 and PON3) are lactonases with overlapping and distinct substrate specificities. *Lipid Res.* 46: 1239–1247.
- [105] Terry-Lorenzo RT, RoadcapDW, Otsuka T (2005) Neurabin/protein phosphatase-1 complex regulates dendritic spine morphogenesis and maturation. *Mol Biol Cell.* 16: 2349–2362.
- [106] Croen LA, Grether JK, Yoshida CK (2005) Maternal autoimmune diseases, asthma, and allergies, and childhood autism spectrum disorders. *Arch Pediatr Adolesc Med.* 159: 151–157.
- [107] Braunschweig D, Ashwood P, Krakowiak P (2008) Autism: maternally derived antibodies specific for fetal brain proteins. *NeuroToxicology.* 29: 226–231.
- [108] Singer HS, Morris CM, Gause CD (2008) Antibodies against fetal brain in sera of others with autistic children. *Neuroimmunol.* 194: 165–172.
- [109] Shen C, Zhao XL, Zhong N. (2010) A proteomic investigation of B lymphocytes in an autistic family: A pilot study of exposure to natural rubber latex (NRL) may lead to autism. *J Mol Neurosci.* 43: 443–452.
- [110] Glessner JT, Wang K, Cai G (2009) Autism genome-wide copy number variation reveals ubiquitin and neuronal genes. *Nature.* 459: 569–573.
- [111] Sakai Y, Shaw CA, Dawson BC (2011) Protein interactome reveals converging molecular pathways among autism disorders. *Sci Transl Med.* 3: 86ra49.
- [112] Voineagu I, Wang X, Johnston P (2011) Transcriptomic analysis of autistic brain reveals convergent molecular pathology. *Nature.* 474: 380–384.