

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Phytochemicals of the Chinese Herbal Medicine *Tacca chantrieri* Rhizomes

Akihito Yokosuka and Yoshihiro Mimaki

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/53668>

1. Introduction

The family Taccaceae is composed of two genera, *Tacca* and *Schizocapsa*, and about 10 species, with most distributed in tropical regions of Asia, the Pacific Islands, and Australia [1]. *Tacca chantrieri* André is a perennial plant that occurs in the southeast region of mainland China, and its rhizomes have been used for the treatment of gastric ulcers, enteritis, and hepatitis in Chinese folk medicine. According to a Chinese herbal dictionary, *T. plantaginea* has also been used for the same purposes as *T. chantrieri* [2]. The chemical constituents of *T. plantaginea* have been extensively examined and a series of highly oxygenated pentacyclic steroids named taccalonolids, which have a γ -enol lactone, have been isolated as characteristic components of the herb [3], but there has been only one report of the secondary metabolites of *T. chantrieri*, in which a few trivial sterols such as stigmasterol and daucosterol, and a diosgenin glycoside were found [4]. Therefore, we focused our attention on the constituents of *T. chantrieri* rhizomes, and a detailed phytochemical investigation of this herbal medicine has been carried out.

In this chapter, we describe the phytochemicals isolated from *T. chantrieri* rhizomes and their biological activities with a focus on cytotoxicity against human cancer cells.

2. Isolation and structural determination

T. chantrieri specimens were collected in Yunnan Province, People's Republic of China. The rhizomes of *T. chantrieri* (fresh weight, 7.3 kg) were extracted with hot MeOH (3 L \times 2). The MeOH extract was concentrated under reduced pressure, and the extract was passed through a polystyrene resin (Diaion HP-20) column eluted with MeOH/H₂O gradients,

EtOH, and EtOAc. The 50% MeOH and MeOH eluate portion was subjected to silica gel and octadecylsilanized silica gel column chromatography to afford a total of 41 compounds, classified into diarylheptanoids (**1** and **2**), diarylheptanoid glucosides (**3–9**), ergostane glucosides (**10–21**), withanolide glucosides (**22** and **23**), spirostan glycosides (**24–28**), furostan glycosides (**29–32**), pseudofurostan glycosides (**33–37**), pregnane glycosides (**38–40**), and a phenolic glucoside (**41**) (Fig.1). Their structures were determined through extensive spectroscopic studies and through chemical transformations followed by chromatographic and spectroscopic analysis.




Figure 1. Extraction, partition, and purification procedures

3. Diarylheptanoids and diarylheptanoid glucosides

Diarylheptanoids consist of two phenyl groups linked by a linear seven-carbon aliphatic chain. Compounds **1** and **2** are diarylheptanoids and **3–9** are diarylheptanoid monoglucosides (Fig. 2) [5].




Figure 2. Structures of **1–9** and their derivatives

Compound **1** was isolated as a viscous syrup, $[\alpha]_D +1.7^\circ$ (MeOH). HREIMS of **1** showed an $[M]^+$ peak at m/z 332.1623, corresponding the empirical molecular formula of $C_{19}H_{24}O_5$, which was also deduced by analysis of its ^{13}C NMR and DEPT spectral data. The IR spectrum suggested the presence of hydroxy groups (3347 cm^{-1}) and aromatic rings (1611 and 1515 cm^{-1}). The UV spectrum showed an absorption maximum due to substituted aromatic rings (281.4 nm). The planar structure of **1** was assigned as 3,5-dihydroxy-1-(3,4-dihydroxyphenyl)-7-(4-hydroxyphenyl)heptane by analysis of the 1D (1H and ^{13}C) and 2D (1H - 1H COSY, HMQC, and HMBC) spectra. The absolute configuration of the 3,5-dihydroxy moieties of the new diarylheptanoids were determined by applying the CD exciton chirality method to acyclic 1,3-dibenzoates [6]. The trimethyl derivative (**1a**) was converted to the corresponding 3,5-bis(*p*-bromobenzoate) (**1b**) and its CD spectrum exhibited positive (237.4 nm , $\Delta\epsilon +29.9$) and negative (253.3 nm , $\Delta\epsilon -20.0$) Cotton effects, which were consistent with a negative chirality. Thus, the absolute configurations were determined as $3R$ and $5R$ (Fig. 3). The

structure of **1** was shown to be (3*R*,5*R*)-3,5-dihydroxy-1-(3,4-dihydroxyphenyl)-7-(4-hydroxyphenyl)heptane. In the same way, the structure of **2** was elucidated as (3*R*,5*R*)-3,5-dihydroxy-1,7-bis(3,4-dihydroxyphenyl)heptane.




Figure 3. Determination of the absolute configurations at C-3 and C-5 of **1**

Compounds **3–9** are diarylheptanoid monoglucosides. Enzymatic hydrolysis of **3–9** with naringinase gave the diarylheptanoid derivatives and D-glucose. Identification of D-glucose, including its absolute configuration, was carried out by direct HPLC analysis of the hydrolysates. In the HMBC spectra, a long-range correlation was observed from each anomeric proton to the C-3 carbon in **3** and **5–9**, and to the C-5 carbon in **4**.

Diarylheptanoids are known to occur in only a limited number species of higher plants belonging to the families Zingiberaceae [7–10], Betulaceae [11], and Aceraceae [12]. This is the first isolation of diarylheptanoids from a plant of the family Taccaceae.

4. Ergostane glucosides

Compounds **10–21** are new ergostane glucosides (Fig. 4) [13–15]. Taccasterosides A–C (**10–12**) are novel bisdesmosideic oligoglucosides of (24*R*,25*S*)-3 β -hydroxyergost-5-ene-26-oic acid (**10a**), whereas **13–20** are those of (24*S*,25*R*)-ergost-5-ene-3 β ,26-diol (**10b**). Compound **21** is an ergostane glucoside with the six-membered lactone on the side chain of the aglycone.

Taccasteroside A (**10**) was obtained as an amorphous solid. Acid hydrolysis of **10** with 1 M HCl in dioxane/H₂O gave D-glucose and a C₂₈-sterol as the aglycone (**10a**). The structure of **10a**, except for the absolute configurations at C-24 and C-25, was identified as 3 β -hydroxyergost-5-en-26-oic acid by analysis of its ¹H, ¹³C, and 2D NMR spectra. In order to determine





Figure 4. Structures of **10-21**

the absolute configuration at C-25, **10a** was reduced with LiAlH₄ to (24*R*,25*S*)-ergost-5-ene-3β,26-diol (**10b**). Then, **10b** was converted to the diastereomeric pairs of (*R*)-MTPA (**10a-R**) and (*S*)-MTPA (**10a-S**) esters with respect to the C-26 primary hydroxy group next to the C-25 chiral center and the differences in the ¹H NMR coupling patterns of the H₂-26 protons

were inspected. The H₂-26 protons of **10a-R** were observed as a doublet-like signal at δ 4.20 ($J = 6.3$ Hz), whereas those of **10a-S** were observed as a doublet of doublets at δ 4.30 ($J = 10.8, 6.6$ Hz) and 4.09 ($J = 10.8, 7.2$ Hz). Application of these spectral data to the empirical rule reported by Yasuhara et al. [17] allowed us to confirm that the C-25 configuration was exclusively *S*. The configuration of C-24 position and other steroidal skeleton were established by the following chemical transformations. Compound **10b** was treated with *p*-toluenesulfonyl chloride to give the 26-*O*-tosylate of **10b** (**10b-T**), which was then reduced with LiAlH₄, affording (24*R*)-ergost-5-ene-3 β -ol, that is, campesterol. The structure of **10a** was determined as (24*R*,25*S*)-3 β -hydroxyergost-5-en-26-oic acid (Fig. 5).



Reagents and conditions: a, LiAlH₄, THF, 0 °C, 5 h; b, (*R*)-MTPA or (*S*)-MTPA, EDC·HCl, 4-DMAP, CH₂Cl₂, r.t., 12 h; c, *p*-TsCl, pyridine, r.t., 6 h; d, LiAlH₄, THF, 0 °C, 5 h

Figure 5. Chemical transformations of **10a**

The severe overlap of the proton signals for the sugar moieties in **10** excluded the possibility of complete assignment in a straightforward way by conventional 2D NMR methods such as the ¹H-¹H COSY, 2D TOCSY, and HSQC spectroscopy. The exact structures of the sugar moieties and their linkage positions of the aglycone were resolved by detailed analysis of the 1D TOCSY and 2D NMR spectra. The ¹H NMR subspectra of individual monosaccharide units were obtained by using selective irradiation of easily identifiable anomeric proton signals, as well as irradiation of other nonoverlapping proton signals in a series of 1D TOCSY experiments [17–19]. Subsequent analysis of the ¹H-¹H COSY spectrum resulted in the sequential assignment of all the proton resonances due to the seven glucosyl units, including identification of their multiplet patterns and coupling constants. The HSQC and HSQC-TOCSY spectra correlated the proton resonances to those of the corresponding one-bond coupled carbons, leading to unambiguous assignments of the carbon shifts. The carbon chemical shifts thus assigned were compared with those of the reference methyl α -D- and β -D-glucosides [20], taking into account the known effects of *O*-glycosylation shifts. The comparison indicated that **10** contained three terminal β -D-glucopyranosyl moieties (Glc', Glc''', Glc'''''), three C-4 substituted β -D-glucopyranosyl moieties (Glc''', Glc''''', Glp'''''), and a C-2 and C-6 disubstituted β -D-glucopyranosyl moiety (Glc''). The β -orientations of the anomeric centers of all the glucosyl moieties were supported by the relatively large J values of their anomeric protons (7.7–8.4 Hz).

In the HMBC spectrum, the anomeric proton of the terminal glucosyl unit (Glc') at δ 5.07 exhibited a long-range correlation with C-3 of the aglycone at δ 78.2, indicating that one glucosyl unit was attached to the C-3 hydroxy group of the aglycone. Consequently, an oligoglucoside composed of six glucosyl units was presumed to be linkage with the C-26 carboxy group of the aglycone. Further HMBC correlations from H-1 of Glc'' at δ 6.30 to C-26 of the aglycone at δ 175.2, H-1 of Glc''' at δ 5.20 to C-2 of Glc'' at δ 82.9, H-1 of Glc'''' at δ 5.17 to C-4 of Glc'''' at δ 80.9, H-1 of Glc'''' at δ 5.16 to C-4 of Glc'''' at δ 81.5, H-1 of Glc'''' at δ 5.13 to C-4 of Glc'''' at δ 80.9, and H-1 of Glc'''' at δ 4.93 to C-6 of Glc'' at δ 69.2 confirmed the hexaglucoside sequence as Glc-(1 \rightarrow 4)-Glc-(1 \rightarrow 4)-Glc-(1 \rightarrow 2)-[Glc-(1 \rightarrow 4)-Glc-(1 \rightarrow 6)]-Glc, which was attached to C-26 of the aglycone (Fig. 6). Accordingly, the structure of **10** was elucidated as (24*R*,25*S*)-3 β -[(β -D-glucopyranosyl)oxy]-ergost-5-en-26-oic acid *O*- β -D-glucopyranosyl-(1 \rightarrow 4)-*O*- β -D-glucopyranosyl-(1 \rightarrow 4)-*O*- β -D-glucopyranosyl-(1 \rightarrow 2)-*O*-[*O*- β -D-glucopyranosyl-(1 \rightarrow 4)- β -D-glucopyranosyl-(1 \rightarrow 6)]- β -D-glucopyranosyl ester.

In the same way, the structures of **11–20** were elucidated as shown in Fig. 4.




Figure 6. HMBC correlations of the sugar moieties of **10**

Phytosterols and their monoglucosides such as campesterol, stigmasterol, and β -sitosterol, and their 3-*O*-glucoside, widely occur in the plant kingdom. However, 10–20 are the first representatives of oligoglucosides of a phytosterol derivative to have sugar moieties with a total of four to seven glucose units. The bisdesmosidic nature of these structures, except for 15, is also notable.

5. Withanolide glucosides

Compounds 22 and 23 are withanolide glucosides, named chantriolides A and B (Fig. 7) [21]. Chantriolides A and B were found to be minor components relative to the other secondary metabolites concomitantly isolated from *T. chantrieri*. However, it is notable that withanolides, which have been isolated almost exclusively from plants of the family Solanaceae previously [22, 23], have now been found in a species of the family Taccaceae in the study.




Figure 7. Structures of 22 and 23

6. Other glycosides

Spirostan glucosides (24–28), furostan glycosides (29–32), pseudofurostan glycosides (33–37), pregnane glycosides (38–40), and a phenolic glucoside (41) were also isolated from *T. chantrieri* rhizomes (Fig. 8) [15, 24–26].

The known naturally occurring 22,26-hydroxyfurostan glycosides exclusively exist in the form of glycoside, bearing a monosaccharide at C-26 [27]. The monosaccharide among the furostan glycosides reported thus far is limited to β -d-glucopyranose, except for one furostan glycoside from *Dracaena afromontana*, which has an α -l-rhamnopyranosyl group at C-26 [28]. Compound 31 is distinctive in carrying a diglucosyl group, *O*-glucosyl-(1→6)-glucosyl, in place of a monoglucosyl unit at C-26.

Compounds **33** is the corresponding $\Delta^{20(22)}$ -furostan glycoside of **29**. This was confirmed by the fact that the peracetate (**33a**) of **33** agreed with the product (**29a**) obtained by treatment of **29** with Ac_2O in pyridine at 110°C for 2.5 h, during which dehydration at C-20 and C-22, as well as the introduction of an acetyl group to all the hydroxy groups of the sugar moieties, occurred (Fig. 9).

The structure of **38**, including the absolute configuration at C-25, was found by the following chemical conversion. When the C-20 and C-22 bond of **33a** was oxidatively cleaved by treating it with CrO_3 in AcOH at room temperature for 2 h, the resultant product was completely consistent with the peracetyl derivative of **38** (**38a**) (Fig. 9).




Figure 8. Structures of **24–41**




Figure 8. Continued.

A few compounds related to **38** and **39** have been isolated [29-31]; however, their C-25 configuration is not clearly presented in all the reports. In this investigation, we unequivocally determined the C-25 configuration of **38** to be *S* by a chemical correlation method. Compounds **38** and **39** could be defined as pregnane glycosides rather than furostan glycosides.




Figure 9. Chemical correlations of the furostan glycosides

7. Biological activity

7.1. Cytotoxic activity against HL-60 cells

The isolated compounds were evaluated for their cytotoxic activity against HL-60 human promyelocytic leukemia cells by a modified MTT assay method [32]. Diarylheptanoids (**1** and **2**), diarylheptanoid glucosides (**3**, **4**, **6**, and **7**), and spirostan glycosides (**24** and **28**) showed moderate cytotoxicity (IC_{50} 1.8–6.4 $\mu\text{g}/\text{mL}$) against HL-60 cells. Compounds **5**, **8–23**, **25–27**, and **29–41** did not show apparent cytotoxic activity against HL-60 cells at a sample concentration of 10 $\mu\text{g}/\text{mL}$.

7.2. Cytotoxic activity and structure–activity relationships of diarylheptanoids and diarylheptanoid glucosides against HL-60 cells, HSC-2 cells, and HGF

The diarylheptanoids and some derivatives, including **9b** prepared by treatment of **9** with CH_2N_2 , were evaluated for their cytotoxic activities against HL-60 cells, HSC-2 human oral squamous carcinoma cells, and normal human gingival fibroblasts (HGF) (Table 1). The diarylheptanoids **1**, **2**, and **7a**, and the diarylheptanoid glucosides **3**, **4**, **6**, and **7**, each of which has three or four phenolic hydroxy groups, showed moderate cytotoxic activity against HL-60 cells with IC_{50} values ranging 1.8 to 6.4 $\mu\text{g}/\text{mL}$, while those possessing two phenolic hydroxy groups (**5**, **5a**, **8**, **8a**, **9**, and **9a**) did not exhibit apparent cytotoxic activity even at a sample concentration of 10 $\mu\text{g}/\text{mL}$. Notably, the diarylheptanoids whose phenolic hydroxy

groups were all masked with methyl groups (**1a**, **2a**, and **9b**) were also cytotoxic. These observations suggest that the number of phenolic hydroxy groups contributes to the resultant cytotoxicity. Compounds **1a**, **2a**, and **9b** showed considerable cytotoxic activity against HSC-2 cells, whereas they had little effect on normal HGF.

compound	IC ₅₀ (μg/mL)		
	HL-60	HSC-2	HGF
1	2.1	54.0	162
1a	5.5	3.9	176
2	1.8	54.0	>250
2a	4.9	6.6	174
3	6.2	158	220
4	5.5	155	>250
5	>10	160	>250
5a	>10	– ^b	– ^b
6	3.0	92.0	189
7	4.5	209	>250
7a	4.1	– ^b	– ^b
8	>10	198	>250
8a	>10	– ^b	– ^b
9	>10	157	213
9a	>10	231	177
9b	6.4	23.0	173
etoposide	0.2	24.0	>200

^aKey: HL-60 (human promyelocytic leukemia cells); HSC-2 (human oral squamous carcinoma cells); and HGF (normal human gingival fibroblasts). ^bnot determined.

Table 1. Cytotoxic activities of compounds **1-9** and their derivatives (**1a**, **4a**, **5a**, **7a-9a**, and **9b**), and etoposide against HL-60 cells, HSC-2 cells, and HGF^a

7.3. Cytotoxic activity and structure–activity relationships of steroidal glycosides against HL-60 cells

Spirostan glycosides (**24** and **28**) showed moderate cytotoxicity (IC₅₀ 1.9 and 1.8 μg/mL) against HL-60 cells. Compounds **25** and **27**, the corresponding C-24 hydroxy derivatives of **24** and **28**, and **26**, the analogue of **24** without the terminal rhamnosyl group linked to C-2 of the inner glucosyl residue, did not show any cytotoxic activity at a sample concentration of 10 μg/mL. Furostan glycosides (**29–32**), pseudofurostan glycosides (**33–37**), and pregnane glycosides (**38–40**) also did not show cytotoxic activity. These data suggest that the structures of both the aglycone and sugar moieties contribute to the cytotoxicity.

7.4. Panel screening in the Japanese Foundation for Cancer Research 39 cell line assay

Diarylheptanoid **2** and spirostan glycosides **24**, which showed significant cytotoxic activity against HL-60 cells, were subjected to the Japanese Foundation for Cancer Research 39 cell line assay [33]. Subsequent evaluation of **2** and **24** showed that the mean concentration required for achieving GI_{50} levels against the panel of cells were 87 μM and 1.8 μM , respectively. Although **2** and **24** exhibited no significant differential cellular sensitivity, some cell lines such as colon cancer HCT-116 (GI_{50} 25 μM), ovarian cancer OVCAR-3 (GI_{50} 36 μM), OVCAR-4 (GI_{50} 39 μM), and stomach MKN-7 (GI_{50} 34 μM) were relatively sensitive to **2**.

8. Conclusion

Our systematic chemical investigations of *T. chantrieri* rhizomes revealed that this plant contains a variety of secondary metabolites, namely, diarylheptanoids, diarylheptanoid glucosides, steroidal glycosides with the aglycone structures of ergostane, withanolide, spirostan, furostan, pseudofurostan, and pregnane, as well as a phenolic glucoside. Some diarylheptanoids and steroidal glycosides showed cytotoxicity against human cancer cells. These compounds may be possible leads for new anticancer drugs.

On the other hand, a number of researchers have reported biological activities of diarylheptanoids and steroidal glycosides other than cytotoxicity. It has been reported that curcuminoids, well-known diarylheptanoid derivatives, showed antioxidant [34, 35], anti-inflammatory [35, 36], estrogenic [37, 38], and anticancer [39] effects. Steroidal glycosides have been shown to have antidiabetic [40, 41], antitumor [42], antitussive [43], antiherpes virus [44], and platelet aggregation inhibitory [45] activities. *T. chantrieri* rhizomes could be applied to treating a wide variety of ailments as an alternative herbal medicine.

Acknowledgements

We are grateful to Dr. Hiroshi Sakagami for evaluating the cytotoxic activities against HSC-2 cells and HGF.

Author details

Akihito Yokosuka and Yoshihiro Mimaki

Tokyo University of Pharmacy and Life Sciences, School of Pharmacy, Tokyo, Japan

References

- [1] Tsukamoto Y. (ed.) The Grand Dictionary of Horticulture Vol. 1. Tokyo: Shogakukan; 1989; pp 148–149.
- [2] Dictionary of Chinese Medicinal Materials Vol. 2. Shanghai: Shanghai Scientific and Technological Press; 1977; pp 1356–1357.
- [3] Chen Z L, Wang B D, Chen M Q. Steroidal bitter principles from *Tacca plantaginea* structures of Taccalonolide A and B. Tetrahedron Letters 1987; 28: 1673–1676.
- [4] Zhou J, Chen C, Liu R, Yang C. Studies on the chemical constituents of *Tacca chantrieri* Andre. Zhiwu Xuebao 1983; 25: 568–573.
- [5] Yokosuka A, Mimaki Y, Sakagami H, Sashida Y. New diarylheptanoids and diarylheptanoid glucosides from the rhizomes of *Tacca chantrieri* and their cytotoxic activity. Journal of Natural Products 2002; 65: 283–289.
- [6] Harada N, Saito A, Ono H, Gawronski J, Gawronska K, Sugioka T, Uda H, Kuriki T. A CD method for determination of the absolute stereochemistry of acyclic glycols. 1. Application of the CD exciton chirality method to acyclic 1,3-dibenzoate systems. Journal of American Chemical Society 1991; 113: 3842–3850.
- [7] Tezuka Y, Gewali M B, Ali M S, Banskota A H, Kadota S. Eleven novel diarylheptanoids and two unusual diarylheptanoid derivatives from the seeds of *Alpinia blepharocalyx*. Journal of Natural Products 2001; 64: 208–213.
- [8] Ali M S, Tezuka Y, Awale S, Banskota A H, Kadota S. Six new diarylheptanoids from the seeds of *Alpinia blepharocalyx*. Journal of Natural Products 2001; 64: 289–293.
- [9] Ali M S, Tezuka Y, Banskota A H, Kadota S. Blepharocalyxins C-E, three new dimeric diarylheptanoids, and related compounds from the seeds of *Alpinia blepharocalyx*. Journal of Natural Products 2001; 64: 491–496.
- [10] Tokawa H, Aiyama R, Ikuta A. A pungent diarylheptanoid from *Alpinia oxyphylla*. Phytochemistry 1981; 20: 769–771.
- [11] Ohta S, Aoki T, Hirata T, Suga T. The structures of four diarylheptanoid glycosides from the female flowers of *Alnus serrulatoides*. Journal of the Chemical Society, Perkin Transactions 1 1984; 1635–1642.
- [12] Nagai M, Kenmochi N, Fujita M, Furukawa N, Inoue T. Studies on the constituents of Aceraceae plants. VI.: Revised stereochemistry of (-)-Centrololol, and new glycosides from *Acer nikoense*. Chemical and Pharmaceutical Bulletin 1986; 34: 1056–1060.
- [13] Yokosuka A, Mimaki Y, Sashida Y. Taccasterosides A–C, novel C₂₈-sterol glucosides from the rhizomes of *Tacca chantrieri*. Chemical and Pharmaceutical Bulletin 2004; 52: 1396–1398.

- [14] Yokosuka A, Mimaki Y, Sakuma C, Sashida Y. New glycosides of the campesterol derivative from the rhizomes of *Tacca chantrieri*. *Steroids* 2005; 70: 257–265.
- [15] Yokosuka A, Mimaki Y. New glycosides from the rhizomes of *Tacca chantrieri*. *Chemical and Pharmaceutical Bulletin* 2007; 55: 273–279.
- [16] Yasuhara F, Yamaguchi S, Kasai R, Tanaka O. Assignment of absolute configuration of 2-substituted-1-propanols by ¹H-NMR spectroscopy. *Tetrahedron Letters* 1986; 27: 4033–4039.
- [17] Kuroda M, Mimaki Y, Ori K, Koshino H, Nukada T, Sakagami H, Sashida Y. Lucilianosides A and B, two novel tetranor-lanostane hexaglycosides from the bulbs of *Chionodoxa luciliae*. *Tetrahedron* 2002; 58: 6735–6740.
- [18] Watanabe K, Mimaki Y, Sakuma C, Sashida Y. Eranthisaponins A and B, two new bisdesmosidic triterpene saponins from the tubers of *Eranthis cilicica*. *Journal of Natural Products* 2003; 66: 879–882.
- [19] Mimaki Y, Harada H, Sakuma C, Haraguchi M, Yui S, Kudo T, Yamazaki M, Sashida Y. Contortisiliosides A–G: isolation of seven new triterpene bisdesmosides from *Enterolobium contortisiliquum* and their cytotoxic activity. *Helvetica Chimica Acta* 2004; 87: 851–865.
- [20] Agrawal P K. NMR spectroscopy in the structural elucidation of oligosaccharides and glycosides. *Phytochemistry* 1992; 31: 3307–3330.
- [21] Yokosuka A, Mimaki Y, Sashida Y. Chantriolides A and B, two new withanolide glucosides from the rhizomes of *Tacca chantrieri*. *Journal of Natural Products* 2003; 66: 876–878.
- [22] Huang Y, Liu J K, Mühlbauer A, Henkel T, Huang Y, Liu J K, Mühlbauer A, Henkel T. Three novel Taccalonolides from the tropical plant *Tacca subflaellata*. *Helvetica Chimica Acta* 2002; 85: 2553–2558.
- [23] Khan P M, Malik A, Ahmad S, Nawaz H R. Withanolides from *Ajuga parviflora*. *Journal of Natural Products* 1999; 62: 1290–1292.
- [24] Yokosuka A, Mimaki Y, Sashida Y. Spirostanol saponins from the rhizomes of *Tacca chantrieri* and their cytotoxic activity. *Phytochemistry* 2002; 61: 73–78.
- [25] Yokosuka A, Mimaki Y, Sashida Y. Two new steroidal glycosides from *Tacca chantrieri*. *Natural Medicines* 2002; 56: 208–211.
- [26] Yokosuka A, Mimaki Y, Sashida Y. Steroidal and pregnane glycosides from the rhizomes of *Tacca chantrieri*. *Journal of Natural Products* 2002; 65: 1293–1298.
- [27] Agrawal P K, Jain D C, Gupta R K, Thakur R S. Carbon-13 NMR spectroscopy of steroidal saponins and steroidal saponins. *Phytochemistry* 1985; 24: 2479–2496.

- [28] Reddy K S, Shekhani M S, Berry D E, Lynn D G, Hecht S M. Afromontoside A new cytotoxic principle from *Dracaena afromontana*. Journal of the Chemical Society, Perkin Transactions 1 1984, 987-992.
- [29] Dong M, Feng X Z, Wang B X, Wu L J, Ikejima T. Two novel furostanol saponins from the rhizomes of *Dioscorea panthaica*. Prain et Burkill and their cytotoxic activity. Tetrahedron 2001; 57: 501-506.
- [30] Dong M, Feng X Z, Wu L J, Wang B X, Ikejima T. Two new steroidal saponins from the rhizomes of *Dioscorea panthaica* and their cytotoxic activity. Planta Medica 2001; 67: 853-857.
- [31] Tran Q L, Tezuka Y, Banskota A H, Tran Q K, Saiki I, Kadota S. New spirostanol steroids and steroidal saponins from roots and rhizomes of *Dracaena angustifolia* and their antiproliferative activity. Journal of Natural Products 2001; 64: 1127-1132.
- [32] Sargent J M, Taylor C G. Appraisal of the MTT assay as a rapid test of chemosensitivity in acute myeloid leukemia. British Journal of Cancer 1989; 60: 206-210.
- [33] Yamori T, Matsunaga A, Sato S, Yamazaki K, Komi A, Ishizu K, Mita I, Edatsugi H, Matsuba Y, Takezawa K, Nakanishi O, Kohno H, Nakajima Y, Komatsu H, Andoh T, Tsuruo T. Potent antitumor activity of MS-247, a novel DNA minor groove binder, evaluated by an in vitro and in vivo human cancer cell line panel. Cancer Research 1999; 59: 4042-4049.
- [34] Masuda T, Hidaka K, Shinohara A, Maekawa T, Takeda Y, Yamaguchi H. Chemical studies on antioxidant mechanism of curcuminoid: Analysis of Radical Reaction Products from Curcumin. Journal of Agricultural and Food Chemistry 1999; 47: 71-77.
- [35] Motterlinia R, Forestia R, Bassia R, Greena C J. Curcumin, an antioxidant and anti-inflammatory agent, induces heme oxygenase-1 and protects endothelial cells against oxidative stress. Free Radical Biology and Medicine 2000; 28: 1303-1312.
- [36] Chan M M Y, Huang H I, Fenton M R, Fong D. In vivo inhibition of nitric oxide synthase gene expression by curcumin, a cancer preventive natural product with anti-inflammatory properties. Biochemical Pharmacology 1988; 55: 1955-1962.
- [37] Suksamrarn A, Ponglikitmongkol M, Wongkrajang K, Chindaduang A, Kittidanairak S, Jankam A, Yingyongnarongkul B, Kittipanumat N, Chokchaisiri R, Khetkam P, Piyachaturawat P. Diarylheptanoids, new phytoestrogens from the rhizomes of *Curcuma comosa*: Isolation, chemical modification and estrogenic activity evaluation. Bioorganic and Medicinal Chemistry 2008; 16: 6891-6902.
- [38] Winuthayanon W, Piyachaturawat P, Suksamrarn A, Ponglikitmongkol M, Arao Y, Hewitt S C, Korach K S. Diarylheptanoid phytoestrogens isolated from the medicinal plant *Curcuma comosa*: Biologic actions in vitro and in vivo indicate estrogen receptor-dependent mechanisms. Environ Health Perspect 2009; 117: 1155-1161.

- [39] Adamsa B K, Ferstlb E M, Davisb M C, Heroldb M, Kurtkayab S, Camalierec R F, Hollingsheadc M G, Kaurc G, Sausvillec E A, Ricklesd F R, Snyderb J P, Liottab D C, Shojia M. Synthesis and biological evaluation of novel curcumin analogs as anti-cancer and anti-angiogenesis agents. *Bioorganic and Medicinal Chemistry* 2004; 12, 3871–3883.
- [40] Nakashima N, Kimura I, Kimura M, Matsuura H. Isolation of pseudoprototimosaponin AIII from rhizomes of *Anemarrhena asphodeloides* and its hypoglycemic activity in streptozotocin-induced diabetic mice. *Journal of Natural Products* 1993; 56: 345–350.
- [41] Choi S B, Park S. A steroidal glycoside from *Polygonatum odoratum* (Mill.) Druce. improves insulin resistance but does not alter insulin secretion in 90% pancreatectomized rats. *Bioscience, Biotechnology, and Biochemistry* 2002; 66: 2036–2043.
- [42] Wu R T, Chiang H C, Fu W C, Chien K Y, Chung Y M, Horng L Y. Formosanin-C, an immunomodulator with antitumor activity. *International Journal of Immunopharmacology* 1990; 12, 777–786.
- [43] Miyata T. Antitussive action of Mai-Men-Dong-Tang: Suppression of ACE inhibitor- and tachykinin-inducing dry cough. *Journal of Traditional Sino-Japanese medicine* 1992; 13: 276–279.
- [44] Ikeda T, Ando J, Miyazono A, Zhu X H, Tsumagari H, Nohara T, Yokomizo K, Uyeda M. Anti-herpes virus activity of Solanum steroidal glycosides. *Biological and Pharmaceutical Bulletin* 2000; 23, 363–364.
- [45] Niwa A, Takeda O, Ishimaru M, Nakamoto Y, Yamasaki K, Kohda H, Nishio H, Segawa T, Fujimaru K, Kuramoto A. Screening test for platelet aggregation inhibitor in natural products. The active principle of *Anemarrhena* Rhizoma. *Yakugaku Zasshi* 1988; 108: 555–561.

