

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Lactic Acid Bacteria in Philippine Traditional Fermented Foods

Charina Gracia B. Banaay, Marilen P. Balolong and Francisco B. Elegado

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/50582>

1. Introduction

The Philippine archipelago is home to a diverse array of ecosystems, organisms, peoples, and cultures. Filipino cuisine is no exception as distinct regional flavors stem from the unique food preparation techniques and culinary traditions of each region. Although Philippine indigenous foods are reminiscent of various foreign influences, local processes are adapted to indigenous ingredients and in accordance with local tastes. Pervasive throughout the numerous islands of the Philippines is the use of fermentation to enhance the organoleptic qualities as well as extend the shelf-life of food.

Traditional or indigenous fermented foods are part and parcel of Filipino culture since these are intimately entwined with the life of local people. The three main island-groups of the Philippines, namely – Luzon, Visayas, and Mindanao, each have their own fermented food products that cater to the local palate. Fermentation processes employed in the production of these indigenous fermented foods often rely entirely on natural microflora of the raw material and the surrounding environment; and procedures are handed down from one generation to the next as a village-art process. Because traditional food fermentation industries are commonly home-based and highly reliant on indigenous materials without the benefit of using commercial starter cultures, microbial assemblages are unique and highly variable per product and per region. Hence the possibility of discovering novel organisms, products, and interactions are likely.

Various microorganisms are involved in common food fermentation processes. In particular, lactic acid bacteria (LAB) in food is a type of biopreservation system. They not only contribute to the flavor of the food but LAB are also able to control pathogenic and spoilage microorganisms through various ways that include, but are not limited to, production of peroxidases, organic acids, and bacteriocins. Traditionally, identification of LAB in foods is largely dependent on culture-based methods; and properties of each isolate are evaluated

under controlled conditions. However, with the advent of molecular techniques, the enumeration of microorganisms missed by culture-dependent methods is now possible. Also, as more LAB metabolites, such as bacteriocins, are being reported, a wider database for identification and comparison with potential novel products are now available.

As the production and consumption of traditional fermented food products become increasingly relevant in the face of rapidly increasing population and food insecurity, more research and development to ensure the safety and nutritional quality of these fermented products is warranted. For a more extensive discussion of the principles and technology of Philippine fermented foods, the readers are directed to Sanchez (2008). This book is a detailed reference based on decades of research. Some data from the book will be presented again here in addition to other data from more recent studies. It is not the intention of this present paper to repeat what has been presented in the book, especially regarding fermentation processes, but only to present, as complete as possible, the data that are available regarding LAB present in indigenous/traditional fermented foods.

This paper aims to briefly review the various lactic acid-fermented indigenous fermented specialties in the different regions of the Philippines. Majority of the discussion will focus on recent data gathered from bacteriocin research and metagenomics studies of Philippine fermented specialties. Lastly, the health applications of the different fermented food products and their development as functional foods will be evaluated.

2. Regional fermented specialties in the Philippines

There are various lactic acid-fermented indigenous food products in the Philippines. Table 1 gives a summary of these different fermented specialties found in the different regions. Although a particular product type can be seen throughout the whole country, the texture, taste, and appearance would vary depending on the local taste, materials used, and process employed. For example, bagoong is a common fermented fish paste found all over the Philippines but the characteristic of the product found in Luzon is different from that found in the Visayas and Mindanao regions. Bagoong also takes on different names; there is bagoong na isda, bagoong alamang, bagoong na sisi, and guinamos (Sanchez, 2008). A product that is processed in a similar manner is dayok; it is made of brined fish entrails. Research indicates that this is also a lactic acid-fermented food but the LAB involved have not been identified yet (Besas and Dizon, 2012). Longanisa is sausage made of beef, pork, or chicken. It also takes on many forms depending on where it is made. The more famous ones are Vigan Longanisa in Northern Luzon, Pampanga Longanisa in Central Luzon, Lucban Longanisa in Southern Luzon, and Cebu Longanisa in the Visayas. The tastes vary from spicy, garlicky, sour, to sweet.

In lactic acid-fermented foods, LAB are important in preventing the growth of spoilage organisms, and altering flavor, aroma, and texture of the product. Although LAB are initially present in low numbers in the raw materials used, they soon proliferate as other organisms are inhibited by the initial addition of salt and as the continuous growth of LAB decreases the pH of the food making it less conducive for growth of other organisms. Recent

studies, however, have shown that there are a lot more benefits that can be derived from LAB in traditional fermented foods.

CATEGORY	PRODUCT NAME	REGION	MAJOR INGREDIENTS	LACTIC ACID BACTERIA INVOLVED (as determined from culture-based methods)	APPEARANCE AND/OR USAGE
Fermented vegetables, fruits	Burong mustasa	Luzon	Mustard leaves, cooked rice and/or rice washings	<i>Leuconostoc mesenteroides</i> , <i>Enterococcus faecalis</i> , <i>Lactobacillus plantarum</i>	Side dish
	Burong pipino	Whole Phil	cucumber	<i>Leu. mesenteroides</i> , <i>L. brevis</i> , <i>Pediococcus cerevisiae</i> , <i>L. plantarum</i>	Side dish
	Burong mangga	Whole Phil	Immature mango	<i>Leu. mesenteroides</i> , <i>L. brevis</i> , <i>P. cerevisiae</i> , <i>L. plantarum</i>	Side dish
	Atchara	Whole Phil	Immature papaya or chayote, or turnip (singkamas)	Unknown	Side dish
Cheese	Kesong puti	Luzon, Visayas	Cow or carabao milk	<i>Lactococcus lactis</i>	White soft cheese
Fermented fish and fishery products	Balao-balao	Luzon	Cooked rice, shrimp, salt	<i>Leu. mesenteroides</i> , <i>P. cerevisiae</i> , <i>L. plantarum</i>	Side dish, condiment
	Burong-isda	Luzon	Freshwater fish, rice, salt	<i>Leu. mesenteroides</i> , <i>E. faecalis</i> , <i>P. cerevisiae</i> , <i>L. plantarum</i> , <i>P. acidilactici</i> , <i>Leu. paramesenteroides</i>	Side dish, condiment
	Tinabal	Visayas	Parrot fish (for tinabal molmol) and frigate fish (for tinabal mangko), salt	<i>P. pentosaceus</i> , <i>S. equinus</i> , <i>Leuconostoc</i> sp., <i>Lactobacillus</i> sp.	Side dish, viand
	Burong talangka	Luzon	Small shore crabs (<i>Varuna litterata</i>)	<i>Leu. mesenteroides</i> , <i>E. faecalis</i> , <i>P. cerevisiae</i> , <i>L. plantarum</i>	Side dish, viand
	Patis	Whole Phil	Small fish, salt	<i>P. halophilus</i> (in mixed fermentation)	Fish sauce (patis), fish paste (bagoong), used as condiment, sauce, flavoring agent, viand
	Bagoong isda	Whole Phil	Small fish, salt		
	Bagoong alamang	Whole Phil	Small shrimps, salt		
	Bagoong na sisi	Visayas	Shell fish, salt		
	Guinamos	Bagoong isda in Visayas, Mindanao	Salt water small fish (dilis/belabid – <i>Stolephorus</i> sp.), salt		Condiment, viand, side dish
Dayok	Visayas, Mindanao	Fish entrails, salt	Unidentified LAB	Condiment, viand, side dish	
Fermented meat, sausages	Longanisa	Whole Phil	Ground pork, beef, or chicken meat, spices and preservatives	<i>P. acidilactici</i> , <i>Lactococcus lactis</i> (together with <i>Micrococcus aurantiacus</i>)	Viand
	Agos-os	Visayas	Sweet potato and ground pig's head	<i>E. faecalis</i>	Viand
	Burong kalabi	Luzon	Cooked rice, ground carabao meat	<i>L. plantarum</i>	Side dish, viand
	Burong babi	Luzon	Cooked rice, ground pork	<i>L. plantarum</i>	Side dish, viand

Fermented rice, cassava, sugar cane, coconut, soya	Puto	Whole Phil	Rice, sugar	<i>L. mesenteroides</i> , <i>E. faecalis</i> , <i>P. cerevisiae</i> (in mixed fermentation with <i>Saccharomyces cerevisiae</i>)	Steamed rice cake
	Bibingka	Whole Phil	Rice, sugar	<i>Saccharomyces cerevisiae</i>	Baked rice cake
	Tapuy	Luzon	Rice, glutinous rice	<i>Leuconostoc</i> , <i>L. plantarum</i> (in mixed fermentation with molds and yeasts)	Wine; beer
	Pangasi	Mindanao	Rice	Unknown	Wine
	Landang	Visayas, Mindanao	Cassava, or buli palm flour	Unknown	Dried jelly pellets, rice substitute
	Puto balanghoy	Mindanao	Cassava	Unknown	Steamed cake
	Basi	Luzon	Sugar cane	Unknown	Wine
	Suka	Whole Phil	Sugar cane juice (for sukang Iloco), palm inflorescence sap (for sukang tuba)	<i>Leuconostoc</i> , <i>Lactobacillus</i> , <i>Streptococcus</i> in the initial fermentation phase only	Vinegar, condiment, flavoring
	Sinamak	Luzon	Sugar cane juice, spices (chilies, onions, garlic)	Unknown	Spiced vinegar, condiment, flavoring
	Pinakurat	Visayas, Mindanao	Coconut sap, chilies, salt, various spices	Unknown	Spiced vinegar, condiment, flavoring
	Tuba	Whole Phil	Coconut sap	Unknown	Wine
	Lambanog	Whole Phil	Coconut sap	Unknown	Wine
Toyo	Whole Phil	Soybeans	<i>P. halophilus</i> , <i>E. faecalis</i> , <i>L. delbrueckii</i> (in mixed fermentation with <i>Aspergillus sojae</i> and <i>Saccharomyces rouxii</i>)	Condiment, flavoring agent, seasoning	

(Sources: Banaay et al., 2004; Besas and Dizon, 2012; Lee, 1999; Olympia et al., 1995; Sanchez, 2008; Tan et al., 2007)

Table 1. Regional Lactic Acid-Fermented Specialties in the Philippines

3. Research initiatives on LAB from Philippine fermented foods

3.1. Bacteriocin research

Bacteriocins are antimicrobial proteins or peptides produced by certain bacterial strains. Unlike the peptide antibiotics they usually have a narrow spectrum of antimicrobial activity, usually inhibiting growth of closely related bacterial species or strains and lacking lethality to the producer strain (Riley and Wertz, 2002).

The bacteriocins of LAB are small, cationic, hydrophobic, or amphiphilic peptides or small proteins, composed of 20 to 60 amino acid residues (Chen & Hoover, 2003). The bactericidal mode of action and biochemical properties depend on the protein moiety that could be specific to a particular LAB strain, *i.e.* the N-terminal amino acids as determinant of receptors in the cell wall of the susceptible strains/species and C-terminal amino acids for the biochemical properties. LAB bacteriocin must have the following desirable properties: “(1) not active and nontoxic to eukaryotic cells, (2) become inactivated by digestive proteases, having little influence on the gut microbiota, (3) low pH and heat-tolerant, (4)

have a relatively broad antimicrobial spectrum, against many food-borne pathogenic and spoilage bacteria, (5) show a bactericidal mode of action, usually acting on the bacterial cytoplasmic membrane: no cross resistance with antibiotics, and (6) have genetic determinants that are usually plasmid-encoded, facilitating genetic manipulation" (Apaga, 2012 as cited from Abriouel et al., 2007).

LAB bacteriocins have attracted attention in recent years because of their generally regarded as safe (GRAS) status and good value as natural biopreservatives which can find applications in the food and cosmetic industries (Cleveland et al., 2001; Daeschel, 1993; Riley and Wertz, 2002). Nisin, produced by strains of *Lactococcus lactis*, has been used in over 50 countries as anti-listerial and anti-clostridium substance. LAB bacteriocins with selective inhibition on food pathogens such as *Listeria monocytogenes*, but no inhibition on important lactic acid bacterial inocula such as the noted probiotic *Lactobacillus paracasei* or *Lactobacillus rhamnosus*; and yogurt-producing *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Lactococcus thermophilus*, may provide advantage over those that have a wider spectrum of antimicrobial activity and would kill these beneficial organisms, including nisin (De Vos, 1993; Jack and Ray, 1995; Nielsen et al., 1990). Hence, efforts on the search for LAB bacteriocins and elucidation of their properties are actively being pursued by several research laboratories. The future holds a wide array of LAB bacteriocins available for various specific applications.

3.2. Isolation and identification of bacteriocin-producing LAB

Some efforts on the isolation of bacteriocin-producing LAB had been started for more than a decade now in two major research institutions in the country namely: University of the Philippines Los Banos (specifically, the National Institutes of Molecular Biology and Biotechnology or BIOTECH-UPLB and the Institute of Biological Sciences or IBS-UPLB) and the Philippine Root Crop Research and Training Center, Visayas State University (VSU). These two institutions branched out knowledge on bacteriocin research through affiliate tutorship, as thesis advisers and as trainers to students and staff from a few other academic institutions which also did bacteriocin researches like University of Santo Tomas (UST), University of the Philippines Manila (UPM), De La Salle University (DLSU) and Ateneo de Manila University (ADMU). BIOTECH-UPLB and IBS-UPLB jointly worked on bacteriocins of *Lactobacillus plantarum* or plantaricins and those of *Pediococcus acidilactici* or pediocins. On the other hand, VSU devoted some efforts on the enterocins of *Enterococcus* spp. (Tan et al., 2001). DLSU also tried isolation of bacteriocin-producing LAB for food applications. UST was able to isolate bacteriocin-like inhibitory substances against medically important pathogens like *K. pneumoniae* (Dedeles et al., 2011). UPM and ADMU worked on human and animal health applications of bacteriocins.

Various fermented food products with proteinaceous components were the major sources of isolated LAB for bacteriocin screening. Such fermented food products are home-grown or produced by small enterprises and are still commercially available from

public markets in Luzon, Philippines and some parts of the Visayas like Leyte island. Examples of Philippine indigenous fermented foods that were good sources of bacteriocin-producing LAB are fermented rice and shrimp (*balao-balao*), fermented rice and fish mixture (*burong kanin at isda*), fermented pork (*burong babi*) in Central Luzon (Elegado et al., 2003; Gervasio and Lim, 2007) and fermented pork and sweet potato (*agos-os*) in Eastern Visayan region (Samar and Leyte). On the other hand, pickled vegetables like mustard leaf (*burong mustasa*) and green papaya (*achara*), fermenting fruits like pickled green mango, *bignay* or mango wine (Samnang 2010), fermented salted fish (*bagoong*), spicy sausages (*longganisa*) may contain some LAB but often times they are not bacteriocinogenic (Gervasio and Lim, 2007). The obvious reasons are the presence of inhibitory substances like salt, spices, alcohol or acid and of course the dearth of proteinaceous materials in the food material.

In one of the first isolation studies for bacteriocinogenic LAB, various proteinaceous fermented foods native to Central and Southern, Philippines were screened for bacteriocin-producing bacterial isolates. Seventy one out of several hundreds of colony-forming unit isolated by agar plate streaking were found antagonistic to the indicator microorganism, *Lactobacillus plantarum* ATCC 14917, through direct assay. By “spot-on-lawn” assay by pH-neutralized culture supernatant, nine (9) isolates were confirmed to be bacteriocin producers (Elegado et al., 2003). Banaay et al. in 2004 also reported on the isolation of 1,100 putative LAB from indigenous fermented foods in Luzon, Philippines. A strain of *Lactobacillus plantarum* was selected as the best bacteriocin producer. In another study, out of the 160 putative LAB obtained from 19 fermented food products from public markets in Central Luzon, 32 LAB isolates were found to be bacteriocinogenic (Gervasio and Lim, 2007). Santiago et al. (2008) were also able to find two LAB isolates, *Lactobacillus fermentum* LBA-19 and *Lactobacillus casei* LTI-21, screened from among several LAB isolates from various fermented food products from different regions in the Philippines.

Being pleomorphic, identification of LAB is quite challenging. A combination of various microbiological and molecular biology tools would help in finding the real identity. Banaay et al. (2004) did a thorough identification of the bacteriocinogenic LAB isolate using conventional morphological, biochemical and physiological methods, chemotaxonomic methods, as well as molecular methods. This is especially relevant to the identification of *Lactobacillus plantarum* which is a known pleomorphic bacteria. Most other Philippine LAB researchers often times directly apply 16S rRNA gene sequencing and homology search for LAB purified through repeated agar streaking and putatively identified as LAB just after determining its acid-forming, Gram positive and catalase negative properties. (Elegado et al., 2003; Gervasio and Lim, 2007; Santiago et al., 2008). Aside from 16S rRNA genes, other conserved genes were used for identification such as phenylalanyl-tRNA synthase (*pheS*) gene (Dedeles et al., 2011). Detection of bacteriocin genes through PCR may also be helpful in confirming the identity of the bacteriocinogenic LAB as well as the probability of producing the bacteriocin (Table 2).

ISOLATE/ STRAIN No.	IDENTIFICATION	(primer) HOMOLOGY to <i>P. acidilactici</i> type strain	REFERENCE	Bacteriocin gene by PCR; fingerprinting; HOMOLOGY
AA-5a	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R)98% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284 (27F) 99% <i>P. acidilactici</i> LAB 001; 99% <i>P. acidilactici</i> DSM20284	Elegado et al. 2003	ped ⁺ ; REP and RAPD
4E2	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 98% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284 (27F) 99% <i>P. acidilactici</i> LAB 001; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺
4E4	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 97% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284 (27F) 98% <i>P. acidilactici</i> 8D2CCH01MX; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺
4E5	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 99% <i>P. acidilactici</i> DSM20284 (27F) 99% <i>P. acidilactici</i> DSM20284	Laxamana et al. (2011)	ped ⁺ ; REP
4E6	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 98% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284; (27F) 99% <i>P. acidilactici</i> 8D2CCH01MX ; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺ ; [99% <i>P.</i> <i>acidilactici</i> bacteriocin genes ; pSMB74]
4E10	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 96% <i>P. acidilactici</i> UL5; 99% <i>P. lolii</i> to NGRI0510Q (27F) 99% <i>P. acidilactici</i> LAB 001 ; 99% <i>P. lolii</i> NGRI 0510Q	Apaga (2012)	ped ⁻
4BL7	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 98% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284 (27F) 99% <i>P. acidilactici</i> 8D2CCH01MX; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺
3G3	API CHL50 ID: <i>Lactobacillus</i> <i>pentosus</i> (doubtful) partial 16S rRNA gene ID: <i>P.</i> <i>acidilactici</i>	(1492R) 99% <i>P. acidilactici</i> IMAU20090 (27F) 98% <i>P. acidilactici</i> DSM20284	Elegado and Perez (2012)	ped ⁺ ; REP; ped ⁺
3G8	partial 16S rRNA gene ID: <i>P.</i> <i>acidilactici</i>	(1492R) 99% <i>P. acidilactici</i> UL5 (27F) 98% <i>P. acidilactici</i> DSM20284	Elegado and Perez (2012)	ped ⁺
3F3	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 95% <i>P. acidilactici</i> UL5 (27F) 98% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺
3F8	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 98% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284 (27F) 99% <i>P. acidilactici</i> LAB 001; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺
3F10	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 97% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284 (27F) 97% <i>P. acidilactici</i> LAB 001; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺ [99% <i>P.</i> <i>acidilactici</i> genomic scaffold];
IG7	partial 16S rRNA gene ID: <i>P.</i> <i>acidilactici</i>	(1492R) 97% <i>P. acidilactici</i> UL5 99% <i>P. acidilactici</i> 8D2CCH01MX; (27F) 98% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺ [100% pediocin operon;PSMB74];

K ₂ A ₂ -3	API: <i>Pediococcus pentosaceus</i> (good) partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 97% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284 (27F) 99% <i>P. acidilactici</i> LAB 001; 99% <i>P. acidilactici</i> DSM20284	Villarante (2011); Elegado and Perez (2012)	ped ⁺ ; plan ⁺ ped ⁺ ; REP
K ₂ A ₂ -1	API: <i>P. acidilactici</i> (doubtful)	-	Abuel (2007)	ped ⁺ ; plan ⁺ ped ⁺
K ₂ A ₂ -5	API: <i>P. acidilactici</i> (doubtful); partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 97% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284 (27F) 99% <i>P. acidilactici</i> LAB 001; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺ [99% <i>P. acidilactici</i> genomic scaffold]; plan ⁺
K ₂ A ₁ -1	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 99% <i>P. acidilactici</i> L94; 99% <i>P. acidilactici</i> DSM20284 (27F) 98% <i>P. acidilactici</i> JS-9-4; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺
K ₃ A ₂ -2	API: <i>Lactococcus lactis</i> (good) partial 16S rRNA gene ID: <i>P. pentosaceus</i>	-		ped ⁺ ; plan ⁺ ; ped ⁺
K ₃ A ₂ -3	partial 16S rRNA gene ID: <i>P. acidilactici</i>	100% <i>P. acidilactici</i> UL5	Elegado and Perez (2012)	ped ⁺
S3	partial 16S rRNA gene ID: <i>P. acidilactici</i>	(1492R) 98% <i>P. acidilactici</i> UL5; 99% <i>P. acidilactici</i> DSM20284 (27F) 97% <i>P. acidilactici</i> LAB 001; 99% <i>P. acidilactici</i> DSM20284	Apaga (2012)	ped ⁺ [99% pediocin operon; pSMB72];

Table 2. Identification and bacteriocin gene determination of putative *Pediococcus acidilactici* through 16S rRNA and pediocin gene PCR amplification and sequencing.

3.3. Purification and characterization of bacteriocins

Purification of bacteriocin peptides or small proteins into homogeneity is necessary in order to fully characterize them, particularly the determination of molecular mass, the primary structure or amino acid sequence and secondary structure. For pediocin, it was found that a simple and rapid method is effective for its purification. This method involves adsorption of pediocin onto the cell wall of the producer cell at pH 6 and 0.05 M NaCl and then subsequent desorption at pH 2.0 and 1 M NaCl (Elegado et al., 1997; Yang et al., 1992). This method seemed more applicable to pediocin but not with the lactococci, nisin or plantaricin. The reason is not clear but it could be related to variation in cell wall properties. The pH-adsorption/desorption method was able to provide materials for pH and temperature tolerance assays, estimation of molecular mass through SDS-PAGE, residual activity determination after protease, amylase and other enzyme actions (Laxamana et al., 2011). Enough amount of semi-purified bacteriocin from pediococci using this method was obtained for further purification through preparative reverse phase HPLC for various characterization studies, including the determination of secondary structures by circular dichroism and confirmation of double bonds through trypsin digestion and electrospray mass spectrometry (Elegado and Kwon, 1998). Other preparative purification methods prior to reverse phase HPLC and spectrometry included ion exchange chromatography and gel

filtration chromatography (Elegado et al., 2003), and hydrophobic interaction chromatography (Villarante et al., 2011). This method could also be applied with bacteriocins of pediococci and lactobacilli. The properties obtained from well characterized bacteriocinogenic LAB are shown in Table 3.

Isolate	Identity	Bacteriocin	Purification mode	Properties
AA5a	<i>Pediococcus acidilactici</i>	pediocin	pH adsorption/desorption Reversed-phase HPLC	Tolerant to pH 2-9 and 121 °C
BS25	<i>Lactobacillus plantarum</i>	plantaricin	Gel filtration chromatography Reversed-phase HPLC	MW = 3,830 Da
K2a2-3	<i>Pediococcus acidilactici</i>	pediocin	Hydrophobic interaction and ion-exchange chromatographies Reversed-phase HPLC	MW = 4,626 Da
K2a2-1	<i>Pediococcus acidilactici</i>	pediocin	pH adsorption/desorption	Optimum pH = 5-7 Resistant to boiling but not to autoclaving
4E5	<i>Pediococcus acidilactici</i>	pediocin	pH adsorption/desorption	Tolerant to pH 2-9; slight loss of activity at 100 °C; loss of activity at 121 °C; tolerates high salt; est. MW = 6,500 Da by SDS-PAGE

Table 3. List of purified and characterized bacteriocins from LAB isolated from Philippine indigenous fermented foods.

3.4. Optimization of bacteriocin production through fermentation kinetics

Bacteriocin production is largely dependent on the nutrients and nitrogen content of the fermentation medium. For instance, increased yeast extract concentration and polypeptone amount increases bacteriocin production. Molasses, raw sugar and sago hydrolyzates of amylase digestion were found to be good carbon sources. Other possible substrate base and supplements are cheese whey, coconut water and rice bran extract. Initial sugar concentration of usually 2 to 3% and inoculation rate of 3% by volume of at least 10^8 cells/mL provides good bacteriocin production (Elegado et al., 2001).

Bacteriocin production is highly dependent on cell or biomass growth. LAB are microaerophilic and most are either mesophilic or slightly thermophilic. The following conditions are applicable to their production: pH= 5.5 to 6.0; temperature = 35 – 40 °C; agitation = 50 rpm; without aeration. Usually, bacteriocin is optimally produced or secreted in the culture broth during the early stationary phase of growth. For *Pediococcus acidilactici*, culturing at 40 °C promotes earlier optimum bacteriocin production of around 10-12 hours. At 37 °C, bacteriocin production is from 14-16 hours (Sagpao et al., 2007).

3.5. Applications

Pediocins and plantaricins are the commonly found bacteriocins in Philippine fermented foods so far studied. Their antimicrobial properties have been investigated in several studies (Banaay et al., 2004; Elegado et al., 2003, 2004, 2007; Marilao et al., 2007). Although pediocins

and plantaricins show promise, their applications are limited at present because it is a well-known fact that other bacteriocins aside from nisin are not yet approved for food use. For pediocins and plantaricins, the most practical use for now would be dermatological and animal health care use. But since the bacteriocin-producing LAB are of GRAS status, those with probiotic properties such as tolerance to acidic pH (2.0 -3.0) and bile (0.3%) and adhesion properties to intestinal mucosa would be an advantage when used as adjunct inocula in fermented food products (Gervasio and Lim, 2007).

Perhaps another importance of bacteriocin-producing LAB is their effectiveness in biomedical applications. In one study, for example, partially-purified pediocin K2a2-3, through pH-mediated bacteriocin extraction method, was found cytotoxic against human colon adenocarcinoma (HT29) and human cervical carcinoma (HeLa) cells *in vitro* as determined by MTT [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] assay (Villarante et al., 2011). Other potential biomedical applications will be discussed in the succeeding section.

4. Probiotics and functional foods

An offshoot of the initial research on bacteriocins of LAB isolated from indigenous fermented foods is the emergence of probiotic research towards developing functional foods for biomedical applications. Probiotics refer to microorganisms that, when administered in adequate amounts, confers health benefits to the host. Although there are many microorganisms that can be considered as probiotics, LAB are the most common types because they produce antimicrobial compounds that inhibit other harmful microorganisms, they are able to tolerate acids and bile present in the digestive system, and they are able to adhere and establish themselves in the gut surfaces.

Many benefits have been ascribed to probiotics. For example, *Lactobacillus casei* (Shirota strain in Yakult®) have been shown effective in preventing diarrhea due to enterotoxigenic *Escherichia coli* (ETEC) and choleraenic vibrios (*V. cholerae* biotype E1 Tor and classical *V. cholerae*) using rats (Jacalne et al., 1990). This may be accounted for by its ability to kill the pathogens and inhibit further growth (Consignado et al., 1994). Because the probiotic used in the two studies mentioned is a commercial strain, current research on probiotics progressed to the search for indigenous LAB for use in the development of locally-produced functional food and investigation of their utility for biomedical applications. Metagenomic approaches to investigating LAB present in fermented foods have shown the diversity of potentially beneficial species present other than those that are readily detected by conventional culture-based methods. The development of functional food products shows potential in disease management. Research using metagenomic analysis in searching for microbial markers for use in functional foods to address certain lifestyle diseases as well as malnutrition is on the way.

4.1. Metagenomic and diversity studies

Traditional culture-based methods have been used for isolating LAB from fermented foods. These studies form the basis for the starter cultures used in food fermentation technologies employed for commercial production. Sanchez (2008) gives detailed information on the different

technologies and cultures used for the production of some traditional as well as developed technologies that have arisen from the culture-based studies conducted in earlier years.

In recent years culture-based approaches in LAB isolation have become more targeted for detection of bacteriocin-producers and those that have potential as probiotics. In one initiative, LAB isolates from fermented foods were screened for bacteriocin production and a PCR-based assay was used to detect specific bacteriocin-encoding genes. Acid and bile tolerance were also determined. Among all the isolates tested, *Lactobacillus fermentum* 4B1 and *Lactobacillus pentosus* 3G3 (later identified as *Pediococcus acidilactici*) have been identified as most promising for the development of new probiotic food products, hence they were chosen for subsequent biomedical application assays (Lim and Gervacio, 2007). In another study, LAB from traditionally fermented wine and vinegar from Visayas and Mindanao were isolated, identified, and tested for inhibitory activity against *Enterococcus faecium*, *Listeria innocua*, and *Staphylococcus aureus*. Five *Lactobacillus paracasei* and one *Lactobacillus brevis* showed antimicrobial properties against the tester strains (Licaros and Bautista, 2009).

With the advent of molecular techniques, the existence of non-culturable microorganisms has been acknowledged especially since the occurrence of culture-bias is already well-accepted. Culture-independent approaches, therefore, have been gaining popularity in microbial diversity studies and this includes researches on microorganisms found in fermented foods. The microbial populations in selected Philippine fermented foods were assessed through Polymerase Chain Reaction followed by Denaturing Gradient Gel Electrophoresis (PCR-DGGE) in two recent studies (Dalmacio et al., 2011; Larcia, 2010). Food samples tested include *burong mustasa* (fermented mustard), *alamang* (fermented shrimp paste), *burong isda* (fermented rice-fish mixture), *balao-balao/burong hipon* (fermented rice-shrimp mixture), *tuba* (sugar cane wine), and *sinamak* (spiced vinegar). Analysis of the 16S rRNA gene sequences revealed the presence of several LAB that have not been reported in these food products before. *Weissella cibaria*, *Lactobacillus plantarum*, *Lactobacillus pontis*, *Lactobacillus panis*, and *Lactobacillus fermentum* were detected in *burong mustasa* (Larcia, 2010). *L. panis* and *L. fermentum* were present in *alamang*; *L. pontis* and *L. plantarum* in *burong isda*; *L. panis*, *L. pontis*, and *L. fermentum* in *burong hipon*; and *W. cibaria*, *L. pontis*, *L. panis*, *L. fermentum* and *L. plantarum* in *burong mustasa* (Dalmacio et al., 2011).

The results of the two studies using molecular approaches in defining diversity of LAB in Philippine fermented foods show that culture-independent approaches are efficient tools for the analysis of microbial populations in fermented foods. Majority of the identified bacteria (LAB and other bacterial groups) have not been reported in culture-dependent studies. As such, the isolated bacterial 16S rRNA genes were cloned to have an initial partial 16S rRNA gene library for Philippine fermented foods (Dalmacio et al., 2011).

4.2. Biomedical applications

1. Anti-Obesity

Obesity is defined as an abnormal or excessive fat accumulation that presents risks to health. Probiotics can help in fighting obesity by reducing lipid absorption through its action on bile

acid metabolism, and by assimilation of cholesterol thus eliminating it from the host's system. Several studies were conducted to examine anti-obesity properties of different probiotic strains.

In one study, oral administration of *Lactobacillus paracasei* K3-4C, isolated from a locally fermented food had significant effect on lowering blood glucose levels (by 46%) and body weight (by 13%) in female BALB/c mice induced to be diabetic and obese through a 28-day high-fat diet (Parungao et al., 2006). In another study, orally administered *L. fermentum* 4B1 reduced adipose cell size, and decreased adipose tissue weight and overall body weight of mice fed with a high-fat diet for 49 days (Bautista et al., 2008). Likewise, oral administration of *P. acidilactici* 3G3 reduced body weight in diet-induced obese female Swiss mice (Parungao et al., 2009). In the last two studies described, the effects of the probiotics were determined to be comparable with the effects of the commercial anti-obesity drug Orlistat based on the parameters measured.

Recently, it has been postulated that the development of obesity may be caused by a shift in the composition of the gut microbiota towards the Firmicutes population (Ley et al., 2005). Firmicutes characterize obese versus lean/non-obese individuals together with a drop or no change in Bacteroidetes (Delzenne and Cani, 2010). Interestingly, Ley et al. (2006) found that a low fat diet had an effect to reverse the shift of Firmicutes/Bacteroidetes proportion. Because of this, dietary manipulation has been seen as a potential means of changing bacterial populations in the colonic microbiota and perhaps treating or at least preventing diseases like obesity. Although the root cause of obesity is excessive caloric intake coupled with a sedentary lifestyle (Blaut and Bischoff, 2010), Ley et al. (2005) proposed in their findings that alteration in the populations of mice gut microflora may have caused or may have been an effect of obesity. Because of this, current researches aim in using probiotics in the treatment of diseases such as obesity.

In two related studies (Arroyo and Fabriculana, 2011; Parungao et al., 2012), the effect of a functional food containing *P. acidilactici* 3G3 on microbial community changes in the gut of obese and non-obese mice was determined through PCR-DGGE. Results of these two preliminary studies showed that obese and non-obese mice had different baseline colonic microbiota. There were also indications that treatment with probiotics shifts the microbiota of obese mice towards the normal non-obese type. As these are preliminary studies, more research is warranted to elucidate the nature of the changes in gut microbiota and how it is related to obesity and the anti-obesity effects of probiotics.

2. Immuno-enhancement

A preliminary *in vitro* study to examine the immune-enhancing properties of viable and heat-killed preparations of two LAB previously isolated from traditional fermented foods (*L. fermentum* 4B1 and *P. acidilactici* 3G3) on murine peritoneal macrophage cells and splenic T-cells showed that isolate 4B1 was able to induce NO production in murine macrophages but, like 3G3, was unable to stimulate murine T-cell proliferation (Tan et al., 2008). Furthermore, this study showed that preparations of *L. fermentum* 4B1 have the ability to induce NO production in murine macrophage cells and its effects were more potent when it was alive.

The study also showed that isolate 4B1 exhibited better immune-enhancing effect than the probiotic species found in a commercial probiotic drink. T-cell proliferation, however, was not observed in any of the treatments in this study and was attributed to the delayed stimulation in cells responding to a first-time exposure to the different probiotic strain preparations used.

3. Reduction of blood glucose levels

A study by Ngo et al. (2008) showed that oral administration of kefir, a common fermented food consumed by the elderly, significantly decreased blood glucose levels and body weight of diabetic obese male Sprague Dawley rats. The results of the study showed lower blood glucose levels (from 198.5 to 105.6 mg/dL) and clinically lower body weights (from 342.9 to 311.5 g) of the treated diabetic-obese rats than the untreated diabetic-obese control group.

4. Prevention of hypercholesterolemia

The effect of *P. acidilactici* 3G3 administration on hypercholesterolemic Swiss Albino mice was determined (Parungao et al., 2009). This strain was able to assimilate cholesterol in the *in vitro* plate assay and decrease HDL, LDL, and total cholesterol in the *in vivo* assay using mice. Strain 3G3 was also shown to adhere well to the duodenum and middle colon. Results suggest the potential of *P. acidilactici* 3G3 in preventing hypercholesterolemia.

4.3. Development of functional foods

The development of functional foods containing known probiotic strains stems from earlier researches on bacteriocins and isolation of potential probiotics from traditional fermented foods. The beneficial effects of probiotic-supplemented chocolate bars (Arroyo et al., 2010; Arroyo and Fabiculana, 2011), fermented mustard leaves (Calapardo et al., 2006), and coffee wine (Parungao, 2007) have been investigated. Initial studies on mango-milk and carrot juice drinks supplemented with probiotic strains have also been conducted (Bugarin et al., 2010; Elegado et al., 2005). These potential functional foods contain probiotic strains, previously isolated from traditional fermented foods such as *P. acidilactici* AA5a (Elegado et al., 2003), *L. plantarum* BS25 (Banaay et al., 2004), and *P. acidilactici* 3G3 (Lim and Gervacio, 2007). Research on functional foods is still in its infancy but this food category shows promise in disease management as well as in contributing to food security in the country. Commercial interest in probiotic food products is increasing due to the growing understanding of its health benefits. This growing industry can derive benefits from the researches conducted on this emerging food category.

5. Future perspectives

Aside from the research works presented earlier in this paper as well as on-going follow-up studies related to them, future goals may include research on a variety of other possible biomedical applications of LAB with potential probiotic properties. The effect of probiotics on *Helicobacter pylori* infections (that may cause peptic ulcers) may be determined. Their ability to modulate inflammatory and hypersensitivity responses as well as their effect on

irritable bowel syndrome and colitis may be investigated. Further research on possible anti-cancer properties of probiotics is warranted as follow-up studies on the work done by Villarante et al. (2011). These studies are very important as these have the potential to address some of the more serious health concerns of our society.

Much is still to be learned about the existing probiotic strains. The molecular biology and genomics of these isolates may be pursued in order to further elucidate their properties and mechanisms of action.

Determination of factors affecting probiotic viability in foods is also important as these will determine if their survival in the food, and therefore their delivery into the host, is maintained. This will constitute a quality control for functional foods.

The potential physiological effects of multiple prebiotic strains, as opposed to a single strain, are also interesting areas of research. The delivery of multiple probiotic strains may help ensure its effectiveness in an environment that contains high diversity of resident microflora. The potential benefits of synbiotics, (combination of probiotic and prebiotic) which have synergistic interaction, may also be investigated. A good combination will greatly enhance the health benefits to humans.

Author details

Charina Gracia B. Banaay

*Institute of Biological Sciences, College of Arts and Sciences,
University of the Philippines Los Baños, Laguna, Philippines*

Marilen P. Balolong

*Department of Biology, University of the Philippines Manila,
Padre Faura Street, Ermita, Manila, Philippines*

Institute of Molecular Biology and Biotechnology, National Institutes of Health,

University of the Philippines Manila, Pedro Gil Street, Ermita, Manila, Philippines

Francisco B. Elegado

*National Institute of Molecular Biology and Biotechnology,
University of the Philippines Los Baños, Laguna, Philippines*

6. References

- Abriouel H, Galvez A, Lopez RL, Omar NB (2007) Bacteriocin-based strategies for food biopreservation. *International Journal of Food Microbiology* 120: 51–70.
- Apaga DLT (2012) Detection of bacteriocin structural genes and bacteriocinogenic activity of several pediococcus isolates. Undergraduate thesis. University of the Philippines Manila. 56 p.
- Arroyo KZO, Garcia JAR, Elegado FB, Calapardo M, Parungao MM (2010) Survival of *Lactobacillus pentosus* 3G3 in home-made chocolate bars. 39th Annual Convention and Scientific Meeting,, PSM, Inc. April 29-30, 2010, City of Naga, Camarines Sur.

- Arroyo MJJ, Fabiculana PRS (2011) Gut microflora assessment of obese and non-obese mice (*Mus musculus* L.) orally-fed with Ip-3g3 chocolate. Department of Biology, University of the Philippines Manila, Manila, Philippines. Unpublished Undergraduate Thesis.
- Banaay CGB, Elegado FB, Dalmacio IF (2004) Identification and characterization of bacteriocinogenic *Lactobacillus plantarum* BS25 isolated from balao-balao, a locally fermented rice-shrimp mixture from the Philippines. *The Philippine Agricultural Scientist* 87(4): 427-438.
- Bautista RLS, Ecarma NCA, Balolong ECJr, Hallare AV, Parungao MM (2008) The Effects of Orally-Administered *Lactobacillus* sp. 4B1 on the Adipose Tissues of Diet-Induced Obese Mice (*Mus musculus* L.). International Symposium on Probiotic from Asian Traditional Fermented Foods for Healthy Gut Function. August 19-20, 2008. Sari-Pan Pacific Hotel, Jakarta, Indonesia.
- Besas JR, Dizon EI (2012) Influence of salt concentration on histamine formation in fermented tuna viscera (Dayok). *Food and Nutrition Sciences* 3:201-206.
- Blaut M and Bischoff SC (2010) Probiotics and Obesity. *Ann Nutr Metab* 2010; 57(Suppl.1):20-23.
- Bugarin MA, Sison AAD, Elegado FB, Calapardo M, Parungao MM (2010) Survival of *Lactobacillus plantarum* BS25 on Varying Ratio of Mango-Milk Substrates and Storage Temperature. 39th Annual Convention and Scientific Meeting, PSM, Inc. April 29-30, 2010, City of Naga, Camarines Sur.
- Calapardo MR, Bueno MOV, Guillermo MKB, Saguibo JD, Parungao MM, Elegado FB (2006) Bile and Acid Tolerance of Bacteriocinogenic *Lactobacillus plantarum* and its Use as Adjunct Inoculum in Pickled Mustard Leaves. Proceedings of the 5th Asia-Pacific Biotechnology Congress and 35th Annual Convention of the Philippine Society for Microbiology Inc. (PSM). "Microbiology and Biotechnology: Roadmaps and Milestones for Enhancing Sustainable Productivity in the Asia-Pacific Region." Bohol Tropics Resort. Tagbilaran City, Bohol, Philippines.
- Chen H, Hoover DG (2003) Bacteriocins and their food applications. <http://www.ift.org/publications/crfsfs> (accessed 22 July 2006)
- Cleveland J, Chikindas ML, Montville TJ, Nes IF (2001) Bacteriocins: safe, natural antimicrobials for food preservation. *International Journal of Food Microbiology* 71: 1-20.
- Consignado GO, Pena AC, Jacalne AV (1994) *In vitro* study on the bacterial activity of *Lactobacillus casei* (commercial Yakult drink) against four diarrhea-causing organisms: enterotoxigenic *E. coli*, *Salmonella enteritidis*, *Shigella dysenteriae*, *Vibrio cholerae*. *Philipp J Microbiol Infect Dis* . 23(2): 50-55.
- Daeschel MA (1993) Applications and interactions of bacteriocins of lactic acid bacteria in food and beverages. In: Hoover, D.G., L.R. Steenson, eds. *Bacteriocins of Lactic Acid Bacteria*. New York: Academic Press Inc. P. 63-70.
- Dalmacio LMM, Angeles AKJ, Larcia LLH, Parungao-Balolong MM, Estacio RC (2011) Assessment of microbial diversity in Philippine fermented food products through polymerase chain reaction- denaturing gradient gel electrophoresis (PCR-DGGE). *Beneficial Microbes*, 2(4): 273-281.

- De Vos WM (1993) Future prospects for research and applications of nisin and other bacteriocins. In: Hoover, D.G., L.R. Steenson, eds. *Bacteriocins of Lactic Acid Bacteria*. New York: Academic Press Inc. P. 249-258.
- Dedeles GR, Caranza MAE, Elegado FB (2011) Bacteriocin-like inhibitory substances (BLIS) from lactic acid bacteria and their potential as biopreservatives in foods. Proceedings of the 6th Asian Conference on Lactic Acid Bacteria and XIII International Congress of Bacteriology and Applied Microbiology. September 6-10, 2011. Sapporo, Japan. 120 p.
- Delzenne NM and Cani PD (2010) Nutritional modulation of gut microbiota in the context of obesity and insulin resistance: potential interest of prebiotics. *Int. Dairy J.* 20: 277-280.
- Elegado FB, Kim WJ, Kwon DY (1997) Rapid purification, partial characterization and antimicrobial spectrum of the bacteriocin, Pediocin ACM from *Pediococcus acidilactici* M. *International Journal of Food Microbiology* 37(1): 1-11.
- Elegado FB, Kwon DY (1998) Primary structure and conformational studies of Pediocin AcM, a bacteriocin from *Pediococcus acidilactici* M. *The Philippine Journal of Biotechnology* 9(1): 19-26.
- Elegado FB, Sonomoto K, Ishizaki A (2001) Molecular characterization of a bacteriocin of *Pediococcus acidilactici* and its production in sago and sugarcane-based substrate. *Biotechnology for Sustainable Utilization of Biological Resources in the Tropics*. Murooka, Y., T. Yoshida, T. Seki, P. matangkasombut, T.M. Espino, U. Soetisna and M.I.A. Karim. eds. JSPS-NRCT/DOST/LIPI/VCC Joint Seminar. Nov. 7-9, 2001. Bangkok, Thailand. 15:173-180.
- Elegado FB, Opina ACL, Banaay CGB, Dalmacio IF (2003) Purification and characterization of novel bacteriocins from lactic acid bacteria isolated from Philippine fermented rice-shrimp or rice-fish mixtures. *The Philippine Agricultural Scientist* 86(1): 65-74.
- Elegado FB, Guerra MARV, Macayan RA, Estolas MT, Lirazan MB (2004) Antimicrobial activity and DNA fingerprinting of bacteriocinogenic *Pediococcus acidilactici* through RAPD-PCR. *The Philippine Agricultural Scientist* 87(2):229-237.
- Elegado FB, Calapardo MR, Fabregas JF, Ona SEN, Parungao MM (2005) Antilisterial Efficacy of Bacteriocinogenic *Pediococcus acidilactici* AA5a in Carrot Juice Fermentation. Proceedings of the 3rd Asian Conference on Lactic Acid bacteria. August 25-27, 2005. Sanur Paradise Hotel, Bali, Indonesia.
- Elegado FB, Abuel BJA, Te JT Jr., Calapardo MR, Parungao MM (2007) Antagonism against *Listeria* spp. and *Staphylococcus aureus* by bacteriocin-producing lactic acid bacteria screened from the intestine of Philippine carabao using polymerase chain reaction. *The Philippine Agricultural Scientist* 90(4): 305-314.
- Elegado FB and Perez MTM (2011) Genetic identification of lactic acid bacteria and bacteriocin structural gene elucidation. National Academy of Science and Technology (NAST) Scientific Meeting. July 11-12, 2012. Manila Hotel, Manila, Philippines.
- Gervasio ATR, Lim VMT (2007) Probiotic characterization of bacteriocinogenic lactic acid bacteria isolated from fermented foods of selected areas. Undergraduate thesis. University of the Philippines Manila. 105 p.
- Jack RJT, Ray B (1995) Bacteriocin of gram positive bacteria. *Microbiological Reviews.* 59(2): 171-200.

- Jacalne AV, Jacalne RR, Hirano H, Suetomi T, Villahermosa CG, Castaneda I (1990) *In-vivo* studies on the use of *Lactobacillus casei* (Yakult Strain) as biological agent for the prevention and control of diarrhea. *Acta Med Philipp.* 26(2): 116-122
- Larcia, LLH (2010) A study of the bacterial profile of Philippine fermented mustard (*burong mustasa*) through culture-independent methods / Levi Letlet Larcia, II. Thesis (MS Biochemistry)--University of the Philippines Manila. 158pp. LG995 2010 B3 L37
- Laxamana FLM, Carillo MCO, Elegado FB (2011) Characterization of a bacteriocin of lactic acid bacteria isolated from fermented rice-fish mixture. Proceedings of the 40th Annual Convention and Scientific Meeting of the Philippine Society for Microbiology, Inc. May 10-14, 2011. Manila, Philippines.
- Lee C (1999) Cereal fermentations in countries of the Asia-Pacific Region. In: FAO, Fermented Cereals: A Global Perspective, FAO Agricultural Services Bulletin Issue 138. Food and Agriculture Organization 144 p.
- Ley RE, Bäckhed F, Turnbaugh P, Lozupone C A, Knight RD, Gordon JI (2005) Obesity alters gut microbial ecology. *Proc Natl Acad Sci USA*, 102(31): 11070-11075.
- Ley RE, Turnbaugh PJ, Klein S, Gordon JI (2006) "Human gut microbes associated with obesity." *Nature* 444:1022-1023.
- Licaros A, Bautista A (2009) Culture-Dependent and Culture-Independent Analysis of Vinegars from Visayas and Mindanao. Department of Biology, University of the Philippines Manila, Manila, Philippines. Unpublished Undergraduate Thesis.
- Lim V, Gervacio TA (2007) Bacteriocin-producing LAB from fermented foods of Central Luzon. Department of Biology, University of the Philippines Manila, Manila, Philippines. Unpublished Undergraduate Thesis.
- Marilao CG, Calapardo MR, Ciron CE, Elegado FB (2007) Antilisterial action of *Pediococcus acidilactici* AA5a in pork sausage fermentation. *The Philippine Agricultural Scientist* 90(1): 40-45.
- Ngo RE, Estrada MZ, Balolong EC Jr., Parungao MM (2008) Orally-administered kefir on lowering blood glucose levels and body weight in diet-induced diabetic obese rats. Project Terminal Report. National Institutes of Health University of the Philippines Manila.
- Nielsen JW, Dickson JS, Crouse JD (1990) Use of a bacteriocin produced by *Pediococcus acidilactici* to inhibit *Listeria monocytogenes* associated with fresh meat. *Applied and Environmental Microbiology* 54: 2349-2353.
- Olympia M, Fukuda H, Ono H, Kaneko Y, Takano M (1995) Characterization of starch-hydrolyzing lactic acid bacteria isolated from a fermented fish and rice food, "Burong Isda", and its amyolytic enzyme. *Journal of Fermentation and Bioengineering* 80(2):124-130.
- Parungao MM (2007) Probiotic Properties of Philippine Coffee Wine. The 4th Asian Conference on Lactic Acid Bacteria & the 3rd International Symposium on Lactic Acid Bacteria and Health. October 17-19, 2007. Shanghai, China.
- Parungao MM, Castillo RL, Mortel MRA (2006) Effects of Oral Administration of *Lactobacillus paracasei* K3-4C on Blood Glucose Levels and Body Weight in Diabetic

- Obese Mice (*Mus musculus* L.). December 15-17, 2006. Proceedings of the 11th Biological Sciences Graduate Congress, Chulalongkorn University, Bangkok, Thailand.
- Parungao MM, Ong C, Laluces N, De Torres K, Usisa J, Calapardo M, Trinidad L, Elegado FB (2009) The Cholesterol-Reducing Ability and Adhesion Properties of *Lactobacillus pentosus* 3G3. Proceedings of the 5th Conference of Federation for Societies of Lactic Acid Bacteria. July 1-4, 2009. National University of Singapore, Singapore.
- Parungao-Balolong M, Libed AAO, Loma K, Villena JPDS, Villafuerte AR, Balolong EC Jr., Dalmacio LMM (2012) Influence of probiotic drinks on the distal gut bacterial flora of mice (*Mus musculus* L.) fed with standard diet or high-fat diet. Terminal report, National Institutes of Health University of the Philippines Manila.
- Riley MA, Wertz JE (2002) Bacteriocins: evolution, ecology and applications. *Annual Review of Microbiology* 56:117-137.
- Sagpao SMN, Elegado FB, Zamora A (2007) Optimization of production and partial purification of pediocin from *Pediococcus acidilactici* PNCM 10289. Poster paper presented at the 29th National Academy of science and Technology (NAST) Philippines Annual Scientific Meeting, Manila Hotel, July 11-12, 2007.
- Samnang (2010) Isolation and identification of lactic acid bacteria from fermenting bignay (*Antidesma bunius* (L.) Spreng) and mango (*Mangifera indica* L.) wines. M.S. thesis. University of the Philippines Los Banos. 242 p.
- Sanchez PC (2008) Philippine fermented foods: principles and technology. The University of the Philippines Press. Diliman, Quezon City, Philippines. 516 p.
- Santiago MR, Lopez CM, Tenorio EL, de Guzman EE (2007) Detection and partial characterization of two bacteriocins produced by *Lactobacillus casei* and *Lactobacillus fermentum* isolated from fermented rice-fish mixtures indigenous in the Philippines. Proceedings of the 37th Annual Convention and Scientific Meeting of the Philippine Society for Microbiology, Inc. May 7-9, 2008. Boracay Philippines. P. 32.
- Tan JD, Galvez FCF, Asano K, Tomita F (2001) Isolation and partial purification of bacteriocin produced by microorganisms from *agos-os*. Biotechnology for Sustainable Utilization of Biological Resources in the Tropics. Murooka, Y., T. Yoshida, T. Seki, P. matangkasombut, T.M. Espino, U. Soetisna and M.I.A. Karim. eds. JSPS-NRCT/DOST/LIPI/VCC Joint Seminar. Nov. 7-9, 2001. Bangkok, Thailand. 15: 104-110.
- Tan JAS, Yalung PM, Evangelista KV, Parungao MM (2008) *In vitro* Study on the Effect of *Lactobacillus* sp. 4B1 and *Lactobacillus* sp. 3G3 on Murine Macrophage Nitric Oxide Production and Splenic T-cell Proliferation. Proceedings of the International Symposium on Probiotic from Asian Traditional Fermented Foods for Healthy Gut Function. August 19-20, 2008. Sari-Pan Pacific Hotel, Jakarta, Indonesia.
- Villarante KI, Elegado FB, Iwatani S, Zendo T, Sonomoto K, de Guzman EE (2010) Purification, characterization and *in vitro* cytotoxicity of the bacteriocin from *Pediococcus acidilactici* K2a2-3 against human colon adenocarcinoma (HT29) and human cervical carcinoma (HeLa) cells. *World Journal of Microbiology and Biotechnology* 27: 975-980.
- Yang R, Johnson MC, Ray B (1992) Novel method to extract large amounts of bacteriocins from lactic acid bacteria. *Appl. Environ. Microbiol.* 58: 3355-3359.