

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Cellular Immunotherapy Using Dendritic Cells in Multiple Myeloma: New Concept to Enhance Efficacy

Je-Jung Lee, Youn-Kyung Lee, Hyun Ju Lee,
Sung-Hoon Jung and Thanh-Nhan Nguyen-Pham

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/54100>

1. Introduction

Multiple myeloma (MM) is a clonal B-cell malignancy that is currently incurable with conventional chemotherapy, even if high-dose chemotherapy with autologous or allogeneic hematopoietic stem cell transplantation (HSCT) and the development of novel molecular target agents have resulted in a marked improvement in overall survival [1, 2]. Allogeneic HSCT, which induces a clinically significant immune-mediated allogeneic graft-versus-myeloma (GVM) effect, has provided the framework for the development of immunotherapeutic strategies [3, 4]. To prolong the survival of patients with MM, who are undergoing allogeneic HSCT, a donor lymphocyte infusion can be used successfully as a salvage therapy, which is based on the GVM effect in some cases of MM that relapse after allogeneic HSCT [5, 6]. A clinically significant immune-mediated GVM effect provides the framework for the development of immune-based therapeutic options that use antigen-presenting cells (APCs) with increased potency, such as dendritic cells (DCs), in MM [6].

DCs are the most potent APCs for initiating cellular immune responses through the stimulation of naive T cells. Because of their ability to stimulate T cells, DCs act as links between innate immunity and adaptive immunity in antitumor immune responses [7]. DCs orchestrate a variety of immune responses by stimulating the differentiation of naïve CD4⁺ T cells into helper T effectors such as Th1, Th2 or Th17 type [8, 9]. Cytokines secreted by DCs at the time of initial T cell stimulation play an important role in the subsequent differentiation of effector T cells. Th1 cells, through interferon-gamma (IFN- γ) production, regulate antigen presentation and immunity against intracellular pathogens [8]. DC-based vaccines have become the most attractive tools for cancer immunotherapy and have been used in more than 20 malignancies; most commonly melanoma, renal cell carcinoma, prostate cancer and colorectal carcinoma

[10]. Cellular immunotherapy using DCs is emerging as a useful immunotherapeutic modality to treat MM [11]. While antigen-specific cytotoxic T lymphocytes (CTLs) and immune response can be induced by DC vaccination in MM patients, clinical responses so far have been largely unsatisfying to be observed only in a minority of treated patients with MM. Progress in understanding DC biology in cancer patients and the recruitment of suppressive cells of the adaptive and innate immune system in antitumor immunity of cellular immunotherapy is leading to new concept which aims at improved immune and clinical outcomes in MM. New concept is developing to generate novel therapeutic targets that could restore DC capacity to prime T cells and trigger effective anticancer responses in combination with other therapies to offset tumor-induced suppression in MM.

2. Dendritic cell in myeloma immunity

DCs have a potent antigen-specific T cell stimulatory capacity and therefore should be considered to the one of the promising antitumor immunotherapeutic options. In tumor-specific immunity, secreted products or fragments from tumor cells enter into DCs through the endosome and are processed and presented on MHC class molecules of DCs [12]. Processed antigens presented on these molecules of DCs are recognized by CD4⁺ T helper cells, which not only enhance to the CD8⁺ T cell response but also facilitate to develop a humoral immune response for surface antigens expressed on the tumor cells. The antigens presented on MHC class I are recognized by CD8⁺ CTLs, which have a direct cytotoxic effect on tumor cells. Unfortunately, patients with MM have basically dysfunctional DCs that are functionally defective, evidenced by the decreased number of circulating precursors of DCs as well as the impaired T cell stimulatory capacities compared with normal controls [13, 14]. The defective functions of DCs in patients with MM are partially attributed to the production of IL-6 and other tumor-derived factors. DCs in MM patients are a target of tumor-associated suppressive factors, such as IL-10, transforming growth factor- beta (TGF- β), vascular endothelial growth factor (VEGF), and IL-6, resulting in their aberrant functions and impaired development of effector functions in tumor-specific lymphocytes [15]. There were only few patients with MM who responded clinically to vaccination with antigen-loaded autologous DCs. There may be several reasons for this failure from MM patients itself. MM is believed to induce immunoparesis that interferes with DC function and hence affects the effective antitumor immune responses in these patients. They are able to escape immune surveillance by down-regulation of immune markers as well as through the production of immunosuppressive cytokines by the tumor cells or by activation of suppressor cells such as regulatory T cells and myeloid cells. Myeloma cells can produce immuno-inhibitory cytokines, such as IL-10, TGF- β , VEGF, and IL-6, which play major roles in the pathogenesis of MM [15]. In addition, the survival and proliferation of myeloma cells are partially facilitated by impaired endogenous immune surveillance against tumor antigens, including the abrogation of DC function, by constitutive activation of the signal transducer and activator of transcription 3 (STAT3) [13]. Impairment in both humoral and cellular immunity in MM is associated with impaired B cell responses; decreased T cell numbers including CD4⁺ T cells and impaired CTL responses; and dysfunction

of natural killer (NK) cells and NKT cells responses [16-19]. In addition, the recruitment and expansion of CD4⁺CD25⁺ regulatory T cells (Tregs) in the suppression of tumor immunity has been reported in MM patients [20, 21]. More recently, the proportion of CD14⁺HLA-DR^{-low} myeloid-derived suppressor cells (MDSCs) and CD4⁺ forkhead box P3 (FoxP3)⁺ Tregs cells was increased in MM patients at diagnosis, resulting in a significant impediment of immune cells related to cancer immunotherapy [22].

3. Current DC vaccination research in MM

Usually, *ex vivo* DCs were generated from circulating blood precursors (i.e. monocytes) or bone marrow progenitor cells and educated them with myeloma-associated antigens prior to vaccination to patients with MM.

3.1. Idiotype-pulsed DCs

Immunoglobulin idiotype (Id) is a tumor-specific antigen that is produced by each B cell tumor clone. Id protein has been used for immunotherapy in patients with MM [23, 24]. Id vaccination could induce immune responses by both antibodies and Id-specific T cells, including CD4⁺ and CD8⁺ T cells, through the presentation of Id protein on the surface of professional APCs [24]. Id-specific CTL lines that kill autologous primary myeloma cells *in vitro* have been generated [25, 26]. Autologous DCs that were generated from MM patients have been shown to efficiently endocytose different classes of Id proteins, and autologous Id-specific CTLs that were generated by Id-pulsed DCs were able to recognize and kill autologous primary myeloma cells *in vitro* [25, 26]. Various studies of DC-based Id vaccination in MM have been reported [27-34]. Although Id-specific CTLs and immune responses could be induced in some patients, clinical responses have rarely been observed after vaccination possibly because Id protein is a weak antigen and immature DCs have been used in some studies [27].

3.2. Myeloma-associated antigens-based DC immunotherapy

In general, the production of DC vaccines using whole tumor antigens has become a promising tool for immunotherapy against MM. There are several types of myeloma-associated antigen for loading onto DCs: loading with myeloma lysates [35, 36], loading with dying myeloma cells [37-39], transfection with myeloma-derived RNA [40], pulsing with myeloma-derived heat shock protein (HSP) gp96 [41, 42], and hybridization with myeloma cells [43, 44]. These techniques have the advantage of allowing the presentation of multiple epitopes to MHC on DCs, therefore inducing polyclonal T cell responses from many potentially unknown tumor-associated antigens (TAAs) and reducing the probability of immune escape by a single TAA.

Various myeloma-associated antigens that may induce immune responses from DC-based vaccines have been identified in MM patients. MUC1-specific CTLs that were induced *in vitro* using peptide-pulsed DCs or plasma cell RNA-loaded DCs efficiently killed not only target cells pulsed with the antigenic peptide but also MM cells [40, 45]. DCs transfected with PTD-NY-ESO-1 protein can induce CD8⁺ cellular antitumor immunity superior to that

achieved with NY-ESO-1 protein alone [46]. Sp17-specific HLA class I-restricted CTLs were successfully generated by DCs that had been loaded with recombinant Sp17 protein and were able to kill autologous tumor cells that expressed Sp17 [47]. The overexpression of hTERT on MM compared to the expression levels in normal cells indicated that this telomerase also could be used as a myeloma-associated antigen. hTERT was capable of triggering antitumor-CTL responses and killing hTERT⁺ tumor cells [48]. Recently, a report demonstrated that activated T lymphocytes were able to successfully kill myeloma cells after stimulation by DCs loaded with hTERT- and MUC1-derived nonapeptides [49]. DKK1, a novel protein that is not expressed in most normal tissues but is expressed in almost all myeloma cells, may be an important antigenic target for anti-myeloma immunotherapy. DKK1-specific CTLs that were generated by DCs pulsed with DKK1 peptides were specifically lysed by autologous primary myeloma cells and DKK1-positive cell lines [50].

4. New concepts to enhance the efficacy of cellular immunotherapy in MM

4.1. How to enhance the efficacy of DC vaccinations

Because of unsatisfied clinical response of DC vaccination trials in MM, a number of groups have looked at whether the DC vaccination may be more effective if better cytokine combinations are used to enhance DC function, effective tumor antigens are investigated to use, suppressive signal transcriptions are blocked to overcome defective DC function, the interaction with immunosuppressor cells is interrupted to avoid the effect of these suppressor cells, or DC vaccines need to be combined with other therapies.

4.2. The next generation of DCs

To improve DC vaccination, the investigators exploit to the microbial activation signals leading to generate potent DCs with high secretion of cytokines such as IL-12p70, which generate strong tumor-specific Th1 response and helper function for the generation of memory T cells, high production of polarizing signals, which help the generation of high avidity in CTLs that may be resistant to tumor microenvironment, and strong costimulation mediated via several costimulatory molecular pathways [51, 52]. This induces to eliminate Tregs and block tumor microenvironment results in the full activity of elicited CTLs and tumor rejection.

The initial phase of DC-based vaccines involving immature or partially-mature “first-generation” DCs has been reported [53, 54]. However, such DCs express suboptimal levels of costimulatory molecules and constitute weaker immunogens than subsequently implemented mature DCs, the “second-generation” of clinically applied standard DCs (sDCs), which induced by cytokine cocktails containing IL-1 β /TNF- α /IL-6/prostaglandin E₂ (PGE₂) [55]. However, to date, sDC vaccines still have some drawbacks, including the mediation of Th2 polarization by increased secretion of the immunosuppressive cytokine IL-10 from DCs and high activity in activating Tregs [56, 57]. Therefore, several investigators, including our group, have tried to develop new generation of potent DC that possess all required features for inducing effective tumor-specific immune responses. We demonstrated the feasibility of

inducing potent α -type 1-polarizing DCs (α DC1s) by exposing immature DCs to α -type 1-polarizing cytokine cocktail containing IL-1 β , TNF- α , IFN- α , IFN- γ , and polyinosinic:polycytidylic acid [poly(I:C)] to generate strong functional CTLs on average 20-fold higher than sDCs [58-62]. Recently, we successfully generated α DC1s from MM patients with high expression of costimulatory molecules, significant production of IL-12p70, and potent generation of myeloma-specific CTLs [37, 38]. Such a novel strategy would provide improved potency of *ex vivo*-generated DCs for cancer immunotherapy.

The other strategy to induce new potent DCs from patients with MM was the use of helper cells to promote type 1-polarization of DCs. Indeed, it has been demonstrated that NK cells play a major immunoregulatory role in the development of protective T cell-mediated immunity against intracellular pathogens and cancers [63]. Such helper activity of NK cells is at least partially mediated by the functional modulation of DCs. This phenomenon depended on the production of IFN- γ and TNF- α from the activated NK cells [63] and was associated with enhanced cross-presentation of tumor antigen and the induction of Th1 and CTL responses [39, 64, 65]. Recent data from our laboratory and other groups has demonstrated that NK-DC interactions promote the subsequent induction of tumor-specific responses in CD4⁺ and CD8⁺ T cells, allowing NK cells to act as helper cells in the development of type 1-polarized DCs in responses against cancer [39, 64, 65]. Resting NK cells that are activated in the presence of toll-like receptor (TLR) agonists, IL-2, and IFN- α can induce potent DCs with enhanced IL-12p70 production *in vitro*, generating strong antigen-specific CTLs against myeloma cells [39].

We also found that the selected combinations of TLR agonists synergistically triggered a Th1-polarizing capacity through production of high amounts of IL-12p70 [66]. However, the major limitation of this combination was the decreased ability of these cells to migrate into lymph nodes compared to that of conventional sDCs. When DCs are activated by individual TLR agonists, such as lipopolysaccharide (LPS) or poly(I:C), or by a combination of 2 TLR agonists, all cells mature and produce high levels of bioactive IL-12p70 in early phase of maturation and after subsequent stimulation with T cell-related DC activating signal CD40L. In addition, the phenotypes of these matured DCs were markedly enhanced when a combination of type I and type II IFN was added. These combinations of stimuli also regulated the expression of CD38 and CD74, markers related to the full activation of DCs [67, 68]. We demonstrated that, at the optimal concentration used to stimulate DCs, the combination of 2 TLR agonists with type I and II IFNs can be used to generate fully mature DCs that have high migratory capacity and can maintain IL-12p70-producing capacity. The regulation of CD38 and CD74 in DCs could in turn enhance the migratory activity of DCs in the presence of a combination of 2 TLR agonists and IFNs [69].

Ursolic acid (URC) is isolated from *Uncaria rhynchophylla* and phytochemically classified as a triterpene. Triterpene compounds have been identified as a unique class of natural products possessing diverse biological activities. Recently, we reported that URC activates human DCs in a fashion that favors Th1 polarization via the activation of TLR2- and/or TLR4-dependent IL-12p70 and induces the production of IFN- γ by CD4⁺ naïve T cells [70]. In addition, combination URC and IFN- γ enhanced the activation of DCs via promotion of IFN- γ -induced Th1

cell polarization that was dependent on the activation of IL-12p70 and independent of TLR4 [71, 72]. The potential of natural products to enhance DC maturation and activation has important implications for the use of DCs as a cancer vaccine.

4.3. New sources of myeloma-associated antigens for DC vaccines

Another important consideration to improve the efficacy of DC vaccination in patients with MM is an effective tumor antigen, instead of using idioype proteins with a weak antigenicity. The use of whole tumor cells, instead of single antigen, may help to enhance antitumor effects to target multiple tumor variants. It is necessary to use purified, optimized myeloma cells, if possible, as a source of tumor antigens for loading onto DCs to generate potent myeloma-specific CTLs [35]. However, it is not only impractical to obtain sufficient amounts of purified autologous myeloma cells for tumor antigens in the clinical setting from patients with MM and it is also unsuitable for those with a lower tumor burden status. As an alternative source of tumor-relevant antigens, allogeneic tumor cells or established cancer cell lines have been used to overcome the limitation in various tumors [37, 38]. DCs loaded with tumor antigens derived from allogeneic myeloma cells could generate myeloma-specific CTLs against autologous myeloma cells in patients with MM [37, 39]. The success of using an allogeneic myeloma cell line as tumor antigens led to the possibility that allogeneic myeloma cells could also be used as a viable source of tumor antigens in the context of appropriate major MHC alleles to autologous CTLs. In addition, autologous DCs loaded with dying myeloma cells of allogeneic matched monoclonal immunoglobulin subtype showed to generate potent myeloma-specific CTLs against autologous myeloma cells in MM patients [38]. These findings suggested that allogeneic myeloma cell lines and allogeneic matched monoclonal immunoglobulin subtype of myeloma were effective tumor antigens capable of inducing functional CTLs against patients' own myeloma cells.

Improved understanding of which specific anticancer agents lead to immunogenic cell death and whether these process can enhance antitumor immunity may facilitate the mechanism how chemotherapy and immunotherapy combination can induce immune responses against cancer. Recently, we have worked to develop strategies that recover dysfunction of DCs caused from loading tumor antigens through treatment of myeloma cells with a combination of the selective JAK/STAT3 inhibitor, JSI-124, and a kind of proteasome inhibitor, bortezomib. We observed that production of inhibitory cytokines, such as IL-10, IL-23, and especially IL-6, which induces DC dysfunction in MM patients, was down-regulated in DCs loaded with dying myeloma tumor cells that induced by these agents. Furthermore, phospho-STAT3 was also down-regulated in the DCs. These DCs displayed a superior ability to induce myeloma-specific responses of CTLs. More recently, we are investigating whether chaetocin could be used to induce dying tumor cells for loading onto DCs to enhance myeloma-specific antitumor responses. We show that anti-myeloma drug-induced dying tumor cells can be used as the source of myeloma antigens to loading onto DCs that could elicit potent anti-myeloma activity of CTLs due to the expression of HSP and cancer testis antigens as a mechanism of immunogenic death of human MM cells.

4.4. Blocking immunosuppressive activity during the loading of tumor antigens for DC vaccines

The suppressive effects of tumor cells during DC generation have been explained previously by the ability of the tumor microenvironment to suppress DC differentiation [73]. This process can influence STAT3 and ERK phosphorylation, resulting in hyperactivation of STAT3 and ERK, which may be responsible for defective generation of DCs [74]. The immune-mediated antitumor effects of DCs are enhanced by inhibition of the JAK2/STAT3 pathway [75], inhibition of p38 or activation of the MEK/ERK or MAP kinase pathways, and neutralization of IL-6 [76]. Recently, we found that when MM-derived DCs were generated by loading tumor lysates from autologous myeloma cells, these DCs showed lower phenotypic maturation, less T cell stimulatory capacity, less CTL activity, and highly abnormal IL-6 and IL-12 secretion compared to the secretion by unloaded DCs. Moreover, the levels of VEGF, phospho-STAT3, and phospho-ERK1/2 in these DCs were significantly higher than in unloaded DCs. After neutralization of VEGF activity, DC functions, signal transduction, and cytokine production were returned to normal level. Therefore, inhibitory factors and abnormal signaling pathways during maturation with tumor antigens in DCs may be responsible for the defective activity of DCs in MM, and these abnormalities may be overcome by neutralizing the signaling that would lead to a suppressed immune response [77].

5. Combination therapy: New concept to enhance efficacy of DC vaccines

Many factors contribute to the limited clinical efficacy of DC vaccines. The tumor microenvironment contains different kinds of inhibitory cells, such as Tregs and MDSC, and inhibitory molecules, such as IL-10, IL-6, TGF- β , and VEGF, all of which prevent the activation of effector T cells in response to DC responses [16-21, 23, 78, 79]. Although DC vaccines showed effective antitumor effect in experimental *ex vivo* systems, they didn't effectively induce strong immune responses that were enough to kill tumors *in vivo*. Therefore, strategies to improve the efficacy of DC vaccines are to overcome the immune tolerance/suppression induced by these cells, which are involved in the use of a combination of DC vaccine with either stimulatory cytokines or the targeting elimination of inhibitory cells and molecules in tumor microenvironment.

5.1. DC vaccine and cytokine combination

Cytokines, such as GM-CSF or IL-2, known to enhance cell-mediated immune responses may be administered as adjuvants with the vaccines aiming to create an environment where specific immune responses are readily induced [80, 81]. To enhance the efficacy of DC vaccination, Id-pulsed DCs were combined with GM-CSF [80, 82-84], with immunogenic carrier molecules such as KLH [27, 28, 31-33, 82, 85], or cytokine IL-2 [80, 83] to improve the effectiveness of these DC vaccines in patients with MM. Recently, a phase I study was performed in patients with MM using autologous DCs/tumor cells fusion in combination with GM-CSF administration at the day of DC vaccination [86]. The expansion of circulating CD4⁺ and CD8⁺ T cells reactive with autologous myeloma cells were detected in 11 of 15 evaluable patients. A majority of

patients with advanced disease demonstrated disease stabilization. In a murine myeloma model, mice were vaccinated with DC-plasmacytoma cell fusions and demonstrated that administration of IL-12 with the vaccine resulted in potentiation of *in vivo* T cell proliferation and cytotoxicity and eradication of established disease [87]. Therefore, the combination of DC vaccine with stimulatory cytokines is a feasible approach to provide a new source of DC-based vaccines for the development of immunotherapy against MM.

5.2. DC vaccine and chemotherapy combination

Chemotherapy can help to reverse the immunosuppression caused by cancers and also further enhance the capacity of DCs to trigger antitumor immunity [88]. Accumulating evidence indicates that conventional chemotherapy as well as radiotherapy selectively eliminates immunosuppressive cells, triggers the activation of DCs, and enhances antigen cross-presentation. Furthermore, specific anticancer agents lead to immunogenic cell death of tumor cells and these processes can enhance antitumor immunity.

Recent studies have shown that chemotherapeutic agents increase the efficacy of active or adoptive antitumor immunotherapies through beneficial immunomodulatory effects [89, 90]. Cyclophosphamide eliminates the activities of tumor-induced suppressor T cells in tumor-bearing hosts [90] and induces the production of immunostimulatory cytokines, such as type I IFN [91]. In addition, low-dose cyclophosphamide has been shown to down-regulate suppressor T cells and to decrease the production of TGF- β and IL-10 while inducing a Th2/Th1 shift in the cytokine profile [92-94]. Low-dose cyclophosphamide may enhance the antitumor efficacy of DC vaccines by increasing the proportion of IFN- γ secreting lymphocytes and suppressing the proportion of CD4⁺CD25⁺FoxP3⁺ Tregs in tumor-bearing mice [95]. The result of a clinical trial using allogeneic DC vaccines combined with low-dose cyclophosphamide has revealed that the combination therapy could induce stronger antitumor responses compared to the DC vaccine alone [96]. Recently, we demonstrated that a single administration of low-dose cyclophosphamide before the first DC vaccination showed to augment antitumor effects of DC vaccines to completely eradicate the tumor and to prolong the survival of vaccinated mice [64].

Lenalidomide is a thalidomide analog that has more potent anti-myeloma effects and less adverse effects [97]. Lenalidomide can induce apoptosis of myeloma cells, inhibit the production of cytokines (IL-6, VEGF, and TNF- α) in bone marrow of myeloma patients, and stimulate T cell and NK cell proliferation, cytotoxicity, and cytokine (IL-2, IFN- γ) production [97]. In addition, lenalidomide can inhibit the frequency and function of Tregs, resulting in inhibition of Treg expansion and FoxP3 expression in cancer patients [98]. Interestingly, this drug can also induce the activation of APC function, resulting in upregulation of CD40, CD80, and CD86 in chronic lymphocytic leukemia [99]. Therefore, lenalidomide can be used as an immunomodulatory drug in order to enhance immune responses against cancer. Our *in vitro* study showed that lenalidomide enhanced the maturation and function of DCs in the presence of LPS, resulting in synergistic stimulation of DCs to increase phenotype expression, IL-12p70 production, T cell stimulation capacities, and CTL activities against myeloma cells, and to suppress the generation of Tregs. Moreover, our *in vivo* mouse myeloma model showed that

a treatment combining the lenalidomide with DC vaccination markedly improved antitumor effect by inhibiting immunosuppressor cells, recovering effector cells, and inducing superior polarization of the Th1/Th2 balance in favor of the Th1 response. This immunomodulatory effect may be a crucial component of the enhancer-like properties of lenalidomide in the context of antitumor immunity against MM.

5.3. Chemotherapeutic agent can induce “immunogenic myeloma-cell death” to trigger activation of DCs and to enhance cross-presentation of DCs

Most of chemotherapeutic agents kill tumor cells by the induction of apoptosis. Previously, chemotherapy and immunotherapy have usually been regarded as unrelated therapy in the treatment of cancers because chemotherapy-induced apoptotic cell death has long been considered as non-immunogenic or inducing immune tolerance. Recently, apoptotic cell death when coupled with inflammatory signals, such as HSPs, is clearly known to induce the activation of DCs and triggers the immune response [100]. Some chemotherapeutic agents could induce a type of tumor cell death that activates efficient antitumor immunity, so it is called “immunogenic tumor-cell death”. Immunogenic tumor-cell death expresses danger signals on the tumor cell surface or secretes immunostimulatory factors, such as HSPs, calreticulin, high mobility group box 1 protein (HMGB1), and ATP, into the tumor microenvironment, thereby promoting DC maturation and stimulating a powerful T cell immune response [88].

Cyclophosphamide is well known as a potent cytotoxic and lymphoablative drug in conventional and high dosages. However, more recent work highlighted as an immunostimulatory and/or antiangiogenic agent at low dosages, opening up novel indication in the field of cancer immunotherapy. In recent reports, cyclophosphamide administration in tumor-bearing mice induced pre-apoptotic surface translocation of calreticulin on tumor cells [101], which serves as an “eat-me” signal for phagocytes [102] and the release of high-mobility group box1 (HMGB1) protein in the extracellular milieu [101], which constitutes a “danger signal” triggering activation of the DC processing machinery [103]. These events are prerequisites for adequate engulfment of tumor apoptotic material and optimal CD8⁺ T cell cross-priming by DCs [102, 103].

HSPs are intracellular chaperones for many proteins, but they can also be expressed on the cell surface or even be released under stress conditions [104, 105]. HSP acts as an adjuvant in initiating the activation of DCs or as protein vehicle to facilitate the presentation of antigen peptides to T cells. Spisek et al. [106] reported that uptake of myeloma cells by DCs after tumor cell death induced by bortezomib leads to the induction of antitumor immunity and enhances DC-mediated tumor immune response, indicating the probability mechanism due to the expression of HSP90 on the surface of dying cells, thereby facilitating the activation of DCs in response to dying tumor cells. Our study also found that HSPs released from dying tumor cells, which were induced by a combination of the selective JAK/STAT3 inhibitor JSI-124 and proteasome inhibitor bortezomib, act on tumor cells to recover DC dysfunction and to induce cytokine and chemokine production from DCs, resulting in generation of potent myeloma-specific CTL response against myeloma cells.

5.4. Possible combination DCs and other approaches

In the presence of regulatory and suppressive environment, it is very difficult to elicit or induce effective immune response after DC vaccination in cancer patients. To improve the clinical outcomes, DC vaccines need to be combined, in particular for patients at advanced stages, with other approaches that offset the suppressive tumor environment [107]. It has been known that the specific depletion of CD4⁺CD25⁺ Treg cells by anti-CD25 antibodies increases the efficiency of the anti-tumor immune response of tumor-bearing animals, although the tumors are not completely rejected [108]. An increased number of CD4⁺CD25⁺FoxP3⁺ regulatory T cells have been demonstrated in patients with MM [22, 109]. Depletion of Treg may have resulted in improved response to tumor vaccine in animal models and a clinical study. In addition, blocking antibodies or soluble receptors were exploited for the blockade of suppressive cytokines in the tumor microenvironment, such as IL-10 [110], IL-13 [111], TGF- β [112] and VEGF [113]. Such strategies can be used to block immune-inhibitory signals in lymphocytes as illustrated by anti-CTLA-4 [114] and/or anti-PD1 [115] or to block their ligands expressed on tumors.

Another strategy to improve DC vaccination is combination approach with other immune cells, including adoptive T cells or NK cells. In adoptive T-cell transfer, one can seek to modulate the number of regulatory T cells, and transfer a population of activated effector cells. The combination of DC vaccination and adoptive T-cell transfer led to a more robust antitumor response than the use of each treatment modality [116]. These findings illuminate a new potential application for DC vaccination in the *in vivo* stimulation of adoptively transferred T cells. Therefore, combining active and passive immunotherapies in the treatment of MM may enhance the efficacy of tumor vaccine in the future.

6. Future perspectives

Progress in understanding DC biology in MM patients and the recruitment of suppressive cells of the adaptive and innate immune system in antitumor immunity of cellular immunotherapy is leading to new concept which aims at improved immune and clinical outcomes in MM. The new generation of DCs may be a potential vaccine therapy for inducing the rate of tumor responses and prolonging survival of patients with MM. Furthermore, information from studies that combine DC vaccine with other therapies, including chemotherapy, radiation therapy, molecular target agents, other immunotherapy (adaptive T cells or NK cells), or adjuvants will have high impact on enhancing therapeutic immunity in MM by simultaneously enhancing the potency of immune responses and offsetting immunoregulatory pathways.

Acknowledgements

This study was financially supported by grant no. 2011-0005285 from General Researcher Program Type II of the National Research Foundation of Korea; grant no. RTI05-01-01 from

the Regional Technology Innovation Program of the Ministry of Commerce, Industry and Energy; grant no. A000200058 from the Regional Industrial Technology Development program of the Ministry of Knowledge and Economy; grant no. 1120390 from the National R&D Program for Cancer Control, Ministry for Health and Welfare; grant no. 2011-0030034 from Leading Foreign Research Institute Recruitment Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology (MEST), Republic of Korea.

Author details

Je-Jung Lee^{1,2,3}, Youn-Kyung Lee³, Hyun Ju Lee^{1,2}, Sung-Hoon Jung^{1,2} and Thanh-Nhan Nguyen-Pham^{1,2}

1 Research Center for Cancer Immunotherapy, Chonnam National University Hwasun Hospital, Jeollanamdo, Republic of Korea

2 Department of Hematology-Oncology, Chonnam National University Hwasun Hospital, Jeollanamdo, Republic of Korea

3 Vaxcell-Bio Therapeutics, Hwasun, Jeollanamdo, Republic of Korea

References

- [1] Kyle RA, Rajkumar SV. Multiple myeloma. *N Engl J Med* 2004; 351: 1860-1873.
- [2] Sirohi B, Powles R. Multiple myeloma. *Lancet* 2004; 363: 875-887.
- [3] Attal M, Harousseau JL. The role of high-dose therapy with autologous stem cell support in the era of novel agents. *Semin Hematol* 2009; 46: 127-132.
- [4] Lonial S, Cavenagh J. Emerging combination treatment strategies containing novel agents in newly diagnosed multiple myeloma. *Br J Haematol* 2009; 145: 681-708.
- [5] Perez-Simon JA, Martino R, Alegre A et al. Chronic but not acute graft-versus-host disease improves outcome in multiple myeloma patients after non-myeloablative allogeneic transplantation. *Br J Haematol* 2003; 121: 104-108.
- [6] Harrison SJ, Cook G, Nibbs RJ, Prince HM. Immunotherapy of multiple myeloma: the start of a long and tortuous journey. *Expert Rev Anticancer Ther* 2006; 6: 1769-1785.
- [7] Banchereau J, Steinman RM. Dendritic cells and the control of immunity. *Nature* 1998; 392: 245-252.

- [8] O'Garra A. Cytokines induce the development of functionally heterogeneous T helper cell subsets. *Immunity* 1998; 8: 275-283.
- [9] Wynn TA. T(H)-17: a giant step from T(H)1 and T(H)2. *Nat Immunol* 2005; 6: 1069-1070.
- [10] Ridgway D. The first 1000 dendritic cell vaccinees. *Cancer Invest* 2003; 21: 873-886.
- [11] Banchereau J, Palucka AK. Dendritic cells as therapeutic vaccines against cancer. *Nat Rev Immunol* 2005; 5: 296-306.
- [12] Albert ML, Sauter B, Bhardwaj N. Dendritic cells acquire antigen from apoptotic cells and induce class I-restricted CTLs. *Nature* 1998; 392: 86-89.
- [13] Brown RD, Pope B, Murray A et al. Dendritic cells from patients with myeloma are numerically normal but functionally defective as they fail to up-regulate CD80 (B7-1) expression after huCD40LT stimulation because of inhibition by transforming growth factor-beta1 and interleukin-10. *Blood* 2001; 98: 2992-2998.
- [14] Ratta M, Fagnoni F, Curti A et al. Dendritic cells are functionally defective in multiple myeloma: the role of interleukin-6. *Blood* 2002; 100: 230-237.
- [15] Yu H, Kortylewski M, Pardoll D. Crosstalk between cancer and immune cells: role of STAT3 in the tumour microenvironment. *Nat Rev Immunol* 2007; 7: 41-51.
- [16] Ogawara H, Handa H, Yamazaki T et al. High Th1/Th2 ratio in patients with multiple myeloma. *Leuk Res* 2005; 29: 135-140.
- [17] Maecker B, Anderson KS, von Bergwelt-Baildon MS et al. Viral antigen-specific CD8+ T-cell responses are impaired in multiple myeloma. *Br J Haematol* 2003; 121: 842-848.
- [18] Dhodapkar MV, Geller MD, Chang DH et al. A reversible defect in natural killer T cell function characterizes the progression of premalignant to malignant multiple myeloma. *J Exp Med* 2003; 197: 1667-1676.
- [19] Jarahian M, Watzl C, Issa Y et al. Blockade of natural killer cell-mediated lysis by NCAM140 expressed on tumor cells. *Int J Cancer* 2007; 120: 2625-2634.
- [20] Prabhala RH, Neri P, Bae JE et al. Dysfunctional T regulatory cells in multiple myeloma. *Blood* 2006; 107: 301-304.
- [21] Banerjee DK, Dhodapkar MV, Matayeva E et al. Expansion of FOXP3high regulatory T cells by human dendritic cells (DCs) in vitro and after injection of cytokine-matured DCs in myeloma patients. *Blood* 2006; 108: 2655-2661.
- [22] Brimnes MK, Vangsted AJ, Knudsen LM et al. Increased level of both CD4+FOXP3+ regulatory T cells and CD14+HLA-DR/low myeloid-derived suppressor cells and decreased level of dendritic cells in patients with multiple myeloma. *Scand J Immunol* 2010; 72: 540-547.

- [23] Kwak LW, Taub DD, Duffey PL et al. Transfer of myeloma idiotype-specific immunity from an actively immunised marrow donor. *Lancet* 1995; 345: 1016-1020.
- [24] Li Y, Bendandi M, Deng Y et al. Tumor-specific recognition of human myeloma cells by idiotype-induced CD8(+) T cells. *Blood* 2000; 96: 2828-2833.
- [25] Butch AW, Kelly KA, Munshi NC. Dendritic cells derived from multiple myeloma patients efficiently internalize different classes of myeloma protein. *Exp Hematol* 2001; 29: 85-92.
- [26] Wen YJ, Barlogie B, Yi Q. Idiotype-specific cytotoxic T lymphocytes in multiple myeloma: evidence for their capacity to lyse autologous primary tumor cells. *Blood* 2001; 97: 1750-1755.
- [27] Lim SH, Bailey-Wood R. Idiotypic protein-pulsed dendritic cell vaccination in multiple myeloma. *Int J Cancer* 1999; 83: 215-222.
- [28] Liso A, Stockerl-Goldstein KE, Auffermann-Gretzinger S et al. Idiotype vaccination using dendritic cells after autologous peripheral blood progenitor cell transplantation for multiple myeloma. *Biol Blood Marrow Transplant* 2000; 6: 621-627.
- [29] Titzer S, Christensen O, Manzke O et al. Vaccination of multiple myeloma patients with idiotype-pulsed dendritic cells: immunological and clinical aspects. *Br J Haematol* 2000; 108: 805-816.
- [30] Yi Q, Desikan R, Barlogie B, Munshi N. Optimizing dendritic cell-based immunotherapy in multiple myeloma. *Br J Haematol* 2002; 117: 297-305.
- [31] Reichardt VL, Milazzo C, Brugger W et al. Idiotype vaccination of multiple myeloma patients using monocyte-derived dendritic cells. *Haematologica* 2003; 88: 1139-1149.
- [32] Bendandi M, Rodriguez-Calvillo M, Inoges S et al. Combined vaccination with idiotype-pulsed allogeneic dendritic cells and soluble protein idiotype for multiple myeloma patients relapsing after reduced-intensity conditioning allogeneic stem cell transplantation. *Leuk Lymphoma* 2006; 47: 29-37.
- [33] Yi Q, Szmania S, Freeman J et al. Optimizing dendritic cell-based immunotherapy in multiple myeloma: intranodal injections of idiotype-pulsed CD40 ligand-matured vaccines led to induction of type-1 and cytotoxic T-cell immune responses in patients. *Br J Haematol* 2010; 150: 554-564.
- [34] Rollig C, Schmidt C, Bornhauser M et al. Induction of cellular immune responses in patients with stage-I multiple myeloma after vaccination with autologous idiotype-pulsed dendritic cells. *J Immunother* 2011; 34: 100-106.
- [35] Lee JJ, Choi BH, Kang HK et al. Induction of multiple myeloma-specific cytotoxic T lymphocyte stimulation by dendritic cell pulsing with purified and optimized myeloma cell lysates. *Leuk Lymphoma* 2007; 48: 2022-2031.

- [36] Wen YJ, Min R, Tricot G et al. Tumor lysate-specific cytotoxic T lymphocytes in multiple myeloma: promising effector cells for immunotherapy. *Blood* 2002; 99: 3280-3285.
- [37] Yang DH, Kim MH, Hong CY et al. Alpha-type 1-polarized dendritic cells loaded with apoptotic allogeneic myeloma cell line induce strong CTL responses against autologous myeloma cells. *Ann Hematol* 2010; 89: 795-801.
- [38] Yang DH, Kim MH, Lee YK et al. Successful cross-presentation of allogeneic myeloma cells by autologous alpha-type 1-polarized dendritic cells as an effective tumor antigen in myeloma patients with matched monoclonal immunoglobulins. *Ann Hematol* 2011; 90: 1419-1426.
- [39] Nguyen-Pham TN, Im CM, Nguyen TA et al. Induction of myeloma-specific cytotoxic T lymphocytes responses by natural killer cells stimulated-dendritic cells in patients with multiple myeloma. *Leuk Res* 2011; 35: 1241-1247.
- [40] Milazzo C, Reichardt VL, Muller MR et al. Induction of myeloma-specific cytotoxic T cells using dendritic cells transfected with tumor-derived RNA. *Blood* 2003; 101: 977-982.
- [41] Qian J, Wang S, Yang J et al. Targeting heat shock proteins for immunotherapy in multiple myeloma: generation of myeloma-specific CTLs using dendritic cells pulsed with tumor-derived gp96. *Clin Cancer Res* 2005; 11: 8808-8815.
- [42] Qian J, Hong S, Wang S et al. Myeloma cell line-derived, pooled heat shock proteins as a universal vaccine for immunotherapy of multiple myeloma. *Blood* 2009; 114: 3880-3889.
- [43] Hao S, Bi X, Xu S et al. Enhanced antitumor immunity derived from a novel vaccine of fusion hybrid between dendritic and engineered myeloma cells. *Exp Oncol* 2004; 26: 300-306.
- [44] Vasir B, Borges V, Wu Z et al. Fusion of dendritic cells with multiple myeloma cells results in maturation and enhanced antigen presentation. *Br J Haematol* 2005; 129: 687-700.
- [45] Brossart P, Schneider A, Dill P et al. The epithelial tumor antigen MUC1 is expressed in hematological malignancies and is recognized by MUC1-specific cytotoxic T-lymphocytes. *Cancer Res* 2001; 61: 6846-6850.
- [46] Batchu RB, Moreno AM, Szmania SM et al. Protein transduction of dendritic cells for NY-ESO-1-based immunotherapy of myeloma. *Cancer Res* 2005; 65: 10041-10049.
- [47] Chiriva-Internati M, Wang Z, Salati E et al. Sperm protein 17 (Sp17) is a suitable target for immunotherapy of multiple myeloma. *Blood* 2002; 100: 961-965.

- [48] Vonderheide RH, Hahn WC, Schultze JL, Nadler LM. The telomerase catalytic subunit is a widely expressed tumor-associated antigen recognized by cytotoxic T lymphocytes. *Immunity* 1999; 10: 673-679.
- [49] Ocadlikova D, Kryukov F, Mollova K et al. Generation of myeloma-specific T cells using dendritic cells loaded with MUC1- and hTERT- driven nonapeptides or myeloma cell apoptotic bodies. *Neoplasma* 2010; 57: 455-464.
- [50] Qian J, Xie J, Hong S et al. Dickkopf-1 (DKK1) is a widely expressed and potent tumor-associated antigen in multiple myeloma. *Blood* 2007; 110: 1587-1594.
- [51] Palucka K, Ueno H, Zurawski G et al. Building on dendritic cell subsets to improve cancer vaccines. *Curr Opin Immunol* 2010; 22: 258-263.
- [52] Palucka K, Ueno H, Banchereau J. Recent developments in cancer vaccines. *J Immunol* 2011; 186: 1325-1331.
- [53] Hsu FJ, Benike C, Fagnoni F et al. Vaccination of patients with B-cell lymphoma using autologous antigen-pulsed dendritic cells. *Nat Med* 1996; 2: 52-58.
- [54] Nestle FO, Alijagic S, Gilliet M et al. Vaccination of melanoma patients with peptide- or tumor lysate-pulsed dendritic cells. *Nat Med* 1998; 4: 328-332.
- [55] Jonuleit H, Kuhn U, Muller G et al. Pro-inflammatory cytokines and prostaglandins induce maturation of potent immunostimulatory dendritic cells under fetal calf serum-free conditions. *Eur J Immunol* 1997; 27: 3135-3142.
- [56] Kalinski P, Vieira PL, Schuitemaker JH et al. Prostaglandin E(2) is a selective inducer of interleukin-12 p40 (IL-12p40) production and an inhibitor of bioactive IL-12p70 heterodimer. *Blood* 2001; 97: 3466-3469.
- [57] Yamazaki S, Inaba K, Tarbell KV, Steinman RM. Dendritic cells expand antigen-specific Foxp3+ CD25+ CD4+ regulatory T cells including suppressors of alloreactivity. *Immunol Rev* 2006; 212: 314-329.
- [58] Mailliard RB, Wankowicz-Kalinska A, Cai Q et al. alpha-type-1 polarized dendritic cells: a novel immunization tool with optimized CTL-inducing activity. *Cancer Res* 2004; 64: 5934-5937.
- [59] Lee JJ, Foon KA, Mailliard RB et al. Type 1-polarized dendritic cells loaded with autologous tumor are a potent immunogen against chronic lymphocytic leukemia. *J Leukoc Biol* 2008; 84: 319-325.
- [60] Wesa A, Kalinski P, Kirkwood JM et al. Polarized type-1 dendritic cells (DC1) producing high levels of IL-12 family members rescue patient TH1-type antimelanoma CD4+ T cell responses in vitro. *J Immunother* 2007; 30: 75-82.
- [61] Giermasz AS, Urban JA, Nakamura Y et al. Type-1 polarized dendritic cells primed for high IL-12 production show enhanced activity as cancer vaccines. *Cancer Immunol Immunother* 2009; 58: 1329-1336.

- [62] Okada H, Kalinski P, Ueda R et al. Induction of CD8+ T-cell responses against novel glioma-associated antigen peptides and clinical activity by vaccinations with {alpha}-type 1 polarized dendritic cells and polyinosinic-polycytidylic acid stabilized by lysine and carboxymethylcellulose in patients with recurrent malignant glioma. *J Clin Oncol* 2011; 29: 330-336.
- [63] Gerosa F, Baldani-Guerra B, Nisii C et al. Reciprocal activating interaction between natural killer cells and dendritic cells. *J Exp Med* 2002; 195: 327-333.
- [64] Pham TN, Hong CY, Min JJ et al. Enhancement of antitumor effect using dendritic cells activated with natural killer cells in the presence of Toll-like receptor agonist. *Exp Mol Med* 2010; 42: 407-419.
- [65] Mailliard RB, Son YI, Redlinger R et al. Dendritic cells mediate NK cell help for Th1 and CTL responses: two-signal requirement for the induction of NK cell helper function. *J Immunol* 2003; 171: 2366-2373.
- [66] Napolitani G, Rinaldi A, Bertonni F et al. Selected Toll-like receptor agonist combinations synergistically trigger a T helper type 1-polarizing program in dendritic cells. *Nat Immunol* 2005; 6: 769-776.
- [67] Frasca L, Fedele G, Deaglio S et al. CD38 orchestrates migration, survival, and Th1 immune response of human mature dendritic cells. *Blood* 2006; 107: 2392-2399.
- [68] Faure-Andre G, Vargas P, Yuseff MI et al. Regulation of dendritic cell migration by CD74, the MHC class II-associated invariant chain. *Science* 2008; 322: 1705-1710.
- [69] Nguyen-Pham TN, Lim MS, Nguyen TA et al. Type I and II interferons enhance dendritic cell maturation and migration capacity by regulating CD38 and CD74 that have synergistic effects with TLR agonists. *Cell Mol Immunol* 2011; 8: 341-347.
- [70] Jung TY, Pham TN, Umeyama A et al. Ursolic acid isolated from *Uncaria rhynchophylla* activates human dendritic cells via TLR2 and/or TLR4 and induces the production of IFN-gamma by CD4+ naive T cells. *Eur J Pharmacol* 2010; 643: 297-303.
- [71] Bae WK, Umeyama A, Chung IJ et al. Uncarinic acid C plus IFN-gamma generates monocyte-derived dendritic cells and induces a potent Th1 polarization with capacity to migrate. *Cell Immunol* 2010; 266: 104-110.
- [72] Kim KS, Pham TN, Jin CJ et al. Uncarinic Acid C Isolated from *Uncaria rhynchophylla* Induces Differentiation of Th1-Promoting Dendritic Cells Through TLR4 Signaling. *Biomark Insights* 2011; 6: 27-38.
- [73] Savill J, Dransfield I, Gregory C, Haslett C. A blast from the past: clearance of apoptotic cells regulates immune responses. *Nat Rev Immunol* 2002; 2: 965-975.
- [74] Kitamura H, Kamon H, Sawa S et al. IL-6-STAT3 controls intracellular MHC class II alphabeta dimer level through cathepsin S activity in dendritic cells. *Immunity* 2005; 23: 491-502.

- [75] Nefedova Y, Gabrilovich DI. Targeting of Jak/STAT pathway in antigen presenting cells in cancer. *Curr Cancer Drug Targets* 2007; 7: 71-77.
- [76] Wang S, Hong S, Yang J et al. Optimizing immunotherapy in multiple myeloma: Restoring the function of patients' monocyte-derived dendritic cells by inhibiting p38 or activating MEK/ERK MAPK and neutralizing interleukin-6 in progenitor cells. *Blood* 2006; 108: 4071-4077.
- [77] Yang DH, Park JS, Jin CJ et al. The dysfunction and abnormal signaling pathway of dendritic cells loaded by tumor antigen can be overcome by neutralizing VEGF in multiple myeloma. *Leuk Res* 2009; 33: 665-670.
- [78] Ostrand-Rosenberg S, Sinha P. Myeloid-derived suppressor cells: linking inflammation and cancer. *J Immunol* 2009; 182: 4499-4506.
- [79] Bergenbrant S, Yi Q, Osterborg A et al. Modulation of anti-idiotypic immune response by immunization with the autologous M-component protein in multiple myeloma patients. *Br J Haematol* 1996; 92: 840-846.
- [80] Massaia M, Borriore P, Battaglio S et al. Idiotype vaccination in human myeloma: generation of tumor-specific immune responses after high-dose chemotherapy. *Blood* 1999; 94: 673-683.
- [81] Shimizu K, Fields RC, Giedlin M, Mule JJ. Systemic administration of interleukin 2 enhances the therapeutic efficacy of dendritic cell-based tumor vaccines. *Proc Natl Acad Sci U S A* 1999; 96: 2268-2273.
- [82] Coscia M, Mariani S, Battaglio S et al. Long-term follow-up of idiotype vaccination in human myeloma as a maintenance therapy after high-dose chemotherapy. *Leukemia* 2004; 18: 139-145.
- [83] Rasmussen T, Hansson L, Osterborg A et al. Idiotype vaccination in multiple myeloma induced a reduction of circulating clonal tumor B cells. *Blood* 2003; 101: 4607-4610.
- [84] Cull G, Durrant L, Stainer C et al. Generation of anti-idiotypic immune responses following vaccination with idiotype-protein pulsed dendritic cells in myeloma. *Br J Haematol* 1999; 107: 648-655.
- [85] Reichardt VL, Okada CY, Liso A et al. Idiotype vaccination using dendritic cells after autologous peripheral blood stem cell transplantation for multiple myeloma--a feasibility study. *Blood* 1999; 93: 2411-2419.
- [86] Rosenblatt J, Vasir B, Uhl L et al. Vaccination with dendritic cell/tumor fusion cells results in cellular and humoral antitumor immune responses in patients with multiple myeloma. *Blood* 2011; 117: 393-402.

- [87] Gong J, Koido S, Chen D et al. Immunization against murine multiple myeloma with fusions of dendritic and plasmacytoma cells is potentiated by interleukin 12. *Blood* 2002; 99: 2512-2517.
- [88] Lake RA, Robinson BW. Immunotherapy and chemotherapy--a practical partnership. *Nat Rev Cancer* 2005; 5: 397-405.
- [89] Mihalyo MA, Doody AD, McAleer JP et al. In vivo cyclophosphamide and IL-2 treatment impedes self-antigen-induced effector CD4 cell tolerization: implications for adoptive immunotherapy. *J Immunol* 2004; 172: 5338-5345.
- [90] North RJ. Cyclophosphamide-facilitated adoptive immunotherapy of an established tumor depends on elimination of tumor-induced suppressor T cells. *J Exp Med* 1982; 155: 1063-1074.
- [91] Proietti E, Greco G, Garrone B et al. Importance of cyclophosphamide-induced bystander effect on T cells for a successful tumor eradication in response to adoptive immunotherapy in mice. *J Clin Invest* 1998; 101: 429-441.
- [92] Berd D, Maguire HC, Jr., Mastrangelo MJ. Potentiation of human cell-mediated and humoral immunity by low-dose cyclophosphamide. *Cancer Res* 1984; 44: 5439-5443.
- [93] Matar P, Rozados VR, Gervasoni SI, Scharovsky GO. Th2/Th1 switch induced by a single low dose of cyclophosphamide in a rat metastatic lymphoma model. *Cancer Immunol Immunother* 2002; 50: 588-596.
- [94] Matar P, Rozados VR, Gonzalez AD et al. Mechanism of antimetastatic immunopotentiality by low-dose cyclophosphamide. *Eur J Cancer* 2000; 36: 1060-1066.
- [95] Liu JY, Wu Y, Zhang XS et al. Single administration of low dose cyclophosphamide augments the antitumor effect of dendritic cell vaccine. *Cancer Immunol Immunother* 2007; 56: 1597-1604.
- [96] Holtl L, Ramoner R, Zelle-Rieser C et al. Allogeneic dendritic cell vaccination against metastatic renal cell carcinoma with or without cyclophosphamide. *Cancer Immunol Immunother* 2005; 54: 663-670.
- [97] Quach H, Ritchie D, Stewart AK et al. Mechanism of action of immunomodulatory drugs (IMiDS) in multiple myeloma. *Leukemia* 2010; 24: 22-32.
- [98] Galustian C, Meyer B, Labarthe MC et al. The anti-cancer agents lenalidomide and pomalidomide inhibit the proliferation and function of T regulatory cells. *Cancer Immunol Immunother* 2009; 58: 1033-1045.
- [99] Ramsay AG, Gribben JG. Immune dysfunction in chronic lymphocytic leukemia T cells and lenalidomide as an immunomodulatory drug. *Haematologica* 2009; 94: 1198-1202.

- [100] Restifo NP. Building better vaccines: how apoptotic cell death can induce inflammation and activate innate and adaptive immunity. *Curr Opin Immunol* 2000; 12: 597-603.
- [101] Schiavoni G, Sistigu A, Valentini M et al. Cyclophosphamide synergizes with type I interferons through systemic dendritic cell reactivation and induction of immunogenic tumor apoptosis. *Cancer Res* 2011; 71: 768-778.
- [102] Obeid M, Tesniere A, Ghiringhelli F et al. Calreticulin exposure dictates the immunogenicity of cancer cell death. *Nat Med* 2007; 13: 54-61.
- [103] Apetoh L, Ghiringhelli F, Tesniere A et al. Toll-like receptor 4-dependent contribution of the immune system to anticancer chemotherapy and radiotherapy. *Nat Med* 2007; 13: 1050-1059.
- [104] Chen T, Cao X. Stress for maintaining memory: HSP70 as a mobile messenger for innate and adaptive immunity. *Eur J Immunol* 2010; 40: 1541-1544.
- [105] Srivastava P. Interaction of heat shock proteins with peptides and antigen presenting cells: chaperoning of the innate and adaptive immune responses. *Annu Rev Immunol* 2002; 20: 395-425.
- [106] Spisek R, Charalambous A, Mazumder A et al. Bortezomib enhances dendritic cell (DC)-mediated induction of immunity to human myeloma via exposure of cell surface heat shock protein 90 on dying tumor cells: therapeutic implications. *Blood* 2007; 109: 4839-4845.
- [107] Dougan M, Dranoff G. Immune therapy for cancer. *Annu Rev Immunol* 2009; 27: 83-117.
- [108] Javia LR, Rosenberg SA. CD4+CD25+ suppressor lymphocytes in the circulation of patients immunized against melanoma antigens. *J Immunother* 2003; 26: 85-93.
- [109] Beyer M, Kochanek M, Giese T et al. In vivo peripheral expansion of naive CD4+CD25^{high} FoxP3⁺ regulatory T cells in patients with multiple myeloma. *Blood* 2006; 107: 3940-3949.
- [110] Moore KW, de Waal Malefyt R, Coffman RL, O'Garra A. Interleukin-10 and the interleukin-10 receptor. *Annu Rev Immunol* 2001; 19: 683-765.
- [111] Terabe M, Matsui S, Noben-Trauth N et al. NKT cell-mediated repression of tumor immunosurveillance by IL-13 and the IL-4R-STAT6 pathway. *Nat Immunol* 2000; 1: 515-520.
- [112] Terabe M, Ambrosino E, Takaku S et al. Synergistic enhancement of CD8⁺ T cell-mediated tumor vaccine efficacy by an anti-transforming growth factor-beta monoclonal antibody. *Clin Cancer Res* 2009; 15: 6560-6569.
- [113] Rabinovich GA, Gabrilovich D, Sotomayor EM. Immunosuppressive strategies that are mediated by tumor cells. *Annu Rev Immunol* 2007; 25: 267-296.

- [114] Peggs KS, Quezada SA, Korman AJ, Allison JP. Principles and use of anti-CTLA4 antibody in human cancer immunotherapy. *Curr Opin Immunol* 2006; 18: 206-213.
- [115] Pilon-Thomas S, Mackay A, Vohra N, Mule JJ. Blockade of programmed death ligand 1 enhances the therapeutic efficacy of combination immunotherapy against melanoma. *J Immunol* 2010; 184: 3442-3449.
- [116] Lou Y, Wang G, Lizee G et al. Dendritic cells strongly boost the antitumor activity of adoptively transferred T cells in vivo. *Cancer Res* 2004; 64: 6783-6790.

IntechOpen