

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Electron Transfer Routes in Oxygenic Photosynthesis: Regulatory Mechanisms and New Perspectives

---

Snježana Jurić, Lea Vojta and Hrvoje Fulgosi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/55339>

---

## 1. Introduction

In nature, the intensity of light changes rapidly, spanning from  $2000 \mu\text{mol}_{\text{PHOTONS}} \text{m}^{-2} \text{s}^{-1}$  during the brightest sunlight, down to  $200\text{-}500 \mu\text{mol}_{\text{PHOTONS}} \text{m}^{-2} \text{s}^{-1}$ , when shaded by clouds, followed by the period of dark, to encircle with the next sunrise of barely  $10 \mu\text{mol}_{\text{PHOTONS}} \text{m}^{-2} \text{s}^{-1}$ . Plants, capable of performing limited movements, like leaf and chloroplast movements, respectively, have been forced to evolve different means of coping with the unpredictable nature. To better understand the acclimatization mechanisms, the intensity of light has been roughly divided into low-light and high-light conditions, shaped by the laboratory conditions of plant growth and the strength of the measuring apparatus available. Moreover, looking at the time scale, it is possible to differentiate between short-term and long-term acclimatization processes in plants. These artificial divisions are needed to simplify the most important process in photosynthesis in vascular plants: the rate and the distribution of excitation energy between two photosystems. Photosystem units are organized into large supercomplexes with peripherally attached antenna complexes, being further assembled into megacomplexes [1]. Two types of peripheral antenna proteins associated to photosystem II (PSII) are known: the major Light-Harvesting Complex II (LHCII), which occurs as a trimeric complex containing the proteins Lhcb1, Lhcb2 and Lhcb3, and three minor monomeric complexes, namely Lhcb4 (CP29), Lhcb5 (CP26), and Lhcb6 (CP24). These peripheral complexes bind variable number of molecules of chlorophyll *a* and *b*, and of some xanthophyll molecules (for physicochemical properties of chlorophylls see the Chapter Kobayashi et al.). In Arabidopsis PSII-LHCII supercomplexes, Lhcb are organized into two rings around the dimeric PSII core complexes, with Lhcb1, Lhcb2, Lhcb4 and Lhcb5 detected in the inner ring, where Lhcb1 and Lhcb2 participate in strongly bound LHCII trimer, while the outer ring consists of Lhcb6 and of moderately bound LHCII trimer (the Lhcb1 and Lhcb3 gene products). Photosystem I (PSI) is associated with the Light-

Harvesting Complex I (LHCI) that binds 10 molecules of chlorophyll *a* or chlorophyll *b* plus a few xanthophylls per one Lhca protein. In green plants, LHCI consists of four polypeptides (Lhca1-Lhca4) from the Lhc protein superfamily. In Arabidopsis, two additional proteins have been identified (Lhca5 and Lhca6), but their contribution to LHCI is still under debate [1]. More distantly related family members are the photoprotective early light-induced stress response proteins (ELIPS), and the component of PSII, PsbS [2].

## 2. The way the plant protects itself: Non-photochemical quenching


The non-photochemical quenching (NPQ) is a short-term response by which plants harmlessly dissipate excess excitation energy into heat under high-light conditions. NPQ is observed in all higher plants, in lower plants, green algae and diatoms [3]. NPQ is also present in oceanic picophytoplankton species (see Chapter Kulk et al.). Basically, during the absorption of sunlight by light-harvesting complexes (LHCs) associated with reaction centres, a chlorophyll *a* molecule shifts from its ground energetic state to its singlet excited state. It can return to its ground state *via* one of several pathways: re-emission of excitation energy in the form of chlorophyll fluorescence; transfer of excitation energy to reaction centres to be utilised in photochemistry reactions; de-excitation by dissipating heat (NPQ); production of triplet excited state, which would be a highly profitable valve for excess excitation, but it indirectly produces a very reactive oxygen species (ROS), singlet oxygen, by transferring energy to the ground-state oxygen [4]. In addition to the dissipation of excitation energy, non-photochemical processes also quench or diminish chlorophyll *a* fluorescence, therefore being mainly observed at PSII. The phenomenon of quenching of chlorophyll *a* fluorescence is usually analysed in terms of three components, based on their relaxation kinetics: state-transitions (qT),  $\Delta$ pH-dependent quenching (qE) and photoinhibition (qI). The majority of NPQ is believed to occur through qE in the PSII antenna pigments bound to the LHCI proteins [5, 6]. State-transitions are considered to be the component of NPQ because the fluorescence yield of PSII diminishes due to the lateral redistribution of the phosphorylated LHCI proteins and their attachment to PSI [5]. Photoinhibition exhibits the slowest relaxation and it is the least defined. qI quenching is proposed to be involved in long-term down-regulation of PSII [4, 7].

### 2.1. $\Delta$ pH-dependent quenching

The process of qE is triggered by acidification of the thylakoid lumen under light-saturating conditions, which activates the interconversion of specific xanthophyll pigments (oxygenated carotenoids), that are mostly bound to the LHC proteins. For comparison, in cyanobacteria, strong blue-green or white light activates the orange carotenoid protein (OCP) which interacts with phycobilisome and dissipates the excess energy in the form of heat [8]. The xanthophyll cycle in plants, green and brown algae consists of the pH-dependent conversion from violaxanthin first to antheraxanthin and then to zeaxanthin. In plants, the reactions towards zeaxanthin are catalysed by the enzyme violaxanthin de-epoxidase, while the relatively slow reactions towards violaxanthin are catalysed by the enzyme zeaxanthin epoxidase [4]. The *npq1* mutants are unable to convert violaxanthin to zeaxanthin but still exhibit qE, demon-

strating that the xanthophyll cycle is not the prerequisite for qE formation [9]. Zeaxanthin was demonstrated to have an additional photoprotective function not connected with NPQ, but rather with thylakoid membrane lipids. It is hypothesised that the function of the non-protein bound zeaxanthin could be the removal of highly deleterious singlet oxygen, working together with the well-known antioxidant tocopherol [10]. The *npq4* mutant, which completely lacks the PsbS protein and qE, can survive under high-light conditions, thus implying that carotenoids may compensate to some extent the deficiency in qE formation [9].

Due to the slower kinetics of formation and relaxation of qE compared to the proton gradient, and taking into account that the light causes changes in charge distribution, which consequently alter the aggregation state of thylakoids, it was proposed that such a conformational change might accompany the qE event [6, 11]. Indeed, reports from at least three independent laboratories confirmed the structural rearrangement of the PSII-LHCII macro-organization during qE [12-14]. Time-correlated single photon counting (TCSPC) measurements revealed an additional far-red fluorescence component in the leaves of the high-light-adapted wild-type, the mutant unable to accumulate zeaxanthin at high-light (*npq1*), and in the mutant overexpressing PsbS, the protein proposed to act as a luminal pH sensor that consequently determines the level of qE [12, 15], respectively. The same component was not observed in the mutant devoid of PsbS, *npq4*, or in the dark-adapted state of abovementioned plant lines [12]. It was concluded that this fluorescence originated from the major LHCII antenna complex detached specifically from PSII, with the required presence of the PsbS protein [12]. Furthermore, it was biochemically demonstrated that the supramolecular complex B4C (Lhcb4, Lhcb6, and moderately bound LHC trimer) dissociates during light exposure, and associates back during dark period [13]. However, *npq4* mutants did not show light-induced B4C dissociation, establishing the role of PsbS as the key player in thylakoid rearrangements [13]. Finally, Ruban group used freeze-fracture electron microscopy to demonstrate that the formation of qE was indeed associated with the reorganisation of PSII and LHCII within the thylakoids [14]. Their experimental data support and update the 20-years-old hypothetical model built to explain the mechanistic of qE, the LHCII aggregation model [16]. According to the model, LHCII antenna could be found in four different states: (i) dark-adapted, unquenched; (ii) dark-adapted, partially aggregated, quenched; (iii) illuminated, partially aggregated, quenched; and (iv) illuminated, aggregated, fully quenched. Illumination causes two events necessary for LHCII aggregation to occur: conversion of violaxanthin to zeaxanthin and the protonation of LHCII, respectively [6]. Precisely, the formation of  $\Delta$ pH triggers a conformational change within the LHCII antenna, which leads to the partial dissociation of LHCII trimers from the PSII-LHCII supercomplexes, and, consequently, to their aggregation (Figure 1). In parallel, deoxidation of violaxanthin to zeaxanthin promotes the LHCII aggregation and the formation of NPQ [14]. The question for debate would be the number and the exact position of quenched complexes. According to Holzwarth group, two quenching centres are formed: detached and aggregated LHCII antenna measured within 1 to 5 minutes, and the minor components of LHCII; still attached to the PSII core, measured within 10 to 15 minutes [12]. As PSII core does not bind any zeaxanthin molecules, it is excluded as a quenching site; however, it is well-covered by formation of the second quenching centre.



**Figure 1.** Structural rearrangement of the PSII-LHCII macro-organization during qE. Composition and arrangement of the LHCII trimers (dark green) and the minor antenna (light green) around the PSII RCs (light grey) in the wild-type plant during the dark period (left panel) and under the high-light conditions (right panel) were presented according to [1]. Pink circles on the left panel denote violaxanthin, while the orange circles on the right panel denote the process of conversion of violaxanthin to zeaxanthin. The change of colour of the minor antenna from light green (left panel) to orange (right panel), also denotes the conversion of violaxanthin to zeaxanthin, respectively. According to [14], in excess light,  $\Delta\text{pH}$  triggers a conformational change within the LHCII antenna, which leads to the partial dissociation of LHCII trimers from the PSII LHCII supercomplexes, and, consequently, to their aggregation. In parallel, de-epoxidation of violaxanthin to zeaxanthin promotes the LHCII aggregation and the formation of NPQ (quenching site 1). Moreover, according to [12], quenching site 2 is also formed within the minor antenna still attached to the PSII.

Acidification of the thylakoid lumen also causes a conformational change in thylakoids that can be monitored at 535 nm ( $\Delta A_{535}$ ). This change is most likely induced by protonation of the lumen-exposed carboxylate side chains in specific PSII proteins [15]. Interestingly, *npq4* mutants, which do not express the PSII PsbS protein, also lack  $\Delta A_{535}$  [17]. It was proposed that PsbS is not necessary for efficient light harvesting and photosynthesis, but it is involved in NPQ by sensing luminal pH and consequently determining the level of qE [15]. Recently, it became clear that PsbS is indispensable for the physical state of the thylakoid membranes. In dark-adapted *npq4* mutants, the formation of ordered semi-crystalline arrays of PSII was increased, while in the plants over-expressing PsbS no arrays could be found, suggesting that PsbS disrupts the ordering and promotes the protein diffusion within the membrane, leading to the NPQ formation [18].

Although the main components for qE are known, the biophysical mechanism of de-excitation of the excited molecules of chlorophyll *a* is still unidentified. It is hypothesised that either xanthophylls act indirectly as allosteric regulators of the LHCs, causing the conformational change that facilitates the de-excitation, or xanthophylls directly de-excite



the excited molecules of chlorophyll *a* [4]. Models presented in Ruban and Holzwarth groups both permit the formation of internal dissipative pigment interactions, whether they occur between chlorophyll molecules or between carotenoid and chlorophyll molecules [19].

Variations in NPQ capacities and processes in different organisms suggest a strong evolutionary pressure to obtain optimized photoprotection [3]. For example, the LHCII proteins may have evolved from ancestors of contemporary stress-responsive proteins such as HLIP and ELIPS, which probably bind only carotenoids and are involved in photoprotection, while LHCII harvest light through high amounts of chlorophyll binding. In plants, PsbS binds pigments minimally and it is primarily involved in photoprotection, while the light-harvesting complex containing fucoxanthin (LHCF) in diatoms binds both chlorophyll and carotenoids in high amounts and contributes equally to the light harvesting and photoprotection. Under high-light conditions, the photosynthetic reaction centres can not accommodate all the electrons coming through electron transport chain, thus entering into the saturated, "closed" status. The already generated excitation energy becomes the burden for the photosynthetic membranes and has to be channeled safely, before destroying the reaction centres, especially the pigment core of PSII, the pair of the most potent oxidizers known to exist in nature, P680. If the vast energy is not diverted, P680 might rest in its prolonged oxidized state, P680<sup>+</sup>, and oxidize the neighbour protein amino-acids and pigments, ultimately leading to the destruction of the PSII protein D1. However, if P680 can not submit the electron to the oxidized plastoquinone, due to the increased number of already reduced plastoquinones, P680 could go into triplet state, interacting with atmospheric triplet oxygen and producing deleterious singlet oxygen. Both of these processes lead to the photoinhibition, the state in which the decreasing of the electron transport is observed [6]. Photoinhibition has a specific signature that can be monitored by a lower oxygen production, analysis of the D1 protein level, formation of uncoupled chlorophyll and of triplet state of chlorophyll. However, photoinhibition provokes the formation of photoprotection processes, where most of the unwanted energy would be harmlessly dissipated as heat. This phenomenon leaves a palpable trace visible as a drop in the fluorescence intensity, measured by TCSPC. One of the approaches to study the origin of the quenching mechanism *in vivo* is the treatment of Arabidopsis with lincomycin, which blocks the synthesis of chloroplast-encoded proteins, such as the reaction centres of PSI and PSII (RCI and RCII, respectively). The thylakoids of treated plants contain diminished amounts of RCs, but are rich in antenna complexes. Although the maximum chlorophyll fluorescence lifetime in isolated PSII-LHCII supercomplex is 4 ns; when complex is an integral part of the thylakoid membranes, fluorescence lifetime decreases to 2 ns [20]. If RCII, when saturated, contribute to the quenching of excitation of LHCII antenna, in the system almost devoid of RCII, the fluorescence lifetimes should be higher than 2 ns. However, spectroscopic measurements of the long-term lincomycin-treated plants did not differ from the control measurements, *i.e.*, the fluorescence lifetime was still 2 ns, suggesting that the LHCII antenna, and not the closed RCs, are sufficient for the quenching to occur [20].

## 2.2. How does the plant repair the damaged D1 protein?

The D1 repair cycle is regulated *via* reversible protein phosphorylation in thylakoid membranes. When plants are exposed to the high-light stress, the PSII protein subunits become heavily phosphorylated and migrate from grana to stroma thylakoids, this process being facilitated by the actions of STN8 [21]. The migration is accompanied by sequential dephosphorylation of the CP43, D2 and D1 proteins, respectively. Turnover of the D1 protein includes degradation of the photo-damaged polypeptide and co-translational insertion of the newly synthesized protein [22]. The current model envisions that the D1 protein is proteolytically processed at both sides of the thylakoid membrane: from the N-terminal end on the stromal side by the FtsH, a member of the ATPases associated with various cellular activities-subfamily (AAA subfamily), and, on the lumen site by Deg, a member of serine proteases family that does not require ATP [23]. Recently, it was confirmed that FtsH and Deg act in a cooperative manner to efficiently cut the D1 protein under photoinhibitory conditions [24]. According to the D1 digestion model, under all light intensities D1 is processed by the FtsH complexes, to be additionally supported by the Deg proteases under photoinhibitory conditions, in the so-called escape pathway [24]. The requirement of D1 dephosphorylation prior to its proteolysis could be explained by low affinity of phosphorylated N-terminal end of D1 towards the FtsH [23].

## 3. Plastoquinone pool-inflicted plant responses

The most promising redox-active components are the pool of plastoquinone (PQ) and the PSI acceptor site (*e.g.* NADPH, thioredoxin, glutathione and glutaredoxin). The PQ pool regulates two temporally distant responses that acclimate the photosynthetic process to the prevailing environment: state-transitions that occur in the order of minutes (short-term response) [25, 26, 27], and photosystem stoichiometry adjustment that requires hours to days (long-term response) [28]. Both responses occur under low-light conditions, in contrast to the high-light provoked responses such as NPQ, D1 repair cycle or various stress-response programmes [29].

### 3.1. State-transitions

State-transitions re-distribute excitation energy between two photosystems, which are electrochemically connected in series, through the supramolecular reorganisation of the photosynthetic membranes. The molecular complexes that cause the reorganisation are light-harvesting proteins, which collect light excitation and channel it to the reaction centres. Already mentioned LHCI is the prototype of large and abundant class of chloroplast trans-membrane proteins that binds roughly half of the total chlorophyll in chloroplasts [2]. The absorption spectra of LHCI is in the far red region owing to the chlorophyll *a* enrichment, while LHCI is enriched in chlorophyll *b* and has the maximum absorption at shorter wavelengths in the red region, around 650 nm [26]. Under PQ reducing conditions, *i.e.* when PSII is

predominantly excited compared to PSI, a redox-sensitive protein kinase acts to phosphorylate the apoproteins of LHCII. Upon phosphorylation, LHCII partially dissociates from PSII and associates with PSI (state I). Under PQ oxidizing conditions, *i.e.* when PSI is predominantly excited compared to PSII, the kinase is inactive, but an activated phosphatase dephosphorylates the mobile LHCII, which moves laterally, and associates with PSII (state II) [29]. Although state-transitions occur only under low-light conditions, they share the same mechanism of LHCII aggregation with the high-light response mechanism, at least in *Chlamydomonas reinhardtii* [30]. This opens an intriguing question of plant capacity to wisely use the same mechanism for different living conditions. In 2003, Jean-David Rochaix group proposed that the thylakoid-associated serine-threonine Stt7 kinase from the alga *C. reinhardtii* is involved in phosphorylation of LHCII and in state-transitions [31]. In 2005, the same group proposed that the homolog of the Stt7 in Arabidopsis, STN7, is involved in state-transitions [32]. Since LHCII phosphorylation is reversible process, an extensive search has been conducted to identify the protein phosphatase(s) that dephosphorylates LHCII. Recently, two groups described the product of the nuclear gene At4g27800 as a long-sought plastid protein phosphatase specifically involved in dephosphorylation of the mobile pool of major LHCII proteins, titled PPH1 [33], and TAP38 [34], respectively. The specificity of PPH1/TAP38 for LHCII supports the hypothesis that several phosphatases must be involved in dephosphorylation of thylakoid phosphoproteins [33, 34]. Recently, a new phosphatase PBCP, capable of *in vivo* dephosphorylation of PSII core subunits CP43, D1, D2 and PsbH, respectively, was identified [35]. It seems that the PBCP phosphatase targets at proteins phosphorylated by the protein kinase STN8 [36], thus forming a pair with opposing effects on phosphorylation of the photosynthetic core proteins [35].

### 3.2. Stoichiometry adjustments

Photosystem stoichiometry adjustment is a long-term response that affects the relative amounts of the two photosystems by changing the expression of photosynthetic genes both in the chloroplast and in the nucleus [29]. It was shown that the redox state of the PQ pool serves as a major signal in the regulation of LHCII photo-acclimation [37]. In order to avoid excess excitation energy and to minimize oxidative damage, the LHCII protein level decreased by approximately 50% in *Lemna perusilla* upon transfer from low-light to high-light conditions [37]. In contrast to the state-transitions, where active redox-regulated thylakoid-associated STN7 kinase phosphorylates LHCII, thus leading to LHCII detachment from PSII and migration towards stroma thylakoids and PSI, under high-light STN7 is inactive, hence the rate of the phosphorylated LHCII decreases and the excess LHCII undergoes proteolytic degradation. The same stands for the PSI and PSII genes: upon reduction of the PQ pool, the expression of the PSI genes is favoured, while upon oxidation of the PQ pool, the expression of the PSII is favoured [29]. In 2005, the coupling of the long-term response of adjusting photosystem stoichiometry and the short-term response of state-transitions by LHCII kinase STN7 was proposed [36].



## 4. Immunophilins in photosynthesis

Immunophilins comprise a superfamily of conserved ubiquitous proteins consisting of two distinct subfamilies; cyclophilins and FKBP (FK506/rapamycin-binding proteins), the targets of the immunosuppressive drugs cyclosporine A (CsA) and FK506/rapamycin, respectively [38]. Despite the lack of structural similarity, all cyclophilins and FKBP share a common enzymatic, so-called PPIase or rotamase activity, catalyzing *cis-trans* isomerization of proline-imidic peptide bonds [39, 40]; (Rotation around peptide bonds is energetically disfavored due to their partial double-bond character. Delocalization of amide nitrogen electrons results in approx. 22 kcal/mol energy barrier to rotation and restrains the peptide bond in either *cis* or *trans* configurations). A more recently discovered third group of proteins (parvulins), which is insensitive to immunosuppressive drugs, also possesses PPIase activity [41]. The molecular masses of cyclophilins, FKBP, and parvulins range normally in the order of 18-21 kDa, 12-13 kDa, and 10-13 kDa, respectively, but several complex immunophilins of higher molecular weight have been detected recently [42, 43, 41]. The biological significance of this group of enzymes is a matter of intense research. Acceleration of protein folding processes by PPIase *in vitro* [44] and *in vivo* [45] has supported a physiological role as folding catalysts facilitating the slow and generally rate-limiting uncatalysed isomerisation around Xaa-Pro peptide bonds. Immunophilins can also perform chaperone functions [46], or cooperate with other chaperone proteins [47, 48, 49].

Complex immunophilins all contain additional protein-protein interaction domains, such as leucine-zipper motifs and/or tetratricopeptide repeat domains. They may be constituents of supramolecular structures, as shown for the human or avian steroid receptor complex [48] or chaperone supercomplexes [50], and may be involved in hsp90-dependent signal transduction [49, 51, 52]. Various members of the immunophilins are involved in phosphorylation processes *via* transient interaction with kinases [53, 51, 52] and phosphatases [54, 55]. It seems likely that, at least for the components of MAP kinase signaling system (Src, Raf, and Mek), hsp90-immunophilin interactions are essential for the kinase regulation [see 52]. The immunophilin FKBP65 together with hsp90 forms a regulatory association with the serine/threonine kinase c-Raf-1 [51]. In mammals, the Ca<sup>2+</sup>/calmodulin-dependent heterodimeric protein phosphatase calcineurin can bind immunophilins with complex immunosuppressive drugs. This interaction inhibits the protein phosphatase activity, resulting in interruption of the signal-transduction cascades required for T-cell activation [see 54]. A unique immunophilin-related protein is the PP5 protein phosphatase. This enzyme is a major constituent of the glucocorticoid receptor-hsp90 complex, with properties of an FK506-binding immunophilin with low affinity FK506 binding activity [56].

Plants possess a substantial number of immunophilins which are localized in cytosol, chloroplasts, nucleus, mitochondria, or associated with secretory pathways. The majority of the cyclophilins are single-domain proteins, 23 isoforms in Arabidopsis [57]. AtCYP20-3, AtCYP20-2, AtCYP26-2, AtCYP28 and AtCYP37 can be found in chloroplasts, mostly in thylakoid lumen. The multidomain cyclophilin isoforms possess unique domain arrangements, as exemplified by AtCYP38. AtCYP40 contains a C-terminal tetratricopeptide repeat module

(TPR). Four additional multidomain cyclophilins contain RNA interaction domains, which implicates their involvement in nuclear RNA processing machinery.

Plant FKBP family encompasses also 23 members in Arabidopsis and this family is one of the largest FKBP family identified to date. These FKBP family members can also be divided into single and multidomain isoforms, consisting of 16 and 7 members, respectively [58]. Interestingly, 11 single-domain FKBP family members appear to be targeted to the thylakoid lumen. Thus, this chloroplast sub-compartment appears to have very important role in immunophilin function, or the processes taking place in lumen require specific activity of this protein family.

The dephosphorylation of D1, D2 and CP43 in spinach is catalysed by a cyclophilin-regulated PP2A-like protein phosphatase, which was found to be associated with, and regulated by, a cyclophilin-like protein, TLP40 [59]. TLP40 is proposed to suppress phosphatase activity, when bound to the lumen-exposed epitope of the protein phosphatase, but to induce its activity, when released to the lumen [59]. However, it remains to be seen if the Arabidopsis PP2C phosphatase PBCP [35] is also under the control of TLP40. Vener group demonstrated that D1 was not only vulnerable to light, but also to high temperature [60]. Raising the temperature from 22 °C to 42 °C resulted in a very rapid dephosphorylation of the D1, D2 and CP43 proteins and in release of TLP40 from membrane into the thylakoid lumen. These events are proposed to trigger an accelerated repair of photodamaged PSII and to initiate other heat-shock responses in chloroplasts.

Higher plant thylakoid lumen prolineisomerases [38], or complex immunophilins, are found to be regulated by light and are responsive to various forms of environmental stress [61]. The structure of TLP40 and its association with the thylakoid membrane system implicates diverse functions and involvement in the intracellular signaling networks [62]. Binding of thylakoid membrane associated phosphatase involved in dephosphorylation of PSII core proteins might occur *via* two putative phosphatase-binding modules on the N-terminal side of TLP40 [59, 60]. In 2002, Baena-González and Aro [63] and in 2005 Aro et al. [64] further suggested that damaged D1 repair cycle includes TLP40 ortholog *AtCYP38*, which lacks peptidyl-prolyl *cis/trans* isomerase (PPIase) activity [65, 66, 67]. It was shown that *AtCYP38* is involved in the assembly of oxygen evolving complex (OEC) [68] and maintenance of PSII [69]. Most recently, the interaction of the E-loop of chlorophyll protein 47 (CP47) with *AtCYP38* was demonstrated [67]. This interaction is mediated through putative cyclophilin domain [67]. Further, *in vivo* role of *AtCYP38* has been investigated in *cyp38* mutant Arabidopsis plants [68, 69, 70], suggesting its primary role in PSII biogenesis and repair.

## 5. Flow and partitioning of photosynthetic electrons

Photosynthetic apparatus has to be able to efficiently convert energy at low light and to avoid over-reduction and damage at excess light. This requires switching between different regulatory mechanisms which keep the cellular ATP pool almost at the constant level. Exposure of plants to higher light intensities than required for efficient photosynthesis results in saturation of photosynthetic electron transport (PET). The over-reduction of PET chain can lead to the

formation of reactive oxygen species and may irreversibly damage the photosystems, as well as the cells and the whole organism [71, 72, 73]. To counteract and reduce the photoinhibitory damage, plants have developed several short- and long-term regulation mechanisms which include processes that modulate the structure and function of antenna complexes, including NPQ, alternative electron transport pathways and movement of chloroplasts, leaves and whole organisms away from intense light [74, 75]. Dissipation of excess light energy to heat in the antenna or in the reaction centre of PSII also counteracts the photoinhibitory damage. Retrograde signaling transduces information on the metabolic state of the organelle and induces many activities in the cytosol, nucleus and mitochondria, inducing alterations in nuclear gene expression of organelle-targeted proteins [76].

### 5.1. Linear, cyclic and pseudo-cyclic electron transfer routes

Various photosynthetic electron transfer routes become turned on and off, according to the need of the plant to adapt to wide-ranging quantities and qualities of light. Three major electron transfer pathways known are linear, cyclic and pseudo-cyclic electron transfer (LEF, CEF and PCEF, respectively). All three pathways are necessary for poised and sustained synthesis of ATP and NADPH and their interplay enables the flexibility of photosynthesis in meeting different metabolic demands [77].

During non-cyclic electron transport or LEF, light drives the conversion of water to oxygen at the level of oxygen-evolving complex of PSII and  $\text{NADP}^+$  to NADPH on the stromal side of thylakoid membranes. The PET chain consists of PSII, the *Cytb6f* complex, PSI, and the free electron carriers plastoquinone (PQ) and plastocyanin (PC). Electron transport includes the two quinone binding sites and two cytochromes b6 (the 'Q cycle') that give two protons translocated for each electron transferred from PSII to PSI. Hydrogen ions are transferred across chloroplast membrane and accumulated on the luminal side of the thylakoids, where they drive ATP synthesis through a membrane ATPase. This way electron transport helps to establish a proton gradient that powers ATP production and also stores energy in the reduced coenzyme NADPH to power the Calvin-Benson cycle to produce sugar and other carbohydrates.

Arnon, who discovered photophosphorylation in isolated chloroplasts in 1954, demonstrated that there is also CEF in the thylakoids, driven solely by PSI. CEF is a light-driven flow of electrons through a photosynthetic reaction centre with the electrons being transferred from PSI to *Cytb6f* complex *via* ferredoxin (Fd), with associated formation of proton gradient. PQ is reduced by Fd or NADPH *via* one or more enzymes collectively called PQ reductase, rather than by PSII, as in LEF. From hydroplastoquinone (PQH<sub>2</sub>), electrons return to PSI *via* the *Cytb6f* complex. Four possible routes of CEF that may operate in parallel have been proposed so far: NAD(P)H dehydrogenase (NDH)-dependent route, Fd-dependent route, Nda2, a type 2 NAD(P)H:PQ oxidoreductase route and *Cytb6f* complex and FNR route [recently reviewed by 78].

CEF around PSI occurs under conditions when acceptor limitation or ATP shortage results in a highly reduced PQ pool and contributes to the formation and maintenance of a pH gradient

across a membrane but does not produce NADPH. The pH gradient generated may drive the production of ATP (cyclic photophosphorylation) or may regulate photosynthesis. When more light is absorbed than can be used for assimilation, the increased  $\Delta\text{pH}$  is a switch to dissipate excess of the light absorbed by chlorophyll molecules of PSII [79]. CEF is diminished when its components are completely reduced. Also, there is no CEF when its components are completely oxidized because there are no electrons to cycle [80]. In an attempt to avoid these two extreme situations, kinetics, post-translational modifications [25, 26, 81] and redox control of reaction-centre gene expression [82] are all employed in maintaining a poised PQ pool. In spite of these control mechanisms, over-reduction easily occurs when the Calvin-Benson cycle is unable to use NADPH, usually due to the lack of ATP. The major physiological significance of CET most probably lies in additional availability of ATP.

LEF in chloroplasts produces a number of reduced components associated with PSI that may subsequently participate in reactions that reduce oxygen. When Fd transfers electrons to molecular oxygen instead of  $\text{NADP}^+$ , PCEF linked with phosphorylation arises.  $\text{O}_2$  is directly reduced to superoxide radical in so-called Mehler reaction. Subsequently, two superoxides dismutate to form  $\text{H}_2\text{O}_2$  and  $\text{O}_2$ , which are by further redox processes converted to water. A reduced state of the FeS pool (Fd and PSI centres) promotes the Mehler reaction. The Mehler reaction leading to PCEF restores the redox poise when the PET chain is over-reduced, thereby allowing CEF to function and to generate ATP for the Calvin-Benson cycle, which will in turn oxidize NADPH and restore LEF [83]. Under high light PCEF could also cover an increased energy demand, as long as the antioxidant systems for  $\text{H}_2\text{O}_2$  removal is sufficiently active (ascorbate and GSH recycling).

A great number of contemporary research topics on photosynthesis aims at elucidating novel, alternative electron transfer routes as the safest pathways for channelling of unwanted electrons. Chlororespiration is a well-known respiratory process that can maintain a trans-thylakoid proton gradient, thus acting as an effective alternative electron sink in preventing over-reduction of the PQ pool and protecting RCIIIs from photodamage under photoinhibitory light conditions [84]. Two enzymes are important for chlororespiratory function: NADH dehydrogenase complex and nucleus-encoded plastid-localized terminal oxidase (PTOX), through which electrons from plastoquinol are transferred to molecular oxygen, forming water in the stroma [85]. In oat leaves incubated at high temperature and under high-light intensities, the amounts of both enzymes were increased, suggesting that, under unfavourable conditions, chlororespiration can act as a protective mechanism [85].

Apart from the Mehler reaction and the chlororespiration, respectively, photorespiration is another efficient mechanism to adjust the ATP/NADPH ratio and to consume excess energy [86]. Although the major part of energy supplied in the form of ATP and NADPH by the light reaction is consumed in the Calvin-Benson cycle, it is also needed for multiple anabolic processes in chloroplasts, such as synthesis of lipids and proteins and of many secondary metabolites. ATP/NADPH ratio in chloroplasts could be increased by indirect export of reducing equivalents in the form of malate through the malate valve [87]. Malate can be used in numerous ways in cytoplasm and mitochondria, providing NADH for nitrate reduction and/or ATP for sucrose synthesis. Also, different stages of tissue growth require different ATP/



NADPH ratios. The ATP/NADPH ratio that is available in the light can vary substantially because of the many possibilities for electron pathways and regulatory mechanisms [88]. Therefore, the cooperation of different electron transport pathways enables optimization of ATP/NADPH stoichiometry [77]. In example, sudden dark-to-light transition induces over-reduction of the PET chain in the first minute, which is relieved by electron transport to O<sub>2</sub> [89], and by the rapid activation of the chloroplast NADP-MDH [90]. Calculating the photosynthetic stoichiometries, it was estimated that ATP/NADPH ratio arising from LEF is about 1.28, which is not sufficient for driving the Calvin-Benson cycle [91, 77]. It is therefore obvious that the optimal operation of the Calvin-Benson cycle requires both LEF and CEF, whose tuning enables adjustment of ATP/NADPH to meet the cellular demands. Green alga *C. reinhardtii* in which intracellular ATP depletion induces a switch from LEF to CEF is a good example for such regulation [92].

## 5.2. Electron partitioning at the ferredoxin hub

In photoautotrophic plants, ferredoxin (Fd) accepts one electron from the stromal side of PSI involving the subunits PsaC, PsaD and PsaE [93]. Fd acts simultaneously as bottleneck and as a hub which distributes high-energy electrons to a multitude of enzymes involved in chloroplast metabolism. The hierarchy of electron distribution and subsequent regulation of channeling of photosynthetically derived electrons into different areas of chloroplast metabolism is still not defined. Electrons are preferentially directed to carbon assimilation, which requires NADPH, and so the majority of Fd is immediately oxidized by the enzyme ferredoxin:NADPH reductase (FNR), which is associated with the thylakoid membrane.

Besides its crucial metabolic role in reducing NADP<sup>+</sup> and thioredoxin (TRX) *via* FNR and ferredoxin-thioredoxin-reductase (FTR) respectively, Fd-dependent enzymatic reactions are also linked to nitrite and sulfur metabolism by ferredoxin-nitrite-reductase and sulfite reductase. Furthermore, Fd is a electron donor for fatty acid desaturase and glutamine-2-oxoglutarate amino transferase. In some green algae, upon transition from dark to the light under anaerobic conditions, Fd transfers transiently electrons to chloroplast hydrogenases, which in turn catalyse the formation of hydrogen [94, 95] dissipating excess reducing power when the Calvin-Benson cycle is not yet fully activated. Fd and NADPH can also act in CEF, returning electrons to the PQ pool *via* the PGR5 (proton gradient regulation 5)-dependent [96] and NDH complex-dependent [97, 98, 99] pathways, respectively. FNR may also act as the direct Fd:PQ reductase, establishing redox regulation and antioxidant defense point. FNR activity seems to represent a critical point in photosynthetic electron partitioning, because it is integral to most of these electron cascades and can associate with several different membrane complexes.

At least four Fd isoforms occur in plants [100]. Fd1 and Fd2 are found in leaves, while Fd3 and Fd4 appear to play roles in non-photosynthetic metabolism [101, 100]. Fd as electron distributing hub is particularly suitable to provide information on the redox state of the system to be transmitted into the regulatory network. Fd contributes to the control of chloroplast energization by feeding electrons into the CEF pathway and thereby controls both phosphorylation potential and reductive power.



### 5.3. TROL-FNR interaction influences the energy conductance

The last step of the photosynthetic electron transfer from Fd to NADP<sup>+</sup> is catalyzed by FNR. There are two evolutionary conserved types of FNR in the chloroplasts of higher plants: predominantly, or exclusively, thylakoid membrane-bound isoproteins and 'soluble', non-tightly bound isoproteins [101, 102]. The membrane-bound FNR is supposed to be involved in electron transport, while the soluble enzyme provides protection against oxidative stress [103]. Two chloroplast-type *FNR* genes have been found in Arabidopsis genome. In 2008, Hanke et al. investigated knock-out mutant of Arabidopsis FNR isoprotein, *fnr1*. The loss of the strong thylakoid binding was observed, which affected the channeling of photosynthetic electrons into NADPH- and Fd-dependent metabolism. Also, these mutants had complex variation in CEF, dependent on light conditions [104]. In *fnr1*, thylakoid NADP<sup>+</sup> photoreduction was greatly reduced even on addition of soluble FNR to rate saturating concentrations, which is consistent with the lack of membrane-bound FNR [104].

NADP<sup>+</sup> photoreduction activity of FNR was shown to be greater when the enzyme is associated with the thylakoid membrane and it has been proposed that binding of FNR to the thylakoid membrane regulates the enzyme activity [105, 106]. Subsequently, interactions of FNR and several photosynthetic protein complexes, such as *Cytb6f* [107, 108], PSI [109] or NDH complex [110] have been shown. However, the factors controlling relative localization of FNR to different membrane complexes have not yet been established. It was shown that maize contains three chloroplast FNR proteins with completely different membrane association and distribution between cells, conducting predominantly CEF in bundle sheath cells and LEF in mesophyll cells [111]. Expression of maize FNRs in Arabidopsis as chimeras and truncated proteins showed that N-terminus determines recruitment of FNR to different membrane complexes, which impacts the photosynthetic electron flow [111].

It was also demonstrated that FNR interacts specifically with two chloroplast proteins, Tic62 (62 kDa component of the translocon at the inner envelope of chloroplasts) and TROL (thylakoid rhodanese-like protein), *via* a conserved Ser/Pro-rich motif [112, 113]. Both Tic62 and TROL seem to act as molecular anchors for FNR, because they form high molecular weight complexes with FNR at the thylakoid membranes. TROL possesses centrally positioned rhodanese-like domain, which is most probably involved in redox regulation of FNR binding and release [113]. We have proposed that such regulation could be important for balancing the redox status of stroma with the membrane electron transfer chain and therefore preventing the over-reduction of any of these two compartments and maintaining the redox poise [113]. TROL-FNR complex was clearly visible during the dark and it disappeared during light periods [112]. FNR-Tic62/TROL interaction is clearly pH-dependent. During high photosynthetic activity in the light, stroma becomes alkaline due to the transport of protons to the thylakoidal lumen. During the dark, stromal pH decreases again, and that is when FNR-Tic62/TROL complexes were found predominantly associated with the thylakoid membrane [112, 113]. The mechanism by which TROL influences the FNR activity could be that during the dark period, FNR is bound to the thylakoids *via* TROL and NADPH production does not occur. This stage could be sustained through the binding of small molecule, possibly oxidized PQ, to the RHO cavity. In conditions of growth-light, FNR is bound to the thylakoids *via* TROL and efficiently

produces NADPH. When the light is saturating, FNR is released from TROL by a signal molecule, possibly reduced PQ that competes for the RHO binding site. Once soluble, FNR acts as NADPH consumer and released protons are passed to an unknown scavenger [78, 113, 114]. It was reported that membrane attachment of FNR is influenced by the stromal redox state (NADP<sup>+</sup>/NADPH ratio), which mimics variations in environmental conditions [115]. Therefore, reversible attachment of FNR to the thylakoid membrane *via* TROL and/or Tic62 provides an elegant way to store redundant molecules, not required when photosynthesis is less active or dormant.

We have already mentioned that reducing equivalents could be exported in the form of malate through the malate valve to increase the ATP/NADPH ratio in chloroplasts [87]. TROL-deficient plants grown under growth-light conditions show significant up-regulation of NADP-malic enzyme 2 that catalyses the oxidative decarboxylation of malate, producing pyruvate, carbon dioxide and NAD(P)H in cytosol [113]. Therefore, the TROL knock-out Arabidopsis mutant lines (*trol*) could act as efficient NADPH producers, fighting the possible hyper-reduction of the thylakoids by exporting the reducing energy in a form of malate to the cytosol.

The *trol* mutants show severely lowered relative electron transport rates at high-light intensities. Also, under high-light conditions, but in a short-term, NPQ amount was higher in the *trol* line, compared to the wild-type [113]. Moreover, TROL is important for NPQ, since the reversion of the plant without TROL to the plant that expresses TROL, even with certain alterations, leads to the restoration of wild-type levels of NPQ (Jurić, Fulgosi, Ruban, unpublished results). We are not dealing with extreme NPQ phenotype, but, the question remains, how is it possible that small changes in the TROL protein could modulate NPQ so effectively? Our unpublished data also suggest the possible enrolment of some LHCII subunits in the TROL-containing complexes, which, in the light of these results, should be thoroughly investigated. In addition, we observed that, after more than two weeks of exposure to the high-light conditions, *trol* plants exhibited NPQ almost at the wild-type levels, suggesting that some sort of long-term acclimation could also be involved.

## 6. Conclusions and perspectives

Photosynthesis is the crucial converter of sunlight into the chemical energy that is subsequently utilised to sustain life on Earth. However, without the layers of regulative pathways, it would be virtually impossible to discuss the efficient photosynthesis. Now, we are aware of the fact that more than 40-years-old “Z-scheme” has been constantly upgraded with many alternative routes allied with still not well-characterised electron sources and sinks. In addition, it seems that the thylakoid membranes are more flexible than anticipated, with PSII antenna involved directly in NPQ origin through detachment and aggregation.

One of the interesting domains in photosynthesis, but not enough explored, would be the involvement of lipids in a complex network of protein interactions. Lipids, especially phospholipids (phosphatidylglycerol; PG) and glycolipids (monogalactosyldiacylglycerol; MGDG,

digalactosyldiacylglycerol; DGDG, and sulfoquinovosyldiacylglycerol; SQDG) are major building blocks of thylakoid membranes, providing the safe docking sites for proteins and protein complexes, respectively. It has been proposed that DGDG and PG are especially involved in the assembly and repair of PSII [116]. Moreover, plants lacking almost 30% of the wild-type amount of PG, caused by the point mutation in one of the genes involved in PG synthesis pathway, displayed pale green leaves and somewhat reduced capacity for photosynthesis [117], while plants that accumulated only 10% of the wild-type PG amount were incapable of surviving on the growth medium without the addition of sucrose [118].

The other attractive domain would be retrograde signalling, *i.e.* the pathways flowing from chloroplasts to the nucleus. At least five classes of chloroplast-derived signals have been studied and discussed: tetrapyrrole biosynthesis, chloroplast gene expression-dependent pathway, redox state of chloroplasts, ROS, and sugar and hormone signalling. While some molecules look more promising than others (sugars and hormones for example), ROS could be more challenging to investigate, due to its non-specificity and a power to respond quickly to different types of stress in different parts of the cell.

The rich past of exploring photosynthetic processes gave us a wealth of knowledge, but, it seems that we have just scratched the surface. In years to come, it would be interesting to fill the missing blanks and to develop some new, “out-of-the-box” perspectives.

## Author details

Snježana Jurić, Lea Vojta and Hrvoje Fulgosi\*

Division of Molecular Biology, Ruđer Bošković Institute, Zagreb, Croatia

## References

- [1] Dekker JP, Boekema EJ. Supramolecular organization of thylakoid membrane proteins in green plants. *Biochim Biophys Acta*. 2005;1706(1-2):12-39.
- [2] Barros T, Kühlbrandt W. Crystallisation, structure and function of plant light-harvesting Complex II. *Biochim Biophys Acta*. 2009;1787(6):753-772.
- [3] Horton P, Ruban A. Molecular design of the photosystem II light-harvesting antenna: photosynthesis and photoprotection. *J Exp Bot*. 2005;56(411):365-373.
- [4] Müller P, Li XP, Niyogi KK. Non-photochemical quenching. A response to excess light energy. *Plant Physiol*. 2001;125(4): 1558-1566.

- [5] Szabó I, Bergantino E, Giacometti GM. Light and oxygenic photosynthesis: energy dissipation as a protection mechanism against photo-oxidation. *EMBO Rep.* 2005;6(7):629-634.
- [6] Ruban AV, Johnson MP, Duffy CD. The photoprotective molecular switch in the photosystem II antenna. *Biochim Biophys Acta.* 2012;1817(1):167-181.
- [7] Eberhard S, Finazzi G, Wollman FA. The dynamics of photosynthesis. *Annu Rev Genet.* 2008; 42:463-515.
- [8] Kirilovsky D, Kerfeld CA. The orange carotenoid protein in photoprotection of photosystem II in cyanobacteria. *Biochim Biophys Acta.* 2012;1817(1):158-166.
- [9] Xiao FG, Shen L, Ji HF. On photoprotective mechanisms of carotenoids in light harvesting complex. *Biochem Biophys Res Commun.* 2011;414(1):1-4.
- [10] Jahns P, Holzwarth AR. The role of the xanthophyll cycle and of lutein in photoprotection of photosystem II. *Biochim Biophys Acta.* 2012;1817(1):182-193.
- [11] Barber J. Influence of surface charges on thylakoid structure and function. *Annu Rev Plant Physiol.* 1982;33:261-295.
- [12] Holzwarth AR, Miloslavina Y, Nilkens M, Jahns P. Identification of two quenching sites active in the regulation of photosynthetic light-harvesting studied by time-resolved fluorescence. *Chem Phys Lett.* 2009;483(4-6):262-267.
- [13] Betterle N, Ballottari M, Zorzan S, de Bianchi S, Cazzaniga S, Dall'osto L, Morosinotto T, Bassi R. Light-induced dissociation of an antenna hetero-oligomer is needed for non-photochemical quenching induction. *J Biol Chem.* 2009;284(22):15255-15266.
- [14] Johnson MP, Goral TK, Duffy CD, Brain AP, Mullineaux CW, Ruban AV. Photoprotective energy dissipation involves the reorganization of photosystem II light-harvesting complexes in the grana membranes of spinach chloroplasts. *Plant Cell.* 2011;23(4):1468-1479.
- [15] Li Z, Wakao S, Fischer BB, Niyogi KK. Sensing and responding to excess light. *Annu Rev Plant Biol.* 2009;60:239-260.
- [16] Horton P, Ruban AV, Rees D, Pascal AA, Noctor G, Young AJ. Control of the light-harvesting function of chloroplast membranes by aggregation of the LHClI chlorophyll-protein complex. *FEBS Lett.* 1991;292(1-2):1-4.
- [17] Li XP, Björkman O, Shih C, Grossman AR, Rosenquist M, Jansson S, Niyogi KK. A pigment-binding protein essential for regulation of photosynthetic light harvesting. *Nature.* 2000;403(6768):391-395.
- [18] Goral TK, Johnson MP, Duffy CD, Brain AP, Ruban AV, Mullineaux CW. Light-harvesting antenna composition controls the macrostructure and dynamics of thylakoid membranes in *Arabidopsis*. *Plant J.* 2012;69(2):289-301.

- [19] Ahn TK, Avenson TJ, Ballottari M, Cheng YC, Niyogi KK, Bassi R, Fleming GR. Architecture of a charge-transfer state regulating light harvesting in a plant antenna protein. *Science*. 2008;320(5877):794-797.
- [20] Belgio E, Johnson MP, Jurić S, Ruban AV. Higher plant photosystem II light-harvesting antenna, not the reaction center, determines the excited-state lifetime-both the maximum and the nonphotochemically quenched. *Biophys J*. 2012;102(12):2761-2771.
- [21] Tikkanen M, Nurmi M, Kangasjärvi S, Aro EM. Core protein phosphorylation facilitates the repair of photodamaged photosystem II at high light. *Biochim Biophys Acta*. 2008;1777(11):1432-1437.
- [22] Vener AV. Environmentally modulated phosphorylation and dynamics of proteins in photosynthetic membranes. *Biochim Biophys Acta*. 2007;1767(6):449-457.
- [23] Kato Y, Sakamoto W. Protein quality control in chloroplasts: a current model of D1 protein degradation in the photosystem II repair cycle. *J Biochem*. 2009;146(4):463-469.
- [24] Kato Y, Sun X, Zhang L, Sakamoto W. Cooperative D1 Degradation in the Photosystem II Repair Mediated by Chloroplastic Proteases in Arabidopsis. *Plant Physiol*. 2012;159(4):1428-1439.
- [25] Allen JF, Forsberg J. Molecular recognition in thylakoid structure and function. *Trends Plant Sci*. 2001;6(7):317-326.
- [26] Wollman FA. State-transitions reveal the dynamics and flexibility of the photosynthetic apparatus. *EMBO J*. 2001;20(14):3623-3630.
- [27] Mullineaux CW, Emlyn-Jones D. State-transitions: an example of acclimation to low-light stress. *J Exp Bot*. 2004;56(411):389-393.
- [28] Allen JF, Pfannschmidt T. Balancing the two photosystems: photosynthetic electron transfer governs transcription of reaction centre genes in chloroplasts. *Philos Trans R Soc Lond B Biol Sci*. 2000; 355(1402):1351-1359.
- [29] Pfannschmidt T, Bräutigam K, Wagner R, Dietzel L, Schröter Y, Steiner S, Nykytenko A. Potential regulation of gene expression in photosynthetic cells by redox and energy state: approaches towards better understanding. *Ann Bot*. 2009;103(4):599-607.
- [30] Iwai M, Yokono M, Inada N, Minagawa J. Live-cell imaging of photosystem II antenna dissociation during state transitions. *Proc Natl Acad Sci USA*. 2010;107(5):2337-2342.
- [31] Depège N, Bellafiore S, Rochaix JD. Role of chloroplast protein kinase Stt7 in LHCII phosphorylation and state transition in *Chlamydomonas*. *Science*. 2003;299(5612):1572-1575.
- [32] Bellafiore S, Barneche F, Peltier G, Rochaix JD. State-transitions and light adaptation require chloroplast thylakoid protein kinase STN7. *Nature*. 2005;433(7028):892-895.



- [33] Shapiguzov A, Ingelsson B, Samol I, Andres C, Kessler F, Rochaix JD, Vener AV, Goldschmidt-Clermont M. The PPH1 phosphatase is specifically involved in LHCII dephosphorylation and state-transitions in Arabidopsis. *Proc Natl Acad Sci USA*. 2010;107(10):4782-4787.
- [34] Pribil M, Pesaresi P, Hertle A, Barbato R, Leister D. Role of plastid protein phosphatase TAP38 in LHCII dephosphorylation and thylakoid electron flow. *PLoS Biol*. 2010;8(1):e1000288.
- [35] Samol I, Shapiguzov A, Ingelsson B, Fucile G, Crèvecoeur M, Vener AV, Rochaix JD, Goldschmidt-Clermont M. Identification of a Photosystem II Phosphatase Involved in Light Acclimation in Arabidopsis. *Plant Cell*. 2012;24(6):2596-2609.
- [36] Bonardi V, Pesaresi P, Becker T, Schleiff E, Wagner R, Pfannschmidt T, Jahns P, Leister D. Photosystem II core phosphorylation and photosynthetic acclimation require two different protein kinases. *Nature*. 2005;437(7062):1179-1182.
- [37] Oelze ML, Kandlbinder A, Dietz KJ. Redox regulation and overreduction control in the photosynthesizing cell: complexity in redox regulatory networks. *Biochim Biophys Acta*. 2008;1780(11):1261-1272.
- [38] Schreiber SL. Chemistry and biology of the immunophilins and their immunosuppressive ligands. *Science*. 1991;251 (4991):283-287.
- [39] Fischer G, Wittmann-Liebold B, Lang K, Kiefhaber T, Schmid FX. Cyclophilin and peptidyl-prolyl *cis-trans* isomerase are probably identical proteins. *Nature*. 1989;337(6206):476-478.
- [40] Takahashi N, Hayano T, Suzuki M. Peptidyl-prolyl *cis-trans* isomerase is the cyclosporin A-binding protein cyclophilin. *Nature*. 1989;337(6206):473-475.
- [41] Rudd KE, Sofia HJ, Koonin EV, Plunkett GI, Lazar S, Rouviere PE. A new family of peptidyl-prolyl isomerases. *Trends Biochem Sci*. 1995;20(1):12-14.
- [42] Gething MJ, Sambrook J. Protein folding in the cell. *Nature*. 1992;355(6355):33-45.
- [43] Kieffer LJ, Seng TW, Li W, Osterman DG, Handschumacher RE, Bayney R. Cyclophilin-40, a protein with homology to the P59 component of the steroid receptor complex. Cloning of the cDNA and further characterization. *J Biol Chem*. 1993;268(17):12303-12310.
- [44] Fruman DA, Burakoff SJ, Bierer BE. Immunophilins in protein folding and immunosuppression. *FASEB J*. 1994;8(6):391-400.
- [45] Matouschek A, Rospert S, Schmid K, Click BS, Schatz G. Cyclophilin catalyzes protein folding in yeast mitochondria. *Proc Natl Acad Sci USA*. 1995;92(14):6319-6323.
- [46] Freskgård PO, Bergenhem N, Jonson BH, Svensson M, Carlsson U. Isomerase and chaperone activity of prolyl isomerase in folding of carbonic anhydrase. *Science*. 1992;258(5081):466-468.

- [47] Chang HC, Lindquist S. Conservation of Hsp90 macromolecular complex in *Sacharomyces cerevisiae*. J Biol Chem. 1994;269(40):24983-24988.
- [48] Johnson JL, Toft DO. A novel chaperone complex for steroid receptors involving heat shock proteins, immunophilins, and p23. J Biol Chem. 1994;269(40):24989-24994.
- [49] Duina AA, Chang HC, Marsh JA, Lindquist S, Gaber RF. A cyclophilin function in Hsp90-dependent signal transduction. Science. 1996;274(5293):1713-1715.
- [50] Freeman BC, Toft DO, Morimoto RI. Molecular chaperone machines: Chaperone activities of the cyclophilin Cyp-40 and the steroid aporeceptors-associated protein p23. Science. 1996;274(5293):1718-1720.
- [51] Coss MC, Stephens RM, Morison DK, Winterstein D, Smith LM, Simek SL. The immunophilin FKBP65 forms an association with the serine/threonine kinase c-Raf-1. Cell Growth Diff. 1998;9(1):41-48.
- [52] Pratt WB. The hsp90-based chaperone system-involvement in signal transduction from variety of hormone and growth factor receptors. Proc Soc Exp Biol Med. 1998;217(4):420-434.
- [53] Owens-Grillo JK, Hoffmann K, Hutchison KA, Yem AW, Deibel MR Jr, Handschumacher RE, Pratt WB. The cyclosporin A-binding immunophilin CyP-40 and the FK506-binding immunophilin hsp56 bind to a common site on hsp90 and exist in independent cytosolic heterocomplexes with the untransformed glucocorticoid receptor. J Biol Chem. 1995;270(35):20479-20484.
- [54] Schreiber SL. Immunophilin-sensitive protein phosphatase action in cell signaling pathways. Cell. 1992;70(3):365-368.
- [55] Wilson LK, Benton BM, Zhou S, Thorner J, Martin GS. The yeast immunophilin Fpr3 is a physiological substrate of the tyrosine-specific phosphoprotein phosphatase Ptp1. J Biol Chem. 1995;270(42):25185-25193.
- [56] Silverstein AM, Galigniana MD, Mei-Shya C, Owens-Grillo JK, Chinkers M, Pratt WB. Protein phosphatase 5 is a major component of glucocorticoid receptor-hsp90 complex with properties of an FK606-binding immunophilin. J Biol Chem. 1997;271(26):16224-16230.
- [57] Romano P, Gray J., Horton P, Luan S. Plant immunophilins: functional versatility beyond protein maturation. New Phytologist. 2005;166(3):753-769.
- [58] He Z, Li L, Luan S. Immunophilins and parvulins. Superfamily of peptidyl prolyl isomerases in Arabidopsis. Plant Physiol. 2004;134(4):1248-1267.
- [59] Vener AV, Rokka A, Fulgosi H, Andersson B, Herrmann RG. A cyclophilin-regulated PP2A-like protein phosphatase in thylakoid membranes of plant chloroplasts. Biochemistry. 1999;38(45):14955-14965.
- [60] Rokka A, Aro EM, Herrmann RG, Andersson B, Vener AV. Dephosphorylation of photosystem II reaction center proteins in plant photosynthetic membranes as an im-

- mediate response to abrupt elevation of temperature. *Plant Physiol.* 2000;123(4):1525-1536.
- [61] Luan S, Albers MW, Schreiber SL. Light-regulated, tissue-specific immunophilins in a higher plant. *Proc Natl Acad Sci USA.* 1994;91(3):984-988.
- [62] Fulgosi H, Vener AV, Altschmied L, Hermann RG, Andersson B. A novel multi-functional chloroplast protein: identification of 1 40 kDa immunophilin-like protein located in the thylakoid lumen. *EMBO J.* 1998;17(6):1577-1587.
- [63] Baena-González E, Aro EM. Biogenesis, assembly and turnover of photosystem II units. *Philos Trans R Soc Lond B Biol Sci.* 2002;357(1426):1451-1460.
- [64] Aro EM, Suorsa M, Rokka A, Allahverdiyeva Y, Paakkarinen V, Saleem A, Battchikova N, Rintamäki E. Dynamics of photosystem II: A proteomic approach to thylakoid protein complexes. *J Exp Bot.* 2005;56(411):347-356.
- [65] Shapiguzov A, Edvardsson A, Vener AV. Profound redox sensitivity of peptidyl-prolylisomerase activity in arabidopsis thylakoid lumen. *FEBS Lett.* 2006;580(15):3671-3676.
- [66] Edvardsson A, Shapiguzov A, Petersson UA, Schroder WP, Vener AV. Immunophilin AtFKBP13 sustains all peptidyl-prolylisomerase activity in the thylakoid lumen from *Arabidopsis thaliana* deficient in AtCYP20-2. *Biochemistry.* 2007;46(33):9432-9442.
- [67] Sirpiö S, Khrouchtchova A, Allahverdiyeva Y, Hansson M, Fristedt R, Vener AV, Scheller HV, Jensen PE, Haldrup A, Aro EM. AtCYP38 ensures early biogenesis, correct assembly and sustenance of photosystem II. *Plant J.* 2008;55(4):639-651.
- [68] Fu A, He Z, Sun Cho H, Lima A, Buchanan BB, Luan S. A chloroplast cyclophilin functions in the assembly and maintenance of photosystem II in *Arabidopsis thaliana*. *Proc Natl Acad Sci USA.* 2007;104(40):15947-15952.
- [69] Lepeduš H, Tomašić A, Jurić S, Katanić Z, Cesar V, Fulgosi H. Photochemistry of PSII in CYP38 *Arabidopsis thaliana* Mutant. *Food Technol Biotechnol.* 2009;47(3):275-280.
- [70] Schreiber U, Bilger W, Neubauer C, Schreiber U, Bilger W, Neubauer C. Chlorophyll fluorescence as a nonintrusive indicator for rapid assessment of in vivo photosynthesis. In: Schulze E-D, Caldwell MM. (eds.) *Ecophysiology of Photosynthesis*. Vol 100. Springer: Berlin Heidelberg New York; 1994. p49-70.
- [71] Asada K. Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol.* 2006;141(2):391-396.
- [72] Sonoike K. Photoinhibition of photosystem I. *Physiol Plant.* 2011;142(1):56-64.
- [73] Vass I. Molecular mechanisms of photodamage in the Photosystem II complex. *Biochim Biophys Acta.* 2012;1817(1):209-217.

- [74] Baker NR, Harbinson J, Kramer DM. Determining the limitations and regulation of photosynthetic energy transduction in leaves. *Plant Cell Environ.* 2007;30(9):1107–1125.
- [75] Rochaix JD. Reprint of: Regulation of photosynthetic electron transport. *Biochim Biophys Acta.* 2011; 1807(8):375–383.
- [76] Barajas-López JD, Blanco NE, Strand A. Plastid-to-nucleus communication, signals controlling the running of the plant cell. *Biochim Biophys Acta.* 2013;1833(2):425–437.
- [77] Allen JF. Cyclic, pseudocyclic and noncyclic photophosphorylation: new links in the chain. *Trends Plant Sci.* 2003;8(1):15–19.
- [78] Vojta L, Fulgosi H. Energy Conductance from Thylakoid Complexes to Stromal Reducing Equivalents, *Advances in Photosynthesis - Fundamental Aspects*, Dr Mohammad Najafpour (Ed.), ISBN: 978-953-307-928-8, InTech, 2012, Available from: <http://www.intechopen.com/books/advances-in-photosynthesisfundamental-aspects/energy-conductance-from-thylakoid-complexes-to-stromal-reducing-equivalents>
- [79] Heber U, Walker D. Concerning a dual function of coupled cyclic electron transport in leaves. *Plant Physiol.* 1992;100(4):1621–1626.
- [80] Whatley FR. Photosynthesis by isolated chloroplasts: the early work in Berkeley. *Photosynth Res.* 1995;46(1-2):17–26.
- [81] Finazzi G, Rappaport F, Furia A, Fleischmann M, Rochaix JD, Zito F, Forti G. Involvement of state transitions in the switch between linear and cyclic electron flow in *Chlamydomonas reinhardtii*. *EMBO Rep.* 2002;3(3):280–285.
- [82] Pfannschmidt T, Nilsson A, Allen JF. Photosynthetic control of chloroplast gene expression. *Nature.* 1999;39:625–628.
- [83] Ort DR, Baker NR. A photoprotective role for O<sub>2</sub> as an alternative electron sink in photosynthesis? *Curr Opin Plant Biol.* 2002;5(3):193–198.
- [84] Ivanov AG, Sane PV, Hurry V, Oquist G, Huner NP. Photosystem II reaction centre quenching: mechanisms and physiological role. *Photosynth Res.* 2008;98(1-3):565–574.
- [85] Quiles MJ. Stimulation of chlororespiration by heat and high light intensity in oat plants. *Plant Cell Environ.* 2006;29(8):1463–1470.
- [86] Scheibe R, Dietz KJ. Reduction-oxidation network for flexible adjustment of cellular metabolism in photoautotrophic cells. *Plant Cell Environ.* 2012;35(2):202–216.
- [87] Scheibe R. Malate valves to balance cellular energy supply. *Physiol Plant.* 2004;120(1):21–26.
- [88] Kramer DM, Evans JR. The importance of energy balance in improving photosynthetic productivity. *Plant Physiol.* 2011;155(1):70–78.

- [89] Steiger HM, Beck E. Formation of hydrogen peroxide and oxygen dependence of photosynthetic CO<sub>2</sub> assimilation by intact chloroplasts. *Plant Cell Physiol.* 1981;22:561-576.
- [90] Scheibe R, Stitt M. Comparison of NADP-malate dehydrogenase activation, QA reduction and O<sub>2</sub> evolution in spinach leaves. *Plant Physiol Biochem.* 1988;26:473-481.
- [91] Joliot P, Joliot A. Cyclic electron transfer in plant leaf. *Proc Natl Acad Sci USA.* 2002;99(15):10209-10214.
- [92] Hemschemeier A, Happe T. Alternative photosynthetic electron transport pathways during anaerobiosis in the green alga *Chlamydomonas reinhardtii*. *Biochim Biophys Acta.* 2011;1807(8):919-926.
- [93] Sétif P, Fischer N, Lagoutte B, Bottin H, Rochaix JD. The ferredoxin docking site of photosystem I. *Biochim Biophys Acta.* 2002;1555(1-3):204-209.
- [94] Zhang L, Happe T, Melis A. Biochemical and morphological characterization of sulfur-deprived and H<sub>2</sub>-producing *Chlamydomonas reinhardtii* (green alga). *Planta.* 2002;214(4):552-561.
- [95] Tolleter D, Ghysels B, Alric J, Petroustos D, Tolstygina I, Krawietz D, Happe T, Auroy P, Adriano JM, Beyly A, Cuiné S, Plet J, Reiter IM, Genty B, Cournac L, Hippler M, Peltier G. Control of hydrogen photoproduction by the proton gradient generated by cyclic electronflow in *Chlamydomonas reinhardtii*. *Plant Cell.* 2011;23(7):2619-2630.
- [96] Munekage Y, Hojo M, Meurer J, Endo T, Tasaka M, Shikanai T. PGR5 is involved in cyclic electron flow around photosystem I and is essential for photoprotection in *Arabidopsis*. *Cell.* 2002;110(3):361-371.
- [97] Burrows PA, Sazanov LA, Svab Z, Maliga P, Nixon PJ. Identification of a functional respiratory complex in chloroplasts through analysis of tobacco mutants containing disrupted plastid *ndh* genes. *EMBO J.* 1998;17(4):868-876.
- [98] Sazanov LA, Burrows PA, Nixon PJ. The chloroplast *Ndh* complex mediates the dark reduction of the plastoquinone pool in response to heat stress in tobacco leaves. *FEBS Lett.* 1998;429(1):115-118.
- [99] Shikanai T, Endo T, Hashimoto T, Yamada Y, Asada K, Yokota A. Directed disruption of the tobacco *ndhB* gene impairs cyclic electron flow around photosystem I. *Proc Natl Acad Sci USA.* 1998;95(16):9705-9709.
- [100] Hanke GT, Hase T. Variable photosynthetic roles of two leaf-type ferredoxins in *Arabidopsis*, as revealed by RNA interference. *Photochem Photobiol.* 2008;84(6):1302-1309.
- [101] Hanke GT, Okutani S, Satomi Y, Takao T, Suzuki A, Hase T. Multiple iso-proteins of FNR in *Arabidopsis*: evidence for different contributions to chloroplast function and nitrogen assimilation. *Plant Cell Environ.* 2005;28(9):1146-1157.



- [102] Okutani S, Hanke GT, Satomi Y, Takao T, Kurisu G, Suzuki A, Hase T. Three maize leaf ferredoxin:NADPH oxidoreductases vary in subchloroplast location, expression, and interaction with ferredoxin. *Plant Physiol.* 2005;139(3):1451-1459.
- [103] Rodriguez RE, Lodeyro A, Poli HO, Zurbriggen M, Peisker M, Palatnik JF, Tognetti VB, Tschiersch H, Hajirezaei MR, Valle EM, Carrillo N. Transgenic tobacco plants overexpressing chloroplastic ferredoxin-NADP(H) reductase display normal rates of photosynthesis and increased tolerance to oxidative stress. *Plant Physiol.* 2007;143(2):639-649.
- [104] Hanke GT, Endo T, Satoh F, Hase T. Altered photosynthetic electron channelling into cyclic electron flow and nitrite assimilation in a mutant of ferredoxin:NADP(H) reductase. *Plant Cell Environ.* 2008;31(7):1017-1028.
- [105] Forti G, Bracale M. Ferredoxin-ferredoxin NADP reductase interaction. *FEBS Lett.* 1984;166(1):81-84.
- [106] Nakatani S, Shin M. The reconstituted NADP photoreducing system by rebinding of the large form of ferredoxin-NADP reductase to depleted thylakoid membranes. *Arch Biochem Biophys.* 1991;291(2):390-394.
- [107] Clark RD, Hawkesford MJ, Coughlan SJ, Bennett J, Hind J. Association of ferredoxin-NADP<sup>+</sup> oxidoreductase with the chloroplast cytochrome b-f complex. *FEBS Lett.* 1984;174(1):137-142.
- [108] Zhang H, Whitelegge JP, Cramer WA. Ferredoxin: NADP<sup>+</sup> oxidoreductase is a subunit of the chloroplast cytochrome b6f complex. *J Biol Chem.* 2001;276(41):38159-38165.
- [109] Andersen B, Scheller HV, Moller BL. The PSI E subunit of photosystem I binds ferredoxin:NADP<sup>+</sup> oxidoreductase. *FEBS Lett.* 1992;311(2):169-173.
- [110] Quiles MJ, Cuello J. Association of ferredoxin-NADP oxidoreductase with the chloroplastic pyridine nucleotide dehydrogenase complex in barley leaves. *Plant Physiol.* 1998;117(1):235-244.
- [111] Twachtmann M, Altmann B, Muraki N, Voss I, Okutani S, Kurisu G, Hase T, Hanke GT. N-Terminal Structure of Maize Ferredoxin:NADP<sup>+</sup> Reductase Determines Recruitment into Different Thylakoid Membrane Complexes. *Plant Cell.* 2012;24(7):2979-2991.
- [112] Benz JP, Stengel A, Lintala M, Lee YH, Weber A, Philippar K, Gugel IL, Kaieda S, Ikegami T, Mulo P, Soll J, Bölter B. *Arabidopsis* Tic62 and ferredoxin-NADP(H) oxidoreductase form light-regulated complexes that are integrated into the chloroplast redox poise. *Plant Cell.* 2009;21(12):3965-3983.
- [113] Jurić S, Hazler-Pilepić K, Tomašić A, Lepeduš H, Jeličić B, Puthiyaveetil S, Bionda T, Vojta L, Allen JF, Schleiff E, Fulgosi H. Tethering of ferredoxin:NADP<sup>+</sup> oxidoreduc-

tase to thylakoid membranes is mediated by novel chloroplast protein TROL. *Plant J.* 2009;60(5):783–794.

- [114] Vojta L, Horvat L, Fulgosi H. Balancing chloroplast redox status – regulation of FNR binding and release. *Periodicum biologorum.* 2012;114(1):25–31.
- [115] Stengel A, Benz P, Balsera M, Soll J, Bölder B. TIC62 redox-regulated translocon composition and dynamics. *J Biol Chem.* 2008;283(11):6656–6667.
- [116] Mizusawa N, Wada H. The role of lipids in photosystem II. *Biochim Biophys Acta.* 2012;1817(1):194–208.
- [117] Xu C, Härtel H, Wada H, Hagio M, Yu B, Eakin C, Benning C. The *pgp1* mutant locus of *Arabidopsis* encodes a phosphatidylglycerophosphate synthase with impaired activity. *Plant Physiol.* 2002;129:594–604.
- [118] Hagio M, Sakurai I, Sato S, Kato T, Tabata S, Wada H. Phosphatidylglycerol is essential for the development of thylakoid membranes in *Arabidopsis thaliana*. *Plant Cell Physiol.* 2002;43(12):1456–1464.