

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Biosynthesis of Lipids and Hydrocarbons in Algae

Masato Baba and Yoshihiro Shiraiwa

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/56413>

1. Introduction

Lipids function as important storage compounds to maintain cellular activities. Lipids store high reducing power and energy since those biosynthetic processes require high amount of reducing cofactors and ATP. Storage lipids do not cause any chemical effect on cellular activity such as osmolarity, pH and ion strength because of its hydrophobicity. Membrane lipids such as phospholipids, carotenoids, and cholesterols play a housekeeping role. In addition, some of lipids function as protein modifiers or signaling molecules.

Recently, plant oils are gathering keen interest as a source of renewable energy according to rapid increase in social demands for establishing a low-carbon-society. However, oil production for biofuels and biorefinery using higher plants and crops is strongly worried for competing with food production and to increase those market prices. Therein, algae came into play a new oil-producing organism since algae do not compete with food production. According to their high productivity per unit area, prokaryotic photoautotrophs such as cyanobacteria and eukaryotic algae such as protists are expected to become a promising feedstock in future (Gong & Jiang, 2011).

Although numerous kinds of lipids exist in nature, main carbon chain of the molecules is almost derived from limited numbers of precursor molecules such as fatty acids and isoprenoids. Interestingly, some parts of the synthetic pathways of lipids are quite different among animals, higher plants, cyanobacteria and some eukaryotic microalgae. Although there is quite few information on lipid biosynthesis and metabolism in algae, it is noteworthy that most of biosynthetic pathways of hydrocarbons such as fatty acid- and isoprene-derived hydrocarbon have been well characterized in microalgae.

In this chapter, we will introduce recent progresses on lipid and hydrocarbon biosynthetic pathways in microalgae: First, unique features of algal lipid synthetic pathways mostly hypothesized by advanced DNA sequencing technique although those are not well proved

experimentally yet. Second, two factors for hydrocarbon biosynthesis in microalgae characterized recently by a combination of expressed sequence tags (EST) analysis and novel enzyme characterization. Those are: (1) decarbonylase to produce fatty acid-derived hydrocarbons in cyanobacteria and (2) isoprene-derived hydrocarbon biosynthetic pathway in a representative oil-producing colonial microalga, *Botryococcus braunii*. Third, the mechanism for carbon flow and energy balance in lipid and hydrocarbon biosynthetic pathways. Finally, we will describe a future perspective of algal lipid biosynthetic pathways and its application to biofuel production.

2. Unique features of algal lipid biosynthetic pathways

Although there is no agreed definition and classification of “lipids” (The AOCS Lipid Library, <http://lipidlibrary.aocs.org/>), here we define a term “lipid” as follows: 1) it is biological component of and derived from organisms; 2) it is basically very soluble in organic solvents but not in water; 3) it contains hydrocarbon group in its structure. We adopt biosynthetic classification to categorize lipids such as fatty acid, isoprene or others of unique lipids as shown in Table 1, instead of a conventional lipid classification such as simple lipid, derived lipid, complex lipid, and so on. Here we used the term “lipids” for compounds composed of only carbon, hydrogen, and oxygen.

Table 1 indicates the list of various lipids and their functions. Although numberless lipids exist in nature, main carbon chain of the molecules is mostly derived from fatty acids, isoprenes and their homologous compounds via some synthetic pathways. Recent progress in genome/transcriptome sequencing technology and its computational analysis on similarity of those base sequences among organisms, namely *in silico* analysis, enable us rapid prediction of metabolic pathway even in uncharacterized organism. Owing to the modern *in silico* analyses, some unique features of algal lipid biosynthesis are starting to be enlightened. Interestingly, some parts of the synthetic pathways are quite different among animals, higher plants, cyanobacteria and eukaryotic algae.

Name	Structure	Function
Fatty acid	C _{2n} -/straight-carbon chain with carboxyl group	Membrane component; Bioactivity
Polyketide	Various carbon chain with polyketone group	Antibiotic; Bioactivity
Glyceride	Ester of fatty acid & glycerol	Common storage lipid
Terpenoid	C _{5n} -/branched-carbon chain; isoprene derivate	Bioactivity
Steroid	Tri-terpenoid derivate	Common hormone
Carotenoid	Tetra-terpenoid derivate; conjugated double bond; absorbent	Pigment

Table 1. Various lipids and their functions

2.1. Fatty acid biosynthesis

Acetyl-coenzyme A (CoA) is a universal carbon donor for fatty acid biosynthesis. Acetyl-CoA is supplied via multiple paths from various origins and then subsequently metabolized into malonyl-acyl carrier protein (ACP) by sequential reactions. One molecule of ATP (1ATP) is used for the carboxylation of acetyl unit to produce one malonyl unit. In general, fatty acid biosynthesis utilizes acetyl-CoA and malonyl-ACP as starting substrates and acetyl unit donors. Primarily, butyryl(C₄)-ACP is synthesized from acetyl(C₂)-CoA and malonyl(C₃)-ACP via sequential reactions of condensation, decarboxylation, and reduction of non-malonyl-ACP derived keto unit. Two molecules of NADPH (2NADPH) is used for the reduction of keto group. Accordingly, 1ATP and 2NADPH are consumed to elongate chain of fatty acid molecule by adding C₂-saturated carbon unit in fatty acid biosynthesis. Acyl-ACP is elongated up to acyl(C_{16 or 18})-ACP. Molecules with C_{2n}-carbon chain are widely distributed among various organisms and those of C_{2n-1}-carbon chain are synthesized from C_{2n}-compounds by carbon-loss (Řezanka & Sigler, 2009). In bacteria, *iso*- and *anteiso*-fatty acids (branched-chain fatty acids) are synthesized from amino-acid-derived precursors with branch (Kaneda, 1991). ACP is subsequently removed from acyl(C_{16 or 18})-ACP to form fatty acid(C_{16 or 18}). The fatty acids synthesized on the plastid envelopes are excreted into cytosol by accompanying probably with the process of binding of CoA (Joyard et al., 2010). Oppositely, fatty acid is necessary to be activated by the binding of ACP for passing through the cell membrane in the cyanobacteria *Synechocystis* species.

In any step, a synthesized carbon chain can be metabolized into various products including glycerolipids, triacylglycerides (TG), phospholipids and glycolipids (Joyard et al., 2010). Fatty acids synthesized excessively are stored as TG in most eukaryotes. Usually prokaryotes do not accumulate TG although *Actinomyces* and a few other bacteria exceptionally synthesize TG (Alvarez & Steinbüchel, 2002).

Same fatty acids as metabolites are widely observed in various organisms but their biosynthetic pathways are different depending on classification. There are four known groups of enzyme(s) for fatty acid biosynthesis; type-I fatty acid synthase (FAS), type-II FAS, particular elongases, and enzymes for catalyzing the reversal of β -oxidation. Typically, animals and fungi possess type-I FAS which is a large multi-functional enzyme with multiple functional domains (Chan & Vogel, 2010; Joyard et al., 2010). Bacteria, plastids and mitochondria have type-II FAS which is composed of four subunit proteins such as β -ketoacyl-ACP synthase (KAS), β -ketoacyl-ACP reductase, β -hydroxyacyl-ACP dehydratase and enoyl-ACP reductase (Chan & Vogel, 2010; Joyard et al., 2010; Hiltunen et al., 2010). A trypanosomatid *Leishmania major* possesses three elongases instead of above mentioned FASs for fatty acid biosynthesis (Lee et al., 2006). In a microalga *Euglena gracilis* fatty acids are synthesized *de novo* via the reversal pathway of β -oxidation under anaerobic conditions (Hoffmeister et al., 2005; Inui et al., 1984). As such, wide variation seems to exist among organisms but detailed information on fatty acid biosynthesis is not well understood in algae.

One of model organism *Chlamydomonas reinhardtii* (Chlorophyta) possesses type-II FAS gene which is homologous to that of land plants (Riekhof et al., 2005). Land plants have both plastidal- and mitochondrial-FASs which exhibit different substrate specificity (Yasuno et al.,

2004). Mitochondrial ACP-type enzymes are well characterized especially in the yeast *Saccharomyces cerevisiae* (Hiltunen et al., 2010). In the green alga *C. reinhardtii*, FAS is thought to be localized both in the plastid and the mitochondrion individually since respective FAS gene is represented as single gene in each organelle (Riekhof et al., 2005). Expressed sequence tags (ESTs) of Chlorophyta *Botryococcus braunii* Bot-88-2 (race A) contained a partial sequence of type-I FAS (accession number: FX056119) which is partially similar to animal FAS (Baba et al., 2012a). However, further information of complete genome sequence is necessary to confirm it. In *E. gracilis* which has triple-layer envelope, there are five fatty acid synthetic pathways located in four subcellular compartments; the type-I FAS system in the cytosol, two type-II FAS systems in the plastid, fatty acid synthesis associated partly with wax ester fermentation in the microsomes (see 3.1.4. *Wax ester*) and a malonyl-ACP independent process located in the mitochondria. *E. gracilis* synthesizes wax esters when cells were grown under either heterotrophic or anaerobic conditions (Hoffmeister et al., 2005). Wax ester fermentation under anaerobic conditions produces ATP in net profit while general malonyl-ACP-dependent fatty acid synthesis consumes ATP (Inui et al., 1982). Fatty acids supplied for wax ester production are *de novo* synthesized by acetyl-CoA condensing reaction which is similar to the reversal pathway of β -oxidation, where *trans*-2-enoyl-CoA reductase contributes instead of acyl-CoA dehydrogenase (mitochondrial β -oxidation enzyme) or acyl-CoA oxidase (peroxisomal β -oxidation enzyme) (Hoffmeister et al., 2005; Inui et al., 1984). The *trans*-2-enoyl-CoA reductase reaction is a key step in the reversal pathway of β -oxidation. The direction of the pathway is O_2 -dependently determined to proceed irreversibly under aerobic conditions since the key enzyme pyruvate:NADP⁺ oxidoreductase is sensitive to O_2 (Tucci et al., 2010). The reversal pathway may be essential to maintain the redox balance in the mitochondria under anaerobic conditions (Tucci et al., 2010). Such hypothesis is supported by another report which demonstrated that β -oxidation can be reversed in genetically engineered *Escherichia coli* (Dellomonaco et al., 2011). In mammalian mitochondria, the reversal pathway functions for only elongation process due to substrate specificity of mammal enoyl-CoA reductase and therefore the pathway contributes to short-chain fatty acid elongation process (Inui et al., 1984).

2.2. Fatty acid elongation

C_{18} -Fatty acid can be further elongated via the fatty acid elongation pathway. Fatty acid elongation process is very similar to that of the fatty acid synthesis although acyl-CoA and malonyl-CoA are used as a substrate. In the process, 1ATP and 2NADPH are required for C_2 -unit elongation of saturated carbon chain since CoA-activation is not essential as suggested by another study (Hlousek-radojic et al., 1998). Fatty acid elongation reaction site was shown to be located in the endoplasmic reticulum (Kunst & Samuels, 2009). In contrast to FAS system, all known elongation systems are basically compatible and functions simultaneously. Fatty acid elongase constitutes an enzyme complex of four subunits which is similar to type-II FAS; namely, β -ketoacyl-CoA synthase (KCS), β -ketoacyl-CoA reductase, β -hydroxyacyl-CoA dehydratase and enoyl-CoA reductase. There are two different KCSs; namely "elongation of very long-chain fatty acid" (ELOVL or merely ELO)-type elongase which contributes to sphingolipid biosynthesis and "fatty acid elongation" (FAE)-type elongase which contributes to plant seed TG or wax biosynthesis (Venegas-Calderón et al., 2010). Typically, animals and

fungi possess ELO-type while land plants possess FAE-type. In some cases, ELO and FAE subunits are inaccurately referred as mere “elongase” since heterologous expression of single gene for KCS often results in successful elongation of acyl-CoA by the help of the other subunit of the host (typically, yeast and land plant *Arabidopsis thaliana*).

Poly unsaturated very long-chain fatty acid (PUVLCFA, PULCA, VLC-PUFA, etc.) is one of elongated fatty acids (e.g. Arachidonic acid, Eicosapentaenoic acid, and Docosahexaenoic acid). PUVLCFA is commonly observed in algae such as Euglenophytes, diatoms (*Phaeodactylum tricornerutum*, *Thalassiosira pseudonana*) and haptophytes (*Emiliania huxleyi*, *Isochrysis galbana*, *Pavlova salina*) which possess desaturase/elongases to produce EPA and DHA from C₁₈ fatty acid and derivatives (Venegas-Calderón et al., 2010). This elongation is shown to be catalyzed by ELO-type fatty acid elongase. On the other hand, contribution and physiological function of FAE-type elongase has not been proved in microalgae yet. Macroalgae produce short and long-chain aldehydes (Moore, 2006). These metabolites are suggested to be produced from PUVLCFA although the detailed synthetic pathway is almost unknown. Long-chain aldehyde producing reaction was reported to be catalyzed by lipooxygenase which functions to oxygenize and cleave PUVLCFA at a specific position to form short-chain metabolites (Moore, 2006).

2.3. Polyketide biosynthesis

Polyketide includes various complex compounds such as antibiotics (e.g. erythromycin, tetracycline, lovastatin) (Staunton & Weissman, 2001). Polyketide biosynthesis is similar to C₄ and longer fatty acid synthesis except successive reduction of *keto*-group and utilizes various ACP/CoA compounds as its substrate. Mycolic acids, extremely large fatty acids (ca. C₉₀) in the cell envelope of mycobacteria (Verschoor et al., 2012), are synthesized by condensation of two VLCFA acyl-CoA molecules via polyketide biosynthesis (Portevin et al., 2004).

There are three types of polyketide syntheses (PKSs): type-I PKS which is a large multifunctional enzyme in the consequence of multiple functional domains, type-II PKS which is composed of monofunctional proteins to form complex and type-III PKS which resembles chalcone synthase catalyzing the committed step in flavonoid biosynthesis in higher plants and some bryophytes (Shen, 2003). Type-I PKSs are further classified into two, namely iterative and non-iterative (modular) types. Bacteria possess type-I to III of PKSs. Fungi and animal typically possess type-I iterative PKS which is closely related each other (Jenke-Kodama & Dittmann, 2009). Interestingly, there is evolutionary connection between PKSs and FASs (Jenke-Kodama & Dittmann, 2009; John et al., 2008; Sasso et al., 2011).

In the genomes of chlorophyta (*C. reinhardtii*, *Chlorella variabilis*, two *Ostreococcus* species, etc.), heterokontophyta (*Aureococcus anophagefferens*), and haptophyceae (*Emiliania huxleyi* and *Chrysochromulina polylepis*), “non-iterative” type-I PKS is coded but not in land plant, rhodophyta (*Cyanidioschyzon merolae* and *Galdieria sulphuraria*), Stramenopiles (*Thalassiosira pseudonana*, *Phaeodactylum tricornerutum*, two *Phytophthora* species), and Euglenozoa (*Trypanosoma brucei*, *Trypanosoma cruzi* and *Leishmania major*) (Sasso et al., 2011). Polyketide synthesis plays a role in biosynthesis of cyanobacterial toxin, microcystin (Jenke-Kodama & Dittmann, 2009). Dinoflagellate toxins are also polyketides although its synthetic pathway remains unknown

because of technical restrictions (Sasso et al., 2011). In addition, biosynthetic processes of polyketides in macroalgae are one of important targets to be urgently elucidated.

2.4. Terpenoid biosynthesis

Terpenoid which is composed of branched C_{5n} carbon unit are synthesized by condensation of C_5 isoprene units (as isopentenyl diphosphate (IPP) and its isomer dimethylallyl diphosphate (DMAPP) *in vivo*) (Bouvier et al., 2005). In general, isoprene unit is supplied via either or both mevalonic acid (MVA) pathway or/and methylerythritol phosphate (MEP) pathway (or non-mevalonic acid pathway). MVA pathway is located in the cytosol of Archaea and eukaryotes or in the peroxisome (Lohr et al., 2012). In MVA pathway, three molecules of acetyl-CoA are condensed into DMAPP through MVA and then IPP by sequential reactions, using 3ATP and 2NADPH. The MEP pathway is known to be located in the cytosol of bacteria including cyanobacteria and in the stroma of plastids in plants and eukaryotic algae (Joyard et al., 2009). In MEP pathway, pyruvate and glyceraldehyde-3-phosphate (GAP) react to form IPP or DMAPP via MEP using energy of three high-energy phosphate bonds on either ATP or CTP and reducing power from at least 1NADPH and four reducing coenzymes (not completely identified yet) (Hunter, 2007). One molecule of CO_2 is released in MVA and MEP pathway respectively. In land plants, isoprene units can be exchanged through chloroplast envelope by that MVA and MEP pathways complement each other. However, any protein for isoprene unit exchange has not been isolated so far (Bouvier et al., 2005; Joyard et al., 2009; Lohr et al., 2012).




Figure 1. Pathway for terpenoid biosynthesis

Primary and terminal molecules are underlined respectively. Substrates multiply used are shown in bold. [1]: putative Isoprene transporter. [2]: a predicted junction from the pentose phosphate pathway to the MEP pathway in cyanobacteria. AACT, acetoacetyl-CoA thiolase; CMK, 4-(cytidine 5'-diphospho)-2-C- methylerythritol kinase; DXR, 1-deoxy-D-xylulose 5-phosphate reductoisomerase; DXS, 1-deoxy-D-xylulose 5-phosphate synthase; HDR, 4-hydroxy-3-methylbut-2-en-1-yl diphosphate reductase; HDS, 4-hydroxy-3-methylbut-2-en-1-yl diphosphate synthase; HMGS, 3-hydroxy-3-methylglutaryl-CoA synthase; HMGR, 3-hydroxy-3-methylglutaryl-CoA reductase; IDI, isopentenyl diphosphate:dimethylallyl diphosphate isomerase; MCT, 2-C-methyl-D-erythritol 4-phosphate cytidyltransferase; MDS, 2-C-methyl-D-erythritol 2,4-cyclodiphosphate synthase; MVD, mevalonate-5-diphosphate decarboxylase; MVK, mevalonate kinase; PMK, 5-phosphomevalonate kinase.

In silico analysis suggested that algae have enzymes for terpenoid biosynthesis which resembles with those in land plants (Lohr et al., 2012; Sasso et al., 2011). On the other hand, MVA pathway genes are often lost in some algae although such algal unique system remains to be understood in future work (Lohr et al., 2012; Sasso et al., 2011). EST analysis on race A and B of *B. braunii* revealed that MEP pathway genes are actively expressed but not MVA pathway genes. Some secondary symbiotic algae possess a mosaic MVA pathway which involves enzymes originated from both primary and secondary hosts (Lohr et al., 2012). MEP pathway connected with the pentose phosphate pathway was observed in a species of cyanobacteria although detailed mechanism is still not clear (Poliquin et al., 2004).

IPP and DMAPP are metabolically conjugated by condensation and dephosphorylation to produce polyterpenoid. No ATP or reducing power is required when isoprene units get into condensation reaction by head-to-tail conjunction (e.g. farnesyl pyrophosphate formation while 1NADPH is required in case of tail-to-tail condensation (e.g. squalene formation). Polyterpenoid is individually or cooperatively synthesized either in the cytosol, plastid or mitochondrion (Bouvier et al., 2005; Joyard et al., 2009; Lohr et al., 2012). Each terpenoid condensation enzyme has particular specific to isoprene molecules such as mono-/sesqui-/di-/tri-/tetra-terpene, respectively. Unlikely land plant, the biosynthesis of isoprene in green macroalgae proceeds via MEP pathway in the plastid (Lohr et al., 2012) and it functions to produce special natural products such as bioactive halogenated poly terpenoid (Moore, 2006). Vanadium bromoperoxidase is an abundant enzyme to produce brominated products in all classes of marine macroalgae and vanadium iodoperoxidase is also identified and characterized (Moore, 2006). However, vanadium chloroperoxidase is not yet identified despite the abundance of chlorinated compounds in algae. These haloperoxidases catalyze both halogenation and cyclization to produce various unique halogenated cyclic terpenoid in macroalgae, but such unique isoprene condensing enzyme is not yet identified in microalgae (Sasso et al., 2011).

3. Hydrocarbon biosynthesis in algae

Table 2 shows a list of lipids and hydrocarbons which can be candidates for renewable energy sources. These compounds are metabolites derived from the elemental lipids shown in Table

Name	Structure	Function
Odd-chain fattyhydrocarbon	Hydrocarbonfrom fatty acid (C_{2n-1})	Unknown
Wax ester	Ester of fatty acid & fatty alcohol	Cuticle component
Alkenones	<i>trans</i> -unsaturated-/straight-carbon chainwith ketone group	Storage lipid?
Heterocyst glycolipid	Alcohol-/ketone-glycoside	Cell wall component
Even-chain fatty hydrocarbon	Hydrocarbon from fatty acid (C_{2n})	Unknown
Olefinichydrocarbon	Hydrocarbon from fatty acidwith multiple double bonds	Unknown
Terpenoid hydrocarbon	Hydrocarbon from terpenoid	Unknown

Table 2. Lipids and hydrocarbons for renewable energy source

1. Their pool sizes of metabolites in cells and production capability largely varies among species and even strains of a certain species. The most extreme example can be seen in a colonial oil-producing green alga *B. braunii*: a certain strain dominantly produces odd-chain fatty hydrocarbons while another produces terpenoid derived hydrocarbons and those strains are classified as race A, B and L.

Microalgal species/strains nominated as oil-producer are simply classified into three groups by their main products: namely, hydrocarbons, TG/free fatty acids and the other lipids. For example, bacteria (Schirmer et al., 2010), a unicellular green alga *Pseudochoricystis ellipsoidea* (Satoh et al., 2010), a colonial green alga *B. braunii* race A (Yoshida et al., 2012) accumulate fatty hydrocarbons. According to a recent review (Yoshida et al., 2012), *B. braunii* race B and L and a heterotrophic Labyrinthulea *Aurantiochytrium sp.* produce terpenoid-derived hydrocarbons. Green algae *C. reinhardtii*, *Chlorella vulgaris*, *Chlorella protothecoides*, diatoms, *Nannochloropsis* spp. usually produce and accumulate TG and free fatty acids. *E. gracilis* Z produces wax ester and accumulate it in the cell (Inui et al., 1982). Haptophytes, but only five species, produce long-chain ketones, called as “alkenones” (Laws et al., 2001; Eltgroth et al., 2005; Toney et al., 2012). Red and green macroalgal species, *Gracilaria salicornia* and *Ulva lactuca*, respectively, contain only limited amounts of lipids including arachidonic acid and Docosapentaenoic acid

ranging 1~2% of dry weight (Tabarsa et al., 2012). Recent big news was that genetic transformation by the particle-gun bombardment method was successfully established in alliphatic hydrocarbon producing photosynthetic eukaryote *Pseudochoricystis ellipsoidea* (Trebouxiophyceae) (Imamura et al., 2012). This achievement may open a new trail toward the genetic manipulation of metabolism for algal biofuel production.

3.1. Fatty hydrocarbons and the other fatty acid derivatives

Alliphatic carbon-chain is a ubiquitous structure which exists in the molecules produced via fatty acid biosynthesis in organisms. In this part we introduce some fatty acid derivatives and their molecular properties and biosynthetic pathways.

3.1.1. Odd-chain fatty hydrocarbon

Bacteria, microalgae and land plants produce odd-chain hydrocarbons (Řezanka & Sigler, 2009; Tornabene, 1981). Plant wax constitutes of odd-chain hydrocarbons without any branching, namely fatty hydrocarbons (Jetter & Kunst, 2008). This type of hydrocarbons is suggested to be produced via the decarbonylation pathway (Jetter & Kunst, 2008; Schirmer et al., 2010). First, acyl-CoA is reduced to form fatty aldehyde using 1NADPH as a reductant cofactor (Schirmer et al., 2010; Willis et al., 2011). In pea, the decarbonylation reaction is catalyzed by a membrane-bound enzyme, fatty acyl-CoA reductase (Cheesbrough & kolattukudy, 1984; Vioque & Kolattukudy, 1997) which is also present in the race A of *B. braunii* (Wang & Kolattukudy, 1995).

A cyanobacterium *Synechococcus elongatus* PCC7942 has fatty acyl-ACP reductase which prefers acyl-ACP, not acyl-CoA, as substrate (Schirmer et al., 2010). Bacterial gene for the aldehyde-forming fatty acyl-CoA reductase was identified, but not eukaryotic gene yet. Fatty aldehyde is decarbonylated to form odd-chain fatty hydrocarbons with a release of carbon monoxide. Such aldehyde decarbonylase activity was successfully determined in land plants (Jetter & Kunst, 2008), a colonial green alga *B. braunii*, and bacteria (Schirmer et al., 2010), but its gene was identified only in bacteria at present (Schirmer et al., 2010). Using microsomal preparations of *B. braunii*, alkane was proved to be synthesized from fatty acid and aldehyde only under anaerobic conditions (Dennis & Kolattukudy, 1991). The aldehyde decarbonylase is a cobalt porphyrin enzyme which was suggested to locate in the microsomes (Dennis & Kolattukudy, 1992). However, it is not known yet how intracellular hydrocarbons are transferred to extracellular space in race A of *B. braunii* (Casadevall et al., 1985; Largeau et al., 1980; Templier et al., 1992). In addition, an enzyme for the synthesis of odd-chain fatty hydrocarbons is unidentified yet since the product of the enzyme was different from native hydrocarbons of *B. braunii* by lacking terminal double bond seen in the natural product. Although decarbonylation reaction does not require any reductant, the reaction of *in vitro* decarbonylation from octadecanal to heptadecane was observed only in the presence of ferredoxin, ferredoxin reductase and NADPH. This result suggest essential requirement of reductant for aldehyde decarbonylase to exhibit activity in the cyanobacterium *Nostoc punctiforme* PCC73102 (Schirmer et al., 2010).

3.1.2. Wax ester

Wax esters consist of fatty acids (acyl-CoAs *in vivo*) and fatty alcohols and are one of components of plant cuticles or seed oils (Jetter & Kunst, 2008; Kunst & Samuels, 2009). *E. gracilis* produces wax esters under either heterotrophic or anaerobic conditions (Inui et al., 1982). Wax esters are produced by condensation of fatty acids and primary alcohols which are synthesized from acyl-CoAs (see also 3.1.5. *Even-chain fatty hydrocarbon*). Wax ester synthase/acyl-CoA:diacylglycerol acyltransferase (WSD1) is a condensation enzyme identified in *A. thaliana* (Li et al., 2008). In *E. gracilis*, both NADH-requiring alcohol-forming fatty acyl-CoA reductase (EgFAR) and wax synthase (EgWS) are already identified and the sequences of those genes showed similarity with those of jojoba (land plant) (Teerawanichpan & Qiu, 2010). EgWS utilizes a broad range of fatty acyl-CoAs and fatty alcohols as substrates with the preference towards myristic acid and palmitoleyl alcohol (Teerawanichpan & Qiu, 2010). Those substrates are suggested to be produced via various fatty acid synthetic pathways in this alga (see 2.1. *Fatty acid biosynthesis*).

3.1.3. Alkenones

At least five species of haptophyceae (*Chrysothile lamellose*, *Emiliania. huxleyi*, *Gephyrocapsa oceanica*, *Isochrysis galbana*, *Pseudoisochrysis* sp.) were reported to accumulate highly alkenes, alkenoates (PUVLCFA-methyl/ethyl esters) and alkenones (PUVLC ketones) (Eltgroth et al., 2005; Laws et al., 2001; Toney et al., 2012). In this section, we call those compounds “alkenones” for our convenience. “Alkenones” are discriminated from the other lipids by 2 to 4 *trans*-carbon double bonds stocked under low temperature conditions, and by its remarkable length (about C₃₈). “Alkenones” are suggested to be synthesized near the chloroplast, and then stored in the intracellular lipid body (Eltgroth et al., 2005). Mechanisms for alkenone biosynthesis and its desaturation are not known yet. Alkenones may act as a storage lipid since behavior of alkenones was shown to be similar to TG in the other algae functioning storage lipids (Eltgroth et al., 2005).

3.1.4. Heterocyst glycolipid

The heterocyst of cyanobacterium *Anabaena* sp. PCC 7120 is surrounded by cell wall involving unique glycolipids as a component (heterocyst glycolipid: HGL) (Bauersachs et al., 2009; Awai & Wolk, 2007). HGL may be a good source of biofuel since such sugar-conjugated molecule (aglycon) is changed to C₂₆₋₂₈ fatty polyhydric alcohol or ketone by removing sugar residues. HGL is known to be biosynthesized from fatty polyhydric alcohol via an unknown pathway which may be composed of fatty acid and/or polyketide synthetic enzymes (Awai & Wolk, 2007; Bauersachs et al., 2009) as its mechanisms including transportation of HGL during heterocyst development has been well studied (Bauersachs et al., 2009; Nicolaisen et al., 2009). Fatty polyhydric alcohol is metabolized to glycoside by the catalysis of glycosyltransferase (HGL formation protein: HglT) (Awai & Wolk, 2007).

3.1.5. Even-chain fatty hydrocarbon

The bacterium *Vibrio furnissi* M1 was suggested to synthesize even-chain fatty hydrocarbons by the reduction of fatty acids through primary fatty alcohols (Park, 2005). Primary fatty alcohols are commonly synthesized from acyl-CoA by either one-step or two-step reductions. The one-step reduction is catalyzed by “alcohol-forming” fatty acyl-CoA reductase of which gene was already identified in bacteria (Willis et al., 2011) and various eukaryotes such as land plants and *Euglena* (Teerawanichpan & Qiu, 2010; Vioque & Kolattukudy, 1997). The two-step reduction reactions are catalyzed by two enzymes: namely, “aldehyde-forming” fatty acyl-CoA reductase and then fatty aldehyde reductase of which gene is identified only in bacteria (Wahlen et al., 2009). In the two-step reductive reactions, 2NADPH are necessary to reduce acyl-CoA to primary fatty alcohol. However, mechanism for consequential reactions to form even-chain fatty hydrocarbon by reducing fatty alcohol has not been identified yet (Park, 2005) but elucidation of the mechanism and knowledge on its distribution among species are very important for the progress of biofuel production.

3.1.6. Olefinic hydrocarbon

Olefinic hydrocarbons contain many unsaturated bonds in the molecule. *Ole* (olefin) *ABCD* is a gene set harbored in bacteria and catalyzes the production of an olefinic hydrocarbon molecule by head-to-head condensation reaction (Beller et al., 2010; Sukovich et al., 2010a; Sukovich et al., 2010b). The head-to-head condensation reaction is summarized as follows: 1) a carboxyl group ($R_1\text{-COOH}$) of fatty acid X is reduced to a carbonyl group ($R_1\text{-CHO}$); 2) the carbonyl (or thioester ($R_1\text{-CO-S-CoA}$ with carbon number x)) group of X reacts with an alpha-carbon of another fatty acid Y ($R_2\text{-H}_2\text{-COOH}$ with carbon number y) to combine by forming a hydroxyl group (-OH) and releasing one carbon molecule; 3) the hydroxyl group changes into a carbon double bond ($R_1\text{-C=}$ R_2); 4) the reduction of the carbon double bond results to produce hydrocarbon molecule with carbon number of $[x + y - 1]$ (Albro & Dittmer, 1969). So, acyl-CoA can react with an alpha-carbon of fatty acid instead of carbonyl unit of the aldehyde (Sukovich et al., 2010a). As the gene set *oleABCD* was recently identified, further kinetic and phylogenetic studies can be performed.

3.2. Hydrocarbon biosynthesis from isoprene: A novel terpenoid hydrocarbon biosynthetic pathway in a colonial green alga *Botryococcus braunii* (race B)

A colonial green alga *B. braunii* produces hydrocarbon up to 75% of its dry weight (Yoshida et al., 2012). *B. braunii* is generally classified into three races by its products: race A produces odd-chain fatty hydrocarbons (alkadiene, alkatriene); race B (triterpene) and L (tetraterpene) produces hydrocarbons from isoprene, namely terpenoid hydrocarbons (Yoshida et al., 2012). *B. braunii* cells are botryoidally-aggregated by a network of covalently and/or non-covalently conjugated hydrocarbon molecules to build colony structure: namely, as structural hydrocarbons (Metzger et al., 1993; Metzger et al., 2007; Metzger et al., 2008). Extracellular space is filled with liquid hydrocarbons being discriminated from the structural hydrocarbons (Weiss et al., 2012). Hydrocarbon biosynthesis in the race A is partially understood but no recent progress

was reported (see 3.1.1. *Odd-chain fatty hydrocarbon*). Properties of the race B hydrocarbons are mostly well-known although the race L is still enigmatic.

The race B hydrocarbons are methylsqualene and botryococcene which are specifically produced by *B. braunii*. The amount of the race B hydrocarbons accumulated in the colony did not decrease in the dark, suggesting that extracellular hydrocarbons are a physiologically inactive storage compounds (Sakamoto et al., 2012). C₅-isoprene unit for synthesizing hydrocarbon molecules is supplied via MEP pathway in *B. braunii* (Sato et al., 2003) (see 2.4. *Terpenoid biosynthesis*). The enzyme 1-deoxy-D-xylulose-5-phosphate synthase catalyzes the first step of the MEP pathway and its three distinct isoforms are well-characterized (Matsushima et al., 2012). ESTs of most genes in the MEP pathway, but never the MVA pathway, are already obtained by transcriptome analysis although those are not completely cloned and characterized yet (Ioki et al., 2012a; Ioki et al., 2012b). Botryococcene is known to be synthesized via a similar pathway to squalene biosynthesis besides of cleavage manner of cyclopropane base in a precursor, presqualene pyrophosphate (PSPP) (Metzger & Largeau, 2005; Banerjee et al., 2002). Squalene synthase, which is widely observed in eukaryotes, catalyzes two step reactions: namely, (1) condensation of two farnesyl diphosphates (FPP, triterpene) to form PSPP and (2) dephosphorylation, cyclopropane cleavage, carbon bond reformation and NADPH-dependent reduction of PSPP to produce squalene (Okada, 2012; Jennings et al., 1991). These two reactions are catalyzed at the domain 3 and 4 for 1st reaction and the domain 5 for 2nd reaction of six domains sequentially (Gu et al., 1998). *Botryococcus* squalene synthase (BSS) was isolated by methods of homology screening of cDNA library based on its putative homologous sequence obtained by the degenerate PCR method to already known squalene synthase (Okada et al., 2000). BSS expressed in *E. coli* produced only squalene but not botryococcene (Okada et al., 2000). Squalene synthase like-1 (SSL-1) is a protein which is homologous to BSS possessing quite different amino acid sequence at the domain 5 (Niehaus et al., 2011). The purified SSL-1 did not function to produce neither botryococcene nor squalene *in vitro*. Instead, SSL-1 stimulates botryococcene production when it was added to *B. braunii* cell extracts. Furthermore, over expression of SSL-1 in FPP accumulating yeast resulted in the accumulation of presqualene alcohol (dephosphorylated PSPP) in the cells. These results suggested that SSL-1 functions as PSPP synthase but subsequent reactions for squalene synthesis are catalyzed by other enzymes (Niehaus et al., 2011). SSL-2 and SSL-3 are proteins which are also homologous to BSS (Niehaus et al., 2011). SSL-2 catalyzes two NADPH-dependent reactions for the production of squalene from PSPP and bisfarnesyl ether with a little squalene from FPP. SSL-3 catalyzes a NADPH-dependent reaction to form botryococcene from PSPP. These results suggest that the pathway for terpenoid hydrocarbon biosynthesis in *B. braunii* is quite unique although cooperation of BSS, SSL-1, SSL-2, and SSL-3 is still unclear (Niehaus et al., 2011). After the synthesis, both squalene and botryococcene are subsequently methylated but the number of methylation is variable. The name of botryococcene was originally designated to methylated botryococcene but now it is used for both compounds. The methyl group is transferred from S-adenosyl methionine by triterpene methyltransferases (TMTs) although completely methylated (tetra-methylated) squalene and botryococcene is not yet produced *in vitro* (Niehaus et al., 2012).


4. Carbon flow and energy balance in lipid and hydrocarbon biosynthetic pathways

In Table 3 and Fig. 1, lipid and hydrocarbon biosynthetic pathways are summarized. All hydrocarbons are produced from precursors (namely acyl-ACP or IPP/DMAPP) which are produced from three primary metabolites; acetyl-CoA, pyruvate and GAP. GAP should be the primary metabolite during photosynthesis and transported into the cytosol. Then acetyl-CoA and pyruvate are sequentially produced from GAP in the glycolysis. On the other hand, acetyl-CoA is primarily produced by the degradation of various lipids via β -, α -, and ω -oxidation (Graham & Eastmond, 2002). Any pathway for hydrocarbon production includes decarboxylation of carbon chain supplied as substrate and consumption of ATP and reducing power (see the MVA/MEP pathway and glycolysis in Fig. 1). GAP production mostly depends on carbon fixation rate by the photosynthetic C_3 cycle and the process seems to be the most effective limiting factor for hydrocarbon production. Gene expression level for fatty acid synthesis is relatively higher in race A (fatty hydrocarbon) than race B (terpenoid hydrocarbon) in *B. braunii* while the expression of isoprene synthetic genes showed opposite trend (Ioki et al., 2012c). Transcriptional regulation network of fatty acid metabolism is well studied in *E. coli* (Fujita et al., 2007) but not yet in the other organisms. Carbon allocation into lipids is known to be affected by environmental change via metabolic regulation; e.g. wax ester fermentation in *E. gracilis* under anaerobic conditions (Tucci et al., 2010), TG accumulation in algae under nutrient deficiency such as nitrogen (Hu et al., 2008; Miao & Wu, 2006) and under cold-stress (Li et al., 2011; Renaud et al., 1995). On the other hand, such environmental changes do not affect carbon allocation into hydrocarbon in *B. braunii* (Baba et al., 2012b; Metzger & Largeau, 2005; Sakamoto et al., 2012).

Supply of inorganic and organic carbon sources, nutrient deficiency and low-temperature are empirically known to be stimulating factors for lipid biosynthesis. Enrichment of CO_2 as inorganic carbon source stimulated lipid biosynthesis and cell growth by accelerating photosynthetic carbon fixation in microalgae (Kumar et al., 2010). Neutral lipid production and accumulation was strongly accelerated in the presence of exogenous organic carbon source by accompanying with abolishing chlorophylls in a unicellular green alga *Chlorella protothecoides* (Miao & Wu, 2006). Nitrogen is the most effective factor for changing carbon/nitrogen metabolism and stimulates neutral lipid accumulation under N-deficient conditions (Hu et al., 2008). However, nitrogen deficiency diminishes whole cellular productivity and therefore the metabolic regulation to stimulating lipid biosynthesis does not always result in the increase in gross productivity of lipids. Lipid accumulation under cold (Li et al., 2011; Renaud et al., 1995) or other stress conditions (e.g., nutrient limitation at the stationary growth phase) also induce change in metabolisms. Either cold or high-salinity stress (Lu et al., 2009) stimulates lipid desaturation catalyzed by various lipid desaturases. Desaturation degree influences properties of lipids such as melting point, reactivity, odor, degradability and so on. Lipid biosynthesis by microalgae was shown to be affected by changing wavelength, namely stimulation of lipid production by red light, via the modulation of nitrogen and carbon metabolism in the cells (Miyachi et al., 1978).

Name	Reactions	Products	Notes
Carbon chain biosynthesis			
Fatty acid biosynthesis (C2)	$\text{acyl}(n)\text{-ACP} + \text{acetyl-CoA} (+ \text{CO}_2) + \text{ATP} + 2\text{NADPH} + 2\text{H}^+ \rightarrow \text{acyl}(n+2)\text{-ACP} (+ \text{CO}_2) + \text{H}_2\text{O} + \text{CoA} + \text{ADP} + \text{Pi} + 2\text{NADP}^+$	Fatty acid(C4~18)	Reference pathway
Fatty acid biosynthesis (C2)	$\text{acyl}(n)\text{-CoA} + \text{acetyl-CoA} + 2\text{NADPH} + 2\text{H}^+ \rightarrow \text{acyl}(n+2)\text{-CoA} + \text{H}_2\text{O} + \text{CoA} + 2\text{NADP}^+$	Fatty acid(C4~16)	<i>E. gracilis</i> mitochondria
Fatty acid biosynthesis (C2)	$\text{acyl}(n)\text{-CoA} + \text{acetyl-CoA} + 2\text{NADPH} + 2\text{H}^+ \rightarrow \text{acyl}(n+2)\text{-CoA} + \text{H}_2\text{O} + \text{CoA} + 2\text{NADP}^+$	Fatty acid(C4~18)	Engineered <i>Escherichia coli</i>
MVA pathway (C5)	$3\text{acetyl-CoA} + \text{H}_2\text{O} + 3\text{ATP} + 2\text{NADPH} + 2\text{H}^+ \rightarrow \text{IPP} + 3\text{CoA} + \text{CO}_2 + 3\text{ADP} + \text{Pi} + 2\text{NADP}^+$; $\text{IPP} \rightleftharpoons \text{DMAPP}$	Isoprene	Reference pathway
MEP pathway (C5)	$\text{pyruvate} + \text{GAP} + \text{ATP} + \text{CTP} + \text{NADPH} + 4\text{e}^- + 5\text{H}^+ \rightarrow \text{IPP (DMAPP)} + \text{CO}_2 + 2\text{H}_2\text{O} + \text{ADP} + \text{CMP} + \text{PPi} + \text{NADP}^+$	Isoprene	Incomplete about redox
Carbon chain elongation			
Fatty acid elongation (C2)	$\text{acyl}(n)\text{-CoA} + \text{acetyl-CoA} (+ \text{CO}_2) + \text{ATP} + 2\text{NADPH} + 2\text{H}^+ \rightarrow \text{acyl}(n+2)\text{-CoA} (+ \text{CO}_2) + \text{H}_2\text{O} + \text{CoA} + \text{ADP} + \text{Pi} + 2\text{NADP}^+$	Fatty acid(C20~28)	Reference pathway
Fatty acid elongation (C2)	$\text{acyl}(n)\text{-CoA} + \text{acetyl-CoA} + 2\text{NADPH} + 2\text{H}^+ \rightarrow \text{acyl}(n+2)\text{-CoA} + \text{H}_2\text{O} + \text{CoA} + 2\text{NADP}^+$	Fatty acid(C6~16)	Human mitochondria
Head-to-tail isoprene condensation (C5)	$(n)\text{isoprene} \rightarrow \text{Poly terpenoid-PP} + (n-1)\text{PPi} (n \geq 2)$	Terpenoid	Reference reaction
Head-to-head isoprene condensation (C5)	$2 \text{isoprene} + \text{NADPH} \rightarrow \text{Poly terpenoid} + 2\text{PPi} + \text{NADP}^+ + \text{H}^+$	Hydro-carbon	Reference reaction
Fatty acid (ACP/CoA) reduction			
Fatty aldehyde formation	$\text{acyl-CoA} + \text{NADPH} + \text{H}^+ \rightarrow \text{fatty aldehyde} + \text{CoA} + \text{NADP}^+$	Fatty-aldehyde	Reference reaction
Fatty aldehyde formation	$\text{acyl-ACP} + \text{NADPH} + \text{H}^+ \rightarrow \text{fatty aldehyde} + \text{ACP} + \text{NADP}^+$	Fatty-aldehyde	Cyanobacteria
Alcohol formation	$\text{acyl-CoA} + 2\text{NADPH} + 2\text{H}^+ \rightarrow \text{fatty alcohol} + \text{CoA} + 2\text{NADP}^+$	Fatty-alcohol	Reference reaction
Alcohol formation	$\text{fatty aldehyde} + \text{NADPH} + \text{H}^+ \rightarrow \text{fatty alcohol} + \text{NADP}^+$	Fatty-alcohol	Reference reaction
Odd-chain fatty hydrocarbon biosynthesis			
Aldehyde decarbonylation	$\text{fatty aldehyde}(n) \rightarrow \text{fatty hydrocarbon}(n-1) + \text{CO}$	Hydro-carbon	Reference reaction
Even-chain fatty hydrocarbon biosynthesis			
Alcohol reduction	$\text{fatty alcohol} (n) \rightarrow \text{fatty hydrocarbon} (n)$	Hydro-carbon	Incomplete about redox
Olefinic hydrocarbon			
Head-to-head acyl-CoA condensation	$2\text{acyl-CoA}(n) \rightarrow \text{alkadiene} (2n-1) + \text{CO}_2 + \text{H}_2\text{O} + 2\text{CoA}$	Hydro-carbon	Incompletely understood
Wax ester			
Fatty acid/alcohol esterification	$\text{fatty acyl-CoA}(X) + \text{fatty alcohol}(Y) \rightarrow \text{wax ester} (X+Y) + \text{CoA}$	Wax ester	Reference reaction

Table 3. Carbon flow, consumption of ATP and reducing power in lipid and hydrocarbon biosynthetic pathways



^aRepresentative localization of pathways in algae and plants.

^bC₁₀, mono-terpene is indicated as a model product which is typically synthesized via GPP *in vivo*.

Figure 2. Carbon flow, consumption of ATP and reducing power in lipid biosynthetic pathways. Black- or blue-chained spheres indicate C-C chain. Orange- or Yellow-colored circles indicate Pi in various compounds including IPP, DMAPP, ATP and so on. Difference in color of box-frames indicates difference in localization of pathways. Fatty acid biosynthesis in engineered *E. coli* and fatty acid (~C₄) elongation in human mitochondria are not shown here since those reactions are same as those in mitochondria of *E. gracilis*; namely, reversal pathway of β -oxidation. Detailed information is written in the text. GPP, geranyl diphosphate.

5. Future perspective of algal lipid biosynthetic pathways

Recent *in silico* analysis suggested the presence of some unique lipid-metabolic pathways in algae although those are not characterized yet (e.g., see Table 3 and 3. *Hydrocarbon biosynthesis in algae*). Therefore, research on algal lipid biosynthetic pathways should be unavoidably worthy task to increase industrial algal-oil production. Study on lipid metabolism is also beneficial for useful and biologically active organic material production to achieve the invention of manufactural lipid synthesis. One of such good examples is the manipulation of β -oxidation to proceed for reversal direction in *E. coli* (Dellomonaco et al., 2011), the construction of recursive “+1” pathway by genetic engineering in contrast to “+2” fatty acid and “+5” isoprene pathways (Marcheschi et al., 2012). Increase in lipid production by eukaryotic algae strongly can be achieved by metabolic manipulation by controlling strict redox status and subcellular compartment of metabolisms.

Genetic engineering in eukaryotic algae is important technology to be established although it still is quite challenging (Gong et al., 2011; Radakovits et al., 2010). It is highly expected that algal oil is efficiently produced with high purity since it is produced by enzymatic reactions in homogenous productive cells. So, characteristics of products, such as chain length and number of double bond in the molecule, can be modified by genetic engineering (Gong & Jiang, 2011; Radakovits et al., 2010). Further, facilitation of lipid extraction (e.g. lipid auto-secretion from cells to the medium) (Cho & Cronan, 1995; Liu et al., 2010; Michinaka et al., 2003; Nojima et al., 1999) and cell precipitation control (Kawano et al., 2011)) are important to be improved since such processes consume vast energy at industrial process of production. It is noteworthy that direct extraction of oil from *B. braunii* was already achieved to reduce energy and cost (Frenz et al., 1989; Kanda et al., in press). In cyanobacteria, trial to generate hydrocarbon tolerant species was just started (Liu et al., 2012).

Finding of limiting step in whole photosynthetic CO₂ fixation process is also important to increase lipid productivity. Algae have evolved by developing ability to facilitate the utilization of ambient level of CO₂ by the action of innate CO₂ concentrating mechanisms (Giordano et al., 2005; Raven, 2010). Exogenous CO₂ supplementation recovers cells from CO₂-limitation when cells are exposed such conditions within few hours. However, the photosynthetic activity quickly changes to optimize their ability to exposed conditions since algal cells possess ability to adapt/acclimate to environmental change. The maximal carbon fixation rate and high-CO₂ tolerance are highly depend on microalgal species/strain and therefore CO₂-enrichment is not so beneficial for the improvement of cost and energy performance of microalgal production (Baba & Shiraiwa, 2012). Further investigation is necessary to produce newly-engineered algal cells which exhibit high and efficient CO₂-utilization and -fixation ability with enhanced photosynthesis and lipid productivity.

Acknowledgements

This work was financially supported, in part, by a Grant-in-Aid for Scientific Research from the Core Research of Evolutional Science & Technology (CREST) program from the Japan Science and Technology Agency (JST) (to YS).

Author details

Masato Baba^{1,2} and Yoshihiro Shiraiwa^{1,2}

1 Faculty of Life and Environmental Sciences, University of Tsukuba, Tsukuba, Ibaraki, Japan

2 CREST, JST, Japan

References

- [1] Albro, P. W, & Dittmer, J. C. (1969). The Biochemistry of Long-chain Nonisoprenoid Hydrocarbons. III. The Metabolic Relationship of Long-chain Fatty Acids and Hydrocarbons and Other Aspects of Hydrocarbon Metabolism in *Sarcina lutea*. *Biochemistry*, May 1969, 0006-2960, 8(5), 1913-1918.
- [2] Alvarez, H. M, & Steinbüchel, A. (2002). Triacylglycerols in Prokaryotic Microorganisms. *Applied microbiology and biotechnology*, December 2002), 0175-7598, 60(4), 367-376.
- [3] Awai, K, & Wolk, C. P. (2007). Identification of the Glycosyl Transferase Required for Synthesis of the Principal Glycolipid Characteristic of Heterocysts of *Anabaena* Sp. Strain PCC 7120. *FEMS Microbiology Letters*, November 2007), 0378-1097, 226, 98-102.
- [4] Baba, M, & Shiraiwa, Y. (2012). High-CO₂ Response Mechanisms in Microalgae, In: *Advances in Photosynthesis- Fundamental Aspects-*. Edited by Najafpour, M., In-Tech, 978-9-53307-928-8Rijeka, Croatia.
- [5] Baba, M, Ioki, M, Nakajima, N, Shiraiwa, Y, & Watanabe, M. M. (2012a). Transcriptome Analysis of an Oil-rich Race A Strain of *Botryococcus Braunii* (BOT-88-2) by de novo Assembly of Pyrosequencing cDNA Reads. *Bioresource technology*, April 2012), 0960-8524, 109, 282-286.
- [6] Baba, M, Kikuta, F, Suzuki, I, Watanabe, M. M, & Shiraiwa, Y. (2012b). Wavelength Specificity of Growth, Photosynthesis, and Hydrocarbon Production in the Oil-producing Green Alga *Botryococcus braunii*. *Bioresource technology*, May 2012), 0960-8524, 109, 266-270.
- [7] Banerjee, A, Sharma, R, Chisti, Y, & Banerjee, U. C. (2002). *Botryococcus braunii*: A Renewable Source of Hydrocarbons and Other Chemicals. *Critical reviews in biotechnology*, 0738-8551, 22(3), 245-279.
- [8] Bauersachs, T, Compaoré, J, Hopmans, E. C, Stal, L. J, Schouten, S, & Sinninghe, J. S. (2009). Phytochemistry Distribution of Heterocyst Glycolipids in Cyanobacteria. *Phytochemistry*, September 2009), 0031-9422, 70(17-18), 2034-2039.
- [9] Beller, H. R, Goh, E, & Keasling, J. D. (2010). Genes Involved in Long-Chain Alkene Biosynthesis in *Micrococcus luteus*. *Applied and environmental microbiology*, February 2010), 0099-2240, 76(4), 1212-1223.
- [10] Bouvier, F, Rahier, A, & Camara, B. (2005). Biogenesis, Molecular Regulation and Function of Plant Isoprenoids. *Progress in lipid research*, November 2005), 0163-7827, 44(6), 357-429.
- [11] Casadevall, E, Dif, D, Largeau, C, Gudin, C, Chaumont, D, & Desanti, O. (1985). Studies on Batch and Continuous Cultures of *Botryococcus braunii*: Hydrocarbon

- Production in Relation to Physiological State, Cell Ultrastructure, and Phosphate Nutrition. *Biotechnology and bioengineering*, March 1985), 0006-3592, 27(3), 286-295.
- [12] Chan, D. I, & Vogel, H. J. (2010). Current Understanding of Fatty Acid Biosynthesis and the Acyl Carrier Protein. *The Biochemical journal*, August 2010), 0264-6021, 430(1), 1-19.
- [13] Cheesbrough, M, & Kolattukudy, P. E. (1984). Particulate Preparation from *Pisum sativum*. *Proceedings of the national academy of sciences of the United States of America*, November 1984), 0027-8424, 81, 6613-6617.
- [14] Cho, H, & Cronan, J. E. Jr. ((1995). Defective Export of a Periplasmic Enzyme Disrupts Regulation of Fatty Acid Synthesis. *The Journal of biological chemistry*, March 1995), 0021-9258, 270(9), 4216-4219.
- [15] Dellomonaco, C, Clomburg, J. M, Miller, E. N, & Gonzalez, R. (2011). Engineered Reversal of the β -oxidation Cycle for the Synthesis of Fuels and Chemicals. *Nature*, August 2011), 0028-0836, 476(7360), 355-359.
- [16] Dennis, M. W, & Kolattukudy, P. E. (1991). Alkane Biosynthesis by Decarbonylation of Aldehyde Catalyzed by a Microsomal Preparation from *Botryococcus braunii*. *Archives of biochemistry and biophysics*, June 1991), 0003-9861, 287(2), 268-275.
- [17] Dennis, M, & Kolattukudy, P. E. Enzyme Converts a Fatty Aldehyde to a Hydrocarbon and CO. *Proceedings of the national academy of sciences of the United States of America*, June 1992), 0027-8424, 89(12), 5306-5310.
- [18] Eltgroth, M. L, Watwood, R. L, & Gordon, V. (2005). Production and Cellular Localization of Neutral Long-Chain Lipids in the Haptophyte Algae *Isochrysis galbana* and *Emiliana huxleyi*. *Journal of phycology*, 0022-3646, 41, 1000-1009.
- [19] Frenz, J, Largeau, C, Casadevall, E, Kollerup, F, & Daugulis, A. J. (1989). Hydrocarbon Recovery and Biocompatibility of Solvents for Extraction from Cultures of *Botryococcus braunii*. *Biotechnology and bioengineering*, September 1989), 0006-3592, 34(6), 755-762.
- [20] Fujita, Y, Matsuoka, H, & Hirooka, K. (2007). Regulation of Fatty Acid Metabolism in Bacteria. *Molecular microbiology*, November 2007), 0095-0382X., 66(4), 829-839.
- [21] Giordano, M, Beardall, J, & Raven, J. A. (2005). CO₂ Concentrating Mechanisms in Algae: Mechanisms, Environmental Modulation, and Evolution. *Annual review of plant biology*, June 2005), 1543-5008ISSN: 1543-5008., 56, 99-131.
- [22] Gong, Y, & Jiang, M. (2011). Biodiesel Production with Microalgae as Feedstock: From Strains to Biodiesel. *Biotechnology letters*, July 2011), 0141-5492, 33(7), 1269-1284.
- [23] Gong, Y, Hu, H, Gao, Y, Xu, X, & Gao, H. (2011). Microalgae as Platforms for Production of Recombinant Proteins and Valuable Compounds: Progress and Prospects.

- Journal of industrial microbiology & biotechnology, December 2011), 1367-5435, 38(12), 1879-1890.
- [24] Graham, I. A, & Eastmond, P. J. (2002). Pathways of Straight and Branched Chain Fatty Acid Catabolism in Higher Plants. *Progress in lipid research*, March 2002), 0163-7827, 41(2), 156-181.
- [25] Gu, P, & Ishii, Y. Spencer, T. a & Shechter, I. ((1998). Function-structure Studies and Identification of Three Enzyme Domains Involved in the Catalytic Activity in Rat Hepatic Squalene Synthase. *The Journal of biological chemistry*, May 1998), 0021-9258, 273(20), 12515-12525.
- [26] Hiltunen, J. K, Chen, Z, Haapalainen, A. M, Wierenga, R. K, & Kastaniotis, A. J. (2010). Mitochondrial Fatty Acid Synthesis- An Adopted Set of Enzymes Making a Pathway of Major Importance for the Cellular Metabolism. *Progress in lipid research*, January 2010), 0163-7827, 49(1), 27-45.
- [27] Hlousek-radojic, A, Evenson, K. J, Jaworski, J. G, & Post-beittenmiller, D. (1998). Fatty Acid Elongation Is Independent of Acyl-Coenzyme A Synthetase Activities in Leek and Brassica napus. *Phytochemistry*, 0031-9422, 116, 251-258.
- [28] Hoffmeister, M, Piotrowski, M, Nowitzki, U, & Martin, W. (2005). Mitochondrial trans-2-enoyl-CoA Reductase of Wax Ester Fermentation from *Euglena gracilis* Defines a New Family of Enzymes Involved in Lipid Synthesis. *The Journal of biological chemistry*, February 2005), 0021-9258, 280(6), 4329-4338.
- [29] Hu, Q, Sommerfeld, M, Jarvis, E, Ghirardi, M, Posewitz, M, Seibert, M, & Darzins, A. (2008). Microalgal Triacylglycerols as Feedstocks for Biofuel Production: Perspectives and Advances. *The Plant journal*, May 2008), 0960-7412, 54(4), 621-639.
- [30] Hunter, W. N. (2007). The Non-mevalonate Pathway of Isoprenoid Precursor Biosynthesis. *The Journal of biological chemistry*, July 2007), 0021-9258, 282(30), 21573-21577.
- [31] Imamura, S, Hagiwara, D, Suzuki, F, Kurano, N, & Harayama, S. (2012). Genetic Transformation of *Pseudochoricystis ellipsoidea*, an Aliphatic Hydrocarbon-producing Green Alga. *Journal of general and applied microbiology*, 0022-1260, 58(1), 1-10.
- [32] Inui, H, Miyatake, K, Nakano, Y, & Kitaoka, S. (1982). Wax Ester Fermentation in *Euglena gracilis*. *FEBS Letters*, December 1982), 0014-5793, 150(1), 89-93.
- [33] Inui, H, Miyatake, K, Nakano, Y, & Kitaoka, S. (1984). Fatty Acid Synthesis in Mitochondria of *Euglena gracilis*. *European journal of biochemistry/FEBS*, July 1984), 0014-2956, 142(1), 121-126.
- [34] Ioki, M, Baba, M, Nakajima, N, Shiraiwa, Y, & Watanabe, M. M. (2012a). Transcriptome Analysis of an Oil-rich Race B Strain of *Botryococcus braunii* (BOT-22) by de novo Assembly of Pyrosequencing cDNA Reads. *Bioresource technology*, April 2012), 0960-8524, 109, 292-296.

- [35] Ioki, M, Baba, M, Nakajima, N, Shiraiwa, Y, & Watanabe, M. M. (2012b). Transcriptome Analysis of an Oil-rich Race B Strain of *Botryococcus braunii* (BOT-70) by de novo Assembly of 5'-end Sequences of Full-length cDNA Clones. *Bioresource technology*, April 2012), 0960-8524, 109, 277-281.
- [36] Ioki, M, Baba, M, Bidadi, H, Suzuki, I, Shiraiwa, Y, Watanabe, M. M, & Nakajima, N. (2012c). Modes of Hydrocarbon Oil Biosynthesis Revealed by Comparative Gene Expression Analysis for Race A and Race B Strains of *Botryococcus braunii*. *Bioresource technology*, April 2012), 0960-8524, 109, 271-276.
- [37] Jenke-kodama, H, & Dittmann, E. (2009). Evolution of Metabolic Diversity: Insights from Microbial Polyketide Synthases. *Phytochemistry*, July 2009), 0031-9422, 70(15-16), 1858-1866.
- [38] Jennings, S. M, Tsay, Y. H, Fisch, T. M, & Robinson, G. W. (1991). Molecular Cloning and Characterization of the Yeast Gene for Squalene Synthetase. *Proceedings of the National Academy of Sciences of the United States of America*, July 1991), 0027-8424, 88(14), 6038-6042.
- [39] Jetter, R, & Kunst, L. (2008). Plant Surface Lipid Biosynthetic Pathways and Their Utility for Metabolic Engineering of Waxes and Hydrocarbon Biofuels. *The Plant journal: for cell and molecular biology*, May 2008), 0960-7412, 54(4), 670-683.
- [40] John, U, Beszteri, B, & Derelle, E. Van de Peer, Y.; Read, B.; Moreau, H. & Cembella, A. ((2008). Novel Insights into Evolution of Protistan Polyketide Synthases Through Phylogenomic Analysis. *Protist*, January 2008), 1434-4610, 159(1), 21-30.
- [41] Joyard, J, Ferro, M, Masselon, C, Seigneurin-berny, D, Salvi, D, Garin, J, & Rolland, N. (2009). Chloroplast Proteomics and the Compartmentation of Plastidial Isoprenoid Biosynthetic Pathways. *Molecular Plant*, November 2009), 1674-2052, 2(6), 1154-1180.
- [42] Joyard, J, Ferro, M, Masselon, C, Seigneurin-berny, D, Salvi, D, Garin, J, & Rolland, N. (2010). Chloroplast Proteomics Highlights the Subcellular Compartmentation of Lipid Metabolism. *Progress in lipid research*, April 2010), 0163-7827, 49(2), 128-158.
- [43] Kanda, H, Li, P, Yoshimura, T, & Okada, S. in press) Wet Extraction of Hydrocarbons from *Botryococcus braunii* by Dimethyl Ether as Compared with Dry Extraction by Hexane. *Fuel*, 0016-2361
- [44] Kaneda, T. (1991). Iso- and Anteiso-fatty Acids in Bacteria: Biosynthesis, Function, and Taxonomic Significance. *Microbiological reviews*, June 1991), 0146-0749, 55(2), 288-302.
- [45] Kawano, Y, Saotome, T, Ochiai, Y, Katayama, M, Narikawa, R, & Ikeuchi, M. (2011). Cellulose Accumulation and a Cellulose Synthase Gene Are Responsible for Cell Aggregation in the Cyanobacterium *Thermosynechococcus vulcanus* RKN. *Plant & cell physiology*, June 2011), 0032-0781, 52(6), 957-966.

- [46] Kumar, A, Ergas, S, Yuan, X, Sahu, A, Zhang, Q, Dewulf, J, Malcata, F. X, & Van Langenhove, H. fixation and biofuel production via microalgae: recent developments and future directions. *Trends in Biotechnology*, July), 0167-7799, 28(7), 371-380.
- [47] Kunst, L, & Samuels, L. (2009). Plant Cuticles Shine: Advances in Wax Biosynthesis and Export. *Current opinion in plant biology*, October 2009), 1369-5266, 12, 721-727.
- [48] Largeau, C, Casadevall, E, & Berkaloff, B. (1980). The Biosynthesis of Long-chain Hydrocarbons in the Green Alga *Botryococcus braunii*. *Phytochemistry*, 0031-9422, 19(6), 1081-1085.
- [49] Laws, E. A, Popp, B. N, Bidigare, R. R, Riebesell, U, Burkhardt, S, & Wakeham, S. G. (2001). Controls on the Molecular Distribution and Carbon Isotopic Composition of Alkenones in Certain Haptophyte Algae. *Geochemistry geophysics geosystems*, January 2001), Paper 0025-3227, 2(2000GC000057)
- [50] Lee, S. H, Stephens, J. L, Paul, K. S, & Englund, P. T. (2006). Fatty Acid Synthesis by Elongases in Trypanosomes. *Cell*, August 2006), 0092-8674, 126(4), 691-699.
- [51] Li, F, Wu, X, Lam, P, Bird, D, Zheng, H, Samuels, L, Jetter, R, & Kunst, L. (2008). Identification of the Wax Ester Synthase/acyl-coenzyme A: Diacylglycerol Acyltransferase WSD1 Required for Stem Wax Ester Biosynthesis in *Arabidopsis*. *Plant physiology*, September 2008), 0032-0889, 148(1), 97-107.
- [52] Li, X, Hu, H, & Zhang, Y. (2011). Growth and Lipid Accumulation Properties of a Freshwater Microalga *Scenedesmus* Sp. Under Different Cultivation Temperature. *Bioresource technology*, February 2011), 0960-8524, 102(3), 3098-3102.
- [53] Liu, X, Brune, D, Vermaas, W, & Curtiss, R. III. ((2010). Production and Secretion of Fatty Acids in Genetically Engineered Cyanobacteria. *Proceedings of the National Academy of Sciences of the United States of America*, July 2010), 0027-8424, 6803(19)
- [54] Liu, J, Chen, L, Wang, J, Qiao, J, & Zhang, W. (2012). Proteomic Analysis Reveals Resistance Mechanism Against Biofuel Hexane in *Synechocystis* Sp. PCC 6803. *Biotechnology for biofuels*, January 2012), 1754-6834, 5(1), 68.
- [55] Lohr, M, Schwender, J, & Polle, J. E. W. (2012). Isoprenoid Biosynthesis in Eukaryotic Phototrophs: A Spotlight on Algae. *Plant science : an international journal of experimental plant biology*, April 2012) 0168-9452, 185-186, 9-22.
- [56] Lu, Y. D, Chi, X, Yang, Q, Li, Z, Liu, S, Gan, Q, & Qin, S. (2009). Molecular cloning and stress-dependent expression of a gene encoding Delta(12)-fatty acid desaturase in the Antarctic microalga *Chlorella vulgaris* NJ-7. *Extremophiles: life under extreme conditions*, November), 1431-0651, 13(6), 875-884.
- [57] Marcheschi, R. J, Li, H, Zhang, K, Noey, E. L, Kim, S, Chaubey, A, Houk, K. N, & Liao, J. C. (2012). A Synthetic Recursive "+1" Pathway for Carbon Chain Elongation. *ACS chemical biology*, April 2012), 1554-8929, 7(4), 689-697.

- [58] Matsushima, D, Jenke-kodama, H, Sato, Y, Fukunaga, Y, Sumimoto, K, Kuzuyama, T, Matsunaga, S, & Okada, S. (2012). The Single Cellular Green Microalga *Botryococcus braunii*, Race B Possesses Three Distinct 1-deoxy-D-xylulose 5-phosphate Synthases. *Plant science: an international journal of experimental plant biology*, April 2012), 0168-9452, 185-186, 309-320.
- [59] Metzger, P, & Largeau, C. (2005). *Botryococcus braunii*: a Rich Source for Hydrocarbons and Related Ether Lipids. *Applied Microbiology*, December 2004), 0003-6919, 66, 486-496.
- [60] Metzger, P, Pouet, Y, Bischoff, R, & Casadeval, E. (1993). An Aliphatic Polyaldehyde from *Botryococcus braunii* (A Race). *Phytochemistry*, 0031-9422, 32(4), 875-883.
- [61] Metzger, P, Rager, M, & Largeau, N. C. ((2007). Polyacetals Based on Polymethylsqualene Diols, Precursors of Algaenan in *Botryococcus braunii* Race B. *Organic geochemistry*, April 2007), 0146-6380, 38(4), 566-581.
- [62] Metzger, P, Rager, M, Fosse, N, & Braunicetals, C. Acetals from Condensation of Macrocyclic Aldehydes and Terpene Diols in *Botryococcus braunii*. *Phytochemistry*, September 2008), 0031-9422, 69(12), 2380-2386.
- [63] Miao, X, & Wu, Q. (2006). Biodiesel Production from Heterotrophic Microalgal Oil. *Bioresource technology*, April 2006), 0960-8524, 97(6), 841-846.
- [64] Michinaka, Y, Shimauchi, T, Aki, T, Nakajima, T, Kawamoto, S, Shigeta, S, Suzuki, O, & Ono, K. (2003). Extracellular Secretion of Free Fatty Acids by Disruption of a Fatty acyl-CoA Synthetase Gene in *Saccharomyces cerevisiae*. *Journal of bioscience and bioengineering*, January 2003), 1347-4421, 95(5), 435-440.
- [65] Miyachi, S, Miyachi, S, & Kamiya, A. (1978). Wavelength effects on photosynthetic carbon metabolism in *Chlorella*. *Plant and Cell Physiology*, 0032-0781, 19(2), 277-288.
- [66] Moore, B. S. (2006). Biosynthesis of Marine Natural Products: Macroorganisms (Part B). *Natural product reports*, August 2006), 0265-0568, 23(4), 615-629.
- [67] Nicolaisen, K, Hahn, A, & Schleiff, E. (2009). The Cell Wall in Heterocyst Formation by *Anabaena* Sp. PCC 7120. *Journal of basic microbiology*, February 2009), 0023-3111X., 49, 5-24.
- [68] Niehaus, T. D, Okada, S, Devarenne, T. P, Watt, D. S, Sviripa, V, & Chappell, J. (2011). Identification of Unique Mechanisms for Triterpene Biosynthesis in *Botryococcus braunii*. *Proceedings of the national academy of sciences of the United States of America*, July 2011), 0027-8424, 108(30), 12260-12265.
- [69] Niehaus, T. D, Kinison, S, Okada, S, & Yeo, Y. Bell, S. a; Cui, P.; Devarenne, T. P. & Chappell, J. ((2012). Functional Identification of Triterpene Methyltransferases from *Botryococcus braunii* Race B. *The Journal of biological chemistry*, March 2012), 0021-9258, 287(11), 8163-8173.

- [70] Nojima, Y, Kibayashi, A, Matsuzaki, H, Hatano, T, & Fukui, S. (1999). Isolation and Characterization of Triacylglycerol-secreting Mutant Strain from Yeast, *Saccharomyces cerevisiae*. *The Journal of general and applied microbiology*, February 1999), 0022-1260, 45(1), 1-6.
- [71] Okada, S, Devarenne, T. P, & Chappell, J. (2000). Molecular Characterization of Squalene Synthase from the Green Microalga *Botryococcus braunii*, Race B. *Archives of biochemistry and biophysics*, January 2000), 0003-9861, 373(2), 307-317.
- [72] Okada, S. (2012). Chapter 7 Elucidation of hydrocarbon synthetic pathway in *Botryococcus braunii*, In: *Technology of Microalgal Energy Production and its Business Prospect*. Edited by Takeyama, H. CMC, 978-4-78130-657-5 Osaka, Japan.
- [73] Park, M. (2005). New Pathway for Long-Chain n-Alkane Synthesis via 1-Alcohol in *Vibrio furnissii* M1. *Journal of bacteriology*, February 2005), 0021-9193, 187(4), 1426-1429.
- [74] Poliquin, K, Ershov, Y. V, & Cunningham, F. X. Jr; Woreta, T. T.; Gantt, R. R. & Gantt, E. & ((2004). Inactivation of *sll1556* in *Synechocystis* Strain PCC 6803 Impairs Isoprenoid Biosynthesis from Pentose Phosphate Cycle Substrates In Vitro. *Journal of bacteriology*, July 2004), 0021-9193, 186(14), 4685-4693.
- [75] Portevin, D. De Sousa-D'Auria, C.; Houssin, C.; Grimaldi, C.; Chami, M.; Daffé, M. & Guilhot, C. ((2004). A Polyketide Synthase Catalyzes the Last Condensation Step of Mycolic Acid Biosynthesis in Mycobacteria and Related Organisms. *Proceedings of the national academy of sciences of the United States of America*, January 2004), 0027-8424, 101(1), 314-319.
- [76] Radakovits, R, Jinkerson, R. E, Darzins, A, & Posewitz, M. C. (2010). Genetic Engineering of Algae for Enhanced Biofuel Production. *Eukaryotic cell*, April 2010), 1535-9786, 9(4), 486-501.
- [77] Raven, J. A. (2010). Inorganic Carbon Acquisition by Eukaryotic Algae: Four Current Questions. *Photosynthesis research*, June 2010), 1573-5079ISSN: 0166-8595., 106(1-2), 123-134.
- [78] Renaud, S. M, Zhou, H. C, Parry, D. L, Thinh, L, & Woo, K. C. (1995). Effect of Temperature on the Growth, Total Lipid Content and Fatty Acid Composition of Recently Isolated Tropical Microalgae *Isochrysis* Sp., *Nitzschia Closterium*, *Nitzschia Paleacea*, and Commercial Species *Isochrysis* Sp. (Clone T.ISO). *Journal of Applied Phycology*, December 1995), 0921-8971, 7(6), 595-602.
- [79] Rezanka, T, & Sigler, K. (2009). Progress in Lipid Research Odd-numbered Very-long-chain Fatty Acids from the Microbial, Animal and Plant Kingdoms. *Progress in lipid research*, March 2009), 0163-7827, 48(3-4), 206-238.

- [80] Riekhof, W. R, Sears, B. B, & Benning, C. (2005). Annotation of Genes Involved in Glycerolipid Biosynthesis in *Chlamydomonas reinhardtii*: Discovery of the Betaine Lipid Synthase BTA1. *Eukaryotic cell*, February 2005), 1535-9786, 4(2), 242-252.
- [81] Sakamoto, K, Baba, M, Suzuki, I, Watanabe, M. M, & Shiraiwa, Y. (2012). Optimization of Light for Growth, Photosynthesis, and Hydrocarbon Production by the Colonial Microalga *Botryococcus braunii* BOT-22. *Bioresource technology*, April 2012), 0960-8524, 110, 474-479.
- [82] Sasso, S, Pohnert, G, Lohr, M, Mittag, M, & Hertweck, C. (2011). Microalgae in the Post-genomic Era: A Blooming Reservoir for New Natural Products. *FEMS microbiology reviews*, (September 2011), 0168-6445, 1-25.
- [83] Sato, Y, Ito, Y, Okada, S, Murakami, M, & Abe, H. (2003). Biosynthesis of the Triterpenoids, Botryococenes and Tetramethylsqualene in the B Race of *Botryococcus braunii* via the Non-mevalonate Pathway. *Tetrahedron letters*, September 2003), 0040-4039, 44(37), 7035-7037.
- [84] Satoh, A, Kato, M, Yamato, K. T, Ikegami, Y, Sekiguchi, H, Kurano, N, & Miyachi, S. (2010). Characterization of the Lipid Accumulation in a New Microalgal Species, *Pseudochorocystis ellipsoidea* (Trebouxiophyceae). *Journal of the Japan institute of energy*, September 2010), 0916-8753, 89(9), 909-913.
- [85] Schirmer, A. Rude, M. a; Li, X.; Popova, E. & Del Cardayre, S. B. ((2010). Microbial Biosynthesis of Alkanes. *Science*, July 2010), 0036-8075, 329(599), 559-562.
- [86] Shen, B. (2003). Polyketide Biosynthesis Beyond the Type I, II and III Polyketide Synthase Paradigms. *Current Opinion in Chemical Biology*, April 2003), 1367-5931, 7(2), 285-295.
- [87] Staunton, J, & Weissman, K. J. (2001). Polyketide Biosynthesis: a Millennium Review. *Natural product reports*, June 2001), 0265-0568, 18(4), 380-416.
- [88] Sukovich, D. J, Seffernick, J. L, Richman, J. E, Hunt, K. A, Gralnick, J. A, & Wackett, L. P. (2010a). Structure, Function, and Insights into the Biosynthesis of a Head-to-head Hydrocarbon in *Shewanella oneidensis* Strain MR-1. *Applied and environmental microbiology*, June 2010), 0099-2240, 76(12), 3842-3849.
- [89] Sukovich, D. J, Seffernick, J. L, Richman, J. E, Gralnick, J. A, & Wackett, L. P. (2010b). Widespread Head-to-Head Hydrocarbon Biosynthesis in Bacteria and Role of OleA. *Applied and environmental microbiology*, June 2010), 0099-2240, 76(12), 3850-3862.
- [90] Tabarsa, M, Rezaei, M, Ramezani, Z, & Waaland, J. R. (2012). Chemical Compositions of the Marine Algae *Gracilaria salicornia* (Rhodophyta) and *Ulva lactuca* (Chlorophyta) as a Potential Food Source. *Journal of the science of food and agriculture*, September 2012), 0022-5142, 92(12), 2500-2506.

- [91] Teerawanichpan, P, & Qiu, X. (2010). Fatty acyl-CoA Reductase and Wax Synthase from *Euglena gracilis* in the Biosynthesis of Medium-chain Wax Esters. *Lipids*, March 2010), 0024-4201, 45(3), 263-273.
- [92] Templier, J, Diesendorf, C, Largeau, C, & Casadevall, E. (1992). Metabolism of n-alkadienes in the A race of *Botryococcus braunii*. *Phytochemistry*, 0031-9422, 31(1), 113-120.
- [93] Toney, J. L, Theroux, S, Andersen, R. a, Coleman, A, Amaral-zettler, L, & Huang, Y. (2012). Culturing of the First 37:4 Predominant Lacustrine Haptophyte: Geochemical, Biochemical, and Genetic Implications. *Geochimica et cosmochimica acta*, February 2012), 0016-7037, 78, 51-64.
- [94] Tornabene, T. G. (1981). Formation of Hydrocarbons by Bacteria and Algae. *Basic life sciences*, January 1981), 0036-8075, 18, 421-438.
- [95] Tucci, S, Vacula, R, Krajcovic, J, Proksch, P, & Martin, W. (2010). Variability of Wax Ester Fermentation in Natural and Bleached *Euglena Gracilis* Strains in Response to Oxygen and the Elongase Inhibitor Flufenacet. *The Journal of eukaryotic microbiology*, December 2010), 1066-5234, 57(1), 63-69.
- [96] Venegas-calerón, M, Sayanova, O, & Napier, J. A. (2010). An Alternative to Fish Oils: Metabolic Engineering of Oil-seed Crops to Produce Omega-3 Long Chain Polyunsaturated Fatty Acids. *Progress in lipid research*, April 2010), 0163-7827, 49(2), 108-119.
- [97] Verschoor]. a; Baird, M. S. & Grooten, J. ((2012). Towards Understanding the Functional Diversity of Cell Wall Mycolic Acids of *Mycobacterium tuberculosis*. *Progress in lipid research*, October 2012), 0163-7827, 51(4), 325-339.
- [98] Vioque, J, & Kolattukudy, P. E. (1997). Resolution and Purification of an Aldehyde-generating and an Alcohol-generating Fatty acyl-CoA Reductase from Pea Leaves (*Pisum sativum* L.). *Archives of biochemistry and biophysics*, April 1997), 0003-9861, 340(1), 64-72.
- [99] Wahlen, B. D, Oswald, W. S, Seefeldt, L. C, & Barney, B. M. (2009). Purification, Characterization, and Potential Bacterial Wax Production Role of an NADPH-dependent Fatty Aldehyde Reductase from *Marinobacter aquaeolei* VT8. *Applied and environmental microbiology*, May 2009), 0099-2240, 75(9), 2758-2764.
- [100] Wang, X, & Kolattukudy, P. E. (1995). Solubilization and Purification of Aldehyde-generating Fatty acyl-CoA Reductase from Green Alga *Botryococcus braunii*. *FEBS letters*, August 1995), 0014-5793, 370(1-2), 15-18.
- [101] Weiss, T. L, Roth, R, Goodson, C, Vitha, S, Black, I, Azadi, P, Rusch, J, Holzenburg, A, Devarenne, T. P, & Goodenough, U. (2012). Colony Organization in the Green Alga *Botryococcus braunii* (Race B) Is Specified by a Complex Extracellular Matrix. *Eukaryotic cell*, August 2012), 1535-9786, 11(12), 1424-1440.
- [102] Willis, R. M, Wahlen, B. D, Seefeldt, L. C, & Barney, B. M. (2011). Characterization of a Fatty acyl-CoA Reductase from *Marinobacter aquaeolei* VT8: a Bacterial Enzyme

Catalyzing the Reduction of Fatty acyl-CoA to Fatty Alcohol. *Biochemistry*, December 2011), 0006-2960, 50(48), 10550-10558.

- [103] Yasuno, R, Von Wettstein-knowles, P, & Wada, H. (2004). Identification and Molecular Characterization of the Beta-ketoacyl-[acyl Carrier Protein] Synthase Component of the Arabidopsis Mitochondrial Fatty Acid Synthase. *The Journal of biological chemistry*, February 2004), 0021-9258, 279(9), 8242-8251.
- [104] Yoshida, M, Tanabe, Y, Yonezawa, N, & Watanabe, M. M. (2012). Energy Innovation Potential of Oleaginous Microalgae. *Biofuels*, November 2012), 1759-7269, 3(6), 761-781.

IntechOpen