

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

**4,800**

Open access books available

**122,000**

International authors and editors

**135M**

Downloads

Our authors are among the

**154**

Countries delivered to

**TOP 1%**

most cited scientists

**12.2%**

Contributors from top 500 universities



**WEB OF SCIENCE™**

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Nod-Factor Signaling in Legume-Rhizobial Symbiosis

---

Sulima Anton Sergeevich, Zhukov Vladimir Alexandrovich,  
Shtark Oksana Yurievna, Borisov Alexey Yurievich and  
Tikhonovich Igor Anatolievich

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/61165>

---

## Abstract

Leguminous plants (or Legumes, family Fabaceae) are known to form symbioses with extremely broad range of beneficial soil microorganisms (BSM), representing examples of almost all plant-microbe mutualistic systems. One of the most ecologically important and well-studied legume beneficial symbioses is root nodule (RN) symbiosis (symbiotic association with nitrogen-fixing bacteria). Compared with other interactions of legumes with BSM, RN symbioses demonstrate high level of genetic and metabolic integrity, which implies, *inter alia*, highly specific mutual recognition of partners. In this chapter, we describe the mechanisms of plant-microbe recognition during initial steps of RN symbiosis using the interaction of model legumes - pea (*Pisum sativum* L.), barrel medic (*Medicago truncatula* Gaertn.) and *Lotus japonicus* (Regel.) K. Larsen - with rhizobia as an example. We paid particular attention to symbiotic system of *P. sativum* since pea, besides its importance as a model object of genetics, is also a valuable crop plant. Hence, in conclusion, we discuss the potential to use obtained knowledge for optimizing the broad spectrum of plant adaptive functions and to improve the sustainability of legume crop production.

**Keywords:** legume-rhizobial symbiosis, Nod factor, plant signaling, genetic control

---

## 1. Introduction

Plants are attached immobile organisms and thus have to adapt to their environment in order to survive and reproduce successfully. Usually, plants experience multiple simultaneous

---

impacts from different sources, so they developed complex signaling pathways to effectively detect these impacts and adequately respond to them [1]. Various microorganisms, which are constantly present in the environment, form one of the major factors affecting the life cycle of plants [2, 3]. Although many plant-associated microbes are pathogens that impair plant growth and reproduction, there are also a lot of beneficial (mutualistic) microorganisms able to provide plants with nutrition and additional defense mechanisms. Cooperation with such microorganisms constitutes the universal and highly effective strategy of plants' ecological adaptation, so they tend to form long-lasting associations, which sometimes grow into highly integrated symbioses where one or both partners can develop novel features useful for their survival. Establishing of such symbiotic relationships involves the complicated developmental programs implemented under the joint control by plant and microbial partners and based on the cross-regulation of their genes.

Leguminous plants (or Legumes, family Fabaceae) are known to form symbioses with extremely broad range of beneficial soil microorganisms (BSM), representing examples of almost all plant-microbe mutualistic systems. One of the most ecologically important and well-studied legume-beneficial symbioses is root nodule (RN) symbiosis (symbiotic association with nitrogen-fixing bacteria). Compared with other interactions of legumes with BSM, RN symbioses demonstrate high level of genetic and metabolic integrity, which implies, *inter alia*, highly specific mutual recognition of partners. As legume plant plays a central role in establishing of RN symbiosis, performing functions of initiation, coordination, and regulation of all developmental processes, it possesses complex receptor system capable of accurate identification of microsymbiotic partner. In this chapter, we describe the mechanisms of plant-microbe recognition during initial steps of RN symbiosis using the interaction of model legumes – pea (*Pisum sativum* L.), barrel medic (*Medicago truncatula* Gaertn.), and *Lotus japonicus* (Regel.) K. Larsen – with rhizobia as an example. We pay particular attention to symbiotic system of *P. sativum* since pea, besides its importance as a model object of genetics, is also a valuable crop plant. Hence, in conclusion, we discuss the potential to use obtained knowledge for optimizing the broad spectrum of plant adaptive functions and to improve the sustainability of legume crop production.

## 2. Legume-rhizobial symbiosis: An example of highly integrated plant-microbe system

Nitrogen is an essential component of all living systems, since it is part of the most important biological molecules – DNA and proteins. Molecular nitrogen ( $N_2$ ) in the atmosphere, despite being abundant, is extremely chemically inert and thus cannot be used by the majority of organisms, causing them to compete for more accessible nitrogen sources. Leguminous plants are able to grow in the soil/substrate without any combined nitrogen due to the fixation of atmospheric nitrogen by their symbiotic nodule bacteria (collectively called rhizobia) [4]. Nitrogen fixation occurs within special plant organs – root nodules (or, in some associations, also stem nodules). Development of these organs represents a well-organized process based on the tightly coordinated expression of specialized symbiotic plant and bacterial genes. The


legume nodules provide an ecological niche for bacteria, as well as structure for metabolic/signal exchange between the partners and for the control of symbionts by the hosts [5].

Family Fabaceae contains about 19,000 species divided between three subfamilies (Caesalpi-  
nioideae, Mimosoideae, and Papilionoideae), with more than 700 genera of worldwide  
distribution [6]. With a single exception (*Parasponia*, family Ulmaceae), the ability for symbio-  
ses with rhizobia is restricted to Fabaceae, although in eight related dicotyledonous families  
(Rosid I clade) an ability to form nodules with the nitrogen-fixing actinomycete *Frankia* is  
known [7].

By contrast to legumes, their nitrogen-fixing microsymbionts do not constitute a taxonomically  
coherent group of organisms. The majority of rhizobia belong to the  $\alpha$ -proteobacteria previ-  
ously assigned to the *Rhizobiaceae* family solely on the basis of their ability to nodulate the  
legumes (e.g., *Azorhizobium*, *Bradyrhizobium*, *Mesorhizobium*, *Rhizobium*, *Sinorhizobium*). In the  
last years, several non-rhizobial symbionts capable of forming nodules and fixing nitrogen in  
legume roots have been documented. According to modern conception, bacteria that can form  
RN symbiotic associations (about 44 species of 12 genera) are not clustered in any common  
lineage, instead being distributed in the classes  $\alpha$ -proteobacteria and  $\beta$ -proteobacteria (close  
to *Burkholderia*, *Cupriavidus* and *Ralstonia*) and dispersed over nine monophyletic groups along  
with taxa that do not contain legume symbionts [8]. Recently, some  $\gamma$ -proteobacteria (belong-  
ing to *Escherichia*, *Enterobacter*, and *Pseudomonas* genera) have been discovered that can also  
form nitrogen-fixing nodules with the legumes [9]. All these bacteria (collectively still referred  
to as rhizobia) vary significantly in their overall genome structure, location of “symbiotic”  
(*sym*) genes, their molecular organization and regulation [10, 11]. However, a particular  
legume plant can find the appropriate rhizobial partner (species, or even strain) due to the  
fine-tuned mechanism of molecular interaction.

The development of nitrogen-fixing nodule is complex process that is traditionally divided  
into three major stages: preinfection, root colonization/nodule morphogenesis, and nitrogen  
fixation. On the first stage, the mutual recognition of partners occurs. The interaction between  
micro- and macrosymbiont begins with the activation of bacterial *nod*-genes under the  
influence of flavonoid molecules secreted by the plant root [12, 13]. *nod*-genes provide the  
synthesis of the main bacterial signaling molecule called Nod-factor (NF), which is crucial for  
identification of microsymbiont [14-16]. After the proper reception of Nod-factor, plant  
activates two parallel processes: bacterial penetration into root hair cells via so-called infection  
thread (IT), and differentiation of nodule from the root cortex. IT is a special structure  
generated by invagination of plant cell membrane, covered with plant-derived cell wall and  
filled with matrix produced by both plant and bacteria. It grows into root hair cell and then to  
the cortex where nodule tissues are formed (Figure 1) [17].

The key stage of nodule development is conversion of bacteria into the form of intracellular  
symbionts through endocytosis-like process. Herein, the distal area of IT transforms into  
structure called infection droplet (ID), which releases membrane vesicles containing bacteria  
into plant cytoplasm. After leaving IT, rhizobia for some time retain their size and shape,  
subsequently differentiating into a specific form called bacteroids [18]. Compared to free-living  
bacteria, bacteroids have significantly (about 3-7 times) increased size and more complex



**Figure 1.** General scheme of RN symbiosis formation and functioning in pea. From left to right: three major stages of symbiosis, namely preinfection, root colonization/nodule morphogenesis, and nitrogen fixation. F – flavonoids excreted by the plant, NF – Nod-factors excreted by nodule bacteria, IT – infection thread, B – bacteroids. *Sym10* and *Sym2*, *Sym37* – stages on which symbiosis is blocked in case of corresponding pea mutants/genotypes.

shape, which can be round, pear-shaped, Y- or X-like, depending on specific symbiotic system. After the aforesaid differentiation, the synthesis of nitrogenase (the enzyme catalyzing reduction of  $N_2$  into  $NH_4^+$ ) and other proteins involved in nitrogen fixation is activated in bacterial cells [19].

Bacteroids are embedded into a membrane structure named symbiosome, which are derived from membrane vesicle originating from ID. They are organelle-like units of plant cell responsible for nitrogen fixation. Symbiosome formation as well as bacteroid differentiation is induced by plant. Peri-bacteroid membrane (PBM) that surrounds bacteroids is an active interface of RN symbiosis where exchange of metabolites between symbionts occurs [19, 20]. Plant cells containing symbiosomes also undergo the deep differentiation, increasing the amount of their membrane structures (endoplasmic reticulum and the Golgi complex), which participate in the development of PBM and biosynthetic processes. Many proteins associated with nitrogen fixation appear in these cells *de novo*.

The developmental program described above is typical only for evolutionary advanced legumes belonging to the inverted repeat-lacking clade (IRLC) of Papilionoideae, such as *Medicago*, *Pisum*, or *Trifolium* (clover). They form so-called “indeterminate” nodules which are characterized by stable apical meristem and division into histological zones with constantly renewed  $N_2$ -fixing zone. Rhizobia in these nodules undergo terminal bacteroid differentiation and cannot revert to free-living form [21, 22]. Other legumes such as *Lotus* or *Phaseolus* (bean),



however, form morphologically more simple “determinate” nodules, where apical meristem exists only for several days, nitrogen-fixing zone is not strongly expressed, and infected ( $N_2$ -fixing) cells intermingle with noninfected ones [21]. Bacteroids in determinate nodules show no sign of terminal differentiation as they usually maintain their normal bacterial size, genome content, and reproductive capacity lacking from those in indeterminate nodules [22].

Several Papilionoideae members, such as *Arachis* and *Stylosanthes*, demonstrate the reductive scheme of nodule development: rhizobia invade roots through the cracks of epidermis, and instead of IT they are brought into cytoplasm by the direct endocytosis from intercellular space [6, 23]. Even more primitive morphology of symbiosis is typical for members of Caesalpinioideae subfamily, as they lack endocytosis step, and nitrogen fixation occurs within modified persistent ITs called “fixation threads” [24]. This is also relevant for evolutionary primitive woody plants from Papilionoideae: *Andira* and *Hymenolobium*, and for *Parasponia*.

Such a complicated system of biological nitrogen fixation will work properly only when suitable partners meet each other in soil. This rendezvous becomes possible owing to reciprocal molecular signal exchange, which is not exhaustively studied to date.

## 2.1. Specificity of legume-rhizobial symbiosis

Root-nodule symbiosis is well known as highly specific plant-microbe interaction. According to the early surveys of symbiotic specificity [25], legumes were suggested to comprise a range of taxonomically restricted cross-inoculation groups (CIG) within which the free cross-inoculation occurs, while the species from different groups do not cross-inoculate.

The best studied examples of this classification are represented by four CIG: “*Trifolium* – *Rhizobium leguminosarum* bv. *trifolii*,” “*Pisum*, *Vicia*, *Lathyrus*, *Lens* – *R. leguminosarum* bv. *viciae*,” “*Galega* – *R. galegae*,” “*Medicago*, *Melilotus*, *Trigonella* – *Sinorhizobium meliloti*, *S. medicae*.” However, it was demonstrated later that such strictly defined specificity is limited to the herbage papilionoid legumes growing in temperate zones and representing the so-called Galegoid complex [26, 27]. Other legumes, including the majority of tropical species, tend to broad their symbiotic specificity, where cross-inoculation is possible between tribes, subfamilies, and even with non-legume plant *Parasponia* [28].

The analysis of CIG structure for both strictly and broadly specific legumes has shown that plant specificity towards rhizobia has good correlation with plant taxonomy on the genus or tribe level. It was also revealed that specificity of nodule formation does not correlate with symbiotic efficiency, i.e., efficiency of nitrogen fixation: several bacterial strains form normal nodules with one plant species, and are inactive (not able to fix nitrogen, Fix<sup>-</sup>) with another [26]. This could be due to the fact that nodulation is an early stage of symbiosis similar (and supposedly related) to pathogenic interaction, and is based on strict cross-activation of plant and bacterial genes (“gene-for-gene” interaction), while nitrogen fixation occurs on the later stages for which “gene-for-gene” interaction is not typical.


Moreover, it is specificity that makes possible the natural selection of effective **symbiotic pairs**, but not the single “symbiotically effective” plant or single “symbiotically effective” microorganism. On the other side, specificity of legume-rhizobial symbiosis should be

somewhat associated with nitrogen-fixing intensity, upon which is based the ecological efficiency of cooperation; otherwise it would not be an evolutionary advantage. The majority of “Galeoid complex” members have both narrow specificity and effective nitrogen fixation, suggesting that these two features are connected, though specificity of recognition is obviously not the only condition required for effective symbiosis.

It is also important to note that the range of potential symbiotic partners can vary for both bacteria and plants. Symbiotic pair *Trifolium–Rhizobium leguminosarum* bv. *trifolii* represents one side of this continuum, as they are the only possible partners for each other. On the opposite side are *Phaseolus vulgaris* and *Vigna unguiculata*, which are able to exchange their symbionts with many unrelated legume species [25]. In bacteria, the *Sinorhizobium fredii* strain NGR234 was shown to interact with more than 120 plant species from all three Fabaceae subfamilies, as well as with *Parasponia*, thus being the most “unscrupulous” strain known so far [29]. The most striking feature of this strain is that its genome, although not particularly large (6.9 Mbp), encodes more different secretion systems than any other known rhizobia and probably most known bacteria [30]. These, among others, include type III and type IV secretion systems which allow bacteria to direct effector proteins or DNA into the cytoplasm of their eukaryotic hosts. There seems to be a correlation between the host range of rhizobia and the number of specialized protein secretion systems they have, as “classic” narrow-host-range rhizobia such as *S. meliloti* and *R. leguminosarum* carry neither type III nor type IV secretion systems. Furthermore, NGR234 is shown to secrete a large family of NFs that are variously 3-O, 4-O, or 6-O carbamoylated, which are N-methylated, and which carry a 2-O-methyl-fucose residue that may be either 3-O sulfated or 4-O acetylated (see below) [29]. Since no other rhizobia synthesize such a large family of NFs, it should be proposed as one of the main aspects contributing to the broad host range of NGR234 [17, 31]. Another possible aspect is that NGR234 not only treats the legume root to a large palette of NFs, but that their concentration is much higher than in even very closely related rhizobia [32].

## 2.2. Initial steps of rhizobium-legume symbiosis

The specificity of legume-rhizobia interactions is expressed mostly during the preinfection stage when rhizobia recognize the roots of appropriate host plants and colonize their surfaces. When the root-excreted signals (in particular, flavonoids) are perceived by bacteria, they activate the bacterial nodulation genes (*nod/nol/noe*) [13]. These genes control the synthesis of lipo-chito-oligosaccharidic (LCO) nodulation factors (Nod-factors) which induce the early stages of RN symbiosis development. NFs represent the unique group of bacterial signal molecules not known outside legume-rhizobia symbiosis. They are among the most potent developmental regulators: their effect is expressed at concentrations merely of  $10^{-8}$  –  $10^{-12}$  M. The core structure of these molecules, common for all rhizobia species, consists of 3-6 residues of N-acetylglucosamine and of a fatty acid (acyl) chain (Figure 2). The type of symbiotic specificity is dependent mainly on the chemical modifications in NF structures [14-16]. However, a sufficient impact to the host specificity of RN symbiosis can also be made by the interactions between bacterial surface molecules (some polysaccharides and proteins) [33, 34] and the lectins located on the root hair surfaces, as well as by means of NFs secretion [35].



$n = 2$  or  $3$ ; regarding Afghan peas, see below (3.2).

**Figure 2.** Example of Nod-factor excreted by *Rhizobium leguminosarum* bv. *viciae* strain TOM nodulating Afghan peas.

Rhizobia possess a wide range of genes involved in the early stages of nodulation, i.e., the NFs production [36]. Genes which are common to all rhizobia – *nodA*, *nodB*, *nodC*, and their regulator *nodD* – are responsible for NF core structure synthesis [37]. The other genes specific for particular species or strains control various modifications of signaling molecule. The difference in the spectrum of hosts possible for microsymbiont to interact with is based on the variety of combinations of different *nod*-genes. For example, presence of gene *nodE*, which encodes protein similar to fatty acid synthase in several genera of rhizobia, provides modification of fatty acid moiety on the nonreducing end of NF molecule, thereby affecting the ability of bacteria to nodulate certain plant species [38, 39]. Genes *nodH*, *nodP*, and *nodQ* found in *Sinorhizobium meliloti* control the specific NF modification – the O-sulfation of reducing end – which makes it recognizable for *Medicago* receptors [40]. Overall, each strain of rhizobia is characterized by specific set of *nod*-genes, which together form the “molecular key” suitable for plant receptor. It is significant to note that most rhizobia secrete an assortment of NFs varying in their structure instead of just one particular kind [41, 42]. Thereby, the symbiotic success of bacteria could be directly connected with diversity of NFs they are able to produce, and “molecular key” rather becomes the “set of lock picks,” with secretion systems and surface molecules being additional tools in it (see above).

By perceiving the NF, plant starts various processes in root tissues. In particular, signaling molecule is required for the activation of plant genes in the epidermis cells and pericycle, as well as for mitotic reactivation of cortical cells and the formation of IT. Genes responsible for proper NF reception were first discovered in mutants of *Lotus japonicus* lacking any response



to NFs [43, 44]. These genes were named *NFR1* and *NFR5*, for Nod-Factor Receptor. Cloning of these genes revealed that they encode receptor-like kinases comprising LysM domains (LysM-RLK). LysM domains occur in a variety of proteins in bacteria and eukaryotes and have been shown to bind glycan-containing ligands (such as chitin) [45]. They consist of a repetition of a small motif typically containing from 44 to 65 amino acid residues – the LysM sequence, or LysM module [46, 47]. One LysM sequence has a  $\beta\alpha\alpha\beta$  secondary structure with the two helices packing onto the same side of an antiparallel  $\beta$  sheet. Multiple LysM modules in a protein are often separated by small Ser-, Thr-, and Asn-rich intervening sequences [48].

Only in plants are LysM domains associated with a kinase-like domain [49] forming two main LysM-RLK gene families: the LYK family and the LYR family. All the LysM-RLKs are predicted to contain three LysM modules, although these modules exhibit a high degree of divergence, both within a protein and between proteins. It is considered that the initial function of LysM-RLKs has been recognition of chitin-based signal molecules produced by hostile microbes (termed as MAMPs (“microbe-associated molecular patterns”) or PAMPs (“pathogen-associated molecular patterns”)), similar to the function of CERK1 receptor-like kinase from *Arabidopsis thaliana* [2]. Based on microsynteny between genomic regions around LysM-RLK genes in legumes and non-legumes (*A.thaliana*, rice) plants, it has been speculated that these genes are the descendants of a common ancestor [50]. Zhang et al. (2007) [51] proposed that in Leguminosae LysM-RLKs have undergone further duplication and diversification, with some LysM-RLKs acquiring the ability to perceive bacterial NFs, leading to mutually beneficial endosymbiosis with rhizobia. One aspect of this diversification is the adaptation of extracellular LysM domains to recognize specific structures of NFs, while another being evolution of the intracellular kinase domains to switch the signals from cascades inducing defense responses to symbiotic gene cascades. Recently, the function of NFRs as NF receptors was confirmed by demonstration of their ability to directly bind NF molecule *in vitro* [52].

In *Medicago* and pea, which belong to IRLC (see above), NF perception seems to be more complicated than in *Lotus*. Genes orthologous to *NFR1* and *NFR5* were identified in *Medicago truncatula* (*LYK3* and *NFP*) and in *Pisum sativum* (*Sym37* and *Sym10*), with careful description of corresponding mutant phenotypes [44, 53-55]. While phenotype of *nfp* and *sym10* mutants (in *Medicago* and pea, respectively) coincided with that of *nfr5* mutants in *Lotus*, mutations in genes *lyk3* and *sym37* (orthologs of *NFR1*) led to significantly different phenotype – successful penetration of bacteria into root hair with subsequent block of IT progress, instead of complete absence of responses to rhizobia [55, 56]. These data support the “two-receptor” model of Nod-factor perception proposed more than 20 years ago [40]. According to this model, which was developed on the base of the infection phenotype of several *S. meliloti nod* mutants, there are two different types of NF receptors – the “recognition” (or “signaling”) receptor inducing early responses with high affinity for Nod-factor and low requirements toward its structure, and the “entry” receptor that controls penetration of bacteria into plant cell and has more stringent demands [40].

It is significant to note that *NFR5* (and its homologs, *NFP* in *Medicago* and *Sym10* in *Pisum*) lacks the independent kinase activity and thus can function properly only in complex with active kinase (which is suggested to be *NFR1*) [52]. It can be assumed, based on the above, that in general the “recognition” receptor (*NFR5*, *NFP* or *Sym10*) perceives NF and afterwards

forms complex with another receptor possessing kinase activity (NFR1, LYK3 or Sym37, respectively), thus constituting the “entry” receptor. Still, results of genome and transcriptome sequencing in *Lotus*, *Medicago* and pea show that legumes possess more than 10 genes of receptor kinases similar by structure to the aforementioned ones. So, the system of NF receptors could be actually much more complicated, suggesting that the overall mechanism of NF perception is probably even more intricate than was thought before.

### 3. Molecular genetics of Nod-factor signaling in legumes

As reviewed in our recent publication [57], plant genes involved in development of RN symbiosis may be divided into two groups, according to approach which was used for the gene identification. The first group, *Sym*-genes, had been identified with the use of formal genetic analysis (started from selection of plant mutants defective in nodule development). The other group of genes called nodulins was identified by molecular genetic methods, through identification of proteins and/or RNAs synthesized *de novo* in root nodules.

The large sizes of genomes of crop legumes (e.g., soybean or pea) in which the formal genetics of symbioses was initially developed, as well as low capability for genetic transformation, complicate greatly the cloning of symbiotic genes, analysis of their primary structures, and gene manipulations. Therefore, in the early 1990s, *Lotus japonicus* [58] and *Medicago truncatula* [59, 60] have been introduced in symbiogenetic studies as model plants. These species are characterized by relatively small genomes (470-500 Mb; [61]) and can be easily genetically transformed [60, 62-64]. In addition, the short life cycle and high seed productivity made them attractive and convenient model objects for studying molecular bases of RN symbioses, as well as other types of plant-microbial symbioses.


The analysis of signaling pathway in RN symbiosis was started with experimental mutagenesis. Large-scale programs of insertion, chemical and X-rays mutagenesis, performed by different research groups, resulted in generation of numerous symbiotic mutants in *L. japonicus* and *M. truncatula* [65, 66] which allowed researchers to identify and characterize a series of *Sym*-genes. The genes involved at the initial stages of nitrogen-fixing symbiosis (named “early *Sym*-genes”) were of primary interest, allowing dissection of the mechanisms by which the NF signal is perceived and transduced by host plants.

#### 3.1. Nod-factor signaling in model legumes

After the first step of NF reception implemented by LysM-receptor kinases (described above), the symbiotic signal is transmitted to the pathway named Common Symbiosis Pathway (CSP), for it shares components with another interaction – arbuscular mycorrhiza (AM) symbiosis, the association with obligate biotrophic fungi of phylum *Glomeromycota*. Arbuscular mycorrhiza is formed by at least 80% of contemporary land plants and is believed to be the most ancient plant-microbe symbiosis which has played a decisive role in plants adaptation for terrestrial life [67-69]. AM is the main source of plants’ phosphoric nutrition, although in many temperate and boreal species it is supplemented or even completely replaced by other forms

of mycorrhiza (ectotrophic, ericoid) with various representatives of the *Ascomycota* and *Basidiomycota*, and for some plants (orchids) fungi supply not only mineral nutrition, but also organic carbon compounds [69, 70]. Being the first beneficial association with microorganisms known for plants (occurred approximately 400 million years ago), AM is considered as an ancestor for other mutualistic plant-microbe interactions, such as RN symbiosis. Therefore, it is supposed that NF signaling evolved on the base of previously existing AM signaling. Intriguingly, arbuscular mycorrhizal fungi excrete a set of chitin-derived Myc-factors structurally similar to Nod-factors [71], which also serve as the signaling molecules. It still remains unknown, however, how exactly the Myc-factors are perceived by plants.

The first player in the CSP was identified more than 10 years ago. It is LRR-receptor kinase, or SymRK (symbiotic receptor kinase) described for *Lotus* as SymRK (Symbiotic Receptor Kinase) and for *Medicago* as NORK (Nodulation Receptor Kinase) [72, 73]. In pea, the gene *Sym19* is orthologous to *SymRK* in *Lotus* and *NORK* (also known as *DMI2*, for Doesn't Make Infections) in *Medicago* [72]. Ligand of this receptor kinase is not known as yet (Figure 3). Interestingly, the activity of SymRK is also required for proper progression of late symbiotic stages, at least for rhizobial infection [74]. SymRK kinase domain has been shown to interact with 3-hydroxy-3-methylglutaryl CoA reductase 1 (HMGR1) from *M. truncatula* [75], and an ARID-type DNA-binding protein [76]. These results suggest that SymRK may form complex with key regulatory proteins of downstream cellular responses. Symbiotic Remorin 1 (SYMREM1) from *M. truncatula* and SymRK-interacting E3 ligase (SIE3) from *L. japonicus* have also been shown to interact with SymRK [77, 78].



From left to right: stages of symbiosis.

**Figure 3.** Receptor kinases of pea participating in nodulation signaling.

The symbiosis receptor kinase SymRK acts upstream of the NF-induced  $\text{Ca}^{2+}$  spiking in the perinuclear region of root hairs within a few minutes after NF application [79]. Perinuclear calcium spiking involves the release of calcium from a storage compartment (probably the

nuclear envelope) through as-yet-unidentified calcium channels. To date, it is known that the potassium-permeable channels might compensate for the resulting charge imbalance and could regulate the calcium channels in plants [80-84]. Also, nucleoporins NUP85 and NUP133 (described only in *Lotus* so far) are required for calcium spiking, although their mode of involvement is currently unknown. Probably, they might be a part of specific nuclear pore subcomplex that plays a crucial role in the signal process requiring interaction at the cell plasma membrane and at nuclear and plastid organelle membranes to induce a  $\text{Ca}^{2+}$  spiking [85-86]. Recently, the third constituent of a conserved subcomplex of the nuclear pore scaffold, NENA, was identified as indispensable component of RN endosymbiotic development [87].

$\text{Ca}^{2+}$  spikes are supposed to activate a calcium- and calmodulin-dependent protein kinase (CCaMK). This kinase contains an autoinhibition domain which, when removed, leads to a spontaneous activation of downstream transcription events and induction of nodule formation even in the absence of rhizobia [88]. Thus, CCaMK appears to be a general “manager” for both RN and AM symbioses and the last member of Common Symbiosis Pathway, because the next steps of nodulation signaling are independent from those of AM: the mutations in downstream *Sym*-genes do not affect the AM symbiotic properties of legume. Interestingly, mutations in any *Sym*-genes do not influence the defense reactions, suggesting that signaling pathways of mutualistic symbioses and pathogenesis are sufficiently different.

The CCaMK is known to form a complex with CYCLOPS, a phosphorylation substrate, within the nucleus [89]. *cyclops* mutants of *Lotus* severely impair the infection process induced by the bacterial or fungal symbionts. During RN symbiosis, *cyclops* mutants exhibit the specific defects in IT initiation, but not in the nodule organogenesis [90], indicating that CYCLOPS acts in an infection-specific branch of the symbiotic signaling network [35]. *Cyclops* encodes a protein with no overall sequence similarity to proteins with known function, but containing a functional nuclear localization signal and a carboxy-terminal coiled-coil domain.

It is supposed that CCaMK with help of CYCLOPS probably phosphorylates the specific transcription factors already present in cell, NSP1 and NSP2, which influence the changes of expression in several genes related to the symbiosis development [91, 92]. The activity of these proteins leads to the transcriptional changes in root tissues, for instance, increasing the level of early nodulins ENOD40, ENOD11, ENOD12, ENOD5, which are known to be the potential regulators of IT growth and nodule primordium formation [93-95]. Also, the changes in cytokinin status of plant are detected, followed by up-regulation of genes encoding for RN symbiosis-specific cytokinin receptors [96-98]. Moreover, transcription regulators NIN and ERN are to be induced specifically downstream of the early NF signaling pathway in order to coordinate and regulate the correct temporal and spatial formation of root nodules [99-102].

The presented genes are responsible for the signal cascade which is aimed to induce the nodulin genes involved in building the symbiotic structures and implementing their biochemical functions. It is supposed that this signaling pathway did not appear *de novo* in legumes when they become able to form nodules, but was developed from already existing system of AM formation into which the novel, nodule-specific genes were recruited. Still, new genes had been involved in RN symbiosis development, especially those encoding the receptors recognizing hormones (e.g., cytokinins) and hormone-like molecules (Nod-factors).



Another important signaling process in RN symbiosis is an autoregulation of nodule formation. It takes place after successful mutual partners' recognition and signal exchange. It is considered that legume host controls the root nodule numbers by sensing the external and internal cues. A major external cue is the concentration of soil nitrate, whereas a feedback regulatory system where nodules formed earlier suppress further nodulation through shoot-root communication is an important internal cue. The latter is known as the autoregulation of nodulation (AON), and is believed to consist of two long-distance signals: a root-derived signal that is generated in infected roots and transmitted to the shoot; and a shoot-derived signal that inhibits nodulation systemically [103-104]. Therefore, AON represents a strategy through which the host plant can balance the symbiotrophic N nutrition with the energetically more "cheap" combined N nutrition.

Recent findings on autoregulation of nodulation suggest that the root-derived ascending signals to the shoot are short peptides belonging to the CLE peptide family [105] [106]. The leucine-rich repeat receptor-like kinase HAR1 of *Lotus* and its homologues in *M. truncatula* and *P. sativum* (SUNN and Sym29, respectively) mediate AON and also the nitrate inhibition of nodulation, presumably by recognizing the root-derived signal [107-110] (Figure 3).

It was suggested that NF signaling induces expression or posttranslation processing of CLE peptides, which likely function as ascending long-distance signals to the shoot [110]. Thus, NF signaling is related to autoregulation as well, but in some indirect way. It is also worth noting that NF signaling pathway appears to work in mature nodules, since aforementioned "early nodulation genes" belonging to CSP, as well as NF receptor kinase genes, are highly expressed in nodule tissues (76, 111). Perhaps the active NF signaling is needed to prevent the induction of defense-like responses and/or to restrict the release of rhizobia into precise cell layers, thus regulating the formation of symbiotic interface [112].

### 3.2. Pea (*Pisum sativum* L.) as a unique example of increased specificity in plant-microbe interaction

Being one of the most ancient crops known to humanity, nowadays garden pea (*Pisum sativum* L.) is widely distributed in the world. According to the recent data, pea is a third most important legume for food industry, following beans and soybeans [113]. It is also the popular model for various genetic and physiological researches, including the studying of symbiosis with nodule bacteria. Despite the fact that work with pea is complicated by the presence of some negative properties, such as relatively large (about 4000 Mb) genome, low seed productivity, and poor transformation capability, the use of this object in study of symbiotic relationships continues and brings significant results.

There are several pea genes known to participate in NFs' reception, with the most interesting of them being *Sym2*. This gene was first described in the 1970s as determinant of "resistance" to nodulation in pea cultivars from Afghanistan and Iran [114, 115]. While being unable to form nodules with the majority of natural *Rhizobium leguminosarum* bv. *viciae* (*Rlv*) strains obtained from European soils, these cultivars have demonstrated the ability to interact normally with strains from the Middle East, such as strain *Rlv* TOM [115]. This feature is controlled by specific recessive allele of *Sym2* named "Afghan allele" (*Sym2<sup>A</sup>*). Presence of



*Sym2<sup>A</sup>* in homozygous state leads to block of infection thread progression in the root hair, similarly to phenotype of *sym37* mutants [55]. Later it was shown that *Rlv* strains able to nodulate “Afghan” cultivars have special gene called *nodX*, which is involved in the modification of NF structure [116, 117]. *NodX* encodes the acetyltransferase providing O-acetylation on reducing end of NF sugar backbone. Thus, only *nodX*-modified NFs can be recognized by plants with *Sym2<sup>A</sup>* allele, although Ovtsyna et al. (2000) [118] show that fucosylation on the same position controlled by *nodZ* gene can also induce nodulation of “Afghan” peas.


More than 20 years ago, *Sym2* was localized on the pea genetic map. Using RAPD (Random Amplification of Polymorphic DNA) markers, Kozik and colleagues [119] created the detailed map of pea I linkage group fragment including *Sym2* and a few other symbiotic genes (such as *Nod3* and *PsENOD7*). Based on the fact that plants with *Sym2<sup>A</sup>* allele show the “Afghan” phenotype then exposed to NF with specific structure, it was suggested that *Sym2* protein could act as an “entry” receptor during preinfection stage (similar to *NFR1* in *Lotus* or *LYK3* in *Medicago*).

When *Pisum* gene *Sym37* was shown to be orthologous for *NFR1* [55], it was at first proposed as a candidate for *Sym2*. This was strongly supported by the fact that the missense mutation in *Sym37* carried by *Pisum* mutant line *RisNod4* led to *Nod<sup>-</sup>* phenotype (the absence of nodulation), which could be suppressed by *Rlv* strain A1 known to produce broad specter of NFs, including *nodX*-modified one [55]. However, the paralogue of *Sym37*, gene *K1*, was discovered shortly after, the similar structure of which indicated a possible involvement in the reception of NF, although the purpose of this additional NF receptor remained unclear.

The comparison of *Sym37* and *K1* nucleotide sequences obtained from “Afghan” (*Sym2<sup>A</sup>*) and “European” pea varieties, as well as amino acid sequences of their corresponding proteins, shows that neither of these genes possesses any features correlating with “Afghan” phenotype [55]. Thus, there must be another determinant corresponding to *Sym2*. Recently, the promising candidate was found – the gene named *LykX* by the authors, which is the second paralogue of *Sym37* localized in the same region of the pea genome (Sulima et al., 2015, in preparation). Analysis of the *LykX* protein sequences revealed that there are amino acid substitutions within first LysM module of receptor domain typical for plants with “Afghan” phenotype [120]. Simultaneously, Li and colleagues [121] compared the sequence of *Sym37* from series of pea genotypes that differ in interaction with rhizobia mutant on *nodE* gene determining the structure of fatty acid on nonreducing end of NF. It was shown that the efficiency of interaction with mutant strain strictly correlates with particular variation of *Sym37*. Similar situation was observed for interaction between *nodX* and *Sym2* (*LykX*) genes: “Afghan” pea varieties requiring NF with additional acetyl group on reducing end of molecule also display characteristic features in structure of receptor protein *LykX*.

We proposed a model, based on the above, according to which the less specific “recognition” receptor (*Sym10*, perhaps in complex with other proteins) perceives the NF signal *per se* and “anchors” NF molecule on the membrane, subsequently “presenting” it to other components of reception complex, with reducing end being tested by *Sym2* (*LykX*), and nonreducing by *Sym37* [122]. Only if all participants in the process react positively will the signal be considered as adequate, and symbiogenesis will start properly (see Figure 4). So, in pea not only one

ortholog of *Lotus NFR1*, but two closely related paralogs – *Sym37* and *Sym2* – are involved in genetic control of Nod-factor reception. This is not surprising, if we take into account the complexity of Nod-factor molecule and the importance of its proper recognition for successful development of symbiosis.



**Figure 4.** Hypothetical model for precise recognition of Nod-factor structure by receptor kinases in pea. The model is proposed by Dr. V.A. Zhukov (ARRIAM, St. Petersburg, Russia). At first step, less specific receptor (probably, *Sym10*) anchors NF molecule onto the membrane; then it presents it to *Sym37*, which tests the structure of the nonreducing end, and to *Sym2*, which tests the structure of reducing end. When both *Sym37* and *Sym2* bind NF, they activate downstream components of signal transduction pathway.

## 4. Conclusion

Among all the multicellular eukaryotes, plants have the greatest need for the beneficial interaction with microorganisms, as they lack active movement and therefore cannot choose more advantageous habitat. That kind of restriction can be compensated by the ability of photosynthesis, as carbon compounds produced by plants are a significant stimulus for various microbes to cooperate with them. As a result of such cooperation, plant acquires an access to the adaptations of microsymbiont, and *vice versa*, according to a principle of genome complementarity that was recently formulated by Prof. I.A. Tikhonovich and Dr. N.A. Provorov (ARRIAM, Russia) [122]. It means that, in spite of lacking the nitrogenase genes in its own genome, plant “exploits” corresponding part of microorganism’s genome in order to implement biological nitrogen fixation, while rhizobia “exploit” plant genes controlling

photosynthetic apparatus, and so forth. Thereby the plant-microbe system acquires an advantage over plants and microbes that compete for survival separately.

The role of symbioses in the evolution of life, and plants in particular, cannot be underestimated. One can state that the tendency to establish mutually beneficial associations with microorganisms is an essential feature of plants, which has a wide variety of manifestations through a long coevolution of symbiotic partners. Photosynthesis itself, the main distinctive feature of plants, is provided by chloroplasts – the descendants of ancient symbiotic cyanobacteria. According to modern conception, plant colonization of land was possible primarily due to the symbiotic association with arbuscular mycorrhiza fungi. AM, in turn, is considered as a basis for the development of highly specific root-nodule symbiosis characteristic for legume plants. The possible path of the AM origin and its connection with RN was largely understood by studying *Geosiphon pyriformis* – the only representative of the phylum *Glomeromycota* that does not form symbiotic association with higher plants. Instead, it contains intracellular symbiotic nitrogen-fixing cyanobacteria of the *Nostoc* genus which are essential for its proper nutrition and development [123, 124]. The intensive exchange of products of nitrogen, carbon, and phosphorus metabolism between partners indicates that mechanisms of reciprocal nutrients' transport probably emerged in symbiotic systems formed by *Geosiphon* and *Nostoc* ancestors and lately have been recruited in the evolution of AM [124, 125]. The transition from *Geosiphon-Nostoc*-type association to AM could occur through an intermediate “triple” symbiosis including plant, common ancestor of AM fungi, and *Geosiphon*, and ancestral forms of *Nostoc*, with subsequent loss of cyanobiont. It should be noted that ancient symbiotic fungi presumably carried additional bacterial symbionts both on the surface and in the cytoplasm. In the cells of modern *Glomeromycota*, including *Geosiphon*, various symbiotic bacteria are found, including those capable of nitrogen-fixation (close to  $\beta$ -proteobacteria of *Burkholderia* genus, some members of which were shown to form the RN symbiosis with legumes; see above) [126]. Thus, the AM symbiosis could be the direct “gateway” for introducing symbiotic bacteria, including the ascendants of modern rhizobia, into plant tissues. This suggestion is also supported by the existence of CSP and the similarity of rhizobial and fungal signal molecules.

Emergence of Nod-factor signaling was among the most important factors that determined the evolutionary success of legume-rhizobial symbiosis. The wide variety of Nod-factors as well as finely tuned receptor system in plants ensure that only specific partners will meet each other in the soil and consequently form a superorganism with high level of genetic and metabolic integration. This appears to be a basis for evolution of the efficiency of symbiotic pairs, instead of single organisms – the results we now observe.

Legumes provide both an important food source for humanity and a unique model for investigation of the evolution and the underlying genetic mechanisms of mutualistic plant-microbe symbioses. Further studies of the genetic bases of signal interactions between plants and microbes can provide more information about evolution of such a mutually beneficial association, as well as about spreading of the legumes across the world. Discovery of genes involved in recognition of partners, transduction of symbiotic signals and overall “management” of symbiosis will also provide a useful tool for agriculture, as the knowledge obtained

from this studying will facilitate the creation of highly-effective specific symbiotic pairs between crop plants and nitrogen-fixing bacteria in field..

## Acknowledgements

A.S. Sulima was financially supported by the grant of Russian Foundation for Basic Research (RFBR) # 14-04-32289\_mol-a. V.A. Zhukov, O.Y. Shtark, A.Y. Borisov, and I.A. Tikhonovich were financially supported by the grant of Russian Science Foundation (RSF) # 14-24-00135. The authors thank Dr. M.N. Povydysh (Saint-Petersburg State Chemical Pharmaceutical Academy, St.Petersburg, Russia) for help in preparation of figures.

## Author details

Sulima Anton Sergeevich, Zhukov Vladimir Alexandrovich\*, Shtark Oksana Yurievna, Borisov Alexey Yurievich and Tikhonovich Igor Anatolievich

\*Address all correspondence to: zhukoff01@yahoo.com

All-Russia Research Institute for Agricultural Microbiology (ARRIAM), St.-Petersburg, Russia

## References

- [1] Mulligan RM, Chory J, Ecker JR. Signaling in plants. *Proc Natl Acad Sci USA*. 1997;94: 2793-2795.
- [2] Nakagawa T, Kaku H, Shimoda Y, Sugiyama A, Shimamura M, Takanashi K, Yazaki K, Aoki T, Shibuya N, Kouchi H. From defense to symbiosis: limited alterations in the kinase domain of LysM receptor-like kinases are crucial for evolution of legume–*Rhizobium* symbiosis. *Plant J* 2011;65: 169-180.
- [3] Hua J. Modulation of plant immunity by light, circadian rhythm, and temperature. *Curr Opin Plant Biol*. 2013;16: 406–413.
- [4] Vance CP. Symbiotic nitrogen fixation and phosphorus acquisition. Plant nutrition in the world of declining renewable resources. *Plant Physiol*. 2001;127(2): 390-397.
- [5] Brewin NJ. Plant cell wall remodeling in the rhizobium-legume symbiosis. *Crit Rev Plant Sci*. 2004;23: 1-24.
- [6] Allen ON, Allen EK. *The Leguminosae. A Source Book of Characteristics, Uses and Nodulation*. Madison: The University of Wisconsin Press; 1981. 806 pp.

- [7] Wall LG. The actinorhizal symbiosis. *J Plant Growth Regul.* 2000;19(2): 167-182.
- [8] Sawada H, Kuykendall LD, Young JM. Changing concepts in the systematics of bacterial nitrogen-fixing legume symbionts. *J Gen Appl Microbiol.* 2003;49(3): 155-179.
- [9] Balachandar D, Raja P, Kumar K, Sundaram SP. Non-Rhizobial nodulation in legumes. *Biotechnol Molec Biol Rev.* 2007;2(2): 49-57.
- [10] Spaink HP, Kondorosi A, Hooykaas PJJ. (eds.) *The Rhizobiaceae. Molecular Biology of Model Plant-Associated Bacteria.* Dordrecht/Boston/London: Kluwer; 1998. 566 pp.
- [11] MacLean AM, Finan T, Sadowsky MJ. Genomes of the symbiotic nitrogen-fixing bacteria of legumes. *Plant Physiol.* 2007;144(2): 615-622.
- [12] Geurts R, Bisseling T. *Rhizobium* nod factor perception and signaling. *Plant Cell.* 2002;14.
- [13] OvtSYna AO, Staehelin C. Bacterial signals required for the *Rhizobium*-legume symbiosis. In: Pandalai SG. (ed.), *Recent Research Developments in Microbiology, Vol 7 (Part II).* Trivandrum, India: Research Signpost; 2003. p 631-648.
- [14] Spaink HP. The molecular basis of infection and nodulation by rhizobia: the Ins and Outs of sympathogenesis. *Ann Rev Phytopathol.* 1995;33: 345-368.
- [15] Schultze M, Kondorosi A. Regulation of symbiotic root nodule development. *Annu Rev Genet* 1998;32: 33-57.
- [16] D'Haeye W, Holsters M. Nod factor structures, responses, and perception during initiation of Nodule development. *Glycobiology* 2002;12(6): 79-105.
- [17] Perret X, Staehelin C, Broughton WJ. Molecular basis of symbiotic promiscuity. *Microbiol Mol Biol. Rev* 2000;64(1): 180-201.
- [18] Oke V, Long SR. Bacteroid formation in the rhizobium-legume symbiosis. *Curr Opin Microbiol* 1999;2(6): 641-646.
- [19] Brewin NJ (1998) Tissue and cell invasion by rhizobium: the structure and development of infection threads and symbiosomes. In: Spaink HP, Kondorosi A, Hooykaas PJJ. (eds.), *The Rhizobiaceae. Molecular Biology of Model Plant-Associated Bacteria.* Dordrecht/Boston/London: Kluwer; 1998. p 417-429.
- [20] Mylona P, Pawlowski K, Bisseling T. Symbiotic Nitrogen Fixation. *Plant Cell.* 1995;7(7): 869-885.
- [21] Brewin NJ. Development of the legume root nodule. *Annu Rev Cell Biol.* 1991;7: 191-226.
- [22] Mergaert P, Uchiumi T, Alunni B, Evanno G, Cheron A, Catrice O, Mausset AE, Barloy-Hubler F, Galibert F, Kondorosi A, Kondorosi E. Eukaryotic Control on Bacterial Cell Cycle and Differentiation in the *Rhizobium*-Legume Symbiosis. *Proc Natl Acad Sci USA.* 2006;103(13): 5230-5235.



- [23] Hirsch AM, LaRue TA. Is the legume nodule a modified root or stem or an organ *sui generis*? *Crit Rev Plant Sci*. 1997;16: 361-392.
- [24] Sprent JI, Raven JA. Evolution of nitrogen-fixing symbiosis. In: Stacey G, Burris RH, Evans HJ. (eds.) *Biological Nitrogen Fixation*. New York, London: Chapman & Hall; 1992. p 461-496.
- [25] Fred EB, Baldwin IL, McCoy E. *Root Nodule Bacteria and Leguminous Plants*. Madison: Univ Wisconsin Stud Sci; 1932. 343 pp.
- [26] Provorov NA. The interdependence between taxonomy of legumes and specificity of their interaction with rhizobia in relation to evolution of the symbiosis. *Symbiosis*. 1994;17: 183-200.
- [27] Broughton WJ, Perret X. Genealogy of legume-rhizobium symbiosis. *Curr Opin Plant Biol*. 1999;2(4): 305-311.
- [28] Lafay B, Bullier E, Burdon JJ. Bradyrhizobia isolated from root nodules of *Parasponia* (Ulmaceae) do not constitute a separate coherent lineage. *Int J Syst Evol Microbiol*. 2006;56(Pt 5): 1013-1018.
- [29] Price NP, Relić B, Talmont F, Lewin A, Promé D, Pueppke SG, Maillet F, Dénarié J, Promé JC, Broughton WJ. Broad-host-range rhizobium species strain NGR234 secretes a family of carbamoylated, and fucosylated, nodulation signals that are O-acetylated or sulphated. *Mol Microbiol*. 1992;6(23): 3575-3584.
- [30] Schmeisser C, Liesegang H, Krysciak D, Bakkou N, Le Quéré A, Wollherr A, Heine-meyer I, Morgenstern B, Pommerening-Röser A, Flores M, Palacios R, Brenner S, Gottschalk G, Schmitz RA, Broughton WJ, Perret X, Strittmatter AW, Streit WR. Rhizobium sp. strain NGR234 possesses a remarkable number of secretion systems. *Appl Environ Microbiol*. 2009;75(12): 4035-4045.
- [31] Broughton WJ, Jabbouri S, Perret X. Keys to symbiotic harmony. *J Bacteriol*. 2000;182: 5641-5652.
- [32] Relić B, Staehelin C, Fellay R, Jabbouri S, Boller T, Broughton WJ. Do Nod-factor levels play a role in host specificity? In: Kiss GB, Endre, G (eds.), *Proceedings of the First European Congress on Nitrogen Fixation*. Officina Press Szed, Szeged, Hungary; 1994. p 69-75.
- [33] Becker A, Pühler A. Production of exopolysaccharides. In: Spaink HP, Kondorosi A, Hooykaas PJJ. (eds.), *The Rhizobiaceae. Molecular Biology of Model Plant-Associated Bacteria*. Dordrecht/Boston/London: Kluwer; 1998. p 87-118.
- [34] Lugtenberg BJJ. Outer membrane proteins. In: Spaink HP, Kondorosi A, Hooykaas PJJ. (eds.), *The Rhizobiaceae. Molecular Biology of Model Plant-Associated Bacteria*. Dordrecht/Boston/London: Kluwer; 1998. p 45-53.
- [35] Jones KM, Sharopova N, Lohar DP, Zhang JQ, VandenBosch KA, Walker GC. Differential response of the plant *Medicago truncatula* to its symbiont *Sinorhizobium meliloti*

- or an exopolysaccharide-deficient mutant. *Proc Natl Acad Sci USA*. 2008;105(2): 704-709.
- [36] Downie JA, Rossen L, Knight CD, Robertson JG, Wells B, Johnston AW. *Rhizobium leguminosarum* genes involved in early stages of nodulation. *J Cell Sci Suppl*. 1985;2: 347-354.
- [37] Spaink HP, Sheeley DM, van Brussel AA, Glushka J, York WS, Tak T, Geiger O, Kennedy EP, Reinhold VN, Lugtenberg BJ. A novel highly unsaturated fatty acid moiety of lipo-oligosaccharide signals determines host specificity of *Rhizobium*. *Nature*. 1991;354(6349): 125-130.
- [38] Demont N, Debellé F, Aurelle H, Dénarié J, Promé JC. Role of the *Rhizobium meliloti* nodF and nodE genes in the biosynthesis of lipo-oligosaccharidic nodulation factors. *J Biol Chem*. 1993;268(27): 20134-20142.
- [39] Spaink HP, Weinman J, Djordjevic MA, Wijffelman CA, Okker RJ, Lugtenberg BJ. Genetic analysis and cellular localization of the *Rhizobium* host specificity-determining NodE protein. *EMBO J*. 1989;8(10): 2811-2818.
- [40] Ardourel M, Demont N, Debellé F, Maillet F, de Billy F, Prome J-C, Denarie J. *Rhizobium meliloti* lipooligosaccharide nodulation factors. *Plant Cell*. 1994;6: 1357-1374.
- [41] Poupot R, Martinez-Romero E, Gautier N, Promé J-C. Wild type *Rhizobium etli*, a bean symbiont, produces acetyl-fucosylated, N-methylated, and carbamoylated nodulation factors. *J Biol Chem*. 1995;270: 6050-6055.
- [42] Yang G-P, Debellé F, Savagnac A, Ferro M, Schiltz O, Maillet F, Promé D, Treillhou M, Vialas C, Lindström K et al. Structure of the *Mesorhizobium huakuii* and *Rhizobium galegae* Nod factors: a cluster of phylogenetically related legumes are nodulated by rhizobia producing Nod factors with  $\alpha$ ,  $\beta$ -unsaturated N-acyl substitutions. *Mol Microbiol*. 1999;34: 227-237.
- [43] Radutoiu S, Madsen LH, Madsen EB, Felle HH, Umehara Y, Grönlund M, Sato S, Nakamura Y, Tabata S, Sandal N, Stougaard J. Plant recognition of symbiotic bacteria requires two LysM receptor-like kinases. *Nature*. 2003;425(6958): 585-592.
- [44] Madsen EB, Madsen LH, Radutoiu S, Olbryt M, Rakwalska M, Szczyglowski K, Sato S, Kaneko T, Tabata S, Sandal N, Stougaard J. A receptor kinase gene of the LysM type is involved in legume perception of rhizobial signals. *Nature*. 2003;425(6958): 637-640.
- [45] Ohnuma T, Onaga S, Murata K, Taira T, Katoh E. LysM domains from *Pteris ryukyuensis* chitinase-A: a stability study and characterization of the chitin-binding site. *J Biol Chem*. 2008;283: 5178-5187.
- [46] Bateman A, Bycroft M. The structure of a LysM domain from *E. coli* membrane-bound lytic murein transglycosylase D (MltD). *J Mol Biol*. 2000;299: 1113-1119.

- [47] Buist G, Steen A, Kok J, Kuipers OP. LysM, a widely distributed protein motif for binding to (peptido)glycans. *Mol Microbiol.* 2008;68: 838-847.
- [48] Visweswaran GRR, van Roosmalen KLM, Kok J, Buist G. Exploiting the peptidoglycan-binding motif, LysM, for medical and industrial applications. *Appl Microbiol Biotechnol.* 2014;98: 4331-4345.
- [49] Bensmihen S, de Billy F, Gough C. Contribution of NFP LysM domains to the recognition of Nod factors during the *Medicago truncatula/Sinorhizobium meliloti* symbiosis. *PLoS One.* 2011;6(11): e26114.
- [50] Zhu H, Riely BK, Burns NJ, Ane JM. Tracing nonlegume orthologs of legume genes required for nodulation and arbuscular mycorrhizal symbioses. *Genetics.* 2006;172: 2491-2499.
- [51] Zhang XC, Wu X, Findley S, Wan J, Libault M, Nguyen HT, Cannon SB, Stacey G. Molecular evolution of lysin motif-type receptor-like kinases in plants. *Plant Physiol.* 2007;144: 623-636.
- [52] Broghammer A, Krusell L, Blaise M, Sauer J, Sullivan JT, Maolanon N, Vinther M, Lorentzen A, Madsen EB, Jensen KJ, Roepstorff P, Thirup S, Ronson CW, Thygesen MB, Stougaard J. Legume receptors perceive the rhizobial lipochitin oligosaccharide signal molecules by direct binding. *PNAS.* 2012;109(34): 13859-13864.
- [53] Limpens E, Franken C, Smit P, Willemse J, Bisseling T, Geurts R. LysM domain receptor kinases regulating rhizobial Nod factor-induced infection. *Science.* 2003;24: 630-633.
- [54] Arrighi JF, Barre A, Ben Amor B, Bersoult A, Soriano LC, Mirabella R, de Carvalho-Niebel F, Journet EP, Ghérardi M, Huguet T, Geurts R, Dénarié J, Rougé P, Gough C. The *Medicago truncatula* lysin [corrected] motif-receptor-like kinase gene family includes NFP and new nodule-expressed genes. *Plant Physiol.* 2006;142(1): 265-279.
- [55] Zhukov V, Radutoiu S, Madsen LH, Rychagova T, Ovchinnikova E, Borisov A, Tikhonovich I, Stougaard J. The pea Sym37 receptor kinase gene controls infection-thread initiation and nodule development. *Mol Plant Microbe Interact.* 2008;21(12): 1600-1608.
- [56] Smit P, Limpens E, Geurts R, Fedorova E, Dolgikh E, Gough C, Bisseling T. *Medicago* LYK3, an entry receptor in rhizobial nodulation factor signaling. *Plant Physiol.* 2007 Sep;145(1): 183-191.
- [57] Zhukov VA, Shtark OY, Borisov AY, Tikhonovich IA. Breeding to improve symbiotic effectiveness of legumes. In: Andersen SB (ed.), *Plant Breeding from Laboratories to Field*. Croatia: InTech; 2013. p 167-207.
- [58] Handberg K, Stougaard J. *Lotus japonicus*, an autogamous, diploid legume species for classical and molecular genetics. *Plant J.* 1992;2: 487-496.

- [59] Barker D, Bianchi S, Blondon F, Dattee Y, Duc G, Essad S, Flament P, Gallusci P, Genier G, Guy P, Muel X, Tourneur J, Denarie J, Huguet T. *Medicago truncatula*, a model plant for studying the molecular genetics of the rhizobium-legume symbiosis. *Plant Mol Biol Rep.* 1990;8: 40-49.
- [60] Cook DR. *Medicago truncatula* – a model in the making. *Curr Opin Plant Biol.* 1999;2(4): 301-304.
- [61] Young ND, Mudge J, Ellis THN. Legume genomes: more than peas in a pod. *Curr Opin Plant Biol.* 2003;6(2): 199-204.
- [62] Cook DR, Vandenbosch K, de Bruijn FJ, Huguet T. Model legumes get the Nod. *Plant Cell.* 1997;9: 275-281.
- [63] Udvardi MK. Legume models strut their stuff. *Mol Plant Microbe Interact.* 2001;14(1): 6-9.
- [64] Stougaard J. Genetics and genomics of root symbiosis. *Curr Opin Plant Biol.* 2001;4(4): 328-335.
- [65] Penmetsa RV, Cook DR. A legume ethylene-insensitive mutant hyperinfected by its rhizobial symbiont. *Science.* 1997;275(5299): 527-530.
- [66] Schauser L, Handberg K, Sandal N, Stiller J, Thykjaer T, Pajuelo E, Nielsen A, Stougaard J. Symbiotic mutants deficient in nodule establishment identified after T-DNA transformation of *Lotus japonicus*. *Mol Gen Genet.* 1998;259(4): 414-423.
- [67] Schüßler A, Schwarzott D, Walker C. A New fungal phylum, the Glomeromycota: phylogeny and evolution. *Mycol Res.* 2001;105: 1413-1297.
- [68] Brundrett MC. Coevolution of roots and mycorrhizas of land plants. *New Phytol.* 2002;154: 275-304.
- [69] Smith SE, Read DJ. *Mycorrhizal Symbiosis* (3rd edn.). London: Academic Press; 2008. 800 pp.
- [70] Bidartondo MI. The evolutionary ecology of myco-heterotrophy. *New Phytol.* 2005;167: 335–352.
- [71] Maillet F, Poinso V, André O, Puech-Pagès V, Haouy A, Gueunier M, Cromer L, Giraudet D, Formey D, Niebel A, Martinez EA, Driguez H, Bécard G, Dénarié J. Fungal lipochitoooligosaccharide symbiotic signals in arbuscular mycorrhiza. *Nature.* 2011;469(7328): 58-63.
- [72] Endre G, Kereszt A, Kevei Z, Mihacea S, Kaló P, Kiss GB. A receptor kinase gene regulating symbiotic nodule development. *Nature.* 2002;417(6892): 962-966.
- [73] Stracke S, Kistner C, Yoshida S, Mulder L, Sato S, Kaneko T, Tabata S, Sandal N, Stougaard J, Szczyglowski K, Parniske M. A plant receptor-like kinase required for both bacterial and fungal symbiosis. *Nature.* 2002;417(6892): 959-962.

- [74] Limpens E, Mirabella R, Fedorova E, Franken C, Franssen H, Bisseling T, Geurts R. Formation of organelle-like N<sub>2</sub>-fixing symbiosomes in legume root nodules is controlled by DMI2. *Proc Natl Acad Sci USA*. 2005; 102(29): 10375-10380.
- [75] Kevei Z, Lougnon G, Mergaert P, Horvath GV, Kereszt A, Jayaraman D, Zaman N, Marcel F, Regulski K, Kiss GB, Kondorosi A, Endre G, Kondorosi, E-Ané JM. 3-Hydroxy-3-methylglutaryl coenzyme A reductase 1 interacts with NORK and is crucial for nodulation in *Medicago truncatula*. *Plant Cell*. 2007;19(12): 3974-3989.
- [76] Zhu H, Chen T, Zhu M, Fang Q, Kang H, Hong Z, Zhang Z. A Novel ARID DNA-binding protein interacts with SymRK and is expressed during early nodule development in *Lotus japonicus*. *Plant Physiol*. 2008;148(1): 337-347.
- [77] Lefebvre B, Timmers T, Mbengue M, Moreau S, Herve C, Tóth K, Bittencourt-Silvestre J, Klaus D, Deslandes L, Godiard L, Murray JD, Udvardi MK, Raffaele S, Mongrand S, Cullimore J, Gamas P, Niebel A, Ott T. A remorin protein interacts with symbiotic receptors and regulates bacterial infection. *Proc Natl Acad Sci USA*. 2010;107(5): 2343-2348.
- [78] Yuan S, Zhu H, Gou H, Fu W, Liu L, Chen T, Ke D, Kang H, Xie Q, Hong Z, Zhang Z. A ubiquitin ligase of symbiosis receptor kinase involved in nodule organogenesis. *Plant Physiol*. 2012; DOI:10.1104/pp.112.199000.
- [79] Wais RJ, Galera C, Oldroyd G, Catoira R, Penmetsa RV, Cook D, Gough C, Dénarié J, Long SR. Genetic analysis of calcium spiking responses in nodulation mutants of *Medicago truncatula*. *Proc Natl Acad Sci USA*. 2000;97(24): 13407-13412.
- [80] Ané JM, Kiss GB, Riely BK, Penmetsa RV, Oldroyd GE, Ajax C, Lévy J, Debelle F, Baek JM, Kalo P, Rosenberg C, Roe BA, Long SR, Dénarié J, Cook DR. *Medicago truncatula* DMI1 required for bacterial and fungal symbioses in legumes. *Science*. 2004;303(5662): 1364-1367.
- [81] Imaizumi-Anraku H, Takeda N, Charpentier M, Perry J, Miwa H, Umehara Y, Kouchi H, Murakami Y, Mulder L, Vickers K, Pike J, Downie JA, Wang T, Sato S, Asamizu E, Tabata S, Yoshikawa M, Murooka Y, Wu GJ, Kawaguchi M, Kawasaki S, Parniske M, Hayashi M. Plastid proteins crucial for symbiotic fungal and bacterial entry into plant roots. *Nature*. 2005;433(7025): 527-531.
- [82] Edwards A, Heckmann AB, Yousafzai F, Duc G, Downie JA. Structural implications of mutations in the pea SYM8 symbiosis gene, the DMI1 ortholog, encoding a predicted ion channel. *Mol Plant Microbe Interact*. 2007;20(10): 1183-1191.
- [83] Riely BK, Lougnon G, Ané JM, Cook DR. The symbiotic ion channel homolog DMI1 is localized in the nuclear membrane of *Medicago truncatula* roots. *Plant J*. 2007;49(2): 208-216.
- [84] Peiter E, Sun J, Heckmann AB, Venkateshwaran M, Riely BK, Otegui MS, Edwards A, Freshour G, Hahn MG, Cook DR, Sanders D, Oldroyd GE, Downie JA, Ané JM.



- The *Medicago truncatula* DMI1 protein modulates cytosolic calcium signaling. *Plant Physiol.* 2007;145(1): 192-203.
- [85] Kanamori N, Madsen LH, Radutoiu S, Frantescu M, Quistgaard EMH, Miwa H, Downie JA, James EK, Felle HH, Haaning LL, Jensen TH, Sato S, Nakamura Y, Tabata S, Sandal N, Stougaard J. A nucleoporin is required for induction of Ca<sup>2+</sup> spiking in legume nodule development and essential for rhizobial and fungal symbiosis. *Proc Natl Acad Sci USA.* 2006;103(2): 359-364.
- [86] Saito K, Yoshikawa M, Yano K, Miwa H, Uchida H, Asamizu E, Sato S, Tabata S, Imaizumi-Anraku H, Umehara Y, Kouchi H, Murooka Y, Szczyglowski K, Downie JA, Parniske M, Hayashi M, Kawaguchi M. NUCLEOPORIN85 is required for calcium spiking, fungal and bacterial symbioses, and seed production in *Lotus japonicus*. *Plant Cell.* 2007;19(2): 610-624.
- [87] Groth M, Takeda N, Perry J, Uchida H, Draxl S, Brachmann A, Sato S, Tabata S, Kawaguchi M, Wang TL, Parniske M. NENA, a *Lotus japonicus* homolog of Sec13, is required for rhizodermal infection by Arbuscular Mycorrhiza fungi and rhizobia but dispensable for cortical endosymbiotic development. *Plant Cell.* 2010;22(7): 2509-2526.
- [88] Gleason C, Chaudhuri S, Yang T, Munoz A, Poovaiah BW, Oldroyd GE. Nodulation independent of rhizobia induced by a calcium-activated kinase lacking autoinhibition. *Nature.* 2006;441(7097): 1149-1152.
- [89] Parniske M. Arbuscular Mycorrhiza: the mother of plant root endosymbioses. *Nat Rev Microbiol.* 2008;6(10): 763-775.
- [90] Yano K, Yoshida S, Muller J, Singh S, Banba M, Vicker K. CYCLOPS, a mediator of symbiotic intracellular accommodation. *Proc Natl Acad Sci USA.* 2008;105(51): 20540-20545.
- [91] Kaló P, Gleason C, Edwards A, Marsh J, Mitra RM, Hirsch S, Jakab J, Sims S, Long SR, Rogers J, Kiss GB, Downie JA, Oldroyd GE. Nodulation signaling in legumes requires NSP2, a member of the GRAS family of transcriptional regulators. *Science.* 2005;308(5729): 1786-1789.
- [92] Smit P, Raedts J, Portyanko V, Debellé F, Gough C, Bisseling T, Geurts R. NSP1 of the GRAS protein family is essential for rhizobial Nod factor-induced transcription. *Science.* 2005;308(5729): 1789-1791.
- [93] Albrecht C, Geurts R, Bisseling T. Legume nodulation and Mycorrhizae formation; two extremes in host specificity meet. *EMBO J.* 1999;18(2): 281-288.
- [94] Heckmann AB, Lombardo F, Miwa H, Perry JA, Bunnewell S, Parniske M, Wang TL, Downie JA. *Lotus japonicus* nodulation requires two GRAS domain regulators, one of which is functionally conserved in a non-legume. *Plant Physiol.* 2006;142(4): 1739-1750.
- [95] Murakami Y, Miwa H, Imaizumi-Anraku H, Kouchi H, Downie JA, Kawasaki KMS. Positional cloning identifies *Lotus japonicus* NSP2, a putative transcription factor of

- the GRAS family, required for NIN and ENOD40 gene expression in nodule initiation. *DNA Res.* 2006;13: 255-265.
- [96] Gonzalez-Rizzo S, Crespi M, Frugier F. The *Medicago truncatula* CRE1 cytokinin receptor regulates lateral root development and early symbiotic interaction with *Sinorhizobium meliloti*. *Plant Cell.* 2006;18(10): 2680-2693.
- [97] Murray JD, Karas BJ, Sato S, Tabata S, Amyot L, Szczyglowski K. A cytokinin perception mutant colonized by rhizobium in the absence of nodule organogenesis. *Science.* 2007;315(5808): 101-104.
- [98] Tirichine L, Sandal N, Madsen LH, Radutoiu S, Albrektsen AS, Sato S, Asamizu E, Tabata S, Stougaard J. A gain-of-function mutation in a cytokinin receptor triggers spontaneous root nodule organogenesis. *Science.* 2007;315(5808): 104-107.
- [99] Schauser L, Roussis A, Stiller J, Stougaard J. A Plant regulator controlling development of symbiotic root nodules. *Nature.* 1999;402(6758): 191-195.
- [100] Borisov AY, Madsen LH, Tsyganov VE, Umehara Y, Voroshilova VA, Batagov AO, Sandal N, Mortensen A, Schauser L, Ellis N, Tikhonovich IA, Stougaard J. The *Sym35* gene required for root nodule development in pea is an ortholog of Nin from *Lotus japonicus*. *Plant Physiol.* 2003;131(3): 1009-1017.
- [101] Marsh JF, Rakocevic A, Mitra RM, Brocard L, Sun J, Eschstruth A, Long SR, Schultze M, Ratet P, Oldroyd GE. *Medicago truncatula* NIN is essential for rhizobial-independent nodule organogenesis induced by Autoactive Calcium/Calmodulin-Dependent Protein Kinase. *Plant Physiol.* 2007;144(1): 324-335.
- [102] Middleton PH, Jakab J, Penmetsa RV, Starker CG, Doll J, Kalo P, Prabhu R, Marsh JF, Mitra RM, Kereszt A, Dudas B, VandenBosch K, Long SR, Cook DR, Kiss GB, Oldroyd GE. An ERF Transcription factor in *Medicago truncatula* that is essential for Nod factor signal transduction. *Plant Cell.* 2007;19(4): 1221-1234.
- [103] Caetano-Anolles G, Gresshoff PM. Plant genetic control of nodulation. *Annu Rev Microbiol.* 1991;45: 345-382.
- [104] Ferguson BJ, Indrasumunar A, Hayashi S, Lin MH, Lin YH, Reid DE, Gresshoff PM. Molecular analysis of legume nodule development and autoregulation. *J Integr Plant Biol.* 2010;52(1): 61-76.
- [105] Okamoto S, Ohnishi E, Sato S, Takahashi H, Nakazono M, Tabata S, Kawaguchi M. Nod factor/nitrate-induced CLE genes that drive HAR1-mediated systemic regulation of nodulation. *Plant Cell Physiol.* 2009;50(1): 67-77.
- [106] Mortier V, Den Herder G, Whitford R, Van de Velde W, Rombauts S, D'Haeseleer K, Holsters M, Goormachtig S. CLE peptides control *Medicago truncatula* nodulation locally and systemically. *Plant Physiol.* 2010;153(1): 222-237.
- [107] Krusell L, Madsen LH, Sato S, Aubert G, Genua A, Szczyglowski K, Duc G, Kaneko T, Tabata S, de Bruijn F, Pajuelo E, Sandal N, Stougaard J. Shoot control of root devel-

- opment and nodulation is mediated by a receptor-like kinase. *Nature*. 2002;420(6914): 422-426.
- [108] Nishimura R, Hayashi M, Wu GJ, Kouchi H, Imaizumi-Anraku H, Murakami Y, Kawasaki S, Akao S, Ohmori M, Nagasawa M, Harada K, Kawaguchi M. HAR1 Mediates systemic regulation of symbiotic organ development. *Nature*. 2002;420(6914): 426-429.
- [109] Schnabel E, Journet EP, de Carvalho-Niebel F, Duc G, Frugoli J. The *Medicago truncatula* SUNN Gene Encodes a CLV1-like leucine-rich repeat receptor kinase that regulates nodule number and root length. *Plant Mol Biol*. 2005;58(6):809-822.
- [110] Staehelin C, Xie ZP, Illana A, Vierheilig H. Long-distance transport of signals during symbiosis: are nodule formation and mycorrhization autoregulated in a similar way? *Plant Signal Behav*. 2011;6(3): 372-377.
- [111] Limpens E, Moling S, Hooiveld G, Pereira PA, Bisseling T, Becker JD, Küster H. Cell- and tissue-specific transcriptome analyses of *Medicago truncatula* root nodules. *PLoS One*. 2013 May 29;8(5): e64377.
- [112] Moling S, Pietraszewska-Bogiel A, Postma M, Fedorova E, Hink MA, Limpens E, Gaddella TW, Bisseling T. Nod factor receptors form heteromeric complexes and are essential for intracellular infection in *Medicago* nodules. *Plant Cell*. 2014;26(10): 4188-4199.
- [113] Bourion V, Rizvi SM, Fournier S, de Larambergue H, Galmiche F, Marget P, Duc G, Burstin J. Genetic dissection of nitrogen nutrition in pea through a QTL approach of root, nodule, and shoot variability. *Theor Appl Genet*. 2010;121(1): 71-86.
- [114] Razumovskaya ZG. Nodule formation in various pea cultivars. *Russian J Microbiol*. 1937;6: 321-328.
- [115] Lie TA. Host genes of *Pisum sativum* L. conferring resistance to European *Rhizobium leguminosarum* strains. *Plant Soil*. 1984;82: 462-465.
- [116] Davis EO, Evans IJ, Johnston AWB. Identification of *nodX*, a gene that allows *Rhizobium leguminosarum* bv. *viciae* strain TOM to nodulate Afghanistan peas. *Mol Gen Genet*. 1988: 531-535.
- [117] Ovtsyna AO, Rademaker GJ, Esser E, Weinman J, Rolfe BG, Tikhonovich IA, Lugtenberg BJ, Thomas-Oates JE, Spaink HP. Comparison of characteristics of the *nodX* genes from various *Rhizobium leguminosarum* strains. *Mol Plant Microbe Interact*. 1999;12(3): 252-258.
- [118] Ovtsyna AO, Schultze M, Tikhonovich IA, Spaink HP, Kondorosi E, Kondorosi A, Staehelin C. Nod factors of *Rhizobium leguminosarum* bv. *viciae* and their fucosylated derivatives stimulate a nod factor cleaving activity in pea roots and are hydrolyzed in vitro by plant chitinases at different rates. *Mol Plant Microbe Interact*. 2000;13(8): 799-807.

- [119] Kozik A, Matvienko M, Scheres B, Paruvangada VG, Bisseling T, van Kammen A, Ellis TH, LaRue T, Weeden N. The pea early gene PsENOD7 maps in the region of linkage group I containing sym2 and leghemoglobin. *Plant Mol Biol.* 1996;31(1): 149-156.
- [120] Zhukov VA, Sulima AS, Porozov YB, Borisov AY, Tikhonovich IA. Polymorphism in gene sequence of LysM receptor kinase is associated with *Sym2*-controlled nodulation in pea (*Pisum sativum* L.). Proceedings of 18th International Conference on Nitrogen Fixation (14–18 October 2013, Myazaki, Japan): 76.
- [121] Li R, Knox MR, Edwards A, Hogg B, Ellis TH, Wei G, Downie JA. Natural variation in host-specific nodulation of pea is associated with a haplotype of the SYM37 LysM-type receptor-like kinase. *Mol Plant Microbe Interact.* 2011;24(11): 1396-1403.
- [122] Tikhonovich IA, Andronov EE, Borisov AY, Dolgikh EA, Zhernakov AI, Zhukov VA, Provorov NA, Roumiantseva ML, Simarov B.V. The Principle of Genome Complementarity in the Enhancement of Plant Adaptive Capacities. *Russian J Genet.* 2015;51(9): 831–846.
- [123] Schüßler A. Molecular phylogeny, taxonomy and evolution of *Geosiphon pyriformis* and arbuscular mycorrhizal fungi. *Plant Soil.* 2002;244: 75-83.
- [124] Schüßler A, Wolf E. *Geosiphon pyriformis* – a glomeromycotan soil fungus forming endosymbiosis with cyanobacteria. In: Declerck S, Strullu DG, Fortin JA. (eds.), *In Vitro Culture of Mycorrhizas*. Berlin–Heidelberg–New York: Springer; 2005. 271-290.
- [125] Kluge M, Mollenhauer D, Wolf E, Schüßler A. The Nostoc-Geosiphon endocytobiosis. In: Rai AN, Bergman B, Rasmussen U. (eds.), *Cyanobacteria in Symbiosis*. Kluwer Academic Publishers; 2003. 19-30.
- [126] Minerdi D, Bianciotto V, Bonfante P. Endosymbiotic bacteria in mycorrhizal fungi: from their morphology to genomic sequences. *Plant Soil.* 2002;244: 211-219.