

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

**4,800**

Open access books available

**122,000**

International authors and editors

**135M**

Downloads

Our authors are among the

**154**

Countries delivered to

**TOP 1%**

most cited scientists

**12.2%**

Contributors from top 500 universities



**WEB OF SCIENCE™**

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Evaluation of Pre- and Postemergence Herbicide Combinations for Broadleaved Weeds in Sugar Beet

---

Irena Deveikyte, Lina Sarunaite and Vytautas Seibutis

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/61437>

---

## Abstract

Lithuania sugar beet growers have few herbicide options available for weed management. Six field trials were conducted at the Institute of Agriculture, Lithuania, in order to evaluate the effects of chemical weed management in sugar beet. Treatments included untreated and hand-weeded control and several rates of phenmedipham plus desmedipham plus ethofumesate, phenmedipham, ethofumesate, triflusaluron, chloridazon, and metamilron. Pre- and postemergence and only postemergence applications similarly affected weed control. Phenmedipham plus desmedipham plus ethofumesate was more effective for controlling weeds when applied in combination with metamilron, triflusaluron, and chloridazon. The significantly lowest efficacy for weed control was phenmedipham combined with ethofumesate and metamilron as compared to the phenmedipham plus desmedipham plus ethofumesate. Reducing the doses of phenmedipham plus desmedipham plus ethofumesate from 114+89+140 g a.i. ha<sup>-1</sup> to 91+71+112 g a.i. ha<sup>-1</sup> and 68+53+84 g a.i. ha<sup>-1</sup> in mixture with triflusaluron resulted in the increase of weed biomass. Full (45 g a.i. ha<sup>-1</sup>) and reduced doses (30 g a.i. ha<sup>-1</sup>) of triflusaluron with phenmedipham plus desmedipham plus ethofumesate similarly affected weeds. The herbicides investigated did not have any negative influence on sugar beet productivity and quality.

**Keywords:** Weeds, herbicides combination, sugar beet

---

## 1. Introduction

Weed competition is one of the major factors which limit sugar beet production in the world [1]. Weed-crop interactions are based on competition for water, nutrients, and light and allelopathic effects may also play a small role. In sugar beet weed interference, all these factors are important too, but light is of prime importance. Weeds may also interfere with harvest operations, making the process less efficient [2]. Due to the fact that a lot of weeds can grow

above the sugar beet canopy and reduce the amount of photosynthetic radiation reaching the crop, these weeds are stronger competitors compared to smaller weeds [3, 4]. In a weed free crop stand, photosynthesis in sugar beet is more efficient and nutrient accumulation in the sugar beet root is higher [5]. Left uncontrolled, weeds may reduce yield, interfere with harvest, reduce the value of the crop, and increase future weed problems. The yield of sugar beet roots and sucrose can be severely decreased by weeds, the extent of the decrease being dependent upon competitive ability, weed density, and the length of time that weeds compete with the crop. The total potential losses from weeds would be between 26 and 100% of the potential crop yield [6-8].

Sugar beet is very sensitive to weed competition from the early stages of growth [9, 10]. Sugar beet is not competitive with emerging weeds until it has at least 8 true leaves [7]. Therefore, effective control of weeds at early stages seems to be more important than that at later developed stages [10]. The length of weed-free period affected yield of sugar beet very markedly [11]. When sugar beet and weeds grow together 30 days after emergence of sugar beet, the root yield is decreased up to 45% [12]. As control of weeds is delayed, the yield lost may be decreased by 1.5% for each day the crop is left unweeded, although sugar beet has some ability to recover from an early check [13]. Understanding the emergence characteristics of weeds can be helpful in determining the optimum time to apply postemergence herbicide [11].

Weed control in sugar beet is accomplished with herbicides, mechanical tillage, cultural practices, and hand labor. Control of weeds with herbicides is generally more profitable than allowing weeds to compete with the crop. Herbicides play an important role for weed control in sugar beet production [14, 15]. For high efficacy of chemical method, the timing of application is very important. Weeds have to be small (cotyledon stage) to ensure successful weed control [16]. The doses of herbicides could be reduced by applying at the early growth stage of the weeds, when the first seed leaves start to appear [14, 15]. The application of lower doses leads to reduction of negative impact of herbicides on environment and cuts expenditures for beet production [17].

In recent years, the use of preplant-applied herbicides has declined and use of postemergence herbicides has increased. The most popular active ingredients are phenmedipham, desmedipham, ethofumesate, metamitron, triflurosulfuron-methyl, lenacil, clopyralid, and chloridazon [7, 18]. The range of weed species controlled by each herbicide is also limited and so mixtures of herbicides are applied [7, 15, 19, 20]. Sugar beet is applied by tank-mix herbicides combinations several times after crop emergence [15, 21, 22]. Mixtures of postemergence, broad-spectrum herbicides have to be applied to control the wide range of weed species in sugar beet crops [23, 24].

Field experiments were carried out in 2004–2005 and 2010–2012 on arable fields located at the Institute of Agriculture in Central Lithuania. The objective of this study was to evaluate the efficacy of different herbicide mixtures used in recommended and reduced doses on broad-leaved weeds applied pre- and postemergence in sugar beet. Treatments included preemergence application of chloridazon (Pyramin Turbo, 520 g ai l<sup>-1</sup>) and metamitron (Goltix SC, 700 g ai l<sup>-1</sup>) and postemergence application of the mixtures of phenmedipham plus desmedipham plus ethofumesate (Betanal Expert, 274 g ai l<sup>-1</sup>) with chloridazon, metamitron, triflurosulfuron-

methyl (Caribou, 500 g ai kg<sup>-1</sup>), ethofumesate (Nortron, 500 g ai l<sup>-1</sup>), and of the mixtures of phenmedipham (Betasana, 160 g ai l<sup>-1</sup>) with ethofumesate, metamitron, mineral oil, and of the mixtures of phenmedipham (Kontakt SC, 320 g ai l<sup>-1</sup>) with ethofumesate, metamitron, rapeseed oil, and of the mixtures of phenmedipham (Betasana) with ethofumesate, metamitron, rapeseed oil. Soil texture was loam consisting of 14.5–17.7% clay, 34.8–39.9% silt, 44.7–51.1% sand. Humus content amounted to 1.6–2.4%, and pH – 6.1–6.9. The field was fertilized with nitrogen, phosphorus, and potassium at the ratio of 105–120:80–120:120–170 kg ha<sup>-1</sup>. Mineral fertilizers were incorporated into the soil during cultivation. Sugar beet was planted with 45 cm row space, at a density of 15 plants m<sup>-2</sup>. The herbicides were tank-mixed and applied postemergence at three different dates. The first application was done at the early cotyledon stage of weed growth. Subsequent applications were applied when the next weed flush emerged or 10–17 days after the first flush. The plot size was 2.5 m x 10 m. The herbicides in the experiment were broadcast-applied. The amount of water was 200 l ha<sup>-1</sup>. Weed dry weight was measured two times: four weeks after herbicide application and before harvest. At the time of assessment a quadrat of 0.20 m x 1.25 m was randomly thrown in each plot. Weed control was assessed by visually estimating the % control relative to the ground cover and vigor of each weed species in the untreated plots. Weed samples were dried at 105°C for 24 h and weighed. Weed density and dry weight data were transformed to  $\sqrt{x+1}$ . The data were analyzed with ANOVA and LSD test.

## 2. Weed flora in sugar beet

In much sugar beet growing areas, dicot weeds of the families Chenopodiaceae, Asteraceae, Brassicaceae, and Polygonaceae are of major importance. The monocots are less important compared to dicot weeds [2, 5]. Broadleaf weeds often grow to a height two to three times that of sugar beet by mid-summer. Annual broad-leaved weeds are usually more competitive than annual grasses [25].

The botanical surveys of species were conducted before herbicide application. Overall, 24 weed species were found. The number of weeds found in 2004–2005 and 2010–2012 was from 41 to 108 weeds m<sup>-2</sup>. In 2011 and 2012, the germination of weeds was lowest in sugar beet; the weed number was 41 and 49 m<sup>-2</sup>, respectively. Weeds abundantly germinated in 2005, the number of weeds was 108 and 106 m<sup>-2</sup>, respectively. The dominant weed species in all years were *Chenopodium album* L. (from 11 to 62 weed m<sup>-2</sup>), *Lamium purpureum* L. (from 3 to 30 weed m<sup>-2</sup>), *Stellaria media* (L.) Vill. (from 2 to 40 weed m<sup>-2</sup>), *Viola arvensis* Murray (from 2 to 18 weed m<sup>-2</sup>), and *Thlaspi arvense* L. (from 1 to 14 weed m<sup>-2</sup>). In Latvia, the most frequent species of annual dicots in sugar beet were: *Tripleurospermum perforatum* (Merat.) M. Lainz, *Chenopodium album*, *Fallopia convolvulus* (L.) Löve, *Capsella bursa-pastoris* (L.) Medik, and *Stellaria media* [26]. *Chenopodium album* was the dominant weed species from the 19–24 species identified. This species accounted for 10–58% of the total weeds documented. According to literature on the population dynamics of a common arable weed, *Chenopodium album*, and its interactions with an arable crop, sugar beet, where *Chenopodium album* and other weeds may also be a considerable problem [7]. Our research data revealed that *Galium aparine* L., *Veronica arvensis* L., and

*Erysimum cheiranthoides* L. were present at a low frequency (Figure 1). Other weeds such as *Tripleurospermum perforatum*, *Fumaria officinalis* L., *Fallopia convolvulus*, *Lapsana communis* L., *Polygonum aviculare* L., *Polygonum persicaria* L., *Capsella bursa-pastoris*, *Sinapis arvensis* L., *Euphorbia helioscopia* L., *Myosotis arvensis* (L.) Hill, *Chaenorhinum minus* (L.) Lange., *Centaurea cyanus* L., *Silene pratensis* (Rafn) Godr., *Anagalis arvensis* L., *Myosurus minimus* L., and *Galeopsis tetrahit* L. were less common species. These species germinated in only a few years of the study.




Figure 1. Weed species in sugar beet before herbicide application data averaged over 2004–2012

### 3. Sensitivity of weeds to phenmedipham, desmedipham, ethofumesate, metamitron, chloridazon, and triflurosulfuron combinations at preemergence application

Weed control in crops is mainly based on the use of herbicides because they are efficient and easily applied [27]. Weed control is one of the most difficult agricultural arrangements in sugar beet growing because of low crop interference with weeds [11]. After herbicide use, significant changes in weed flora were noted in terms of abundance and share of some weed species on total weed community [28, 29]. Herbicides for control of dicots can only be used until the crop starts to develop true leaves and their efficacy decreases as the weeds grow [30].

Weed control programs in sugar beet include both pre- and postemergence herbicide treatments [31]. The effectiveness of preemergence residual herbicides decreases with reductions in rainfall or soil moisture content [32]. Preemergence application of soil herbicides is used limitedly because it strongly depends on soil moisture [33]. Therefore, less than 10% of the total sugar beet crop is treated with preemergence herbicides. The remaining 90% depends solely on a selection of postemergence herbicides to maintain season-long weed control [34].

The advantage of soil applied residual herbicides is that they reduce the number of weeds that emerge with the crop and often sensitize survivors to subsequent postemergence sprays. When

residual herbicides are used after sowing, they must be applied to the soil surface before sugar beet seedlings emerge or crop damage may result. Preemergence herbicides are important for the subsequent postemergence applications and provide some flexibility with timing and selection of postemergence treatments [35].

The main preemergence residual broad-leaved weed control herbicides used on sugar beet crops are chloridazon and metamiltron. Chloridazon is a pyridazinone herbicide with pre-emergence and postemergence activity. This herbicide is usually applied prior to emergence of beet and weeds, and may also be applied postemergence to control common lambsquarters in combination with other herbicides [36]. Metamiltron is a 1, 2, 4-triazinone herbicide which is absorbed predominantly by the roots, but also the leaves. This herbicide is applied predrilling incorporated, pre- and postemergence. Metamiltron is applied in tank-mix with other herbicides postemergence [37].

Our research data revealed that the efficacy of herbicides varied from 35.0 to 100% (Table 1,2). In 2010, the efficacy of herbicides was higher than in 2011 because the growing season of 2010 started later than normal and the spring rainfall was higher than the perennial average. Total amount of rain was significantly higher and amounted to 20 and 80%, respectively, as compared to long-term average. In April and May of 2011, dry weather prevailed. The amount of precipitation was 42 and 90% of that as the long-term average, respectively. Air temperature, soil moisture, and relative humidity affected herbicide efficacy [38].

Treatment	Efficacy in 1 month after DAA, %				
	CHEAL	POLCO	STEME	LAMPU	EPHHE
Weedy check	0	0	0	0	0
Metamiltron, 2100 g a.i. ha <sup>-1</sup> – predrilling (T <sub>0</sub> ); Metamiltron, 700 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> ; Raps oil, 0.5 l ha <sup>-1</sup> – T <sub>0</sub> , T <sub>1</sub> , T <sub>3</sub>	98.5b	89.0b	100.0a	98.3b	98.8b
Metamiltron, 1400 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; Metamiltron, 1050 g a.i. ha <sup>-1</sup> – T <sub>2</sub> , T <sub>3</sub> ; Raps oil, 0.5 l ha <sup>-1</sup> – T <sub>0</sub> , T <sub>1</sub> , T <sub>3</sub>	100.0a	95.3a	99.8a	99.5a	100.0a
Metamiltron + phenmedipham + ethofumesate, 1400+160+35 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; Metamiltron + phenmedipham + ethofumesate, 1050+160+35 g a.i. ha <sup>-1</sup> – T <sub>2</sub> , T <sub>3</sub> ; Raps oil, 0.5 l ha <sup>-1</sup> – T <sub>0</sub> , T <sub>1</sub> , T <sub>3</sub>	100.0a	99.8a	100.0a	100.0a	100.0a

Note. The means followed by the same letter within a line are not significantly different according to Fisher’s Protected LSD test (P<0.05).

**Table 1.** Effect of the herbicide combinations on weeds in sugar beet, 2010

The tank mixture of metamiltron at 1050 g ai ha<sup>-1</sup> or 1400 g ai ha<sup>-1</sup> with phenmedipham at 160 g ai ha<sup>-1</sup> and ethofumesate at 35 g ai ha<sup>-1</sup> and raps oil at 0.5 l ha<sup>-1</sup> significantly reduced *Chenopo-*

*dium album* (CHEAL), *Fallopia convolvulus* (POLCO), *Lamium purpureum* (LAMPU), and *Euphorbia helioscopiai* (EPHHE) as compared with pre- and postemergence application of metamiltron (Table 1). The higher efficacy (95.3–100.0%) on weeds was achieved when metamiltron at 1050 g ai ha<sup>-1</sup> or 1400 g ai ha<sup>-1</sup> with raps oil at 0.5 l ha<sup>-1</sup> was applied postemergence.

In 2011, in dry years, the efficacy of metamiltron alone was lower (35.0–62.5%) than when in combination with other herbicides (Table 2). Preemergence application of metamiltron provided significantly lower efficacy on *Chenopodium album*, but significantly higher efficacy on *Galium aparine* (GALAP) than postemergence application of this herbicide. In other studies, metamiltron controlled *Chenopodium album* up to two weeks after application thoroughly. One month after application *Chenopodium album* regenerated [38]. The combination of metamiltron with phenmedipham plus desmedipham plus ethofumesate plus raps oil resulted excellent control of weeds (>96%).

Treatment	Efficacy in 1 month after DAA, %				
	CHEAL	POLCO	STEME	GALAP	VIOAR
Weedy check	0	0	0	0	0
Metamiltron, 2100 g a.i. ha <sup>-1</sup> – predrilling (T <sub>0</sub> ); Metamiltron, 700 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> ; Raps oil, 0.5 l ha <sup>-1</sup> , T <sub>0</sub> , T <sub>1</sub> , T <sub>3</sub>	56.3de	43.8bc	48.8bc	56.3b	37.5b
Metamiltron, 1400 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; Metamiltron, 1050 g a.i. ha <sup>-1</sup> – T <sub>2</sub> , T <sub>3</sub> ; Raps oil, 0.5 l ha <sup>-1</sup> – T <sub>0</sub> , T <sub>1</sub> , T <sub>3</sub>	62.5bc	42.5bc	43.8cd	46.3c	35.0b
Metamiltron + phenmedipham plus thofumesate, 1400+160+35 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; Metamiltron + phenmedipham + ethofumesate, 1050+160+35 g a.i. ha <sup>-1</sup> – T <sub>2</sub> , T <sub>3</sub> ; Raps oil, 0.5 l ha <sup>-1</sup> , T <sub>0</sub> , T <sub>1</sub> , T <sub>3</sub>	98.0a	98.5a	100.0a	97.8a	96.8a

Note. The means followed by the same letter within a line are not significantly different according to Fisher's Protected LSD test (P<0.05).

**Table 2.** Effect of the herbicide combinations on weeds in sugar beet, 2011

Herbicides can interact with each other in tank-mixed and can cause damage or reduce crop populations [35]. The visual crop injury symptoms included deformation and yellowing of leaves, growth reduction, and thinning (Figure 2). Statistical analysis of the data on visual injury showed that the effect of year with treatments was significant. The visual injury in metamiltron-treated plots ranged from 64% of preemergence and 0% of postemergence when herbicides were applied at low doses (Table 3). Sugar beet recovered from metamiltron injury even at high doses [39]. Other studies also have reported no or less injury of sugar beet plants with the application of herbicides at reduced doses compared to full dose application [40]. No visible symptoms of phytotoxicity on sugar beet plants were noticed after postemergence metamiltron and this herbicide tank-mixed with phenmedipham plus desmedipham plus

ethofumesate plus raps oil application. The phytotoxicity of herbicides decreased with time. To avoid injury, growth depressions, or leaf damage of sugar beet plants, herbicide use has to be carefully adjusted especially to the prevailing weather conditions [41].

Treatment	2010			2011		
	7 DAT	14 DAT	28 DAT	7 DAT	14 DAT	28 DAT
Weedy check	0	0	0	0	0	0
Metamitron, 2100 g a.i. ha <sup>-1</sup> – predrilling (T <sub>0</sub> ); Metamitron, 700 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> ; Raps oil, 0.5 l ha <sup>-1</sup> , T <sub>0</sub> , T <sub>1</sub> , T <sub>3</sub>	64.0**	61.3**	61.3**	0	0	0
Metamitron, 1400 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; Metamitron, 1050 g a.i. ha <sup>-1</sup> – T <sub>2</sub> , T <sub>3</sub> ; Raps oil, 0.5 l ha <sup>-1</sup> , T <sub>0</sub> , T <sub>1</sub> , T <sub>3</sub>	0	0	0	0	0	0
Metamitron + phenmedipham + thofumesate, 1400+160+35 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; Metamitron + phenmedipham + ethofumesate, 1050+160+35 g a.i. ha <sup>-1</sup> – T <sub>2</sub> , T <sub>3</sub> ; Raps oil, 0.5 l ha <sup>-1</sup> , T <sub>0</sub> , T <sub>1</sub> , T <sub>3</sub>	0	0	0	0	0	0

Note. \*\*differences are statistically significant as compared to the control at  $P < 0.01$ . T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, and T<sub>4</sub> – first, second, third, and fourth application.

**Table 3.** Visual injury on sugar beet treated with pre- and postemergence herbicides



**Figure 2.** Sugar beet injury from preemergence application of metamitron: (a) yellowing, (b) thinning

The infestation of *Chenopodium album* (CHEAL), *Fallopia convolvulus* (POLCO), *Galium aparine* (GALAP), *Stellaria media* (STEME), and *Lapsana communis* (LAPCO) were noted (Table 4). After herbicide application, significant changes were noted in the weed flora. When chloridazon was applied preemergence or postemergence, the herbicidal activity was very high. Preemergence



application of chloridazon at 2080 g a.i. ha<sup>-1</sup> and postemergence application of tank-mixed phenmedipham plus desmedipham plus ethofumesate with metamitron resulted in excellent control of *Chenopodium album*, *Fallopia convolvulus*, *Galium aparine*, and *Stellaria media* (99–100%) and provided good control of *Lapsana communis* (91 %).

Treatment	CHEAL	POLCO	GALAP	STEME	LAPCO
Weedy check	259.0b	6.8b	9.9b	6.4b	10.8c
Chloridazon, 2080 g a.i. ha <sup>-1</sup> – predrilling;					
Phenmedipham + desmedipham + ethofumesate + metamitron, 91+71+114+700 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub>	0.0a	0.01ab	0.0a	0.0a	1.0ab
Phenmedipham + desmedipham + ethofumesate + chloridazon 91+71+112+650 g ha <sup>-1</sup> a.i. – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub>	0.9a	0.0a	0.8ab	0.0a	2.0abc
Phenmedipham + desmedipham + ethofumesate + metamitron, 91+71+112+700 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub>	5.0a	0.1ab	0.4ab	0.0a	0.5a

Note. The means followed by the same letter within a line are not significantly different according to Fisher's Protected LSD test ( $P < 0.05$ ).

**Table 4.** Biomass of prevailing weeds species (g m<sup>-2</sup>) in sugar beet 1 month after DAA, data averaged over 2004–2005

Postemergence application of chloridazon with phenmedipham plus desmedipham plus ethofumesate resulted in a similar effect on weeds as with the preemergence application. There was no significant difference when comparing both applications. The combination of chloridazon with phenmedipham plus desmedipham plus ethofumesate and metamitron with phenmedipham plus desmedipham plus ethofumesate provided a similar reduction of weed biomass. At the final assessment (3 month after DAA), weed density and biomass decreased compared with first assessment, respectively 42.3 and 25.7% (Table 5).

Treatment	Density, weed m <sup>-2</sup>		Weed biomass, g m <sup>-2</sup>	
	1 month after DAA	3 month after DAA	1 month after DAA	3 month after DAA
Weedy check	96.9	55.9	424.7	315.4
Chloridazon, 2080 g a.i. ha <sup>-1</sup> – predrilling;				
Phenmedipham + desmedipham + ethofumesate + metamitron, 91+71+114+700 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub>	10.6**	3.2*	6.6**	13.1*
Phenmedipham + ethofumesate + chloridazon 91+71+112+650 g ha <sup>-1</sup> a.i. – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub>	5.1**	1.8**	2.3**	1.5*
Phenmedipham + desmedipham + ethofumesate + metamitron, 91+71+112+700 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub>	13.0**	3.2*	11.3**	5.7*

Note. \*differences are statistically significant as compared to the control at  $P < 0.05$ , \*\*-at  $P < 0.01$ . T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>, and T<sub>4</sub> – first, second, third, and fourth application.

**Table 5.** Effect of the herbicide combinations on weed density and biomass in sugar beet; data averaged over 2004–2005

The results showed that combination of herbicides significantly affected weed control. A preemergence application of chloridazon at 2080 g a.i. ha<sup>-1</sup> and two postemergence applications of phenmedipham plus desmedipham plus ethofumesate with metamiltron resulted in a similar effect on weeds as a postemergence application of tank-mix of phenmedipham plus desmedipham plus ethofumesate with metamiltron and chloridazon. Chloridazon did not influence effectivity. The addition of chloridazon and metamiltron similarly affected efficacy of phenmedipham plus desmedipham plus ethofumesate.

#### **4. Combinations of phenmedipham, desmedipham, ethofumesate, metamiltron, chloridazon, and triflusulfuron at postemergence application on weeds and sugar beet**

Often sugar beets are treated with postemergence herbicides two or more times [16, 20, 28, 40]. Sometimes, more herbicide applications may be necessary [40]. Herbicides are applied at the cotyledon growth stage at 5–14-day intervals [42–45]. The major herbicides are phenmedipham, desmedipham, ethofumesate, chloridazon, metamiltron, clopyralid, lenacil, and triflusulfuron-methyl [7, 46–48]. Individual sugar beet herbicides seldom have a wide enough weed control spectrum or residual activity to control all weeds [49], and tank-mixes of different herbicides are commonly used in order to provide a broad spectrum of weed control [35]. The optimization of herbicide application in the sugar beet protection system can be achieved by using mixtures of appropriate components and their selected doses [30, 49]. Mixing compatible herbicides can have benefits such as consumption reduction, increased weed control, economization of the number of applications, release of fewer chemicals into the ecosystem with using their synergistic effects, decrease in residue of herbicide in soil and crops in low concentrations and reduced occurrence of herbicide resistance in weeds [50]. Weed control is often higher from tank-mixed herbicides than from a single herbicide [20, 38, 41, 47, 50, 51]. The herbicides phenmedipham, desmedipham, and ethofumesate are commonly tank-mixed with metamiltron, while chloridazon and triflusulfuron are used for broad-leaved weed control in sugar beet [37, 38, 43, 45, 52].

The tank-mix of phenmedipham plus desmedipham plus ethofumesate at 1029 g ai ha<sup>-1</sup> controlled *Chenopodium album* better than the combination of this herbicide at 822 g a.i. ha<sup>-1</sup> with triflusulfuron, but the efficacy was lower on *Tripleurospermum perforatum*. Other studies have shown a good control of *Chenopodium album* with phenmedipham plus desmedipham plus ethofumesate [53]. The effect of herbicide treatments on density and biomass of weeds was not significant (Table 6). The addition of triflusulfuron increased the effectiveness of phenmedipham plus desmedipham plus ethofumesate. Results of root yield showed that the combination of herbicides used had no significant effect on root yield as compared to the control.

At the first assessment 1 month after application (DAA), all combinations of herbicides similarly controlled weed density, except where phenmedipham plus desmedipham plus ethofumesate with metamiltron and ethofumesate and triflusulfuron were applied (Table 7). At the final

Treatment	Density, weed m <sup>-2</sup>		Weed biomass, g m <sup>-2</sup>		Root yield, t ha <sup>-1</sup>
	1 month after	3 month after	1 month after	3 month after	
	DAA	DAA	DAA	DAA	
Control (cleaned manually)	7.5	1.1	2.1	2.2	75.8
Phenmedipham + desmedipham + ethofumesate, 114+89+140 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (1029 g)	1.3	5.5**	59.0**	70.6**	76.1
Phenmedipham + desmedipham + ethofumesate, 91+71+112 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (822 g); Triflurosulfuron, 5 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ; 10 g a.i. ha <sup>-1</sup> – T <sub>3</sub>	9.7	4.3**	50.2**	64.1**	75.0

Note. \*\*differences are statistically significant as compared to the control at  $P < 0.01$ .

T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub> – first, second, and third application.

**Table 6.** Effect of the herbicide combinations on weeds and sugar beet; data averaged over 2010–2012

assessment, at the 3-month DAA, all treatments resulted in similar effect on weed density. The significantly lowest efficacy on biomass of weeds was the combination of phenmedipham with ethofumesate and metamitron (544+500+700 g a.i. ha<sup>-1</sup>) and raps oil as compared to the phenmedipham plus desmedipham plus ethofumesate (Control II) and other herbicides treatments. Phenmedipham plus desmedipham plus ethofumesate at 1029 g a.i. ha<sup>-1</sup> and phenmedipham plus desmedipham plus ethofumesate at 822 g a.i. ha<sup>-1</sup> with triflurosulfuron at 15 g a.i. ha<sup>-1</sup> decreased weed biomass similarly. The biomass of weeds was significantly lower after application of tank-mixed phenmedipham plus desmedipham plus ethofumesate with metamitron, ethofumesate, and triflurosulfuron (319+249+492+10 g a.i. ha<sup>-1</sup>) as compared to other herbicide combinations. Other studies also have reported that phenmedipham plus desmedipham plus ethofumesate was more effective for controlling weeds by applying in a mixture with metamitron than by applying alone phenmedipham plus desmedipham plus ethofumesate [54, 55]. The combination of herbicides decreased sugar beet root yield as compared to the hand-weeded check (Control I). Similar results were reported elsewhere [34, 49]. Only application of phenmedipham with ethofumesate and metamitron (544+500+700 g a.i. ha<sup>-1</sup>) and raps oil significantly decreased root yield as compared to control I.

## 5. Sensitivity of weeds to low rates of phenmedipham, desmedipham, ethofumesate, metamitron, chloridazon, and triflurosulfuron

In older systems used for weed control in sugar beets, herbicides were applied at a high, single dose. Herbicides are often applied at rates higher than required for weed control under ideal conditions [44]. A single full-rate of phenmedipham and/or desmedipham controlled weeds better and caused less sugar beet injury than half-rate application [56]. By testing the efficacy

Treatment	Density, weed m <sup>-2</sup>		Weed biomass, g m <sup>-2</sup>		Root yield, t ha <sup>-1</sup>
	1 month after DAA	3 month after DAA	1 month after DAA	3 month after DAA	
Control I (cleaned manually)	1.3	0.9	4.8	2.8	80.6
Control II. Phenmedipham + desmedipham + ethofumesate, 114+89+140 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (1029 g)	16.2	6.4	86.7	96.4	76.8
Phenmedipham + desmedipham + ethofumesate + metamitron, 91+71+112+700 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; Phenmedipham + desmedipham + ethofumesate + ethofumesate, 114+89+140+100 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ;	7.4**	3.5	14.9**	17.1**	77.2
Phenmedipham + desmedipham + ethofumesate + triflusulfuron, 114+89+140 +10 g a.i. ha <sup>-1</sup> – T <sub>3</sub> (319+249+492+10 g)					
Phenmedipham + ethofumesate + metamitron 160+100+700 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; Phenmedipham + ethofumesate 224+150 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ;	20.4	10.5	148.0*	249.7**	72.3**
Phenmedipham + ethofumesate 160+250 g a.i. ha <sup>-1</sup> – T <sub>3</sub> (544+500+700 g) Raps oil 0.5 l ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub>					
Phenmedipham + desmedipham + ethofumesate, 91+71+112 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (822 g); Triflusulfuron 5 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ; 10 g a.i. ha <sup>-1</sup> – T <sub>3</sub> (15 g)	14.2	6.0	75.2	94.0	75.9

Note. \*differences are statistically significant as compared to the control at  $P < 0.05$ , \*\*-at  $P < 0.01$ .

T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub> – first, second, and third application.

**Table 7.** Effect of the herbicide combinations on weed density and biomass in sugar beet; data averaged over 2011–2012

of a herbicide over a wide range of rates, growers will have better information to determine the appropriate weed management program that maximizes net returns and minimizes loading of herbicides into the environment [57]. Reducing the recommended dose of herbicides is one of the important instruments in weed management systems. Reduced herbicide applications could be achieved either by reducing the dosages or the number of treatments [53]. The exploitation of competitiveness factors might favor the development of reduced herbicide use strategies for sugar beet [9]. Numerous research studies have indicated a few reasons for the potential successful use of reduced dose: 1) registered doses are set to ensure adequate control over a wide spectrum of weed species, weed densities, growth stages, and environmental conditions; 2) maximum weed control is not always necessary for optimal crop yields; and 3) combining reduced doses of herbicides with other management practices, such

as tillage or competitive crops, can markedly increase the odds of successful weed control [30, 58]. Another researcher has shown that it is possible to reduce herbicide doses in sugar beet [38, 44, 45, 50, 59, 60]. For example, Goleblowska and Domaradzki [48] reported that a 50% and 67% dose of Betanal Progress + Goltix + Safari and Betanal Progress + Venzar + Safari consistently produced 94–97% weed control. The half dose of herbicides reduced weed biomass significantly [38]. The lower and frequent doses of herbicide reached comparable or better results in comparison with the traditional system of application [34].

The weed spectrum was similar in both years. The results showed that the efficacy of phenmedipham plus desmedipham plus ethofumesate (1029 g a.i. ha<sup>-1</sup>) was lower on *Chenopodium album* (CHEAL), *Tripleurospermum perforatum* (MATIN), *Polygonum aviculare* (POLCO), *Thlaspi arvense* (THLAR), and *Viola arvensis* (VIOAR) (Table 8). The additions of metamiltron (1050 g) and triflusalufuron (15 g) increased efficacy of phenmedipham plus desmedipham plus ethofumesate. Similar cases of metamiltron effectiveness have been reported by many authors [59, 61].

Treatment	CHEAL	MATIN	POLAV	THLAR	VIOAR
Control II. Phenmedipham + desmedipham + ethofumesate, 114+89+140 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (1029 g)	33.4b	4.4b	6.3b	1.5b	0.2b
Phenmedipham + desmedipham + ethofumesate, 91+71+112 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (822 g); Triflusalufuron, 5 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ; 10 g a.i. ha <sup>-1</sup> – T <sub>3</sub> (15 g)	12.2ab	1.7ab	2.0ab	0.3ab	0.2ab
Phenmedipham + desmedipham + ethofumesate, 68+53+84 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (615 g); Metamiltron 350 g ha <sup>-1</sup> a.i. – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (1050 g)	18.2ab	1.4ab	0.4ab	0.3ab	0.02ab

Note. The means followed by the same letter within a line are not significantly different according to Fisher's Protected LSD test (P<0.05).

**Table 8.** Biomass of prevailing weed species (g m<sup>-2</sup>) in sugar beet 1 month after DAA; data averaged over 2010–2011

All herbicide treatments had similar effects on weed density, except treatments where combination of phenmedipham plus desmedipham plus ethofumesate (822 g a.i. ha<sup>-1</sup>) with triflusalufuron were applied (Table 9). The least biomass of weeds was recorded for phenmedipham plus desmedipham plus ethofumesate (1029 g ha<sup>-1</sup> – full dose). Reducing the doses of phenmedipham plus desmedipham plus ethofumesate by 20% with triflusalufuron and by 40% with metamiltron, their effectiveness significantly reduced at final assessment. Metamiltron with tank-mixes of phenmedipham plus desmedipham plus ethofumesate had similar effect on weeds compared to triflusalufuron with phenmedipham plus desmedipham plus ethofumesate. Effect of combination herbicides was not significant on sugar beet root yield as compared with control I.

The postemergence trials showed that commercial mixture of phenmedipham plus desmedipham plus ethofumesate (1029 g ha<sup>-1</sup> – full dose) effectively decreased the biomass of

Treatment	Density, weed m <sup>-2</sup>		Weed biomass, g m <sup>-2</sup>		Root yield, t ha <sup>-1</sup>
	1 month after	3 month after	1 month after	3 month after	
	DAA	DAA	DAA	DAA	
Control I (cleaned manually)	1.8	1.4	2.5	2.5	83.0
Control II. Phenmedipham + desmedipham + ethofumesate, 114+89+140 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (1029 g)	11.1	5.5	46.8	44.4	82.4
Phenmedipham + desmedipham + ethofumesate, 91+71+112 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (822 g); Triflurosulfuron, 5 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ; 10 g a.i. ha <sup>-1</sup> – T <sub>3</sub> (15 g)	7.2	2.2*	21.2	14.0*	81.1
Phenmedipham + desmedipham + ethofumesate, 68+53+84 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (615 g); Metamitron 350 g ha <sup>-1</sup> a.i. – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (1050 g)	8.2	3.5	18.7	14.2*	81.7

Note. \*differences are statistically significant as compared to the control at  $P < 0.05$ .

T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub> – first, second, and third application.

**Table 9.** Effect of the herbicide combinations on weeds and sugar beet; data averaged over 2010–2011

*Chenopodium album* (CHEAL), *Veronica arvensis* (VERAR), and *Galium aparine* (GALAP), but the differences were not statistically significant. In the treatment where by reducing dose of phenmedipham plus desmedipham plus ethofumesate by 40% with triflurosulfuron at 30 and 45 g ha<sup>-1</sup> was applied, the biomass of *Veronica arvensis* (VERAR) was recorded to be higher as compared to that of full dose of phenmedipham plus desmedipham plus ethofumesate. The herbicide combination did not have significant influence on weight of botanical composition of weed flora.

All herbicide combinations similarly affected weed density, except phenmedipham plus desmedipham plus ethofumesate (615 g a.i. ha<sup>-1</sup>) with triflurosulfuron at 30 g a.i. ha<sup>-1</sup> (Table 11). In this mixture, dose of phenmedipham plus desmedipham plus ethofumesate were the lowest. When the dose of phenmedipham plus desmedipham plus ethofumesate in a herbicide mixture was reduced by 40% and addition of triflurosulfuron at reducing dose by 33% (30 g a.i. ha<sup>-1</sup>) was used, the effectiveness of phenmedipham plus desmedipham plus ethofumesate was not reduced. At the first assessment (1 month after DAA), different herbicide treatments had no significant effect on biomass of weeds. At the final assessment, triflurosulfuron with tank-mixes of phenmedipham plus desmedipham plus ethofumesate had a greater effect on biomass of weeds than phenmedipham plus desmedipham plus ethofumesate. When the dose of phenmedipham plus desmedipham plus ethofumesate in this herbicide combination was reduced by 40% the biomass of weeds significantly decreased as compared to phenmedipham plus desmedipham plus ethofumesate of reducing dose by 20%. Weed control from herbicide combinations of phenmedipham plus desmedipham plus ethofumesate with full dose (45 g

Treatment	CHEAL	MATIN	VERAR	POLCO	GALAP
Phenmedipham + desmedipham plus ethofumesate, 114+89+140 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (1029 g)	0.00a	0.93ab	0.00ab	2.27b	0.00ab
Phenmedipham + desmedipham + ethofumesate, 91+71+112 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (822 g); Trilussulfuron, 10 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ; 20 g a.i. ha <sup>-1</sup> – T <sub>3</sub> (30 g)	0.00a	0.05ab	0.00ab	0.00ab	0.10ab
Phenmedipham + desmedipham + ethofumesate, 68+53+84 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (615 g); Trilussulfuron, 15 g a.i. ha <sup>-1</sup> – T <sub>2</sub> , T <sub>3</sub> (30 g)	0.25c	1.91b	0.66ab	0.00ab	0.07ab
Phenmedipham + desmedipham + ethofumesate, 68+53+84 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (615 g); Trilussulfuron, 10 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; 15 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ; 20 g a.i. ha <sup>-1</sup> – T <sub>3</sub> (45 g)	0.01abc	0.00ab	0.77b	0.00ab	0.10b

Note. The means followed by the same letter within a line are not significantly different according to Fisher's Protected LSD test ( $P < 0.05$ ).

**Table 10.** Biomass of prevailing weeds species (g m<sup>-2</sup>) in sugar beet 1 month after DAA; data averaged over 2011–2012

a.i. ha<sup>-1</sup>) of triflussulfuron was the highest. Sugar beet yield was not significantly different between herbicide treatments. All herbicide treatments produced lower sugar beet yields than the hand-weeded check. Similar results were reported elsewhere [49, 62].

Treatment	Density, weed m <sup>-2</sup>		Weed biomass, g m <sup>-2</sup>		Root yield, t ha <sup>-1</sup>
	1 month after DAA		3 month after DAA		
	DAA	DAA	DAA	DAA	
Control. Phenmedipham + desmedipham + ethofumesate, 114+89+140 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (1029 g)	1.5	3.8	3.6	19.1	74.6
Phenmedipham + desmedipham + ethofumesate, 91+71+112 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (822 g); Trilussulfuron, 10 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ; 20 g a.i. ha <sup>-1</sup> – T <sub>3</sub> (30 g)	0.5	1.0	0.2	4.4	70.2
Phenmedipham + desmedipham + ethofumesate, 68+53+84 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (615 g); Trilussulfuron, 15 g a.i. ha <sup>-1</sup> – T <sub>2</sub> , T <sub>3</sub> (30 g)	6.8*	2.0	3.1	4.2*	70.8
Phenmedipham + desmedipham + ethofumesate, 68+53+84 g a.i. ha <sup>-1</sup> – T <sub>1</sub> , T <sub>2</sub> , T <sub>3</sub> (615 g); Trilussulfuron, 10 g a.i. ha <sup>-1</sup> – T <sub>1</sub> ; 15 g a.i. ha <sup>-1</sup> – T <sub>2</sub> ; 20 g a.i. ha <sup>-1</sup> – T <sub>3</sub> (45 g)	3.2	1.0	1.2	1.2*	69.0

Note. \*differences are statistically significant as compared to the control at  $P < 0.05$ .

T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub> – first, second, and third application.

**Table 11.** Effect of the herbicide combinations on weeds and sugar beet; data averaged over 2011–2012

## 6. Conclusion

All herbicide combinations acted similarly on reduction of the following weed species: *Chenopodium album*, *Thlaspi arvense*, *Tripleurospermum perforatum*, *Polygonum aviculare*, *Veronica arvensis*, *Stellaria media*, and *Lapsana communis*. Postemergence application of chloridazon with phenmedipham plus desmedipham plus ethofumesate resulted in a similar effect on weeds compared to preemergence applications. The efficacy of phenmedipham plus desmedipham plus ethofumesate was similar in action as compared to that applied in tank-mixes with chloridazon, metamiltron, and triflusalufuron. There were no significant differences on weight of weeds. The addition of chloridazon, metamiltron, and triflusalufuron controlled weeds similarly. The significantly lowest efficacy on weeds resulted from a combination of phenmedipham with ethofumesate and metamiltron as compared to the phenmedipham plus desmedipham plus ethofumesate. Two reduced doses (by 20% and 40%) of phenmedipham plus desmedipham plus ethofumesate in tank-mix had a significant effect on weeds compared to that of all doses of phenmedipham plus desmedipham plus ethofumesate. Full and reduced doses (by 33%) of triflusalufuron with phenmedipham plus desmedipham plus ethofumesate similarly affected weeds. The herbicides investigated did not have any negative influence on sugar beet productivity and quality.

## Acknowledgements

This study has been supported by the UAB "Nordic Sugar Kedainiai."

## Author details

Irena Deveikyte\*, Lina Sarunaite and Vytautas Seibutis

\*Address all correspondence to: [irenad@lzi.lt](mailto:irenad@lzi.lt)

Institute of Agriculture, Lithuanian Research Centre for Agriculture and Forestry, Instituto al., Akademija, Kedainiai distr., Lithuania

## References

- [1] Heidari G, Nasab AD, Javanshor A, Khoie FR, Moghaddam M. Influence of redroot pigweed (*Amaranthus retroflexus* L.) emergence time and density on yield and quality of two sugar beet cultivars. *J Food Agric Environ* 2006; 5(3-4):261-266.
- [2] Zoschke A, Quadranti M. Integrated weed management: Quo vadis. *Weed Biologic Manag* 2002; 1-10.



- [3] Schäufele WR. Einfluß niedrigwaschsender Unkräuter zwischen den Reihen auf ertrag von Zuckerrüben. *Gesunde Pflanze* 1991; 43:175-179.
- [4] Mittler S, Petersen J, Koch HJ. Bekämpfungsschwellen bei der Unkrautbekämpfung in Zuckerrüben. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz. Sonderheft* 2002; XVIII:499-509.
- [5] Roos H, Brink A. Optimierung von Produkteigenschaften durch Formulierungsentwicklung am Beispiel Rübenherbicide. *Mitteilungen aus der Biologischen Bundesanstalt für Land-und Fortwirtschaft* 1996; 321:63.
- [6] Kemmer A. Zusammenhänge zwischen Selektivität von Rübenherbiziden und Ertrag. *BASF AG* 1997; 25-29, 31-35.
- [7] May M. Crop protection in sugar beet. *Pestic Outlook* 2001;12(5) 188-191.
- [8] Schweizer EE, Dexter AG. Weed control in sugarbeets (*Beta vulgaris*) in North America. *Rev Weed Sci* 1987; 3:1133.
- [9] Paolini R, Principi M, Froud-Wiiliams RJ, Del Puglia S, Biancardi E. Competition betwvn sugar beet and *Sinapis arvensis* and *Chenopodium album*, as affected by timing of nitrogen fertilization. *Weed Res* 1999; 39(6):425-440.
- [10] Salehi F, Esfandiari H, Mashhadi HR. Critical period of weed control in sugar beet in Shahrekord region. *Iran J Weed Sci* 2006; 2(2):1-12
- [11] Jursík M, Holec J, Soukup J, Venclová V. Competitive relationships between sugar beet and weeds in dependence on time of weed control. *Plant Soil Environ* 2008; 54(3): 108-116.
- [12] Soroka SV, Gadzhieva GJ. State of weed infestation and features of sugar beet protection in Belarus. *Matica Srpska Proc Natur Sci* 2006; 110:165-172.
- [13] Montemurro P, Castrignano A, Fracchiolla M, Lasorella C. The critical period for weed control in sugar beet. *Proc 11th EWRS Sympos* 28 June–1 July 1999; Basel, Switzerland; 1999 p. 67.
- [14] Sysmans J, D'Hollander R, Schoonejans T, Tossens H, Vincinaux C. Research on herbicide efficiency and tolerance of 'low dose systems' for weed control in beet. *Mededelingen van de Faculteit Landbouwtwenschappen* 1991; 56:617-631.
- [15] Lajos K, Lajos M. Unkrautbekämpfung mit verringerten Aufwandmengen in Zuckerrüben in Ungarn. *Zeitschrift für Pflanzenkrankheiten und Pflanzenschutz* 2000; XVII: 623-627.
- [16] Dale TM, Renner KA, Kravchenko AN. Effect of herbicides on weed control and sugar beet (*Beta vulgaris*) yield and quality. *Weed Technol* 2006; 20(1):150-156.

- [17] Van Njenhuis JH, Haverkamp HC. Economic view on the reduction of the use of herbicides in sugarbeet. *Mededelingen van de Faculteit Landbouwtwenschappen* 1992; 57(3): 1013-1019.
- [18] Wilson RG. Response of nine sugarbeet (*Beta vulgaris*) cultivars to post-emergence herbicide applications. *Weed Technol* 1999;13 25-29.
- [19] Wilson RG, Smith JA, Yonts CD. Repeated reduced rates of broadleaf herbicides combination with methylated seed oil for post emergence weed control in sugar beet (*Beta vulgaris*). *Weed Technol* 2005; 19(4):855-860.
- [20] Deveikyte I, Seibutis V. The influence of post-emergence herbicides combinations on broad-leaved weeds in sugar beet. *Zemdirbyste-Agriculture* 2008; 95(3):43-49.
- [21] Bauer H. Neue Strategien gegen spezielle Rübenukräuter. Die landwirtschaftliche *Zeitschrift für Management Produktion und Technik*. 1997; 1:58-63.
- [22] Dexter AG, Luecke JL. Weed control in transgenic sugarbeet in North Dakota and Minnesota. *Proc N Central Weed Sci Soc*; 1997; 52:142-143.
- [23] Scepanovic M, Baric K, Galzina N, Ostojic Z. Effect of low-rate herbicide treatments on weed biomass and yield of sugar beet. *Proc 14th EWRS Sympos* 18-21 June 2007; Norway. Oslo; 2007 p. 66.
- [24] Deveikyte I, Seibutis V. Broadleaf weeds and sugar beet response to phenmedipham, desmedipham, ethofumesate and triflusaluron-methyl. *Agron Res* 2006; 4 159-162.
- [25] Schweizer EE, May MJ. Weeds and weed control. In: Cooke DA, Scott RK, editors. *The Sugar Beet Crop*. Chapman & Hall; 1993. p. 485-519.
- [26] Vanaga I. Dynamics of the flora of arable fields in central Latvia. *Agonomijas Vestis* 2004; 7:176-182.
- [27] Lodovichi MV, Blanco AM, Chantre GR, Bandoni JA, Sabbatini MR, Vigna M, López R, Gigón R. Operational planning of herbicide-based weed management. *Agric Sys* 2013; 121(2):117-129.
- [28] Smatana J, Macák M, Demjanová E, Dalović I. Herbicide weed control in sugar beet. *Acta Herbologica* 2008; 17(2):131-135.
- [29] Smatana J, Macák M, Demjanová E, Djalović I. Weed control in canopy of sugar beet. *Res J Agric Sci* 2009; 41(1):199-302.
- [30] Strandberg B, Pedersen MB, Elmegaard N. Weed and arthropod populations in conventional and genetically modified herbicide tolerant fodder beet fields. *Agric Ecosys Environ* 2005; 105(2):243-253.
- [31] Coyette B, Tencalla F, Brants I, Fichet Y, Rouchouze D. Effect of introducing glyphosate – tolerant sugar beet on pesticide usage in Europe. *Pestic Outlook* 2002; 13:219-223.

- [32] Kayva R, Buzluk S. Integrated weed control in sugar beet through combinations of tractor hoeing and reduced dosages of a herbicide mixture. *Turk J Agric Forest* 2006; 30:137-144.
- [33] Urban J, Pulkrábek J, Valenta J, Bečková L, Kvíz Z. Influence of lower and more frequent doses of herbicides on yield and technological quality of sugar beet. *Proc 43rd Croatian and 3rd Int Sympos Agric*; 18-21 February 2008; Croatia. Opatija: 2008. p. 646-649.
- [34] Mitchell B. Weed control in sugar beet. *Crop Protect* 2005; 23 April, 40-43.
- [35] Cioni F, Maines G. Weed control in sugarbeet. *Sugar Tech* 2010; 12:3-4:243-255. DOI: 10.1007/s12355-010-0036-2.
- [36] Robinson DE, McNaughton KE, Bilyea D. Comparison of sequential preemergence-postemergence and postemergence-alone weed management strategies for re beet (*Beta vulgaris* L.). *Can J Plant Sci* 2013; 93(5):863-870. DOI: 10.4141/CJPS2012-327
- [37] Tomlin CDS, editor. *The Pesticide Manual*. 11th edn. British Crop Protection Council; 1997. 1606 p.
- [38] Najafi H, Bazoubandi M, Jafarzadeh N. Effectiveness of repeated reduced rates of selective broadleaf herbicides for postemergence weed control in sugar beet (*Beta vulgaris*). *World J Agric Res* 2013; 1(2):25-29.
- [39] Abbaspoor M, Teicher HB, Streibig JC. The effect of root-absorbed PSII inhibitors on Kautsky curve parameters in sugar beet. *Weed Res* 2006; 46(3):226-235
- [40] Wujek B, Kucharski M, Domaradzki K. Weed control programs in sugar beet (*Beta vulgaris* L.): Influence on herbicidal residue and yield quality. *J Food Agric Environ* 2012; 10(3-4):606-609.
- [41] Gummert A, Ladewig E, Märländer B. Guidelines for integrated pest management in sugar beet cultivation – weed control. *Journal für Kulturpflanzen* 2012; 64(4):105-111.
- [42] Konstantinović BI, Meseldžija MU. Occurrence spread and possibilities of invasive weeds control in sugar beet. *Proc Natur Sci Matca Sprska* 2006; 110:173-178
- [43] Odero DC, Meshab AO, Miller SD. Economics of weed management systems in sugarbeet. *Econ Weed Manag* 2008; 45(1&2):49-63.
- [44] Kucharski M. Changes in application system – influence on herbicides residue in soil and sugar beet roots. *J Plant Protect Res* 2009; 49(4):421-425.
- [45] Domaradzki K. Skuteczność mikrodawek herbicydów w systemach chemicznej ochrony buraka cukrowego. *Progr Plant Protect* 2011; 51(4):1683-1688.
- [46] Panjehkeh N, Alamshahi L. Influence of separate and tank-mixed application of some broadleaf herbicides on sugarbeet weeds and their effects on crop productivity. *Austr J Basic Appl Sci* 2011; 5(7):332-335.

- [47] Dewar AM, Champion GT, May MJ, Pidgeon JD. The UK farm scale evaluations of GM crops – a post script. *Outlooks on Pest Management. FEBS Lett.* 2005; 1-10. DOI: 10.1564/16aug00/Odeec53bf77cd564fb000000.
- [48] May M, Wilson RG. *Weeds and weed control*, In: Drycott AP, editor. *Sugar Beet*. Blackwell Publishing Ltd; 2006. p. 359-386.
- [49] Abdollahi F, Ghardiri H. Effect of separate and combined applications of herbicides on weed control and yield of sugar beet. *Weed Technol* 2004;18(4) 968-976.
- [50] Majidi M, Heidari G, Mohammadi K. Management of broad leaved weeds by combination of herbicides in sugar beet production. *Adv Environ Biol* 2011;5:10 3302-3306
- [51] Domaradzki K. Organiczenie nakładów na odchwaszczanie buraka cukrowego poprzez optymalizację dawkowania herbicydów w zabiegach systemowych. *Progr Plant Protect* 2009; 49(4):1790-1797.
- [52] Goleblowska H., Domaradzki K. Systemy chemicznej regulacji zachwaszczenia upraw rolniczych w aspekcie rolnictwa zrównoważonego. *Fragmenta Agronomica* 2010; 27(1):32-43.
- [53] Jursík M, Soukup J, Venclová V, Holec J. POST herbicide combinations for velvetleaf (*Abutilon theophrasti*) control in sugarbeet. *Weed Technol* 2011; 25(1):14-18.
- [54] Chitband AA, Ghorbani R, Mohassel Rashed MH, Abbaspoor M, Abbasi R. Evaluation of broadleaf weeds control with selectivity of post-emergence herbicides in sugar beet (*Beta vulgaris* L.). *Notulae Scientia Biologicae* 2014; 6(4):491-497.
- [55] Deveikyte I, Seibutis V. Effects of the phenmedipham, desmedipham, ethofumesate, Metamitron and Triflusulfuron-methyl on weeds and sugar beet. *Lucrari stiintifice Universitatea de stipinte agricole si medicina veterinara Ion Ionescu de la Brad. Seria Agronomia* 2008; 51: 278-286.
- [56] Deveikyte I. Optimising weed control in sugar beet. *Proc Scient Conf Int Particip* 25-26 September 2003; Slovenia. Nitra; 2003. p. 489.
- [57] Dexter AG. History of sugar beet (*Beta vulgaris* L.) herbicide rate reduction in North Dakota and Minnesota. *Weed Technol* 1994; 8(2):334-337.
- [58] Nurse RE, Hamill AS, Swanton CJ, Tardif FJ, Sikkema PH. Weed control and yields response to foramsulfuron in corn. *Weed Technol* 2007; 21(2):453-458.
- [59] Blackshaw RE, O'Donovan JT, Harker KN, Clayton GW, Stougaard RN. Reduced herbicide doses in field crops: a review. *Weed Biol Manag* 2006; 6(1):10-17.
- [60] Deveikyte I, Seibutis V. Broadleaf weeds and sugar beet response to phenmedipham, desmedipham, ethofumesate and triflusulfuron-methyl. *Agron Res* 2006; 4:159-162.
- [61] Zargar M, Rostami R. Response of broad leaf weeds to chemical and non-chemical management methods in sugar beet. *J Agric Environ Sci* 2011; 11(3):392-397.

- [62] Deveikyte I. Sensitivity of *Tripleurospermum perforatum* and *Chenopodium album* on low rates of phenmedipham, desmedipham, ethofumesate, metamiltron and chloridazon. *Lucrari stiintifice Universitatea de stiinte agricole si medicina veterinara Ion Ionescu de la Brad. Seria Agronomia* 2005; 48:386-204.
- [63] Alford CM, Nelson KK, Miller SD. Plant population, row spacing and herbicide effects on weeds and yield in sugar beets. *Int Sugar J* 2003; 105(1254):283-285.

IntechOpen

IntechOpen