

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Abiotic Stress Alleviation with Brassinosteroids in Plant Roots

Sevgi Marakli and Nermin Gozukirmizi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/61336>

Abstract

This chapter covers the advances in establishment and optimization of brassinosteroids (BRs) in the alleviation of abiotic stresses such as water, salinity, temperature, and heavy metals in plant system, especially roots. Plant roots regulate their developmental and physiological processes in response to various internal and external stimuli. Studies are in progress to improve plant root adaptations to stress factors. BRs are a group of steroidal hormones that play important roles in a wide range of developmental phenomena, and recently they became an alleviation agent for stress tolerance in plants. This review is expected to provide a resource for researchers interested in abiotic stress alleviation with BRs.

Keywords: Water stress, salt stress, temperature stress, heavy metal stress

1. Introduction

Abiotic stress responses in plants occur at various organ levels among which the root-specific processes are of particular importance. Under normal growth condition, root absorbs water and nutrients from the soil and supplies them throughout the plant body, thereby playing pivotal roles in maintaining cellular homeostasis. However, this balanced system is altered during the stress period when roots are forced to adopt several structural and functional modifications. Examples of these modifications include molecular, cellular, and phenotypic changes such as alteration of metabolism and membrane characteristics, hardening of cell wall, and reduction of root length [1, 2]. The root system has the crucial role of extracting nutrients and water through a complex interplay with soil biogeochemical properties and of maintaining these functions under a wide range of stress scenarios to ensure plant survival and reproduction [3].

Water stress is characterized by a reduction of water content and leaf water potential, closure of stomata, and decreased growth. Severe water stress may result in the arrest of photosynthesis, disturbance of metabolism, and finally the death of plant [4]. This water loss causes a loss of turgor pressure that may be accompanied by a decrease in cell volume depending on the hardness of the cell wall [5]. The cells of the root must activate processes to limit water loss and mitigate its harmful effects.

Salinity also affects plant growth, activity of major cytosolic enzymes by disturbing intracellular potassium homeostasis, causing oxidative stress and programmed cell death, reducing nutrient uptake, genetic and epigenetic effects, metabolic toxicity, inhibition of photosynthesis, decreasing CO₂ assimilation, and reducing root respiration [6, 7, 8, 9]. Salt stress affects the root in all developmental zones. Cell division decreases in the meristematic zone and cell expansion attenuates in the elongation zone, resulting in reduced overall growth [10]. Cells also expand radially in the elongation zone [11], and root hair outgrowth suppresses in the differentiation zone [12]. Salt stress additionally results in agravitropic growth [13] as well as reduced lateral root number under high-salt conditions and enhanced lateral root number under moderate-salt conditions [14, 15]. Salt stress develops from excessive concentrations of salt, especially sodium chloride (NaCl) in soil. Root is the primary organ of exposure and hence responds rapidly [16]. Salt stress is known to increase Na⁺/K⁺ ratio in the root that leads to cell dehydration and ion imbalance [17, 18, 19].

High temperature increases the permeability of plasma membrane [20], and also reduces water availability [21]. Moreover, low temperature (chilling and frost stress) is also a major limiting factor for productivity of plant indigenous to tropical and subtropical climates [22]. Chilling stress has a direct impact on the photosynthetic apparatus, essentially by disrupting the thylakoid electron transport, carbon reduction cycle, and stomatal control of CO₂ supply, together with an increased accumulation of sugars, peroxidation of lipids, and disturbance of water balance [23].

Heavy metal contamination in soil could result in inhibition of plant growth and yield reduction and even poses a great threat to human health via food chain [24]. Among heavy metals, Cadmium (Cd) in particular causes increasingly international concern [25]. Cd-contaminated soil results in considerable accumulation of Cd in edible parts of crops, and then it enters the food chain through the translocation and accumulation by plants [26, 27]. Another metal, chromium (Cr III or VI), is not required by plants for their normal plant metabolic activities. On the contrary, excess of Cr (III or VI) in agricultural soils causes oxidative stress for many crops. Reactive oxygen species (ROS), like hydrogen peroxide (H₂O₂), hydroxyl radical (OH[•]), and superoxide radical (O₂⁻) generated under Cr-stress, are highly reactive and cause oxidative damages to DNA, RNA, proteins, and pigments [28, 29]. Nickel (Ni) is one of the most abundant heavy metal contaminants of the environment due to its release from mining and smelting practices. It is classified as an essential element for plant growth [30]. However, at higher concentrations, nickel is an important environmental pollutant. Ni²⁺ ions bind to proteins and lipids such as specific subsequences of histones [31] and induce oxidative damage. Copper (Cu) is also an essential micronutrient for most biological organisms. It is a cofactor for a large array of proteins involved in diverse physiological processes, such as

photosynthesis, electron transport chain, respiration, cell wall metabolism, and hormone signaling [32, 33]. Cu has emerged as a major environmental pollutant in the past few decades because of its excessive use in manufacturing and agricultural industries [34]. Zinc (Zn) is one of the other essential microelement, the second most abundant transition metal, and plays roles in many metabolic reactions in plants [35, 36]. However, high concentrations of Zn are toxic, induce structural disorders, and cause functional impairment in plants. At organism level, Zn stress reduces rooting capacity, stunted growth, chlorosis, and at cellular level alters mitotic activity [37, 38].

The key to find out abiotic stress tolerance resides in understanding the plant's capacity to accelerate/maintain or repress growth. Most plant hormones play a role in development and have been implicated in abiotic stress responses. One of these hormones, BRs, are a group of steroidal hormones that play significant roles in a wide range of developmental phenomena including cell division and cell elongation in stems and roots, photo-morphogenesis, reproductive development, leaf senescence, and also in stress responses [39]. Mitchell et al. [40] discovered BRs which were later extracted from the pollen of *Brassica napus* by Grove et al. [41]. To date, more than 70 BR-related phytosteroids have been identified in plants [42].

BRs increase adaptation to various abiotic stresses such as light [43, 44], low or high temperature [45], drought [46, 47, 48], salt stress [9], and heavy metal stress [49, 50]. BRs may be applied/supplied to plants at different stages of their life cycle such as meiosis stage [51], anthesis stage [52], and root application [9, 53].

In this chapter, the potential role of BRs in alleviating the adverse effects of water, salt, low/high temperature stresses, and heavy metals on plants, especially roots, were discussed.

2. Water stress

Water shortage is predicted as one of the most important environmental problem for the 21st century that limits crop production [54]. Although drought stress inhibites the plant-water relations, exogenous application of BRs maintains tissue-water status [55] by stimulating the proton pumping [56], activating nucleic acid and protein synthesis [57] and regulation of genes expressions [58]. It has been shown that 24-epibrassinolide (24-epiBL)-treated *Arabidopsis* and *B. napus* seedlings had a higher survival rate when subjected to drought [59], and in another study BR-treated sorghum (*Sorghum vulgare*) showed increased germination and seedling growth under osmotic stress [60].

Root nodulation is a fundamental developmental event in leguminous crops, and is sensitive to water shortage [61, 62, 63]. As endogenous hormones play an important role in the organogenesis and initial growth of nodules in roots, attempts have been made to increase root nodulation by growth regulator treatments [64, 65]. The potential of BRs in the improvement of root nodulation and yield have been reported in groundnut [66]. Upreti and Murti [67] also studied the effects of two BRs, epibrassinolide (EBL) and homobrassinolide (HBL), on root nodulation and yield in *Phaseolus vulgaris* L. cv. Arka Suvidha under water stress. They

concluded that water stress negatively influenced nodulated root, but BRs increased tolerance to water stress and EBL was relatively more effective than HBL.

Several researchers have found that increased proline levels can protect plants from water stress. BR treatment increased the contents of proline and protein under water stress [68]. Zhang et al. [69] also indicated that BR treatment promoted the accumulation of osmoprotectants, such as soluble sugars and proline. It may be due to the fact that BRs activated the enzymes of proline biosynthesis, which caused an additive effect on the proline content [70].

Drought stress causes increment in H_2O_2 due to decrease in antioxidative enzyme activities [71]. Plants have improved various defense mechanisms to respond and adapt to water stress [72]. Vardhini et al. [73] studied with sorghum seedlings grown under PEG-imposed water stress and investigated the effects of HBL and 24-epiBL on the activities of four oxidizing enzymes: superoxide dismutase (SOD), glutathione reductase (GR), IAA oxidase, and polyphenol oxidase (PPO). They found that supplementation of both the BRs resulted in enhanced SOD and GR but lowered IAA oxidase and PPO. Li and Feng [68] also reported that treatment of brassinolide significantly increased peroxidase (POD), catalase (CAT), and ascorbate peroxidase (APX) activities of seedlings under normal water and mild water stress. Therefore, increment in enzyme activities provided tolerance of *Xanthoceras sorbifolia* seedlings to drought stress. It has been found that BRs can induce the expression of some antioxidant genes and enhance the activities of antioxidant enzymes such as SOD, POD, CAT, and APX [74, 75].

3. Salt stress

Salinity stress is one of the most serious abiotic stress factors. It causes morphological, biochemical, cytogenetic, and molecular changes in plants [9, 76, 77, 78]. Root lengths, shoot lengths, and root numbers decrease in plants exposed to salt stress [7]. Moreover, salinity also induces oxidative stress in plants due to production ROS [79, 80]. These ROS are produced in the cell and interacted with a number of vital cellular molecules and metabolites, thereby leading to a number of destructive processes causing cellular damage [81].

BRs reduce impacts of salt stress on ROS, gene expression, mitotic index, nutrient uptake, and growth [9, 82, 83–88]. There are lots of studies to analyse alleviation of salt stress by using BRs. In these studies, different parameters have been investigated to understand the mechanism of BRs on salt stress (Table 1).

References	[9]	[84]	[85]	[86]	[87]	[88]
Effects on gene expression	---	---	---	---	Increased <i>Cu/Zn-SOD</i> , <i>APX</i> , <i>CAT</i> , <i>GR</i> and <i>OsBRI</i> expressions	---

					but reduced <i>Fe-SOD</i> and <i>Mn-SOD</i> , <i>OsDWF4</i> and <i>SalT</i>	
Effects on protein content	Increased	---	Increased	---	Increased	---
Effects on enzyme activities	Increased SOD and CAT activities	Increased CAT, GR, POX and SOD activities	Increased POX and SOD activities	Increased SOD and POD activities	Shown varying results depending on 24-epiBL concentration for SOD, APX, CAT, GR	Increased CAT, POX and SOD activities
Effects on growth and/or cell division	Increased	Increased	Increased	Increased	Increased	Increased
Methods of salt and BRs applications	Seeds were grown under both 150–250 mM salt concentrations and 0.5 and 1 μM HBR at 48 h and 72 h.	Plants received 100 mM NaCl as well as 0.01 μM of HBL during 18 days after sowing	25, 50, 100, and 150 mM NaCl were applied and then sprayed twice with 0.05 ppm brassinolide during 25 days. At 45 days from sowing, the plants were collected	Seedlings were exposed to 90 mM NaCl with 0, 0.025, 0.05, 0.10, and 0.20 mg dm ⁻³ 24-epiBL for 10 days	Seeds were soaked for 8 h in different concentrations of 24-epiBL (10 ⁻¹¹ , 10 ⁻⁹ and 10 ⁻⁷ M). After 24-epiBL application, the seeds were sown in autoclaved sand moistened with different concentrations of NaCl (0, 75, 100, 125 mM) during 12 days	The 15-day-old plants were exposed to 100 mM NaCl and they were subsequently treated by exogenous 24-epiBL (10 ⁻⁸ M). The plants were harvested after 30 days of growth
Plant Species	<i>Hordeum vulgare</i> L.	<i>Vigna radiata</i> L.	<i>Vigna sinensis</i> L.	<i>Solanum melongena</i> L.	<i>Oryza sativa</i> L. var. Pusa Basmati-1 cv. indica	<i>Cucumis sativus</i> L.

Table 1. Effects of BRs on plants subjected to salt stress Dashes indicate that there are no results in study.

4. Temperature stress

4.1. High temperature

In general, a transient elevation in temperature (usually 10–15°C above environment) causes heat shock or heat stress [89]. High-temperature effects can be seen at the biochemical and molecular level in plant organs (especially leaves). Heat stress induces decrease in duration of developmental phases, leading to fewer organs, smaller organs, reduce light perception over the shortened life cycle, and finally play an important role in losing the product [90, 91, 92].

High-temperature stress often induces the overproduction of ROS [93] which can cause membrane lipid peroxidation, protein denaturation, and nucleic acid damage [94, 95]. Many studies have demonstrated that ROS scavenging mechanisms play an important role in protecting plants from high-temperature stress [96, 97]. BRs applications decrease ROS levels and increase antioxidant enzyme activities to provide thermotolerance to elevated temperatures [98].

4.2. Low temperature

Chilling and frost stresses affect growth, development, survival, and crop productivity in plants [99, 100, 101]. However, BRs treatments enhance seedling tolerance to chilling stress [101] and increase the height, root length, root biomass, and total biomass of rice under low-temperature conditions [102, 103]. In another study, Krishna [104] reported the same results in maize. They postulated that treatments with BRs promoted growth recovery of maize seedlings following chilling treatment (0–3°C).

Chilling stress increases the proline, betaine, soluble protein, soluble sugar contents of plants [79, 105]. Studies showed that BRs treatment enhanced proline content and therefore increased plant chilling resistance and cell membrane stability [99, 100, 106, 107].

Chilling stress could trigger the production of antioxidant enzymes in plants to prevent the chilling injury [108]. In the previous investigations, it was reported that treatment with BRs further increased the activities of antioxidant enzymes under chilling stress as well [99, 100, 107, 109]. The enhanced activities of the antioxidative enzymes as a result of BRs applications may occur with increasing de novo synthesis or activation of the enzymes, which is mediated through transcription and/or translation of specific genes to gain tolerance [57].

5. Heavy metal stresses

5.1. Cd stress

Cd toxicity has emerged as one of the major agricultural problems in many soils around the world [110]. It has been shown to interfere with the uptake, transport, and utilization of essential nutrients and water, change enzyme activities, cause symptoms (chlorosis, necrosis), decrease in fresh and dry mass of root and shoot and also their lengths [110, 111, 112].

There are lots of studies to investigate the effects of BRs on Cd stress in plant species [110, 113, 114]. In these studies, results showed that BRs change different parameters such as germination, plant dry biomass, protein content, and antioxidant enzyme activities (Table 2). It is proposed that the changes induced by BRs are mediated through the repression and/or de-repression of specific genes [58]. Microarray experiments evaluating gene expression changes in *Arabidopsis* roots and shoots under Cd stress were performed [115]. Moreover, studies showed that gene expression in response to Cd mimics a BR increase, and Cd exposure most probably induces an activation of the BR signaling pathway in *Arabidopsis* [116].

5.2. Cr stress

Cr (III or VI) is not required by plants for their normal plant metabolic activities [117]. The entry of Cr into a plant system occurs through roots via using the specialized uptake systems of essential metal ions required for normal plant metabolism [118]. On the contrary, excess of Cr (III or VI) in agricultural soils causes oxidative stress to many crops. Reduced seed germination, disturbed nutrient balance, wilting, and plasmolysis in root cells and thus effects on root growth of plants have been documented in plants under Cr stress [118, 119].

Choudhary et al. [120] reported that EBL treatment improved seedlings growth under Cr (VI) stress. Ability of EBL to increase seedling growth under this metal stress could be attributed to the capacity of BRs to regulate cell elongation and divisional activities, by enhancing the activity of cell wall loosening enzymes (xyloglucan transferase/hydrolase, XTH) [121]. Studies also indicated that increment in antioxidant activities as a result of BRs application (Table 2) provide plant tolerance to grow under Cr stress.

5.3. Ni stress

The heavy metals that affect (either positively or negatively) plants include Fe, Cu, Zn, Mn, Co, Ni, Pb, Cd, and Cr, but out of them, nickel has recently been defined as an essential micronutrient, because of its involvement in urease activity in legumes [122]. Excess Ni causes different problems. These symptoms include the inhibition in root elongation, photosynthesis and respiration, and interveinal chlorosis [123]. Moreover, the toxic concentration of Ni also inhibits enzyme activities and protein metabolism [124]. This metal also accelerates the activities of antioxidative enzymes [125, 126].

BRs effect on Ni stress in plants has been studied to understand the relationship between BRs and this stress (Table 2). One of these studies was carried out by Yusuf et al. [49]. They showed that seed germination and seedling growth were significantly reduced by Ni treatment, but HBL treatment enhanced germination percentage as well as shoot and root lengths in Ni-stressed seedlings. BRs confer tolerance against heavy metals either by reducing their uptake or by stimulating the antioxidative enzymes in *B. juncea* [127, 128]. The exogenous application of BRs in nickel-stressed *R. sativus* L., and *Triticum aestivum* L. plants enhanced the pool of antioxidant enzyme activity, thus alleviating the toxic effects of this stress [129].

5.4. Cu stress

Among the pollutants of agricultural soils, Cu has become increasingly hazardous due to its involvement in fungicides, fertilizers, and pesticides [130]. In addition, Cu present in excess has been known to decrease root biomass and alter plant metabolism [131, 132]. Sharma and Bhardwaj [127] demonstrated decrease in growth parameters of *Brassica juncea* grown under Cu stress. The reduction in growth parameters due to the Cu stress occurred as a result of decreasing mitotic activity and cell elongation [133, 134]. Moreover, Chen et al. [130] suggested a different opinion. They concluded that Cu-induced inhibition in root growth of rice seedlings was due to the stiffening of the cell wall. Moreover, excess of Cu ion leads to the generation of harmful ROS via the formation of free radicals [135].

Effects of exogenous application of BRs were studied on *Raphanus sativus* seedlings under Cu stress. It was found that 24-epiBL promoted the shoot and root growth by overcoming the Cu toxicity [136]. The growth-promoting effects of BRs on seedlings under Cu stress may be linked to the general ability of BRs to promote cell elongation and cell cycle progression [137, 138] as well as the stimulation of genes encoding xyloglucanases and expansins [139]. BRs applications also increase antioxidant enzyme activities [140, 141]. Increasing all parameters as a result of BRs application improves plant tolerance against Cu stress, and finally plant development (Table 2).

5.5. Zn stress

Zn is an essential microelement, the second most abundant transition metal after iron (Fe), and has a role in many metabolic reactions in plants [35, 36]. However, high concentrations of Zn are toxic, induce structural disorders, and cause functional problems in plants. At organism level, Zn stress causes reduced rooting capacity, growth, and at cellular level alters mitotic activity [37, 38]. It induces oxidative stress by promoting ROS production as a result of indirect consequence of Zn toxicity [142].

Application of BRs on plants alleviates Zn stress via increasing protein content and antioxidant enzyme activities (Table 2). Çağ et al. [143] reported that EBL application effectively enhanced the protein content in *Brassica oleracea* cotyledons. Sharma et al. [144] also reported that pre-sowing treatments of HBL lowered the uptake of metal and enhanced the activities of antioxidant enzymes and protein concentration of *B. juncea* seedlings under Zn stress. Moreover, Ramakrishna and Rao [145] also reported that the application of 24-epiBL significantly alleviated the Zn-induced oxidative stress.

References	[114]	[146]	[30]	[147]	[141]	[145]
Effects on gene expression	---	---	---	---	---	---
Effects on protein content	---	Increased	Decreased protein content	---	Showed varying results	---

Effects on enzyme activities	Increased CAT, POX and SOD activities	Decreased CAT level	Decreased SOD but increased POD activities	Increased SOD, CAT and POX activities	Showed varying results	Increased
Effects on growth and/or cell division	Alleviated	Improved	Improved	Improved	Increased	Increased
Methods of metal and BRs applications	Soil amended with 50 µM Cd and foliage sprayed with 10 ⁻⁸ M HBL at 20 days after sowing. The plants were sampled at 30 days after sowing	Seeds were treated with eEBL (10 ⁻⁹ M) and 1.2 mM Cr(VI) (K ₂ CrO ₄) solution at 7 days	Seeds were soaked for 8 h in different concentrations of 24-epiBL (0, 10 ⁻⁷ , 10 ⁻⁹ and 10 ⁻¹¹ M). Then, seeds were grown under different (0, 0.5, 1.0, 1.5, and 2.0 mM) Ni concentrations	Seeds were grown in different levels of Cu ²⁺ (50 or 100 mg kg ⁻¹ of soil. At 15 and 20 days stage, 10 ⁻⁵ mM HBL was applied. At 45 days, plants were collected	Seeds were grown under both 30 µM – 40 µM Cu and 2 µM HBR at 48 h and 72 h	EBR (0.5, 1, and 2 µM) and 5 mM of Zn were applied to seeds. Seven day old seedlings were collected
Plant Species	<i>Triticum aestivum</i> L.	<i>Raphanus sativus</i> L.	<i>B. juncea</i> L.	<i>Vigna radiata</i> L.	<i>Helianthus annuus</i> L.	<i>Raphanus sativus</i> L.
	Cd Stress	Cr Stress	Ni Stress	Cu Stress		Zn Stress

Table 2. Effects of BRs on plants subjected to Cd, Cr, Ni, Cu and Zn stresses. Dashes indicate that there are no results in study

6. Conclusion

Roots are very important plant organs whose architecture is determined by endogenous and environmental conditions to adjust water and nutrient uptake from soil [148, 149]. BRs, one of the plant hormones, have both positive and negative effects on root growth related to hormone concentrations [150]. Experimental condition is one of the most important factors for analysing BRs effects on root development. The procedures using BRs to alleviate abiotic systems generally are easy, time saving, and one of the most reliable systems [53]. Therefore, BRs open up new approaches for plant tolerance against hazardous environmental conditions [151]. Morphological, biochemical, and molecular analyses have been performed to analyse the effects of BRs. However, detailed analyses should be performed to investigate the relationship between abiotic stresses and BRs, especially gene expression studies will provide knowledge about interaction at molecular level in plants [152]. We tried to cite as many papers as possible. Yet we apologize to authors whose works are gone unmentioned in this chapter.

Acknowledgements

We are grateful to the Research Fund of Istanbul University for financial support (Projects 39824, 43403, and UDP54481).

Author details

Sevgi Marakli* and Nermin Gozukirmizi

*Address all correspondence to: sevgi.marakli@istanbul.edu.tr

Istanbul University, Faculty of Science, Department of Molecular Biology and Genetics,
Vezneciler, Istanbul, Turkey

References

- [1] Gowda VRP, Henry A, Yamauchi A, Shashidhar HE, Serraj R. Root biology and genetic improvement for drought avoidance in rice. *Field Crops Res* 2011;122:1–13. DOI: 10.1016/j.fcr.2011.03.001
- [2] Atkinson NJ, Urwin PE. The interaction of plant biotic and abiotic stresses: from genes to the field. *J Exp Bot* 2012; 63:3523–3543. DOI: 10.1093/jxb/ers100
- [3] Ahmadi N, Audebert A, Bennett MJ. The roots of future rice harvests. *Rice* 2014;7:29. DOI: 10.1186/s12284-014-0029-y
- [4] Jaleel CA, Manivannan P, Lakshmanan GMA, Gomathinayagam M, Panneerselvam R. Alterations in morphological parameters and photosynthetic pigment responses of *Catharanthus roseus* under soil water deficits. *Colloids Surfaces B: Biointerf* 2008;61:298–303. DOI: 10.1016/j.colsurfb.2007.09.008
- [5] Verslues PE, Agarwal M, Katiyar-Agarwal S, Zhu J, Zhu, JK. Methods and concepts in quantifying resistance to drought, salt and freezing, abiotic stresses that affect plant water status. *Plant J* 2006;45:523–39. DOI: 10.1111/j.1365-313X.2005.02593.x
- [6] Abogadallah GM. Antioxidative defense under salt Stress. *Plant Signal Behavior* 2010;5:369-374. DOI: 10.4161/psb.5.4.10873
- [7] Demirkiran A, Marakli S, Temel A, Gozukirmizi N. Genetic and epigenetic effects of salinity on *in vitro* growth of barley. *Genet Molecul Biol* 2013;36(4):566–70. DOI: 10.1590/S1415-47572013000400016
- [8] Liu J, Gao H, Wang X, Zheng Q, Wang C, Wang X, Wang Q. Effects of 24-epibrassinolide on plant growth, osmotic regulation and ion homeostasis of salt stressed canola. *Plant Biol* 2014;16(2):440–50. DOI: 10.1111/plb.12052
- [9] Marakli S, Temel A, Gozukirmizi N. Salt stress and homobrassinosteroid interactions during germination in barley roots. *Not Bot Horti Agrobo* 2014;42:446–52. DOI: 10.15835/nbha4229461

- [10] West G, Inze D, Beemster GTS. Cell cycle modulation in the response of the primary root of *Arabidopsis* to salt stress. *Plant Physiol* 2004;135:1050–8. DOI: 10.1104/pp.104.040022
- [11] BursSENS S, Himanen K, van de Cotte B, Beeckman T, Van Montagu M, Inze D, Verbruggen N. Expression of cell cycle regulatory genes and morphological alterations in response to salt stress in *Arabidopsis thaliana*. *Planta* 2000;211:632–40. DOI: 10.1007/s004250000334
- [12] Halperin SJ, Gilroy S, Lynch JP. Sodium chloride reduces growth and cytosolic calcium, but does not affect cytosolic pH, in root hairs of *Arabidopsis thaliana* L. *J Exp Bot* 2003;54:1269–80. DOI: 10.1093/jxb/erg134
- [13] Sun F, Zhang W, Hu H, Li B, Wang Y, Zhao Y, Li K, Liu M, Li X. Salt modulates gravity signaling pathway to regulate growth direction of primary roots in *Arabidopsis*. *Plant Physiol* 2008;146:178–188. DOI: 10.1104/pp.107.109413
- [14] Wang Y, Li K, Li X. Auxin redistribution modulates plastic development of root system architecture under salt stress in *Arabidopsis thaliana*. *J Plant Physiol* 2009;166:1637–45. DOI: doi:10.1016/j.jplph.2009.04.009
- [15] Zolla G, Heimer YM, Barak S. Mild salinity stimulates a stress-induced morphogenic response in *Arabidopsis thaliana* roots. *J Exp Bot* 2010;61:211–224. DOI: 10.1093/jxb/erp290
- [16] Ghosh D, Xu J. Abiotic stress responses in plant roots: a proteomics perspective, *Front Plant Sci* 2014;5:6. DOI: 10.3389/fpls.2014.00006
- [17] Tester M, Davenport R. Na⁺ tolerance and Na⁺ transport in higher plants. *Ann Bot* 2003;91:503–27. DOI: 10.1093/aob/mcg058
- [18] Cavalcanti FR, Santos Lima JPM, Ferreira-Silva SL, Viegas RA, Silveira JAG, Roots and leaves display contrasting oxidative response during salt stress and recovery in cowpea. *J Plant Physiol* 2007;164:591–600. DOI: 10.1016/j.jplph.2006.03.004
- [19] Munns R, Tester M. Mechanisms of salinity tolerance. *Ann Rev Plant Biol* 2008;59:651–81. DOI: 10.1146/annurev.arplant.59.032607.092911
- [20] Zhang JH, Huang WD, Liu YP, Pan QH. Effects of temperature acclimation pretreatment on the ultrastructure of mesophyll cells in young grape plants (*Vitis vinifera* L. cv. Jingxiu) under cross-temperature stresses. *J Integrat Plant Biol* 2005;47:959–70. DOI: 10.1111/j.1744-7909.2005.00109.x
- [21] Simoes-Araujo JL, Rumjanek NG, Margis-Pinheiro M. Small heat shock proteins genes are differentially expressed in distinct varieties of common bean. *Brazil J Plant Physiol* 2003;15:33–41. DOI: 10.1590/S1677-04202003000100005

- [22] Salveit ME. Chilling injury is reduced in cucumber and rice seedlings in tomato pericarp discs by heat-shocks applied after chilling. *Postharvest Biol Technol* 2001;21:169–77. DOI: doi:10.1016/S0925-5214(00)00132-0
- [23] Allen DJ, Ort DR. Impact of chilling temperatures on photosynthesis in warm-climate plants. *Trends Plant Sci* 2001;6:36–42. DOI: doi:10.1016/S1360-1385(00)01808-2
- [24] Lux A, Martinka M, Vaculik M, White PJ. Root responses to cadmium in the rhizosphere: a review. *J Exp Bot* 2011;62:21–37. DOI: 10.1093/jxb/erq281
- [25] Mulligan CN, Yong RN, Gibbs BF. Remediation technologies for metal-contaminated soils and groundwater: an evaluation. *Eng Geol* 2001;60:193–207. DOI: doi:10.1016/S0013-7952(00)00101-0
- [26] Uraguchi S, Mori S, Kuramata M, Kawasaki A, Arai T, Ishikawa S. Root-to-shoot Cd translocation via the xylem is the major process determining shoot and grain cadmium accumulation in rice. *J Exp Bot* 2009;60:2677–88. DOI: 10.1093/jxb/erp119
- [27] Cai Y, Cao F, Wei K, Zhang G, Wu F. Genotypic dependent effect of exogenous glutathione on Cd-induced changes in proteins, ultrastructure and antioxidant defense enzymes in rice seedlings. *J Hazard Mater* 2011;192:1056–66. DOI: 10.1016/j.jhazmat.2011.06.011
- [28] Ali S, Bai P, Zeng F, Cai S, Shamsi IH, Qui B, Wu F, Zhang G. The ecotoxicological and interactive effects of chromium and aluminum on growth, oxidative damage and antioxidant enzymes on two barley genotypes differing in Al tolerance. *Environ Exp Bot* 2011;70:185–91. DOI: 10.1016/j.envexpbot.2010.09.002
- [29] Sharma I, Pati PK, Bhardwaj R. Effect of 28-homobrassinolide on antioxidant defence system in *Raphanus sativus* L. under chromium toxicity. *Ecotoxicol* 2011;20:862–74. DOI: 10.1007/s10646-011-0650-0
- [30] Kanwar MK, Bhardwaj R, Chowdhary SP, Arora P, Sharma P, Kumar S. Isolation and characterization of 24-Epibrassinolide from *Brassica juncea* L. and its effects on growth, Ni ion uptake, antioxidant defense of *Brassica* plants and in vitro cytotoxicity. *Acta Physiol Plant* 2013;35:1351–62. DOI: 10.1007/s11738-012-1175-8
- [31] Bal W, Kasprzak KS. Induction of oxidative damage by carcinogenic metal. *Toxicol Lett* 2002;127:55–62. DOI: doi:10.1016/S0378-4274(01)00483-0
- [32] Bhakuni G, Dube B.K, Sinha P, Chatterjee C. Copper stress affects metabolism and reproductive yield of chickpea. *J Plant Nutri* 2009;32:703–11. DOI: 10.1080/01904160902743258
- [33] Andre CM, Larondelle Y, Evers D. Dietary antioxidants and oxidative stress from a human and plant perspective: a review. *Curr Nutri Food Sci* 2010;6:2–12. DOI: 10.2174/157340110790909563
- [34] Bouazizi H, Jouili H, Geitmann A, El Ferjani E. Cell wall accumulation of Cu ions and modulation of lignifying enzymes in primary leaves of bean seedlings exposed

- to excess copper. *Biol Trace Element Res* 2011;139:97–107. DOI: 10.1007/s12011-010-8642-0
- [35] Gayor A, Srivastava PS, Iqbal M. Morphogenic and biochemical responses of *Bacopa monniera* cultures to zinc toxicity. *Plant Sci* 1999;143:187–93. DOI: 10.1016/S0168-9452(99)00032-1
- [36] Vaillant N, Monnet F, Hitmi A, Sallanon H, Coudret A. Comparative study of responses in four datura species to zinc stress. *Chemosphere* 2005;59:1005–13. DOI: 10.1016/j.chemosphere.2004.11.030
- [37] Castiglione S, Franchin C, Fossat T, Lingua G, Torrigiani P, Biondi S. High zinc concentrations reduce rooting capacity and alter metallothionein gene expression in white poplar (*Populus alba* L. cv. Villafranca). *Chemosphere* 2007;67:1117–26. DOI: 10.1016/j.chemosphere.2006.11.039
- [38] Tewari RK, Kumar P, Sharma PN. Morphology and physiology of zinc-stressed mulberry plants. *J Plant Nutri Soil Sci* 2008;171:286–94. DOI: 10.1002/jpln.200700222
- [39] Choudhary SP, Yu JQ, Yamaguchi-Shinozaki K, Shinozaki K, Phan Tran L-S. Benefits of brassinosteroid crosstalk. *Trends Plant Sci* 2012;17:594–605. DOI: 10.1016/j.tplants.2012.05.012
- [40] Mitchell JW, Mandava NB, Worley JE, Plimmer JR, Smith MV. Brassins: a family of plant hormones from rape pollen. *Nature* 1970;225:1065–66. DOI: 10.1038/2251065a0
- [41] Grove MD, Spencer GF, Rohwededer WK, Mandava N, Worley JF, Warthen JR JD, Steffens GL, Flippen-Anderson JL, Cook JR JC. Brassinolide, a plant promoting steroid isolated from *Brassica napus* pollen. *Nature* 1979;281:216–7. DOI: 10.1038/281216a0
- [42] Zhao B, Li J. Regulation of brassinosteroid biosynthesis and inactivation. *J Integrat Plant Biol* 2012;54:746–59. DOI: 10.1111/j.1744-7909.2012.01168.x.
- [43] Wang M, Jiang WJ, Yu HJ. Effects of exogenous epibrassinolide on photosynthetic characteristics in tomato (*Lycopersicon esculentum* Mill) seedlings under weak light stress. *J Agricult Food Chem* 2010;58:3642–5. DOI: 10.1021/jf9033893
- [44] Kurepin LV, Joo SH, Kim S-K, Pharis RP, Back TG. (2012). Interaction of brassinosteroids with light quality and plant hormones in regulating shoot growth of young sunflower and *Arabidopsis* seedlings. *J Plant Growth Regulat* 2012;31:156–64. DOI: 10.1007/s00344-011-9227-7
- [45] Wang Q, Ding T, Gao L, Pang J, Yang N. Effect of brassinolide on chilling injury of green bell pepper in storage. *Sci Horticult* 2012;144:195–200. DOI: 10.1016/j.scienta.2012.07.018

- [46] Anjum SA, Wang LC, Farooq M, Hussain M, Xue LL, Zou CM. Brassinolide application improves the drought tolerance in maize through modulation of enzymatic antioxidants and leaf gas exchange. *J Agronomy Crop Sci* 2011;197:177–85.
- [47] Li YH, Liu YJ, Xu XL, Jin M, An LZ, Zhang H, Effect of 24-epibrassinolide on drought stress-induced changes in *Chorispora bungeana*. *Biol Plant* 2012;56:192–6. DOI: 10.1007/s10535-012-0041-2
- [48] Mahesh B, Parshavaneni B, Ramakrishna B, Rao SSR. Effect of brassinosteroids on germination and seedling growth of radish (*Raphanus sativus* L.) under PEG-6000 induced water stress. *Am J Plant Sci* 2013;4:2305–13. DOI: 10.4236/ajps.2013.412285
- [49] Yusuf M, Fariduddin Q, Hayat S, Hasan SA, Ahmad A. Protective responses of 28-Homobrassinolide in cultivars of *Triticum aestivum* with different levels of nickel. *Arch Environ Contam Toxicol* 2011;60:68–76. DOI: 10.1007/s00244-010-9535-0
- [50] Yusuf M, Fariduddin Q, Ahmad A. 24-Epibrassinolide modulates growth, nodulation, antioxidant system, and osmolyte in tolerant and sensitive varieties of *Vigna radiata* under different levels of nickel: a shotgun approach. *Plant Physiol Biochem* 2012;57:143–53. DOI: 10.1016/j.plaphy.2012.05.004
- [51] Saka H, Fujii S, Imakawa A.M, Kato N, Watanabe S, Nishizawa T, Yonekawa S. Effect of brassinolide applied at the meiosis and flowering stages on the levels of endogenous plant hormones during grain filling in rice plant (*Oryza sativa* L.). *Plant Prod Sci* 2003;6:36–42. DOI: 10.1626/ppp.6.36
- [52] Liu H, Guo T, Zhu Y, Wang C, Kang G. Effects of Epibrassinolide (epi-BR) application at anthesis on starch accumulation and activities of key enzymes in wheat grains. *Acta Agronom Sinica* 2006;32:924–30.
- [53] Kartal G, Temel A, Arican E, Gozukirmizi N. Effects of brassinosteroids on barley root growth, antioxidant system and cell division. *Plant Growth Regulat* 2009;58:261–7. DOI: 10.1007/s10725-009-9374-z
- [54] Yuan G-F, Jia C-G, Li Z, Sun B, Zhang L-P, Liu N, Wang Q-M. Effect of brassinosteroids on drought resistance and abscisic acid concentration in tomato under water stress. *Sci Horticult* 2010;126:103–8. DOI: 10.1016/j.scienta.2010.06.014
- [55] Farooq M, Wahid A, Basra SMA, Islam-ud-Din. Improving water relations and gas exchange with brassinosteroids in rice under drought stress. *J Agron Crop Sci* 2009;195:262–9. DOI: 10.1111/j.1439-037X.2009.00368.x
- [56] Khripach VA, Zhabinski VN, Khripach NB. New practical aspects of brassinosteroids and results of their ten year agricultural use in Russia and blakanes. In: Hayat S, Ahmad A, editors. *Brassinosteroids; Bioactivity and Crop Productivity*. Kluwer Academic Publisher, Dordrecht; 2003. pp. 189–230.

- [57] Bajguz A. Effect of brassinosteroids on nucleic acid and protein content in cultured cells of *Chlorella vulgaris*. *Plant Physiol Biochem* 2000;38:209–15. DOI: 10.1016/S0981-9428(00)00733-6
- [58] Felner M. Recent progress in brassinosteroid research: hormone perception and signal transduction. In: Hayat S, Ahmad A, editors. *Brassinosteroids: Bioactivity and Crop Productivity*. Kluwer Academic Publishers, Dordrecht; 2003. pp. 69–86.
- [59] Kagale S, Divi UK, Krochko JE, Keller WA, Krishna P. Brassinosteroid confers tolerance in *Arabidopsis thaliana* and *Brassica napus* to a range of abiotic stresses. *Planta* 2007;225:353–64. DOI: 10.1007/s00425-006-0361-6
- [60] Vardhini BV, Rao SSR. Amelioration of osmotic stress by brassinosteroids on seed germination and seedling growth of three varieties of sorghum. *Plant Growth Regulat* 2003;41:25–31. DOI: 10.1023/A:1027303518467
- [61] Bordeleau LM, Prevost D. Nodulation and nitrogen fixation in extreme environments. *Plant Soil* 1994;161:115–25. DOI: 10.1007/BF02183092
- [62] Ramos MLG, Gordon AJ, Minchin FR, Sprent JI, Parsons R. Effect of water stress on nodule physiology and biochemistry of a drought tolerant cultivar of common bean (*Phaseolus vulgaris* L.). *Annals Bot* 1999;83:57–63. DOI: 10.1006/anbo.1998.0792
- [63] Serraj R, Sinclair TR, Purcell LC. Symbiotic N₂ fixation response to drought. *J Exp Bot* 1999;50:143–55. DOI: 10.1093/jxb/50.331.143
- [64] Singh T, Kumar V. Nodulation and plant growth as influenced by growth regulators in some legumes. *Acta Bot Ind* 1989;17:177–81.
- [65] Fedorova EE, Zhiznevskaya GY, Al'zhapparova ZK, Izmailov SF. Phytohormones in nitrogen-fixing nodules of leguminous plants. *Fiziologiya I Biokhimiya Kul'Turnykh Rastenii*. 1991;23:426–38.
- [66] Vardhini BV, Rao SSR. Effect of brassinosteroids on nodulation and nitrogenase activity in groundnut (*Arachis hypogaea* L.). *Plant Growth Regulat* 1999;28:165–7. DOI: 10.1023/A:1006227417688
- [67] Upreti KK, Murti GSR. Effects of brassinosteroids on growth, nodulation, phytohormone content and nitrogenase activity in French bean under water stress. *Biologia Plant* 2004;48:407–11. DOI: 10.1023/B:BIOP.0000041094.13342.1b
- [68] Li KR, Feng CH. Effects of brassinolide on drought resistance of *Xanthoceras sorbifolia* seedlings under water stress. *Acta Physiol Plant* 2011;33:1293–300. DOI: 10.1007/s11738-010-0661-0
- [69] Zhang M, Zhai Z, Tian X, Duan L, Li Z. Brassinolide alleviated the adverse effect of water deficits on photosynthesis and the antioxidant of soybean (*Glycine max* L.). *Plant Growth Regulat* 2008;56:257–64. DOI: 10.1007/s10725-008-9305-4
- [70] Fariduddin Q, Khanam S, Hasan SA, Ali B, Hayat S, Ahmad A. Effect of 28-homobrassinolide on the drought stress-induced changes in photosynthesis and antioxi-

- dant system of *Brassica juncea* L. *Acta Physiol Plant* 2009;31:889–97. DOI: 10.1007/s11738-009-0302-7
- [71] Reddy AR, Chaitanya KV, Vivekanandan M. Drought induced responses of photosynthesis and antioxidant metabolism in higher plants. *J Plant Physiol* 2004;161:1189–1202. DOI:10.1016/j.jplph.2004.01.013
- [72] Xiong L, Schumaker KS, Zhu JK. Cell signaling during cold, drought, and salt stress. *Plant Cell* 2002;14:S165–83. DOI: 10.1105/tpc.000596
- [73] Vardhini BV, Sujatha E, Rao SSR. Brassinosteroids: alleviation of water stress in certain enzymes of sorghum seedlings. *J Phytol* 2011;3(10):38–43.
- [74] Mazorra LM, Nunez M, Hechavarria M, Coll F, Sanchez-Blanco MJ. Influence of brassinosteroids on antioxidant enzymes activity in tomato under different temperatures. *Biol Plant* 2002;45:593–6. DOI: 10.1023/A:1022390917656
- [75] Cao S, Xu Q, Cao Y, Qian K, An K, Zhu Y, Binzeng, H, Zhao H, Kuai B. Loss of function mutation in *DET2* gene lead to an enhanced resistance to oxidative stress in *Arabidopsis*. *Physiol Plant* 2005;123:57–66. DOI: 10.1111/j.1399-3054.2004.00432.x
- [76] Eraslan F, Inal A, Gunes A, Alpaslan A. Impact of exogenous salicylic acid on the growth, antioxidant activity and physiology of carrot plants subjected to combined salinity and boron toxicity. *Sci Horticult* 2007;113:120–8. DOI: 10.1016/j.scienta.2007.03.012
- [77] Tuteja N. Mechanisms of high salinity tolerance in plants. *Methods Enzymol* 2007;428:419–38. DOI: 10.1016/S0076-6879(07)28024-3
- [78] Munns R, Tester M. Mechanisms of salinity tolerance. *Ann Rev Plant Biol* 2008;59:651–81. DOI: 10.1146/annurev.arplant.59.032607.092911
- [79] Ashraf M, Foolad MR. Improving plant abiotic-stress resistance by exogenous application of osmoprotectants glycinebetaine and proline. *Environ Exp Bot* 2007;59:206–16.
- [80] Daneshmand F, Arvin MJ, Kalantari KM. Acetylsalicylic acid ameliorates negative effects of NaCl or osmotic stress in *Solanum stoloniferum* in vitro. *Biol Plant* 2010;54:781–4. DOI: 10.1007/s10535-010-0142-8
- [81] Ashraf M. Biotechnological approach of improving plant salt tolerance using antioxidants as markers. *Biotechnol Adv.* 2009;27:84–93. DOI: 10.1016/j.biotechadv.2008.09.003
- [82] Talaat NB, Shawky BT. 24-Epibrassinolide alleviates salt-induced inhibition of productivity by increasing nutrients and compatible solutes accumulation and enhancing antioxidant system in wheat (*Triticum aestivum* L.). *Acta Physiol Plant* 2013;35:729–40. DOI: 10.1007/s11738-012-1113-9

- [83] Divi UK, Rahman T, Krishna P. Gene expression and functional analyses in brassinosteroid-mediated stress tolerance. *Plant Biotechnol J* 2015, DOI: 10.1111/pbi.12396.
- [84] Hayat S, Hasan SA, Yusuf M, Hayat Q, Ahmad A. Effect of 28-homobrassinolide on photosynthesis, fluorescence and antioxidant system in the presence or absence of salinity and temperature in *Vigna radiata*. *Environ Exp Bot* 2010;69:105–12. DOI: 10.1016/j.envexpbot.2010.03.004
- [85] El-Mashad AAA, Mohamed HI. Brassinolide alleviates salt stress and increases antioxidant activity of cowpea plants (*Vigna sinensis*). *Protoplasma* 2012;249:625–35. DOI: 10.1007/s00709-011-0300-7
- [86] Ding H-D, Zhu X-H, Zhu Z-W, Yang S-J, Zha D-S, Wu X-X. Amelioration of salt-induced oxidative stress in eggplant by application of 24-epibrassinolide. *Biol Plant* 2012;56(4):767–70. DOI: 10.1007/s10535-012-0108-0
- [87] Sharma I, Ching E, Saini S, Bhardwaj R, Pati PK. Exogenous application of brassinosteroid offers tolerance to salinity by altering stress responses in rice variety Pusa Basmati-1. *Plant Physiol Biochem* 2013;69:17–26. DOI: 10.1016/j.plaphy.2013.04.013
- [88] Fariduddin Q, Mir BA, Yusuf M, Ahmad A. 24-epibrassinolide and/or putrescine trigger physiological and biochemical responses for the salt stress mitigation in *Cucumis sativus* L. *Photosynthetica* 2014;52(3):464–74. DOI: 10.1007/s11099-014-0052-7
- [89] Bajguz A. Brassinosteroid enhanced the level of abscisic acid in *Chlorella vulgaris* subjected to short-term heat stress. *J Plant Physiol* 2009;166:882–6. DOI: 10.1016/j.jplph.2008.10.004
- [90] Stone P. The effects of heat stress on cereal yield and quality. In: Basra AS. (Ed.) *Crop Responses and Adaptations to Temperature Stress*. Binghamton, NY: Food Products Press; 2001. p. 302.
- [91] Rane J, Chauhan H. Rate of grain growth in advanced wheat (*Triticum aestivum*) accession under late-sown environment. *Ind J Agricult Sci* 2002;72:581–5.
- [92] Hussain SS, Mudasser M. Prospects for wheat production under changing climate in mountain areas of Pakistan: an econometric analysis. *Agricult Sys* 2007;94:494–501. DOI: 10.1016/j.agry.2006.12.001
- [93] Janeczko A, Oklestkova J, Pocięcha E, Koscielniak J, Mirek M. Physiological effects and transport of 24-epibrassinolide in heat-stressed barley. *Acta Physiol Plant* 2011;33:1249–59. DOI: 10.1007/s11738-010-0655-y
- [94] Yin H, Chen QM, Yi MF. Effects of short-term heat stress on oxidative damage and responses of antioxidant system in *Lilium longiflorum*. *Plant Growth Regulat* 2008;54:45–54. DOI: 10.1007/s10725-007-9227-6

- [95] Bartwal A, Mall R, Lohani P, Guru SK, Arora S. Role of secondary metabolites and brassinosteroids in plant defense against environmental stresses. *J Plant Growth Regulat* 2012;32:216–32. DOI: 10.1007/s00344-012-9272-x
- [96] Hu WH, Xiao YA, Zeng JJ, Hu XH. Photosynthesis, respiration and antioxidant enzymes in pepper leaves under drought and heat stresses. *Biol Plant* 2010;54:761–5. DOI: 10.1007/s10535-010-0137-5
- [97] Asthir B, Koundal A, Bains NS. Putrescine modulates antioxidant defense response in wheat under high temperature stress. *Biol Plant* 2012;56:757–61. DOI: 10.1007/s10535-012-0209-1.
- [98] Wu X, Yao X, Chen J, Zhu Z, Zhang H, Zha D. Brassinosteroids protect photosynthesis and antioxidant system of eggplant seedlings from high-temperature stress. *Acta Physiol Plant* 2014;36:251–61. DOI: 10.1007/s11738-013-1406-7
- [99] Liu Y, Zhao Z, Si J, Di C, Han J, An L. Brassinosteroids alleviate chilling-induced oxidative damage by enhancing antioxidant defense system in suspension cultured cells of *Chorispora bungeana*. *Plant Growth Regulat* 2009;59:207–14. DOI: 10.1007/s10725-009-9405-9
- [100] Liu YJ, Jiang HF, Zhao ZG, An LZ. Abscisic acid is involved in brassinosteroids-induced chilling tolerance in the suspension cultured cells from *Chorispora bungeana*. *J Plant Physiol* 2011;168:853–62. DOI: 10.1016/j.jplph.2010.09.020
- [101] Wang B, Zeng G. Effect of epibrassinolide on the resistance of rice seedlings to chilling injury. *J Plant Physiol Molecul Biol* 1993;19:38–42.
- [102] Kim KS, Sa JG. Effects of plant growth regulator, brassinolide, on seedling growth in rice (*Oryza sativa* L). *Res Rep Rural Develop Admin Rice* 1989;31(1):49–53.
- [103] Hirai K, Fujii S, Honjo K. The effect of brassinolide on ripening of rice plants under low temperature condition. *Japan J Crop Sci* 1991;60(1):29–35.
- [104] Krishna P. Brassinosteroid-mediated stress responses. *J Plant Growth Regulat* 2003;22:289–97. DOI: 10.1007/s00344-003-0058-z
- [105] Burbulis N, Jonytiene V, Kupriene R, Blinstrubiene A. Changes in proline and soluble sugars content during cold acclimation of winter rapeseed shoots in vitro. *J Food, Agricult Environ* 2011;9:371–4.
- [106] Hu WH, Wu Y, Zeng JZ, He L, Zeng QM. Chill-induced inhibition of photosynthesis was alleviated by 24-epibrassinolide pretreatment in cucumber during chilling and subsequent recovery. *Photosynthetica* 2010;48:537–44. DOI: 10.1007/s11099-010-0071-y
- [107] Fariduddin Q, Yusuf M, Chalkoo S, Hayat S, Ahmad A. 28-homobrassinolide improves growth and photosynthesis in *Cucumis sativus* L. through an enhanced antiox-

- idant system in the presence of chilling stress. *Photosynthetica* 2011;49(1):55–64. DOI: 10.1007/s11099-011-0022-2
- [108] Oidaira H, Sano S, Koshiha T, Ushimaru T. Enhancement of antioxidative enzyme activities in chilled rice seedlings. *J Plant Physiol* 2000;156:811–3. DOI: 10.1016/S0176-1617(00)80254-0
- [109] Ahammed GJ, Yuan HL, Ogweno JO, Zhou YH, Xia XJ, Mao WH, Shi K, Yu JQ. Brassinosteroid alleviates phenanthrene and pyrene phytotoxicity by increasing detoxification activity and photosynthesis in tomato. *Chemosphere* 2012;86:546–55. DOI: 10.1016/j.chemosphere.2011.10.038
- [110] Hasan SA, Hayat S, Ahmad A. Brassinosteroids protect photosynthetic machinery against the cadmium induced oxidative stress in two tomato cultivars. *Chemosphere* 2011;84:1446–51. DOI: 10.1016/j.chemosphere.2011.04.047
- [111] López-Millán AF, Sagardoy R, Solanas M, Abadía A, Abadía J. Cadmium toxicity in tomato (*Lycopersicon esculentum*) plants grown in hydroponics. *Environ Exp Bot* 2009;65:376–85. DOI: 10.1016/j.envexpbot.2008.11.010
- [112] Hayat S, Alyemini MN, Hasan SA. Foliar spray of brassinosteroid enhances yield and quality of *Solanum lycopersicum* under cadmium stress. *Saudi J Biol Sci* 2012;19:325–35. DOI: 10.1016/j.sjbs.2012.03.005
- [113] Cao F, Liu L, Ibrahim W, Cai Y, Wu F. Alleviating effects of exogenous glutathione, glycinebetaine, brassinosteroids and salicylic acid on cadmium toxicity in rice seedlings (*Oryza sativa*). *Agrotechnology* 2013;2:107. DOI:10.4172/2168-9881.1000107
- [114] Hayat S, Khalique G, Wani AS, Alyemini MN, Ahmad A. Protection of growth in response to 28-homobrassinolide under the stress of cadmium and salinity in wheat. *Int J Biol Macromolecul* 2014;64:130–6. DOI: 10.1016/j.ijbiomac.2013.11.021
- [115] Herbette S, Taconnat L, Hugouvieux V, Piette L, Magniette M-LM, Cuine S, Auroy P, Richaud P, Forestier C, Bourguignon J, Renou J-P, Vavasseur A, Leonhardt N. Genome-wide transcriptome profiling of the early cadmium response of *Arabidopsis* roots and shoots. *Biochimie* 2006;88:1751–65. DOI: 10.1016/j.biochi.2006.04.018
- [116] Villiers F, Jourdain A, Bastien O, Leonhardt N, Fujioka S, Tichtinck G, Parcy F, Bourguignon J, Hugouvieux V. Evidence for functional interaction between brassinosteroids and cadmium response in *Arabidopsis thaliana*. *J Exp Bot* 2012;63:1185–200. DOI: 10.1093/jxb/err335
- [117] Gill A, Sagoo MIS. Mutagenic potential and nutritive quality of turnip plants raised over chromium amended soils. *Int J Bot* 2010;6:127–31.
- [118] Shanker AK, Cervantes TC, Loza-Tavera H, Avudainayagam S. Chromium toxicity in plants. *Environ Int* 2005;31:739–53. DOI: 10.1016/j.envint.2005.02.003

- [119] Panda SK, Choudhury S. Chromium stress in plants. *Brazil J Plant Physiol* 2005;17(1): 95–102. DOI: 10.1590/S1677-04202005000100008
- [120] Choudhary SP, Kanwar M, Bhardwaj R, Gupta BD, Gupta RK. Epibrassinolide ameliorates Cr (VI) stress via influencing the levels of indole-3-acetic acid, abscisic acid, polyamines and antioxidant system of radish seedlings. *Chemosphere* 2011;84:592–600. DOI: 10.1016/j.chemosphere.2011.03.056
- [121] Sun Y, Veerabomma S, Abdel-Mageed HA, Fokar M, Asami T, Yoshida S, Allen RD. Brassinosteroid regulates fiber development on cultured cotton ovules. *Plant Cell Physiol* 2005;46:1384–91. DOI: 10.1093/pcp/pci150
- [122] Welch RM. Micronutrient nutrition of plants. *Crit Rev Plant Sci* 1995;14:49–82. DOI: 10.1080/07352689509701922
- [123] Marschner H. *Mineral Nutrition of Higher Plants*. 2nd Ed. Academic Press, London 1995.
- [124] Kevrešan S, Petrovič N, Popovič M, Kandrač M. Effect of heavy metals on nitrate and protein metabolism in sugar beet. *Biol Plant* 1998;41:235–40. DOI: 10.1023/A:1001818714922
- [125] Schickler H, Caspi H. Response of antioxidative enzymes to nickel and cadmium stresses in hyperaccumulator plants of the genus *Alyssum*. *Physiol Plant* 1999;105:39–44. DOI: 10.1034/j.1399-3054.1999.105107.x
- [126] Prasad SM, Dwivedi R, Zeeshan M. Growth, photosynthetic electron transport, and antioxidant responses of young soybean seedlings to simultaneous exposure of nickel and UV-B stress. *Photosynthetica* 2005;43:177–85. DOI: 10.1007/s11099-005-0031-0
- [127] Sharma P, Bhardwaj R. Effects of 24-epibrassinolide on growth and metal uptake in *Brassica juncea* L. under copper metal stress. *Acta Physiol Plant* 2007;29:259–63. DOI: 10.1007/s11738-007-0032-7
- [128] Kanwar MK, Bhardwaj R, Arora P, Choudhary SP, Sharma P, Kumar S. Plant steroid hormones produced under Ni stress are involved in the regulation of metal uptake and oxidative stress in *Brassica juncea* L. *Chemosphere* 2012;86:41–9. DOI: 10.1016/j.chemosphere.2011.08.048
- [129] Sharma I, Pati PK, Bhardwaj R. Effect of 24-epibrassinolide on oxidative stress markers induced by nickel-ion in *Raphanus sativus* L. *Acta Physiol Plant* 2011;33:1723–35. DOI: 10.1007/s11738-010-0709-1
- [130] Chen LM, Lin CC, Kao CH. Copper toxicity in rice seedlings: changes in antioxidative enzyme activities, H₂O₂ level and cell wall peroxidase activity in roots. *Bot Bull Acad Sin* 2000;41:99–103.

- [131] Sheldon AR, Menzies NW. The effect of copper toxicity on the growth and root morphology of Rhode grass (*Chloris gayana* Knuth.) in resin buffered solution culture. *Plant Soil* 2005;278:341–9. DOI: 10.1007/s11104-005-8815-3
- [132] Chatterjee C, Sinha P, Dube BK, Gopal R. Excess copper induced oxidative damages and changes in radish physiology. *Commun Soil Sci Plant Anal* 2006;37:2069–76. DOI: 10.1080/00103620600770425
- [133] Alaoui-Sossé B, Genet P, Vinit-Dunand F, Toussaint ML, Epron D, Badot PM. Effect of copper on growth in cucumber plants (*Cucumis sativus*) and its relationships with carbohydrate accumulation and changes in ion contents. *Plant Sci* 2004;166:1213–8. DOI: 10.1016/j.plantsci.2003.12.032
- [134] Maksymiec W, Krupa Z. Effects of methyl jasmonate and excess copper on root and leaf growth. *Biol Plant* 2007;51:322–6. DOI: 10.1007/s10535-007-0062-4
- [135] Ducic T, Polle A. Transport and detoxification of manganese and copper in plants. *Brazil J Plant Physiol* 2005;17:103–12. DOI: 10.1590/S1677-04202005000100009
- [136] Choudhary SP, Bhardwaj R, Gupta BD, Dutt P, Gupta RK, Kanwar M, Biondi S. Enhancing effects of 24-epibrassinolide and Putrescine on the antioxidant capacity and free radical scavenging activity of *Raphanus sativus* seedlings under Cu ion stress. *Acta Physiol Plant* 2011;33:1319–33. DOI: 10.1007/s11738-010-0665-9
- [137] Guo H, Li L, Ye H, Yu X, Algreen A, Yin Y. Three related receptor-like kinases are required for optimal cell elongation in *Arabidopsis thaliana*. *Proc Nat Acad Sci USA* 2009;106:7648–53. DOI: 10.1073/pnas.0812346106
- [138] Gonzalez-Garcia MP, Vilarrasa-Blasi J, Zhiponova M, Divol F, Mora-Garcia S, Russinova E, Cano-Delgado AI. Brassinosteroids control meristem size by promoting cell cycle progression in *Arabidopsis* roots. *Development* 2011;138:849–59. DOI: 10.1242/dev.057331
- [139] Gudesblat GE, Russinova E. Plants grow on brassinosteroids. *Curr Opin Plant Biol* 2011;14:530–7. DOI: 10.1016/j.pbi.2011.05.004
- [140] Filová A, Sytar O, Krivosudská E. Effects of brassinosteroid on the induction of physiological changes in *Helianthus annuus* L. under copper stress. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis* 2013;61:623–9. DOI: 10.11118/actaun201361030623
- [141] Betul Burun K, Marakli S, Gozukirmizi N. Alleviation of copper stress with brassinosteroid in germinating sunflower roots. *J Animal Plant Sci* 2015; in press.
- [142] Vázquez MN, Guerrero YR, González LM, de la Noval WT. Brassinosteroids and plant responses to heavy metal stress. An overview. *Open J Metal* 2013;3:34–41. DOI: 10.4236/ojmetal.2013.32A1005.
- [143] Çağ S, Gören-Sağlam N, Çingil-Bariş Ç, Kaplan E. The effect of different concentration of epibrassinolide on chlorophyll, protein and anthocyanin content and peroxi-

- dase activity in excised red cabbage (*Brassica oleraceae* L.) cotyledons. *Biotechnol Biotechnol Equip* 2007;21:422–5. DOI: 10.1080/13102818.2007.10817487
- [144] Sharma P, Bhardwaj R, Arora N, Arora HK. Effect of 28-homobrassinolide on growth, Zn metal uptake and antioxidative enzyme activities in *Brassica juncea* L. seedlings. *Brazil J Plant Physiol* 2007;19(3):203–10. DOI: 10.1590/S1677-04202007000300004
- [145] Ramakrishna B, Rao SSR. 24-Epi brassinolide alleviated zinc-induced oxidative stress in radish (*Raphanus sativus* L.) seedlings by enhancing antioxidative system. *Plant Growth Regulat* 2012;68:249–59. DOI: 10.1007/s10725-012-9713-3
- [146] Choudhary SP, Kanwar M, Bhardwaj R, Yu J-Q, Pan Tran L-S. Chromium stress mitigation by polyamine-brassinosteroid application involves phytohormonal and physiological strategies in *Raphanus sativus* L. *PLoS ONE* 2012;7(3):e33210. DOI: 10.1371/journal.pone.0033210
- [147] Fariduddin Q, Khan TA, Yusuf M. Hydrogen peroxide mediated tolerance to copper stress in the presence of 28-homobrassinolide in *Vigna radiata*. *Acta Physiol Plant* 2014;36:2767–78. DOI: 10.1007/s11738-014-1647-0
- [148] Malamy JE. Intrinsic and environmental response pathways that regulate root system architecture. *Plant Cell Environ* 2005;28:67–77.
- [149] Vriet C, Russinova E, Reuzeau C. Boosting crop yields with plant steroids. *Plant Cell* 2012;24:842–57. DOI: 10.1105/tpc.111.094912
- [150] Müssig C, Shin G-H, Altmann T. Brassinosteroids promote root growth in *Arabidopsis*. *Plant Physiol* 2003;133:1261–71. DOI: 10.1104/pp.103.
- [151] Bajguz A, Hayat S. Effects of brassinosteroids on the plant responses to environmental stresses. *Plant Physiol Biochem* 2009;47:1–8. DOI:10.1016/j.plaphy.2008.10.002
- [152] Shu H, Ni W, Guo S, Gong Y, Shen X, Zhang X, Xu P, Guo Q. Root-applied brassinolide can alleviate the NaCl injuries on cotton. *Acta Physiol Plant* 2015;37:75. DOI 10.1007/s11738-015-1823-x