

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Transcriptional Modulation of Inositols and Raffinose Family Oligosaccharides Pathways in Plants — An (A)Biotic Stress Perspective

José Ribamar Costa Ferreira Neto, Amanda Cordeiro de Melo Souza, Manassés Daniel da Silva, Ana Maria Benko-Iseppon, Valesca Pandolfi, Antônio Félix da Costa and Ederson Akio Kido

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/61341>

Abstract

Among the multifunctional molecules that participate in processes of plant tolerance/resistance to stresses, inositol (Ins) and its derivatives (phosphorylated, methylated, oxygenated, and Raffinose Family Oligosaccharides) have attracted the attention of researchers. These compounds represent versatile and dynamic signaling molecules and osmolytes in all eukaryotes. Due to the impacts related to Ins and its derivatives in a plant cell, assays have been conducted to understand how these biomolecules affect plant physiology. Thus, overexpression or knockout of Ins-related genes has been shown as interesting strategies for generating more efficient plants capable of growing under stress conditions. In this chapter, studies using molecular tools are presented, and the impacts of their results are discussed based on the plant stress tolerance/resistance. Furthermore, an informative panel is provided with transcriptional modulation of genes related to Ins and its derivatives expressed in plants under stress. There is a gap involving about two dozen enzymes associated with the synthesis of Ins-related compounds that have not been adequately studied, and they represent an area of high biotechnological potential.

Keywords: Transgeny, tolerance, resistance, biotechnology

1. Introduction

To survive and integrate in the niche in which they germinate, plants constantly regulate their internal environment to external fluctuations encompassing soil, climate, and biological interactions. Thus, along its evolutionary processes, plants were selected through the need of

molecular mechanisms for physiological adjustments to inadequate conditions for development, resulting from adverse conditions. In this way, plants have a diverse and active cellular machinery at different stratified levels, covering perception, signaling, transcriptional control of key metabolic pathways and synthesis of molecules responsive to stresses [1].

Among the molecules functioning in more than one of the aforementioned levels, inositol (Ins; $C_6H_{12}O_6$) is a biomolecule of great interest. It is a cyclic carbohydrate (polyalcohol) that anchors in each of the six carbons forming the ring, a hydroxyl group. Along with their derivatives, Ins has multiple effects on plant metabolism. They act from the production of secondary messengers to the synthesis of osmolytes and antioxidants (more details in the reviews of [2, 3]). Phosphorylated Ins-derivatives [(poly)phosphoinositides and inositol (poly)phosphates] are versatile and dynamic signaling molecules in all eukaryotes, particularly in plants [4]. These two classes of compounds [highlighted in red and orange respectively, in Figure 1] are interdependent. While (poly)phosphoinositides are used in the synthesis of inositol (poly)phosphates through the action of phospholipases; the breaking of inositol (poly)phosphates produces inositol, which is a substrate for the synthesis of (poly)phosphoinositides. Moreover, according to Ins metabolism, shown in Figure 1, another branch realizes the synthesis of methylated derivatives (highlighted in green). These compounds act as important osmoregulators during periods of unfavorable conditions [5]. Additionally, oxygenated Ins-derivatives are observed (highlighted in yellow in Figure 1), which are involved in increasing plant tolerance to stresses by decreasing oxidative damage [6]. Still associated with Ins is the metabolism of the Raffinose Family Oligosaccharides (RFOs) [7]. In this biosynthetic pathway, the galactinol synthase (GolS; EC 2.4.1.123) uses myo-inositol and UDP-galactose to produce galactinol, which serve as galactose donors for subsequent synthesis of RFO members ([8, 9]; highlighted in red in Figure 2). Recent reports indicate that RFOs may assist in the reactive oxygen species (ROS) cleaning process. In periods of stress, ROS accumulation favors the physiological imbalance of plants [10].

Due to the impacts related to the Ins and its derivatives in a plant cell, assays have been conducted to understand how these biomolecules affect the physiology of plants. Thus, overexpression or knockout of genes present in these pathways has been shown as interesting strategy for generating more efficient plants capable of growing under abiotic stress conditions. In this chapter, studies using molecular genetic tools will be presented, which affect the above-mentioned metabolic pathways and the studied organisms.

2. The use of Ins and phosphorylated Ins-derivatives in plant cells under stress

Although there are several articles addressing the involvement of Ins and its derivatives in plant stress responses, so far no report has described the overall transcriptional orchestration of these components covering the metabolic pathways related to them. Information is available only in particular assays covering few genes and their expression modulations, some individual gene knockout analyses or a specific GMO (genetically modified organism) assay. In most

cases, Ins and its phosphorylated derivatives have positive impact in plant tolerance/resistance to several stresses, promoting a biotechnological interest in these compounds.

Among the Ins-derivatives, those that are phosphorylated [(poly)phosphoinositides (highlighted in red in Figure 1) and inositol (poly)phosphates (highlighted in orange)] are the most discussed in the literature. This fact reflects the importance of these compounds in plant physiology in signaling activities. An example is a work developed by Hunt et al. [11] with transgenic tobacco plants (*Nicotiana tabacum* cv. Wisconsin 38) expressing reduced levels of a gene encoding phospholipase C (PI-PLC; EC 3.1.4.11). This enzyme catalyzes the hydrolysis of phosphatidylinositol 4,5-bisphosphate to InsP3 [inositol (1,4,5)-trisphosphate; (Figure 1)], affecting from this point on the rest of the pathway. The obtained transgenic plants showed a partial inhibition of stomatal opening due to the action of ABA (abscisic acid phytohormone). A possible role of PI-PLC enzyme and InsP3 in ABA-dependent signaling pathway was also reported, thus suggesting that a complete response to ABA phytohormone in guard cells requires PI-PLC. However, other calcium-mobilizing pathways could also help in ABA guard cell signaling. The stomatal movement is a critical process for the plant under drought conditions by enabling better use of its water supply.

Mills et al. [12] provide more details on this ABA-mediated stomatal regulation in transgenic plants obtained by Hunt et al. [11]. A three-day assay under drought and in dark-adapted conditions, to reopen the stomata in response to light, was carried out. The results showed that transgenic plants with reduced PI-PLC as compared to control plants (with the empty vector without the transgene insert) have a greater increase in stomatal conductance. Thus, there is a strengthening of the role of inhibition due to PI-PLC in ABA-mediated stomatal opening. Further analysis indicated tobacco PI-PLC acting on the inhibition of stomatal opening by ABA, but not in promoting ABA-induced stomatal closure.

There are also reports of the involvement of Ins-derivatives in ABA-independent mechanisms during periods of drought. Perera et al. [13] obtained *A. thaliana* plants transformed with human type I gene for inositol polyphosphate 5-phosphatase (InsP5-ptase; EC 3.1.3.56; Figure 1). This enzyme hydrolysis InsP3, which is an essential element of the signal transduction pathway in general response to stresses. Looking at the impact on plant response to drought, the authors observed that transgenic plants showed increased stress tolerance after 12 days of watering suspension. After this period of stress, wild and controls plants (with the empty vector without the transgene insert) became brown and dry, while InsP5-ptase transgenic lines remained green and turgid. Furthermore, transgenic plants under drought conditions showed reduced levels of ABA compared with wild plants in the same condition as well as no induction of several genes regulated by the phytohormone. The analyses of stomatal responses in transgenic plants observed that guard cells are less responsive to the inhibition of opening stomata promoted by ABA. Nevertheless, there is an increase in sensitivity to the closing of the stomata, induced by the phytohormone. The transgenic plants showed irregular behavior in coordinated processes via ABA. Despite this, these plants showed a compensatory overexpression of an ABA-independent pathway involving the transcription factor (TF) DREB2A (dehydration-responsive element-binding protein 2A) and a subset of genes regulated by this TF. In this way, the drought tolerance of Ins5-ptase plants was mediated in part via DREB2A-




Figure 1. Inositol phosphate metabolism from the KEGG Pathway database. The stereoisomers (highlighted in blue); the phosphorylated derivatives [inositol (poly)phosphates (highlighted in orange) and (poly)phosphoinositides (highlighted in red)]; the methylated derivatives (highlighted in green); the oxygenated derivatives (highlighted in light green); and others (highlighted in purple).

dependent and that constitutive dampening of InsP3 signal revealed unforeseen interconnections between signaling pathways.

In a similar assay, Khodakovskaya et al. [14] reported physiological consequences in transgenic lines of *Solanum lycopersicum* (cv. Micro-Tom) overexpressing a human type I InsP5-ptase gene. The transgenic lines presented a content of 15–30% of InsP3 observed in the wild-type plants. This reduction led to increases in: (1) the total vegetative biomass (two- to fourfold) with an increased ratio of root:shoot dry weight; (2) lycopene levels (in fruit); and (3) the hexose concentration (on fruits and leaves). After 13 days of water stress, the leaf water potential in transgenic plants was about -0.4 MPa higher than in control plants. Drought tolerance in transgenic lines was associated with increased hexoses in the leaves. This would contribute to maintaining a greater potential for water in transgenic leaves under drought. Furthermore, increases in the number of root biomass may have contributed to this improved performance.

The involvement of InsP3 in other stress tolerance processes, beyond drought, has also been demonstrated. Alimohammadi et al. [15] obtained transgenic tomato plants (*Lycopersicon esculentum* cv. MicroTom,) overexpressing a human type I InsP5-ptase gene. These plants presented a decreased level of InsP3 and supported a continuous exposure to light longer than wild plants. Prolonged exposure to light causes oxidative stress in plant cells and can result in irreversible damage. However, the molecular mechanism involved in this tolerance process was not reported, but these transgenic lines were characterized in more detail by Alimoham-




Figure 2. The Galactose metabolism from the KEGG Pathway database. Highlighted in red are the enzymatic reactions associated with the metabolism of Raffinose Family Oligosaccharides.

mediated et al. [16]. These authors observed that under stress conditions, the transgenic plants maintained high chlorophyll content and accumulated low levels of hydrogen peroxide. This fact was attributed to the induction of genes related to multiple antioxidants [LeAPX1 (*L. esculentum* ascorbate peroxidase 1), SICAT2, LeSOD (*L. esculentum* superoxide dismutase)] during continuous exposure to light. Other effects included overexpression of the LePHYB photoreceptor (*L. esculentum* phytochrome B) and a key enzyme [LeCHS1 (*L. esculentum* chalcone synthase)] in the biosynthesis pathway of flavonoids, which are plant nonenzymatic antioxidants. There was also an overexpression of the SIMYB12 transcription factor, leading to an increase in flavonoids in tomato plants by up-regulation of the LeCHS1 expression. A relationship was established between change in phosphoinositol signaling pathway and increases tolerance to continuous exposure to light, through the activation of ROS-scavenging enzymes, and up-regulation of molecular activators of non-enzymatic antioxidants.

The biotechnological potential through the manipulation of compounds shown in Ins-related metabolic pathways may also be seen in the work of Ahmad et al. [17]. These authors performed a comprehensive analysis of *A. thaliana* genome, using the activation tagging technique in dedifferentiated calli, to identify salt-tolerant mutants (NaCl 150 mM). To this end, plants were modified with pRi35ADen4 binary vector. Such vector contains four copies of the 339 bp long cauliflower mosaic virus (CaMV) 35S enhancer in the construct that induces the expres-

sion of adjacent genes after proper insertion. Of the 18 potential tolerant mutants (150 mM NaCl), a line (*stc1*; salt-tolerant callus 1) presented the gene for myo-inositol-1-P-synthase-1 [At4g39800; MIPS1; EC 5.5.1.4; Figure 1] with induced expression in callus, with or without salt. This gene expressed 45 and 15 times higher compared to wild-type under the control condition. MIPS catalyzes the first step in the biosynthesis of inositol from glucose-6-P (Figure 1). The referred induction was greater in the mutant line than in the wild type (approximately, 260 times higher), both under stress. The *mips* gene transcription in the wild type reduced dramatically under stress condition. The tolerance analysis revealed that the mutant plants regenerated from calli showed salt tolerance in germination and growth. However, the mechanism involved was not disclosed, but the authors suggested that MIPS protect the calli and the plants from salt stress as osmolytes or by providing a precursor in the regulation of signal transduction pathways.

Kusuda et al. [18] went beyond the study of transformed lines overexpressing enzymes from the Ins-related pathways. They analyzed the differences among wild type and transformants lines in regard to salt tolerance in 3.5 days in medium with concentrations up to 250 mM NaCl. They also sought for differences by mining the metabolomes (the fourth leaf tissue harvested at 0, 6, and 12 h after NaCl stress induction) of the studied plants. To this end, a rice cultivar (*Oryza sativa* cv. Kitaake) was transformed with the construction Act::RINO1. The RINO1 gene encodes an MIPS (EC 5.5.1.4, Figure 1). It has been demonstrated that the constitutive overexpression of rice MIPS when compared to a wild type, results in greater tolerance to salt stress. Furthermore, it leads to a range of metabolic changes, with increased production of various metabolites (such as inositol, raffinose, ascorbate, amino acids). These handle the protection of plants from abiotic stresses. Additionally, activation of basal metabolisms such as glycolysis, the pentose phosphate pathway, and the tricarboxylic acid cycle has been observed during induction of the Ins metabolism in those plants overexpressing MIPS.

Ins metabolism and phosphorylated Ins-derivatives are also associated with response to biotic stresses in plants. This fact shows the plurality of actions of these compounds. Murphy et al. [19] report evidence in this direction. They obtained transgenic potato [*S. tuberosum* L. (cv. Desiree)] and *A. thaliana* lines, synthesizing low levels of phytic acid (1-D-myo-inositol-P6 or InsP6, Figure 1). The transgenic potato lines were obtained by: (1) constitutive expression of an antisense sequence of the myo-inositol 3-phosphate synthase gene. This enzyme (IPS, EC 5.5.1.4, Figure 1) catalyzes the first step in the InsP6 biosynthesis; (2) plants expressing the *Escherichia coli* polyphosphate kinase (PPK, EC 2.7.4.1; Figure 1). PPK inserts inorganic phosphate into chains of phosphate residues linked by phospho-anhydride bonds, and this decreases the phosphate available to InsP6 biosynthesis. Genetically modified potatoes presenting both (1) and of (2) showed increased susceptibility to avirulent pathogen potato virus Y and the virulent pathogen tobacco mosaic virus (TMV). In relation to *A. thaliana*, the authors obtained three loss-of-function mutants [two (*atips1* and *atips2*) involving the gene for IPS (EC 5.5.4.1, Figure 1); and one (*atipk1*) involving the gene encoding inositol polyphosphate kinase (IPK1; EC 2.7.1.158, Figure 1)]. IPK1 catalyzes the final step in InsP6 and without this enzyme, very little InsP6 is done. The disruption of InsP6 biosynthesis in *A. thaliana* resulted in an increased susceptibility to viruses (tobacco mosaic virus), bacteria [*Pseudomonas syringae* pv tomato (Pst) DC3000 and Pst DC3000 AvrB], and fungus (*Botrytis cinerea*). The

increased susceptibility only occurred for *atips2* and *atipk1* mutants. For *atips1* mutants showing also InsP6 depletion, the resistance to pathogens was not compromised. This suggests either that a particular pool of InsP6 regulates defense against pathogens in *A. thaliana*, or IPS1 and IPs2 are differentially regulated, and one cannot compensate for the other. It was observed with regard to the typical responses of both species that levels of salicylic acid (SA), a key molecular signal for establishing local and systemic acquired resistance, were not diminished. Therefore, SA was not responsible for increased susceptibility to pathogens analyzed. Furthermore, a relationship between the inositol phosphate metabolism and basal resistance to pathogens (fungi, bacteria, and viruses) has been established. This indicates that InsP6 is also required as an essential operation signal for regulation and defense in plants, together with several other well-known defensive signals (NO, cADPR, SA and Ca²⁺).

Recently, Meng et al. [20] found that *A. thaliana* *ips1* loss-of-function mutant (*atips1*; IPS; EC 5.5.4.1, Figure 1) exhibits spontaneous cellular death and increased resistance to oomycete *Hyaloperonospora arabidopsis* pathogen. This result, together with the above, supports the premise that a particular InsP6 pool may regulate defense pathways, since *atips1* mutants did not show increased susceptibility to the variety of testes pathogens (Tobacco mosaic virus; *Pseudomonas syringae* pv. tomato (Pst) DC3000 and Pst. DC3000 AvrB; and *Botrytis cinerea*). Additionally, it was observed that spraying the *atips1* mutant plants with Ins could suppress the formation of spontaneous lesions, indicating that Ins production and not AtIPS1 protein is required to prevent the lesion formation. The authors also found evidence for the role of Ins (or Ins-derivatives) in the regulation of programmed cell death.

Reports presented in Table 1 also show the broad roles of Ins and its phosphorylated derivatives in plant cells. From the wide spectrum of analyzed genes, a range of effects on plants at different levels was observed. These effects have shown associations with hormone signaling pathways, such as ABA [12], influences in photosystems [21, 22], with reactive oxygen species (ROS; [23]), with relative water content, with osmotic adjustment [24], among others (Table 1).

3. Methylated Ins-derivatives in plant cell and the biotechnological use to increase stress tolerance

Some plants use Ins as precursor of compatible solutes such as D-ononitol and D-pinitol, which act as osmoprotectants (small molecules that act as osmolytes and help organisms survive in extreme osmotic stress [35]). In halophyte ice plant (*Mesembryanthemum crystallinum*), which is considered highly tolerant to drought, salinity, and cold, Ins is methylated to D-ononitol and subsequently epimerized to D-pinitol [36]. The myo-inositol O-methyl transferase gene (IMT1; EC 2.1.1.40; Figure 1) is transcriptionally induced by osmotic stress, whereas neither transcriptional nor enzyme activities is detectable in ice plants under normal growth conditions [37, 38]. Despite the positive influence of these metabolites in plant physiology under abiotic stress conditions, there are less available data for these compounds compared with phosphorylated Ins-derivatives.

EC# Number	Gene origin (specie)	Transformant (specie)	Gene Modulation	Analysed Condition	Impact on Tolerance	Authors	Notes
5.5.1.4	<i>Pc</i>	<i>Os and Bj</i>	Overexp.	HS	Raise	[25]	a.
5.5.1.4	<i>Pc</i>	<i>Nt</i>	Overexp.	HS	Raise	[21]	b.
5.5.1.4	<i>As</i>	<i>At</i>	Overexp.	HS	Raise	[22]	c.
3.1.3.25	<i>Ca</i>	<i>At</i>	Overexp.	HS, P, PEG, and HT	Raise	[26]	d.
2.7.1.140; 2.7.1.151	<i>At</i>	<i>Nt</i>	Overexp.	HS and OS	Raise	[27]	e.
2.7.1.140; 2.7.1.151	<i>Th</i>	<i>Bn</i>	Overexp.	HS, D, and OS	Raise	[28]	f.
3.1.4.11	<i>Zm</i>	<i>Zm</i>	Overexp.	D	Raise	[29]	g.
2.7.1.159; 2.7.1.134	<i>Os</i>	<i>Nt</i>	Overexp.	HS	Decrease	[24]	h.
3.1.3.57	<i>At</i>	<i>At</i>	Knockout	HS, F, and D	Decrease	[30]	i.
2.7.1.137	<i>At</i>	<i>At</i>	Knockout	HS	Decrease	[31]	j.
2.7.1.67	<i>At</i>	<i>At</i>	Overexp.	HS and ABA	Raise	[23]	l.
2.7.8.11	<i>Zm</i>	<i>Zm</i>	Overexp.	D	Raise	[32]	m.
3.1.3.8	<i>At</i>	<i>At</i>	Overexp.	HS and OSM	Raise	[33]	n.
2.7.1.149	<i>At</i>	<i>At</i>	Overexp.	HS, D, and ABA	**	[34]	-

†Enzyme Commission; *According to KEGG Database Pathway; - Not observed; Overexp. (Overexpression); **Legend:** HS (high salinity); P (paraquat); PEG (polyethylene glycol); HT (high temperature); OS (oxidative stress); D (drought); DH (dehydration); ABA (ABA hormone); OSM (osmotic stress); F (freezing). **a.** Albeit to a variable extent, overexpression of this gene confers salt-tolerance to diverse evolutionary organisms (from prokaryotes to eukaryotes), including crop plants; **b.** Transgenic individuals presenting retention of approximately 40–80% of the photosynthetic competence under analyzed stress condition; **c.** Transgenic individuals retained more chlorophyll and carotenoid by protecting the photosystem II; **d.** Improving seed germination and seedling growth in transgenic individuals under stress conditions; **e.** Expression patterns of various stress responsive genes were enhanced, and the activities of antioxidative enzymes were elevated in transgenic plants; **f.** The transcripts of various stress-responsive genes are increased in ThIPK2 transgenic plants under salt stress condition; **g.** The sense transgenic plants had higher relative water content, better osmotic adjustment, increased photosynthesis rates, lower percentage of ion leakage and less lipid membrane peroxidation, higher grain yield than the wild type; **h.** The 1,3,4-trisphosphate 5/6-kinase is a negative regulator of osmotic stress signaling in tobacco; **i.** The genetic evidence indicating that phosphoinositols mediate ABA and stress signal transduction in plants, and their turnover is critical for attenuating ABA and stress signaling; **j.** Salt stress responses, such as increased plasma membrane endocytosis and the intracellular production of ROS, are coordinated by phospholipid-regulated signaling pathways; **l.** AtPI4K γ 3 is activated by DNA demethylation and regulates the ROS accumulation induced by high salt treatment or ABA treatment; **m.** ZmPIS regulates the plant response to drought stress through altering membrane lipid composition and increasing ABA synthesis in maize; **n.** AtPAP15 (3-PHYTASE) may modulate AsA levels by controlling the input of myoinositol into this branch of AsA biosynthesis in *Arabidopsis thaliana*. *At*: *A. thaliana*; *Pc*: *P. coarctata*; *Sa*: *S. alterniflora*; *Ca*: *C. arietinum*; *Th*: *T. halophila*; *Zm*: *Z. mays*; *Os*: *O. sativa*; *Nt*: *N. tabacum*; *Bn*: *B. napus*; *Bj*: *B. juncea*.

Table 1. Transgenic and knockout plant assays available in the literature related to (poly)phosphoinositides and inositol (poly)phosphates. Relevant information: EC number of the enzyme coded by the studied gene, plant donor species, the genetically modified organisms (transformants), the modulation of the studied gene, the analyzed stress condition, the impact on plant tolerance and physiology (additional details, please see the legends).

Sheveleva et al. [39] were one of the first to report the biotechnological potential of methylated Ins-derivatives. In their work, the authors superexpressed O-methyltransferase (IMT1; EC

2.1.1.40; Figure 1) of *Mesembryanthemum crystallinum* in tobacco (*Nicotiana tabacum* cv. SRI). The transgenic plant increased its tolerance to abiotic stresses [drought and high salinity (50–250 mM NaCl)] when compared to the wild control line. An accumulation of methylated inositol D-ononitol in amounts of fresh weight exceeding 35 $\mu\text{mol g}^{-1}$ in the transformed lines was observed. Besides, the photosynthetic CO_2 fixation was less inhibited in those plants during drought and salt stress. Further, transformed plants recovered faster than the wild type after rehydration. In turn, Sengupta et al. [40] observed an increase of the D-pinitol synthesis in a wild-type rice (*Porteresia coarctata*) with halophilic characteristics when subjected to high salt environment (400 mM NaCl). An increment also occurred in both transcriptional and proteomic level of IMT1, not observable in domesticated rice under the same condition. The authors also reported an increase in the expression of L-myo-inositol 1-phosphate synthase (PcMIPS1; EC 5.5.1.4; Figure 1), along with the expression of IMT1. According to the authors, this suggests that the accumulation of D-pinitol would be a mechanism regulated by salt stress.

Recently, Zhu et al. [41] used a similar strategy as the one developed by Sheveleva et al. [39] to express in *A. thaliana*, a gene coding IMT1, from *Mesembryanthemum crystallinum*. The transformed plants showed higher growth compared to the wild control line and increased tolerance to cold stress (4°C). This increment in tolerance was attributed to different factors: (1) the electrolyte leakage content in the transgenic plants was significantly lower than that of the wild-type plants after freezing stress, showing less damage to the membranes of those plants; (2) transgenic plants showed lower MDA content than wild-type plants, not only in normal conditions but also after stress; and (3) a higher proline content presented in transgenic lines than in wild type, after application of stress.

4. Oxygenated Ins-derivatives in plant cell and the plant strategy to tackle stress

So far, the myo-inositol oxygenase (MIOX; EC 1.13.99.1; Figure 1) is the only enzyme known by the oxidation of Ins [42]. Its importance in plants stood out from the statement in the Ins metabolism as a precursor in Ascorbic Acid (AsA) biosynthesis in Arabidopsis. In this sense, Lorence et al. [43] observed the expression of a myo-inositol oxygenase (miox4) increasing the content of AsA in leaves (approximately two- to threefold). By that, they anticipated a potential use of the gene by genetic engineering, enhancing levels of this important antioxidant in plants. Further analysis indicated that D-GlcUA (Figure 1), a derivative from MIOX reaction (EC 1.13.99.1, Figure 1), plays a negligible role for AsA biosynthesis [44]. However, MIOX can control the metabolite level of myo-inositol in plants [44].

Nevertheless, the metabolic consequences of MIOX action are still unclear. In order to contribute with information from gene regulation and catalytic activity of this enzyme, Duan et al. [6] performed a functional characterization in rice (*Oryza sativa*), observing its predominant expression in root, with induced transcription under drought stress (20% PEG6000 solution), H_2O_2 , high salt (200 mM NaCl), cold (4°C) and Abscisic Acid (100 μM). Transgenic rice lines overexpressing an MIOX gene showed a higher survival rate than a wild line, when in contact with 20% PEG6000 solution. In the same way, the authors also reported for the

transgenic lines, transcription levels significantly induced for genes coding enzymes associated with ROS scavenging, suggesting an MIOX activity in reducing oxidative stress.

Moreover, Alford et al. [45] reported MIOX enzymes responding to growing conditions of *A. thaliana* in low energy/nutrient environment. Their data supported the hypothesis that MIOX2 and MIOX4 enzymes would be encoded by multiple genes. Furthermore, on plants growing in low energy conditions/nutrients environments, MIOX2 enzyme would have a significant role in providing Ins to many different tissues, whereas MIOX4 would act in supplemental form at some tissues. An analysis of promoters was presented, considering multiple lines of MIOX2p:GUS and MIOX4p:GUS seedlings growing on: (I) no nutrients (agar); (II) low nutrients (agar + 0.5× MS salts); and (III) optimal nutrients (agar + 0.5× MS salts + 3% glucose) in low light (40 μE) during seven days. As a result, only MIOX2p:GUS was expressed abundantly in condition I, whereas in condition II, both MIOX2p:GUS and MIOX4p:GUS showed moderate expression, having been more expressed previously.

5. Raffinose Family Oligosaccharides (RFOs) and plant strategy to address stresses

RFOs are a class of compatible solutes coming from Ins metabolism ramifications. As mentioned before, the enzyme GolS (EC 2.4.1.123, highlighted in red in Figure 2) connects the metabolism of these compounds, producing galactinol (highlighted in red in Figure 2), which serves as galactose donor for further synthesis of RFO members [8, 9]. To date, structural genomics data and global transcriptome analysis concerning RFOs are only available for corn [46]. For this crop, the authors have performed a genomic identification of genes associated with raffinose metabolism, together with an expression analysis using data-mining from GEO (<http://www.ncbi.nlm.nih.gov/geo>) and PLEXdb databases (<http://www.plexdb.org>). Additionally some transgenic lines overexpressing specific gene isoforms related to RFO pathway, under particular growth conditions, are available for some species. These studies showed a positive impact in some crops.

Taji et al. [47], for example, analyzed the expression of seven genes encoding GolS in *A. thaliana* under different stresses. From those genes, only three (AtGolS1, 2 and 3) were stress-responsive. AtGolS1 and AtGolS2 were induced by drought and high salt (250 mM NaCl), but not by low temperature (4°C). On the other hand, AtGolS3 was induced by low temperatures, but not high salinity or drought. The AtGolS2 overexpression in *A. thaliana* was associated with an increase of raffinose and galactinol and resulted in a reduction of leaf transpiration with a greater drought tolerance. In turn, Dos Santos et al. [48] analyzing three *Coffea arabica* galactinol synthase isoforms (CaGolS1, CaGolS2, CaGolS3) observed a mainly tissue-specific expression but differentiated regulation depending on the applied stress (drought, heat, and high salinity). This reinforces the observation by Taji et al. [47], indicating that different galactinol sets can be necessary for response to various stresses. The GOLs (BnGOLS-1) activity was also positively correlated with desiccation tolerance in cabbage seeds (*Brassica napus*) during the vegetative growth period [8]. The tolerance was observed around 21-24 days after flowering cabbage, coinciding with the accumulation of raffinose and stachyose. The

BnGOLS-1 transcripts accumulation was concomitant with the formation of these two RFO members [8].

In *Arabidopsis thaliana* leaves overexpressing HsfA2 (Heat-Shock Transcription Factor A2; [49]) was also found highly induced GolS1, -2, -4 and Raffinose Syntase 2 (RS2; EC 2.4.1.82, highlighted in red in Figure 2) transcriptions. The galactinol and raffinose levels in the transgenic plants were higher compared to the wild-type lines, both in the control condition. These higher levels were positively correlated with an increase in plant tolerance when exposed to the studied stresses [mevalonate (50 mM), high salt (100 mM NaCl) and low temperatures (4°C)].

Latter, Pennycooke et al. [50] studied the expression of α -galactosidase gene (EC 3.1.2.22 highlighted in red in Figure 2) from petunia (*Petunia x hybrida* "Mitchell"), monitoring acclimated plants to low temperatures (4°C) and in response to increasing temperature (25°C). Transcripts induction were observed after one hour of desacclimation occurring together with an increase in enzymatic activity and decreased raffinose content, suggesting that the rise in temperature can regulate the RFO catabolism of certain members, through gene regulation that encoding α -galactosidase.

Thus, the diversity of functions performed by compounds presenting in the Ins metabolism was shown in the described works. Also, studies of distinct isoforms showed positive correlations with plant responses to various abiotic stresses. In this way, the identification of new transcripts, as well as the understanding of its regulation (spatial and temporal) in plants under unfavorable conditions for the development may lead to the discovery of new genes with biotechnological potential.

According to the Kyoto Encyclopedia of Genes and Genomes (KEGG) Pathway database (<http://www.genome.jp/kegg/pathway.html>), which provides diagrams of various metabolic processes, at least 45 enzymes are associated with the metabolic pathways described here. Of these enzymes, 21 (highlighted in green boxes in Figures 3A and 3B) have been studied in previous works addressing the transcriptional expression of their genes or effects on plants under stress. Therefore, there are at least 24 enzymes (in red boxes in Figures 3A and 3B) that have not been targets of these analyses, with significant potential for further research in biotechnology.

6. Ins and its derivatives in humans: Antinutrients *versus* disease prevention

Once Ins and its derivatives are present in vegetables and these are part of the daily diet of large populations around the world, it is essential to analyze their potential effects on consumers. The Ins and related metabolites play a heterogeneous physiological role, depending on the concerned organism, plant or animal (including human). In plants, as already mentioned, such compounds help regulate plant homeostasis during periods of stress. In animals (including humans), their influence has very diverse physiological repercussions. Initially, they were only seen as harmful agents because some representatives when present in certain

plants could act as antinutritional factors, thereby reducing the bioavailability of important nutrients and the nutritional value of the food. According to Kokhar and Apenten [51], this effect is present a result of a selected adaptive mechanism due to a "chemical warfare" between higher plants and herbivorous pests.




Figure 3. KEGG Pathway database diagrams presenting: (A) Inositol Metabolism; (B) Galactose Metabolism, showing the Raffinose Family Oligosaccharides (highlighted in red). Green Box: enzymes with transcriptional modulation data available from stress assays. Red Box: enzymes without transcriptional modulation data available in the literature. Yellow Box: enzymes not covered in this review.

Among the various Ins-derivatives, phytic acid (1-D-myo-inositol-P6; Figure 1) is the most studied, concerning the impacts on human and animal health. Its unique structure provides the ability to chelate cations such as iron, zinc, potassium, magnesium, and copper, forming

insoluble salts denominated phytate. These salts adversely affect animal's ability to absorb and digest nutrients [52, 53]. Phytates can form complexes with proteins, changing their structures and their enzymatic activities and characteristics of solubility and proteolytic digestibility [54]. However, there are reports that show positive aspects in phytates consumption. The presence of these in the diet of patients with diabetes has positive effects in reducing the level of blood glucose due to decreased starch digestion rate and slowing of gastric evacuation [55]. There are also reports of activity against HIV replication, kidney stones prevention, reduction of cholesterol and triglycerides levels, as well as assistance in prevention of heart diseases (for review see [54]). Studies also indicate that both Ins [56] and phytic acid [56, 57] have anticancer properties. With regard to RFOs, besides the fact that they are potential antinutritional factors, there are indications that they may act as important immunostimulants in animals (including humans). Also, RFOs' involvement is suggested in universal mechanisms of oxidative balance in several taxa [58].

7. Concluding remarks and perspectives

Experimentally, mutants and transgenic analyses are being successfully carried out to uncover the various roles played by Ins-related compounds. It is known today that some phosphorylated derivatives of inositol are connected with a large number of signaling procedures which are regulated by both abiotic and biotic stress. Methylated and oxygenated Ins-derivatives, including RFOs, have also proven to be active agents in the process of plant acclimatization to unfavorable conditions, involved in a number of functions. However, there is a gap to be filled. About two dozen enzymes associated with the synthesis of these compounds have not been adequately studied and they represent an area of high biotechnological potential.

Acknowledgements

The authors thank CNPq (National Council of Technological and Scientific Development – process number 454112/2014-9) for financial support. JRCFN thanks to CAPES (InterSis Network – Biocomp Project) for the fellowship granted. EAK thanks to CNPq for the fellowship granted.

Author details

José Ribamar Costa Ferreira Neto^{1*}, Amanda Cordeiro de Melo Souza¹,
Manassés Daniel da Silva¹, Ana Maria Benko-Iseppon¹, Valesca Pandolfi¹,
Antônio Félix da Costa² and Ederson Akio Kido¹

*Address all correspondence to: netocostaferreira@gmail.com

1 Department of Genetics, Federal University of Pernambuco, Brazil

2 Agronomic Institute of Pernambuco State (IPA), Brazil

References

- [1] Bartels D, Souer E. Molecular Responses of Higher Plants to Dehydration. In: Hirt H, Shinozaki K, editors. *Plant Responses to Abiotic Stress*. 1st edn. Springer; 2004. p. 9–38. DOI: 10.1007/978-3-540-39402-0_2
- [2] Loewus, FA, Murthy PPN. Myo-Inositol Metabolism in Plants. *Plant Sci*. 2000; 150: 1–19. DOI: doi:10.1016/S0168-9452(99)00150-8
- [3] Majumder AL, Biswas BB. *Biology of Inositols and Phosphoinositides: Subcellular Biochemistry*. 2nd edn. Springer; 2006. DOI: 10.1007/0-387-27600-9
- [4] Munnik T, Vermeer JEM. Osmotic Stress-Induced Phosphoinositide and Inositol Phosphate Signalling in Plants. *Plant Cell Environ*. 2010;33:655–669. DOI: 10.1111/j.1365-3040.2009.02097.x
- [5] Flowers TJ, Colmer TD. Salinity Tolerance in Halophytes. *New Phytol*. 2008;179:945–963. doi: 10.1111/j.1469-8137.2008.02531.x
- [6] Duan J, Zhang M, Zhang H, Xiong H, Liu P, Ali J, Li J, Li Z. OsMIOX, a Myo-Inositol Oxygenase Gene, Improves Drought Tolerance through Scavenging of Reactive Oxygen Species in Rice (*Oryza Sativa* L.). *Plant Sci*. 2012;196:143–151. DOI: 10.1016/j.plantsci.2012.08.003
- [7] Keller F, Pharr DM. Metabolism of Carbohydrates in Sinks and Sources: Galactosyl-Sucrose Oligosaccharides. In: Zamski E, Schaffer AA, editors. *Photoassimilate Distribution in Plants and Crops*. 1st edn. New York: Marcel Dekker; 1996. p. 115–184.
- [8] Liu JJJ, Krenz DC, Galvez AF, De Lumen BO. Galactinol Synthase (GS): Increased Enzyme Activity and Levels of mRNA due to Cold and Desiccation. *Plant Sci*. 1998;134:11–20. DOI: 10.1016/S0168-9452(98)00042-9
- [9] Sprenger N, Keller F. Allocation of Raffinose Family Oligosaccharides to Transport and Storage Pools in *Ajuga Reptans*: The Roles of Two Distinct Galactinol Synthases. *Plant J*. 2000;21:249–258. DOI: 10.1046/j.1365-313X.2000.00671.x
- [10] Elsayed A I, Rafudeen MS, Gollack D. Physiological Aspects of Raffinose Family Oligosaccharides in Plants: Protection against Abiotic Stress. *Plant Biol*. 2014;16:1–8. DOI: 10.1111/plb.12053
- [11] Hunt L, Mills LN, Pical C, Leckie CP, Aitken FL, Kopka J, Mueller-Roeber B, McAinsh MR, Hetherington AM, Gray JE. Phospholipase C Is Required for the Con-

- trol of Stomatal Aperture by ABA. *Plant J.* 2003;34:47–55. DOI: 10.1046/j.1365-313X.2003.01698.x
- [12] Mills LN, Hunt L, Leckie CP, Aitken FL, Wentworth M, McAinsh MR, Gray JE, Hetherington AM. The Effects of Manipulating Phospholipase C on Guard Cell ABA-Signalling. *J Exp Bot.* 2004;55:199–204. DOI: 10.1093/jxb/erh027
- [13] Perera IY, Hung C-Y, Moore CD, Stevenson-Paulik J, Boss WF. Transgenic Arabidopsis Plants Expressing the Type 1 Inositol 5-Phosphatase Exhibit Increased Drought Tolerance and Altered Abscisic Acid Signaling. *Plant Cell.* 2008;20:2876–2893. DOI: 10.1105/tpc.108.061374
- [14] Khodakovskaya M, Sword C, Wu Q, Perera IY, Boss WF, Brown CS, Sederoff WH. Increasing Inositol (1,4,5)-Trisphosphate Metabolism Affects Drought Tolerance, Carbohydrate Metabolism and Phosphate-Sensitive Biomass Increases in Tomato. *Plant Biotechnol J.* 2010;8:170–183. DOI: 10.1111/j.1467-7652.2009.00472.x
- [15] Alimohammadi M, De Silva K, Ballu C, Ali N, Khodakovskaya MV. Reduction of Inositol (1,4,5)-Trisphosphate Affects the Overall Phosphoinositol Pathway and Leads to Modifications in Light Signalling and Secondary Metabolism in Tomato Plants. *J Exp Bot.* 2012;63:825–835. DOI: 10.1093/jxb/err306
- [16] Alimohammadi M, Lahiani MH, Khodakovskaya MV. Genetic Reduction of Inositol Triphosphate (InsP3) Increases Tolerance of Tomato Plants to Oxidative Stress. *Planta.* 2015;242:123–135. DOI: 10.1007/s00425-015-2289-1
- [17] Ahmad A, Niwa Y, Goto S, Kobayashi K, Shimizu M, Ito S, Usui Y, Nakayama T, Kobayashi H. Genome-Wide Screening of Salt Tolerant Genes by Activation-Tagging Using Dedifferentiated Calli of Arabidopsis and Its Application to Finding Gene for Myo-Inositol-1-P-Synthase. *PLoS One.* 2015;10: e0115502. DOI: 10.1371/journal.pone.0115502
- [18] Kusuda H, Koga W, Kusano M, Oikawa A, Saito K, Hirai MY, Yoshida KT. Ectopic Expression of Myo-Inositol 3-Phosphate Synthase Induces a Wide Range of Metabolic Changes and Confers Salt Tolerance in Rice. *Plant Sci.* 2015;232:49–56. DOI: 10.1016/j.plantsci.2014.12.009
- [19] Murphy, AM, Otto, B, Brearley, Ca, Carr, JP, Hanke, D.E. A Role for Inositol Hexakisphosphate in the Maintenance of Basal Resistance to Plant Pathogens. *Plant J.* 2008;56:638–652. DOI: 10.1111/j.1365-313X.2008.03629.x
- [20] Meng PH, Raynaud C, Tcherkez G, Blanchet S, Massoud K, Domenichini S, Henry Y, Soubigou-Taconnat L, Lelarge-Trouverie C, Saindrenan P, Renou JP, Bergounioux C. Crosstalks between Myo-Inositol Metabolism, Programmed Cell Death and Basal Immunity in Arabidopsis. *PLoS One.* 2009;4. DOI: 10.1371/journal.pone.0007364
- [21] Majee M, Maitra S, Dastidar KG, Pattnaik S, Chatterjee A, Hait NC, Das KP, Majumder AL. A Novel Salt-Tolerant L-Myo-Inositol-1-Phosphate Synthase from *Porteresia Coarctata* (Roxb.) Tateoka, a Halophytic Wild Rice. *Molecular Cloning, Bacterial*

- Overexpression, Characterization, and Functional Introgression into Tobacco-Confering Salt Tolerance Phenotype. *J Biol Chem.* 2004;279:28539–28552. DOI: 10.1074/jbc.M310138200
- [22] Joshi R, Ramanarao MV, Baisakh N. Arabidopsis Plants Constitutively Overexpressing a Myo-Inositol 1-Phosphate Synthase Gene (SaINO1) from the Halophyte Smooth Cordgrass Exhibits Enhanced Level of Tolerance to Salt Stress. *Plant Physiol Biochem.* 2013;65:61–66. DOI: <http://dx.doi.org/10.1016/j.plaphy.2013.01.009>
- [23] Akhter S, Uddin MN, Jeong IS, Kim DW, Liu X-M, Bahk JD. Role of Arabidopsis At-PI4K γ 3, a Type II Phosphoinositide 4-Kinase, in Abiotic Stress Responses and Floral Transition. *Plant Biotechnol J.* 2015:1–16. DOI: 10.1111/pbi.12376
- [24] Niu X, Chen Q, Wang X. OsITL1 Gene Encoding an Inositol 1,3,4-Trisphosphate 5/6-Kinase is a Negative Regulator of Osmotic Stress Signaling. *Biotechnol Lett.* 2008;30:1687–1692. DOI: 10.1007/s10529-008-9730-5
- [25] Das-Chatterjee A, Goswami L, Maitra S, Dastidar KG, Ray S, Majumder AL. Introgression of a Novel Salt-Tolerant L-Myo-Inositol 1-Phosphate Synthase from *Porteresia Coarctata* (Roxb.) Tateoka (PcINO1) Confers Salt Tolerance to Evolutionary Diverse Organisms. *FEBS Lett.* 2006;580:3980–3988. DOI: 10.1016/j.febslet.2006.06.033
- [26] Saxena SC, Salvi P, Kaur H, Verma P, Petla BP, Rao V, Kamble N, Majee M. Differentially Expressed Myo-Inositol Monophosphatase Gene (CaIMP) in Chickpea (*Cicer Arietinum* L.) Encodes a Lithium-Sensitive Phosphatase Enzyme with Broad Substrate Specificity and Improves Seed Germination and Seedling Growth under Abiotic Stresses. *J Exp Bot.* 2013;64:5623–5639. DOI: 10.1093/jxb/ert336
- [27] Yang L, Tang R, Zhu J, Liu H, Mueller-Roeber B, Xia H, Zhang H. Enhancement of Stress Tolerance in Transgenic Tobacco Plants Constitutively Expressing AtIpk2, an Inositol Polyphosphate 6-/3-Kinase from Arabidopsis Thaliana. *Plant Mol Biol.* 2008;66:329–343. DOI: 10.1007/s11103-007-9267-3
- [28] Zhu JQ, Zhang JT, Tang RJ, Lv QD, Wang QQ, Yang L, Zhang HX. Molecular Characterization of ThIPK2, an Inositol Polyphosphate Kinase Gene Homolog from *Thelungiella Halophila*, and its Heterologous Expression to Improve Abiotic Stress Tolerance in *Brassica Napus*. *Physiol Plant.* 2009;136: 407–425. DOI: 10.1111/j.1399-3054.2009.01235.x
- [29] Wang C-R, Yang A-F, Yue G-D, Gao Q, Yin H-Y, Zhang J-R. Enhanced Expression of Phospholipase C 1 (ZmPLC1) Improves Drought Tolerance in Transgenic Maize. *Planta.* 2008;227:1127–1140. DOI: 10.1007/s00425-007-0686-9
- [30] Xiong L, Lee B, Ishitani M, Lee H, Zhang C, Zhu J. FIERY1 Encoding an Inositol Polyphosphate 1-Phosphatase Is a Negative Regulator of Abscisic Acid and Stress Signaling in Arabidopsis. *Gene Dev.* 2001;15:1971–1984. DOI: 10.1101/gad.891901.netic
- [31] Leshem Y, Seri L, Levine A. Induction of Phosphatidylinositol 3-Kinase-Mediated Endocytosis by Salt Stress Leads to Intracellular Production of Reactive Oxygen Spe-

- cies and Salt Tolerance. *Plant J.* 2007;51:185–197. DOI: 10.1111/j.1365-313X.2007.03134.x
- [32] Liu X, Zhai S, Zhao, Y, Sun B, Liu C, Yang A, Zhang J. Overexpression of the Phosphatidylinositol Synthase Gene (ZmPIS) Conferring Drought Stress Tolerance by Altering Membrane Lipid Composition and Increasing ABA Synthesis in Maize. *Plant Cell Environ.* 2013;36:1037–1055. DOI: 10.1111/pce.12040
- [33] Zhang W, Gruszewski HA, Chevone BI, Nessler CL. An Arabidopsis Purple Acid Phosphatase with Phytase Activity Increases Foliar Ascorbate. *Plant Physiol.* 2008;146:431–440. DOI: 10.1104/pp.107.109934
- [34] Mikami K, Katagiri T, Iuchi S. A Gene Encoding Phosphatidylinositol-4-Phosphate 5-Kinase is Induced by Water Stress and Abscisic Acid in Arabidopsis Thaliana. *Plant J.* 1998;15:563–568. DOI: 10.1046/j.1365-313X.1998.00227.x
- [35] Lang F. Mechanisms and significance of cell volume regulation. *J Am Coll Nutr.* 2007;26(5 Suppl):613S–623S. DOI: 10.2174/1389202911314030001
- [36] Taji T, Takahashi S, Shinozaki K. Inositols and Their Metabolites in Abiotic and Biotic Stress Responses. In: Majumder AL, Biswas, BB, editors. Spring;2006. p. 239–264. DOI: 10.1007/0-387-27600-9_10
- [37] Rammesmayer G, Pichorner H, Adams P, Jensen RG and Bohnert HJ. Characterization of IMT1, Myo-Inositol O-Methyltransferase, from Mesembryanthemum Crystallinum. *Arch Biochem Biophys.* 1995;322:183–188. DOI: doi:10.1006/abbi.1995.1450
- [38] Vernon DM, Bohnert HJ. Increased Expression of a Myo-Inositol Methyl Transferase in Mesembryanthemum Crystallinum is Part of a Stress Response Distinct from Crasulacean Acid Metabolism Induction. *Plant Physiol.* 1992;99:1695–1698. DOI: 10.1104/pp.99.4.1695
- [39] Sheveleva E, Chmara W, Bohnert HJ, Jensen RC. Increased Salt and Drought Tolerance by D-Ononitol Production in Transgenic Nicotiana Tabacum L. *Plant Physiol.* 1997;115:1211–1219. DOI: <http://dx.doi.org/10.1104/pp.115.3.1211>
- [40] Sengupta S, Patra B, Ray S, Majumder AL. Inositol Methyl Transferase from a Halophytic Wild Rice, Porteresia Coarctata Roxb. (Tateoka): Regulation of Pinitol Synthesis under Abiotic Stress. *Plant Cell Environ.* 2008;31:1442–1459. DOI: 10.1111/j.1365-3040.2008.01850.x
- [41] Zhu B, Peng RH, Xiong AS, Xu J, Fu XY, Zhao W, Jin XF, Meng XR, Gao JJ, Cai R, Yao QH. Transformation with a Gene for Myo-Inositol O-Methyltransferase Enhances the Cold Tolerance of Arabidopsis Thaliana. *Biol Plant.* 2012;56:135–139. DOI: 10.1007/s10535-012-0029-y
- [42] Arner RJ, Prabhu KS, Thompson JT, Hildenbrandt GR, Liken AD, Reddy CC. Myo-Inositol Oxygenase: Molecular Cloning and Expression of a Unique Enzyme That

- Oxidizes Myo-Inositol and D-Chiro-Inositol. *Biochem J.* 2001;360:313–320. DOI: 10.1042/0264-6021:3600313
- [43] Lorence A, Chevone BI, Mendes P, Nessler CL. Myo-Inositol Oxygenase Offers a Possible Entry Point into Plant Ascorbate Biosynthesis. *Am Soc Plant Biol.* 2004;134: 1–338. DOI: 10.1104/pp.103.033936.sapiens
- [44] Endres S, Tenhaken R. Myoinositol Oxygenase Controls the Level of Myoinositol in Arabidopsis, but Does Not Increase Ascorbic Acid. *Plant Physiol.* 2009;149:1042–1049. DOI: 10.1104/pp.108.130948
- [45] Alford SR, Rangarajan P, Williams P, Gillaspay GE. Myo-Inositol Oxygenase Is Required for Responses to Low Energy Conditions in Arabidopsis Thaliana. *Front Plant Sci.* 2012;3:1–11. DOI: 10.3389/fpls.2012.00069
- [46] Zhou ML, Zhang Q, Zhou M, Sun ZM, Zhu XM, Shao JR, Tang YX, Wu YM. Genome-Wide Identification of Genes Involved in Raffinose Metabolism in Maize. *Glycobiology.* 2012;22:1775–1785. DOI: 10.1093/glycob/cws121
- [47] Taji T, Ohsumi C, Iuchi S, Seki M, Kasuga M, Kobayashi M, Yamaguchi-Shinozaki K, Shinozaki K. Important Roles of Drought and Cold-Inducible Genes for Galactinol Synthase in Stress Tolerance in Arabidopsis Thaliana. *Plant J.* 2002;29:417–426. DOI: 10.1046/j.0960-7412.2001.01227.x
- [48] dos Santos TB, Budzinski IGF, Marur CJ, Petkowicz CLO, Pereira LFP, Vieira LGE. Expression of Three Galactinol Synthase Isoforms in Coffea Arabica L. and Accumulation of Raffinose and Stachyose in Response to Abiotic Stresses. *Plant Physiol Biochem.* 2011;49:441–448. DOI: 10.1016/j.plaphy.2011.01.023
- [49] Nishizawa-Yokoi A, Yoshida E, Yabuta Y, Shigeoka S. Analysis of the Regulation of Target Genes by an Arabidopsis Heat Shock Transcription Factor, HsfA2. *Biosci Biotechnol Biochem.* 2009;73:890–895. DOI: 10.1271/bbb.80809
- [50] Pennycooke JC, Vepachedu R, Stushnoff C, Jones ML. Expression of an A-Galactosidase Gene in Petunia Is Upregulated during Low-Temperature Deacclimation. *JASHS.* 2004;129:491–496.
- [51] Khokhar S, Apenten, RKO. Antinutritional Factors in Food Legumes and Effects of Processing. In: Squires VR, editor. *The Role of Food, Agriculture, Forestry and Fisheries in Human Nutrition.* 1st edn. EOLSS; 2003. p. 82–116.
- [52] Raboy V, Young KA, Dorsch JA, Cook A. Genetics and Breeding of Seed Phosphorus and Phytic Acid. *J Plant Physiol.* 2001;158:489–497. DOI: 10.1078/0176-1617-00361
- [53] Raboy V. Seeds for a Better Future: “Low Phytate” Grains Help to Overcome Malnutrition and Reduce Pollution. *Trends Plant Sci.* 2001;6:458–462. DOI: 10.1016/S1360-1385(01)02104-5

- [54] Kumar V, Sinha AK, Makkar HPS, Becker K. Dietary Roles of Phytate and Phytase in Human Nutrition: A Review. *Food Chem.* 2010;120:945–959. DOI: 10.1016/j.foodchem.2009.11.052
- [55] Thompson LU. Potential Health Benefits and Problems Associated with Antinutrients in Foods. *Food Res Int.* 1993;26:131–149. DOI: 10.1016/0963-9969(93)90069-U
- [56] Vuverick I, Shamsuddin AM. Cancer Inhibition by Inositol Hexaphosphate (IP6) and Inositol: From Laboratory to Clinic. *J Nutr.* 2003;133:3830–3836.
- [57] Shamsuddin AM. Anti-Cancer Function of Phytic Acid. *Int J Food Sci Technol.* 2002;37:769–782. DOI: 10.1046/j.1365-2621.2002.00620.x
- [58] Van den Ende W. Multifunctional Fructans and Raffinose Family Oligosaccharides. *Front. Plant Sci.* 2013;4:247. DOI: 10.3389/fpls.2013.00247

