

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Dissipative Solitons in Fibre Lasers

---

Vladimir L. Kalashnikov and Sergey V. Sergeyev

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/61713>

---

## Abstract

Interdisciplinary concept of dissipative soliton is unfolded in connection with ultrafast fibre lasers. The different mode-locking techniques as well as experimental realizations of dissipative soliton fibre lasers are surveyed briefly with an emphasis on their energy scalability. Basic topics of the dissipative soliton theory are elucidated in connection with concepts of energy scalability and stability. It is shown that the parametric space of dissipative soliton has reduced dimension and comparatively simple structure that simplifies the analysis and optimization of ultrafast fibre lasers. The main destabilization scenarios are described and the limits of energy scalability are connected with impact of optical turbulence and stimulated Raman scattering. The fast and slow dynamics of vector dissipative solitons are exposed.

**Keywords:** Ultrafast fibre laser, mode-locking, dissipative soliton, non-linear dynamics, vector solitons, optical turbulence, stimulated Raman scattering

---

## 1. Introduction

Over the last decades, ultrafast fibre laser technologies have demonstrated a remarkable progress. By definition [1–4], these technologies concern generation, manipulation and application of optical pulses from a fibre laser or a laser-amplifier system with (i) peak power  $P_0$  exceeding substantially an average laser power  $P_{ave}$  and (ii) pulse widths  $\tau$  which are much lesser than a laser round-trip period  $T$ . Such a definition can be re-interpreted in terms of a laser *mode-locking*, which means a phase-locked interference of the  $M$  laser eigenmodes producing an equidistant train of ultrashort pulses. Then,  $\tau \propto 1/M\delta\omega$  ( $M$  is a number of locked eigenmodes and  $\delta\omega$  is an inter-mode frequency interval defined by  $T \propto 1/\delta\omega$ ) and  $P_0 \propto MP_{ave}$  [5,6]. It means that a mode-locked laser generates a comb of equidistant optical frequencies comprising the broad spectral range  $\Delta \propto M\delta\omega$ . It is clear that the substantial enhancement of  $P_0$  (by the factor of  $M \propto 1/\tau\delta\omega \sim 10^5 \div 10^6$ , i.e. upto over-MW level [7,8]) and  $\tau$ -reduction ( $\sim T/M$ ,

i.e. down to sub-100 fs level [9]) promise an outlook for different applications [10] including non-linear and ultrasensitive laser spectroscopy [11–14], biomedical applications [15–19], micromachining [20,21], high-speed communication systems [22], metrology [23] and many others [24,25]. The extraordinary peak powers in combination with the drastic pulse width decrease bring the high-field physics on tabletops of a mid-level university lab [26–29]. Moreover, the over-MHz pulse repetition rates  $\delta\omega$  provide the signal rate improvement factor of  $10^3 \div 10^4$  in comparison with that of classical chirped-pulse amplifiers [26]. As a result, the signal-to-noise ratio enhances substantially, as well.

Another aspect of the ultrafast laser applications is connected with studying non-linear phenomena [30]. Ultrafast lasers became an effective platform for investigation of general non-linear processes such as instabilities and rogue waves [31,32], self-similarity [33] and turbulence [34]. A coherent self-organization in such non-linear systems [35,36] is the keystone of this review, and it will be considered below in detail. But here, we have to point at the multidisciplinary context of our topic. The ultrafast fibre lasers can be treated as an ideal playground for exploring of non-linear system phenomenology as a whole [37]. Such a playground spans gravity and cosmology [38], condensed-matter physics and quantum field theory [39–41], biology, neurosciences and informatics [42,43]. The advance of ultrafast laser technology is that the theoretical insights promise to become directly testable, controllable and, on the other part, the theory can be urged by new precise measurable experimental challenges.

To date, the solid-state lasers allowed generating shortest pulses with highest peak powers directly from an oscillator with high repetition rates ( $\delta\omega > 1$  MHz) [44–49]. The main advantages of solid-state laser systems are (i) broad gain bands (i.e. very large  $M$ ) allowing generation of extremely short pulses ( $\tau$  approaches one optical cycle for such media as Ti:Sp or Cr:chalcogenides) and (ii) covering the spectral range from visible (Ti:Sp) through infra-red (Ti:Sp, Cr:forsterite and Cr:YAG) to mid-infrared (Cr:chalcogenides) wavelengths, as well as (iii) possibility of independent and precise dispersion [50] and non-linearity [46,48] control. Nevertheless, fibre lasers have unprecedented prospects [51] due to (i) possibility of mean power scaling provided by large gain, (ii) high quality of laser mode, (iii) reduced thermo- and environment-sensitivity, (iv) compactness and integrity of laser setup. Additionally, one has to point at broader gain bands of fibre media in comparison with the energy-scalable, thin-disk, solid-state oscillators operating within analogous wavelength ranges [9,52]<sup>1</sup> and possibility to break into deep-UV and mid-IR optical spectral ranges [55,56].

In this review, we will concern the concepts of a mode-locking and a dissipative soliton in a nutshell.


## 2. Mode-Locking

The concept of mode-locking is universal and closely connected with a principle of synchronisation of coupled oscillators [57–64]. A laser is, in fact, the interferometer which possesses a

---

<sup>1</sup> The width of a gain band is not a decisive factor per se because both pulse width and its spectrum are affected by various factors including higher-order dispersions, non-linearity, etc. [53,54].

set of eigenmodes (longitudinal modes) separated by  $\delta\omega = 2\pi/T$ . Simultaneously, it is an *active* resonator, which means an amplification  $\Delta A$  of mode amplitude  $A$  during the resonator round-trip in the vicinity of the maximum gain frequency  $\omega_0$  as  $\Delta A(\omega) \approx (g(\omega_0) - \ell)A(\omega) - \alpha(\omega - \omega_0)^2 A(\omega)$ , where  $g(\omega_0)$  is a gain at the frequency  $\omega_0$ ,  $\ell$  is a net-loss coefficient and  $\alpha \propto g(\omega_0) / \delta\Omega^2$  takes into account a frequency-dependence of gain coefficient in the vicinity of  $\omega_0$  defined by a gain bandwidth  $\delta\Omega$ . In such an oversimplified model, only *one* mode with the maximum net-gain  $\sigma \equiv g(\omega - \omega_0) - \ell = 0$  is generated because the gain coefficient is energy-dependent, that is, it decreases with  $A$  (i.e. a gain is *saturable* that results in a mode competition or *mode selection*, Figure 1).



**Figure 1.** Comb of frequencies generated by a laser with the repetition rate  $\delta\omega$ . The gain band is centred at  $\omega_0$ . Mode-locking consists in excitation and synchronisation of  $\omega_0 \pm n\delta\omega$  sidebands.

In the case of *active mode-locking*, a periodic external modulation with the frequency of  $\delta\omega$  excites the  $\pm\delta\omega$  sidebands for each mode in the comb so that the modes  $A(\omega)$ ,  $A(\omega \pm \delta\omega)$  become coupled. In the framework of our oversimplified model, a steady-state regime  $\Delta A = 0$  is described by the equation in time domain [59–61]:

$$(g - \ell)A(t) + \alpha \frac{\partial^2 A(t)}{\partial t^2} - \nu t^2 A(t) = 0, \quad (1)$$

which is the classical equation for an oscillator in the potential defined by  $\nu \propto \delta\omega^2$ . This equation has a trivial solution in the form of a Gaussian pulse [59–63]:  $A(t) = A_0 \exp(-2\ln(2)t^2/\tau^2)$ , where the pulse amplitude  $A_0$  is defined by the condition of energy balance of  $\sigma \propto \nu\tau^2$  (the saturable gain coefficient is energy, i.e.  $A_0^2$ -dependent) and the pulse width  $\tau \propto \sqrt{\alpha/\nu}$ . In the general case, the excitation of  $A(t)$  in the form of Hermitian–Gaussian solutions of Equation (1) is possible. Since the pulse width is defined by  $\nu$  so that  $\tau \propto 1/\sqrt{\delta\Omega\delta\omega}$ , the minimum pulse widths of  $\sim 1/\delta\Omega$  are hardly reachable due to limitation imposed by  $\delta\omega$ -value. The situation can be changed in the presence of the self-phase modulation (SPM) [65] and the dynamic gain saturation. Then  $\tau \propto 1/\delta\Omega\sqrt{\sigma}$  [66].

Using the non-linear processes such as SPM, loss and gain saturation allows generating the ultrashort pulses due to mechanism of the so-called *passive mode-locking* [60]. Periodical perturbations caused by transitions through non-linear laser elements such as saturable absorber or gain medium enrich the spectrum with new components  $\omega_0 \pm m\delta\omega$  ( $m = 1, 2, \dots, M$ ), which becomes locked through non-linear interaction [64]:

$$\frac{\partial A(\omega_1, z)}{\partial z} = -\frac{i\pi\omega_1}{2cn(\omega_1)} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} d\omega_2 d\omega_3 d\omega_4 \chi^3(\omega_1, \omega_2, \omega_3, \omega_4) \times A(\omega_2, z) A(\omega_3, z) A(\omega_4, z) e^{i\Delta k z} \delta(\omega_1 - \omega_2 - \omega_3 - \omega_4), \quad (2)$$

where a four-wave non-linear process defined by non-linear susceptibility  $\chi^3$  mixes the frequencies  $\omega_1, \omega_2, \omega_3$  and  $\omega_4$  during the propagation through a non-linear medium along the  $z$ -coordinate ( $\Delta k = k(\omega_1) - k(\omega_2) - k(\omega_3) - k(\omega_4)$  is the difference of wavenumbers,  $c$  is the speed of light,  $n$  is the frequency-dependent refractive index).

Both active and passive mode-locking concepts can be easily united from the point of view of space-time duality [64,67–69]. For instance, let's consider *heat diffusion* equation:

$$\frac{\partial u}{\partial t} = \sigma u + \zeta \frac{\partial^2 u}{\partial x^2} - \nu x^2 u, \quad (3)$$

where a heat radiated at  $x = 0$  by the point source  $\sigma$  diffuses along  $x$ -axis and is absorbed by cooler with the parabolic 'cooling potential'. The replacements  $t \rightarrow z$  and  $x \rightarrow t$  result in an equation for 'diffusion' of light describing an *active amplitude mode-locking* (see Eq. (1)):

$$\frac{\partial A}{\partial z} = \sigma A + \alpha \frac{\partial^2 A}{\partial t^2} - \nu t^2 A. \quad (4)$$

Eq. (4) is clearly understandable in the Fourier domain:  $\frac{\partial \tilde{A}}{\partial z} = \sigma \tilde{A} + \nu \frac{\partial^2 \tilde{A}}{\partial \omega^2} - \alpha \omega^2 \tilde{A}$ , where an external (i.e. active) modulation defined by the  $\nu$ -coefficient 'diffuses' (i.e. broadens) a field spectrum  $\tilde{A}$  and such diffusion is compensated by spectral dissipation defined by the  $\alpha$ -coefficient. In the time-domain, the spectrum broadening corresponds to a light pulse shortening due to parabolic potential action which is balanced by spectral dissipation causing a pulse widening.

The space-time duality can be extended further with the help of *diffraction-dispersion* duality:

$$\frac{\partial A}{\partial z} = -\frac{i}{2k} \frac{\partial^2 A}{\partial x^2} \leftrightarrow \frac{\partial A}{\partial z} = \frac{i}{2} \beta_2 \frac{\partial^2 A}{\partial t^2}, \quad (5)$$

where  $k$  and  $\beta_2$  are the wave number and group-delay dispersion coefficients, respectively. Both processes describe the beam/pulse spreading with propagation which is accompanied by phase  $\phi$  profile distortion, i.e. by appearance of the *chirp*  $Q \propto d^2\phi / dx^2$  (or  $d^2\phi / dt^2$ ). The *active phase modulation* from this point of view

$$\frac{\partial A}{\partial z} = \sigma A + \alpha \frac{\partial^2 A}{\partial t^2} - i\nu t^2 A \quad (6)$$

(compare with (4)) looking as  $\frac{\partial \tilde{A}}{\partial z} = \sigma \tilde{A} + i\nu \frac{\partial^2 \tilde{A}}{\partial \omega^2} - \alpha \omega^2 \tilde{A}$  in the Fourier domain describes a ‘diffraction’ (dispersion) in the frequency domain inspired by phase modulator which is balanced by spectral dissipation. The main difference from (4) is that the phase modulation in (6) distorts the phase and thereby produces chirp like the action of thin lens but in the time domain. In other words, the phase modulation in (6) pushes the spectral components out of the point of stationary phase  $\partial\phi / \partial t = 0$ , adding the frequency shift  $\propto \nu t$  (Doppler shift) which enhances the spectral dissipation on the pulse wings and thereby forms a pulse like the active amplitude modulator. But the phase profile  $\phi(t)$  is parabolic in this case.

The transition to a *passive mode-locking* looks straightforward, but one has to be careful in this case. The space-time duality suggests a simple way to realize the temporal focusing like that in space domain: combination of phase modulation (‘time lens’) from Eq. (6) with dispersion (‘time diffraction’) from Eq. (5) allows compressing a pulse. Therefore, a replacement of time focusing by a time self-focusing (SPM) would provide a laser pulse self-trapping like the effect of laser beam self-trapping:

$$\frac{\partial A}{\partial z} = \frac{i}{2} \beta_2 \frac{\partial^2 A}{\partial t^2} - i\gamma |A|^2 A, \quad (7)$$

which is the famous *non-linear Schrödinger equation* describing propagation of optical solitons in a fiber ( $\beta_2 < 0$  corresponds to an anomalous dispersion,  $\gamma$  is a SPM-coefficient) [35,70,71].

It is appropriate to mention here that the space-time duality  $x \rightarrow t$  allows extending the physical context of consideration beyond scopes of optics. For instance,  $A$ ,  $E = \int dx |A|^2$ , and  $\phi$  can be related for a mean-field amplitude, number of particle (mass of condensate) and momentum (wave number) for a Bose–Einstein condensate [39]. Then, it is clear that the dispersion and SPM terms in Eq. (7) describe the kinetic energy and four-particle interaction potential for gas of bosons. Such an interpretation opens a road to a quantum theory of solitons [72–74].

Following the same procedure for Eq. (4), describing the active amplitude mode-locking results in the simplest version of equation for a passive mode-locking, so-called *cubic non-linear Ginzburg–Landau equation* [35,36,75]:

$$\frac{\partial A}{\partial z} = \sigma A + \alpha \frac{\partial^2 A}{\partial t^2} + \kappa |A|^2 A. \quad (8)$$

This equation describes a combined action of saturated net-gain ( $\sigma$ ), spectral dissipation ( $\alpha$ ) and non-linear gain ( $\kappa$ ). The last term results from loss saturation in a non-linear absorber with the response time much lesser than the pulse width. As will be shown below, such an assumption is valid for a broad class of fibre mode-locking mechanisms. Physics of passive mode-locking resembles that of active one: self-focusing in time domain causes a spectrum broadening which is balanced by spectral dissipation. Loss and energy-dependent gain are required for developing and stabilizing the mode-locking (all these factors are included in  $\sigma$ -term which is  $< 0$  for a steady-state pulse). Eqs. (7) and (8) have a similar solution  $A(t) \propto \text{sech}(t/\tau)$  but the mathematical structures of these equations differ substantially that created discrepancies between concepts of the ‘true’ [76] and *dissipative solitons* (DSs, see next section) [36]. Combining Eqs. (7) and (8) gives the famous complex cubic non-linear Ginzburg–Landau equation (cubic CNGLE) [35–37,42]:

$$\frac{\partial A}{\partial z} = i \left( \frac{\beta_2}{2} \frac{\partial^2 A}{\partial t^2} - \gamma |A|^2 A \right) + \sigma A + \alpha \frac{\partial^2 A}{\partial t^2} + \kappa |A|^2 A, \quad (9)$$

which is a playground for study of DSs. Equation (9) allows a number of further generalizations such as: (i) description of non-distributed evolution due to dependence of the equation coefficients on  $z$  [77]; (ii) generalization of non-linearity type aimed first of all to adequate description of different mode-locking mechanisms (see below); (iii) taking into account the higher-order dispersions, i.e.  $\omega$ -dependence of  $\beta_2$  [68]; (iv) taking into account the vector nature of light, i.e. transition to a system of coupled two-component CNGLEs [68,78–81], etc.


Now let us consider the mode-locking mechanisms for fibre lasers in more detail. Active mode-locking can be utilized for DS generation from a fibre laser [82–84], but the widespread mechanism is based on the *non-linear polarization rotation* (NPR) which uses the effect of intensity-dependent polarization mode coupling in a fibre [85–88]. There is voluminous literature concerning the experimental realization of NPR mode-locking in fibre lasers; therefore, our selection of references is rather subjective and concerns the DS context [95–111].

It is known [70] that an ideal single-mode fibre supports two degenerate orthogonally polarized modes. However, a real fibre has inherent birefringence caused by core asymmetry or mechanical stress (Figure 2).

Since SPM as well as cross-phase modulation (XPM) contribute to refractivity index with the strength defined by field intensity, such a contribution will change the state of polarization (SOP, Figure 3) [60,70,89] that can be described by coupled equations for two orthogonal ( $x$  and  $y$ ) polarization components [70]:



**Figure 2.** Fibre cross-sections with cylindrical symmetry broken by manufacturing process that leads to fibre birefringence.



**Figure 3.** Block scheme of a self-amplitude modulator (SAM) based on NPR. Both linear birefringence and NPR contribute to the change of state of polarization (SOP, rotation angle  $\theta$ ) that leads to intensity-dependent transmission of this scheme.

$$\begin{aligned}
 & \frac{\partial A_x}{\partial z} + \beta_{1x} \frac{\partial A_x}{\partial t} - \frac{i\beta_2}{2} \frac{\partial^2 A_x}{\partial t^2} - \sigma A_x - \alpha \frac{\partial^2 A_x}{\partial t^2} = \\
 & = -i\gamma \left( |A_x|^2 + \frac{2}{3} |A_y|^2 \right) A_x - \frac{i\gamma}{3} A_x^* A_y^2 e^{-2i\Delta\beta z}, \\
 & \frac{\partial A_y}{\partial z} + \beta_{1y} \frac{\partial A_y}{\partial t} - \frac{i\beta_2}{2} \frac{\partial^2 A_y}{\partial t^2} - \sigma A_y - \alpha \frac{\partial^2 A_y}{\partial t^2} = \\
 & = -i\gamma \left( |A_y|^2 + \frac{2}{3} |A_x|^2 \right) A_y - \frac{i\gamma}{3} A_y^* A_x^2 e^{2i\Delta\beta z},
 \end{aligned} \tag{10}$$


where the dissipative factors from Eq. (9) are taken into account and  $\Delta\beta = 2\pi / L_b$  describes a ‘strength’ of linear birefringence ( $L_b$  is a beat-length). As was shown in [81,91–94,179], the multi-scale averaging technique allows reducing Eq. (10) to the modified scalar non-linear Ginzburg–Landau equation (so-called *sinusoidal Ginzburg–Landau equation*) in which the self-amplitude modulation term (SAM, last term in Eq. (9)) is replaced by  $\propto \log Q(|A|^2)A$ , where  $Q$  is a complex function defined by birefringence and settings of laser wave plates and polarizer.



Such an approach opens a way to multi-parametrical optimization of fibre lasers mode-locked by NPR.

Despite its relative simplicity in principle as well as possibility of all-fibre-integrity of a laser, NPR in the form presented in Figure 3 is too sensitive to laser setup, uncontrollable perturbations and requires a precise manual tuning. The modified SAM setup, which can utilize both NPR and scalar SPM, is shown in Figure 4. It is the so-called *non-linear optical loop mirror* or *figure eight laser* (Figure 4) [112,113,265]. In principle, this setup is an all-fibre realization of additive-mode-locking [82,114] with inherently adjusted linear optical propagation lengths for counter-propagating beams. The main control parameter here is the beam splitting ratio  $\rho$  controlling the mutual intensities of counter-propagating beams.

The unique property of this SAM setup is its ability to utilize different types of non-linearities for mode-locking (e.g. see [115–118]). Different modifications of this mode-locking mechanism have been used in DS fibre lasers [119–125]. Nevertheless, a fibre loop defining SAM remains environment- and tuning-sensitive.



**Figure 4.** Block-scheme of SAM based on a non-linear optical loop mirror. Splitter splits input laser beam into two counter-propagating ones (red and green arrows) with some splitting ratio  $\rho$ . The beams interfere after round-trip and partially return into a laser. The result of interference is intensity-dependent due to NPR or/and SPM within a loop.

There is a class of alternative approaches utilizing non-fibre well-controllable non-linearities for mode-locking by the cost of broken fibre-integrity of a laser. Such an alternative was provided by development of high-non-linear *semiconductor saturable absorber mirror* (SESAM) [126–135]. The point is to put a semiconductor layer into a composed multi-layer mirror with the well-controllable spectral characteristics as well as with the adjustable intensity concentration of penetrating field within a semiconductor layer. In fact, it is an advanced non-linear Fabry–Perot interferometer with the reflectivity coefficient depending on the incident intensity (or energy) [136]. Interaction of light with a semiconductor layer can be characterized roughly as excitation of carriers from a valence band of semiconductor to its conduction band. Excited carriers thermalize inside a conduction band with the character time  $\sim 100$  femtoseconds. This

time defines a fastest response of SESAM to a laser radiation. Then, the thermalized carriers can relax into valence band or intra-band trapping states with the characteristic times from picoseconds to nanoseconds. Thus, SAM due to SESAM is *slow* in comparison with that due to NPR or SPM because the response times of the lasts are defined by intra-atom polarization dynamics, i.e. these times belong to femtosecond diapason. Additionally, the *spectral diapason* of SESAM response is substantially squeezed in comparison with that of pure electronic non-linearities due to resonant character of SESAM non-linearity. This can trouble the mode-locking within a spectral range exceeding the SESAM bandwidth. But the reverse side of the SESAM-band squeezing is that a non-linear response of SESAM becomes resonantly enhanced. This means that SESAM can provide more easily starting, stable and controllable mode-locking. The key characteristics of SESAM are [127] *loss saturation fluence*  $E_s = hv / 2\sigma_a$  ( $hv$  is a photon energy,  $\sigma_a$  is an absorption cross section), *modulation depth*  $\mu_0 = \sigma_a N$  ( $N$  is a density of states in semiconductor), *relaxation (recovery) time*  $T_r$ , *unsaturable loss*, *saturable loss bandwidth* and level of *two-photon absorption*.

Akin mode-locking methods providing full fibre-integrity, broadband absorption, sub-picosecond response time and avoiding a complex multi-layer mirror weaving use nanotube and graphene saturable absorbers [30,137–143] and other low-dimensional structures [144].

From the theoretical point of view, the response of saturable absorber (SESAM or other quantum-size structures) to a laser field can be very complicate. In principle, one has to take into account finite loss bandwidth, its dispersion, dependence of refractive index on carrier's (or exciton's) density (so-called linewidth enhancement), complex kinetics of excitation and relaxation, etc. However, the praxis demonstrated that a simple model of two-level absorber is well working [145]:

$$\frac{d\mu}{dt} = -\frac{\mu - \mu_0}{T_r} - \mu \frac{|A|^2}{E_s} \quad (11)$$

with some possible modifications (e.g. see [146]). Since DSs, as a rule, have over-picosecond widths (see next section), one may use an adiabatic approximation for (11) so that the expression for SAM coefficient in the last term in Eq. (9) has to be replaced:

$$\kappa \leftrightarrow (-1) \times \mu = \frac{\mu_0}{1 + |A|^2 / P_s}, \quad (12)$$


where  $P_s = E_s / T_r$  is a loss saturation power.

One may propose a hypothesis that an analogue of Kerr-lens mode-locking, which is a basic mechanism for generation of femtosecond pulses from solid-state lasers [60,85,147], can be realized in a fibre laser as well. Such an insight is based on possible enhancement of the laser beam spatial-trapping induced by non-linearity in a medium with spatially inhomogeneous

gain/loss or refractivity [148–152]. The model for analysis of such phenomena can be based on extension of dimensionality of Eq. (9), with taking into account the diffraction and transverse inhomogeneity of gain, loss or/and refractive index (the last can work as SAM due to the waveguide leaking loss) [153]:

$$\frac{\partial A}{\partial z} = i \left( \frac{\beta_2}{2} \frac{\partial^2 A}{\partial t^2} - \frac{1}{2k} \frac{\partial^2 A}{\partial x^2} - \gamma |A|^2 A \right) + \sigma A + \alpha \frac{\partial^2 A}{\partial t^2} - \kappa x^2, \quad (13)$$

where cylindrical symmetry is assumed,  $x$  is a radial coordinate and  $\kappa$  is a coefficient (complex in general case) which describes a transverse inhomogeneity of a fibre. Figure 5 shows the net-gain profiles (a) and the intra-laser pulse energies (b) as function of an effective aperture size obtained on the basis of variational approach for Eq. (13) [153]. The results demonstrate a principal feasibility of the Kerr-lens mode-locking regime for a DS fibre laser.



**Figure 5.** (a) Transverse net-gain profiles for different transverse parabolic distributions of net-gain coefficient which can be realized by inhomogeneous doping of fiber or by impact of waveguide leaking loss. (b) The dependence of intra-laser DS energy for a DS Yb-fibre laser with the 14 nm filter bandwidth and the average GDD of 330 fs<sup>2</sup>/cm [153]. DS collapses for large energies and cannot start for large aperture sizes (here SAM has inverse sign and the continuous-wave generation prevails).

All these mode-locking techniques are realizable for both soliton proper and DS fibre lasers (excluding the Kerr-lens mode-locking which requires sufficiently high pulse energies provided by only a DS laser). Now let's consider the DSs fibre lasers proper.

### 3. DS concept: Theory and experiment

A 'classical' soliton can be formally defined as a solution of non-linear evolution equation belonging to discrete spectrum of the inverse scattering transform [71,76,154]. The non-linear


equations, which can be solved by inverse scattering transform, are ‘exactly integrable’. This means that they are akin to linear equations in some sense. In particular, they obey the superposition principle and, as a result, can be canonically quantized [155,156]. One has to note that integrability of a non-linear evolution equation and non-dissipative (non-Hamiltonian) character of the latter are not equivalent because there are both non-integrable Hamiltonian systems and integrable dissipative ones [36]. The point is that the DS concept is not connected with ‘integrability’; therefore, DSs are not ‘true’ solitons in a mathematical sense. However, many properties of DSs, in particular, their stable localization, robustness in the processes of scattering and interaction, well-organized internal structure, etc., resemble the properties of ‘true’ solitons. Formally, one may define *DS* as a *localized and stable structure emergent in a non-linear dissipative system far from the thermodynamic equilibrium* [36]. DSs are abundant in the different natural systems ranging from optics and condensed-matter physics to biology and medicine. In this sense, one may paraphrase that DSs “are around us. In the true sense of the word they are absolutely everywhere” [157]. Therefore, the concept of DS became well established in the last decade [36,37,42,158].

Stability of a DS under condition of strong non-equilibrium can be achieved only due to well-organized energy exchange with environment and subsequent energy redistribution within a DS. It results in energy flux inside a DS and, thereby, in DS phase inhomogeneity [36]. For a simplest case of Eq. (9), which has a DS solution in the form of  $A(t)=A_0\text{sech}(t/\tau)^{1+i\psi}$  ( $\psi \propto Q\tau^2 = \tau^2 \partial^2 \phi / \partial t^2$  is a dimensionless chirp parameter) [85,159,160], the DS energy generation [36]

$$E = 2\sigma|A|^2 + 2\kappa|A|^4 - 2\alpha \left| \frac{\partial A}{\partial t} \right|^2 + \alpha \frac{\partial^2 |A|^2}{\partial t^2} \quad (14)$$


as well as the spectrum  $|\tilde{A}(\omega)|^2$  are shown in Figure 6 in dependence of  $\beta_2$  (the data are based on an approach of [160]). One can see that the spectrum broadening transfers the action of spectral dissipation on a pulse ‘in whole’ into well-structured energy exchange: inflow at pulse centre and outflow on its wings. The key characteristic of a dissipation inhomogeneity is a *chirp*, i.e. an inhomogeneity of phase. In absence of the chirp, the spectral dissipation acts on a pulse in whole that, in particular, induces a multi-pulse instability [161]. However, a power-dependent chirp causes inhomogeneity of energy transfer (Figure 7). Energy flows in the region closer to central wavelength where the gain is maximal. This region is located in the vicinity of pulse maximum. Energy flows out from the spectrum wings which are located on the wings of pulse, that is, the pulse localization is supported by spectral dissipation through non-linear mechanism of chirping [42,160,162]. One has to note that a direction of energy fluxes inside a DS depends on parameters and can be inversely related to the direction shown in Figures 6,7 (i.e. energy can flow from wings to centre). The corresponding structure was named *dissipative anti-soliton* [210].

Thus, an additional mechanism of SAM (in addition to mechanisms considered in the previous section) appears, which provides unique robustness of DSs (i.e. DS exists within a broad range of laser parameters [163,164]).



**Figure 6.** (a) Profile of energy generation and (b) logarithm of spectral power in dependence on GDD (normal dispersion range) for a DS of [160].

Below, we will consider a chirp as the essential characteristic of DS [210]. One of the reasons is that the chirp allows DS to accumulate energy  $E \propto \psi$ , which means that DS is *energy-scalable* [37,164–167]. The last statement does not mean that a chirp-free pulse is not energy-scalable. However, the energy-scalability of such pulses can be provided by only fine-tuned and separated control of SPM and GDD that can be achieved in solid-state oscillators [26,168] or in large mode area (LMA) fibre lasers [169]. For fibre lasers such an approach entails issues of full-fibre integrability, higher-order mode control [170,171]<sup>2</sup> and thermo-effects impact [172].



**Figure 7.** (a) Profile of energy generation and (b) power in dependence on chirp for a DS of Eq. (9):  $\sigma = -0.01$ ,  $\kappa = 0.025$ ,  $\alpha = 0.05$ ,  $A_0 = 1$ ,  $\tau = 1$ .

<sup>2</sup> However, namely LMA and photonic-crystal fibres could realize a Kerr-lens mode-locking in a fibre laser [152,153].

In the terms of space-time duality (see above), the mechanism of formation of time window, within which a DS is localized, resembles a phenomenon of total internal reflection from some 'borders' created by phase discontinuity. Such borders are formally defined by the equivalence of the wave number of out-/in-going radiation  $k(\omega) = \beta_2 \omega^2 / 2$  (wave number of dispersive linear wave) and the DS wave number  $q = \gamma P_0$ :  $k(\pm\Delta) = q$ , where  $P_0 = |A_{\max}|^2$  is a DS peak power and a DS spectral width is  $\Delta = \sqrt{2\gamma P_0 / \beta_2}$  [173]. Since a system is dissipative, the above phase equilibrium has to be supplemented by loss compensation condition: spectral loss  $\propto \alpha \Delta^2$  has to be compensated by non-linear gain  $\kappa P_0$ . Combination of above criteria gives a definition of the parametric limits for DS [44,173]:


$$\frac{2\alpha\gamma}{\beta_2\kappa} \leq \begin{cases} 2/3, E \rightarrow \infty \\ 2, E \rightarrow 0 \end{cases} \quad (15)$$

where  $E$  is a DS energy. Eq. (15) is valid for the *cubic-quintic CNGLE* in which SAM has a form of  $(\kappa |A|^2 - \kappa\zeta |A|^4)A$ , that is, a non-linear gain is saturable (despite the unsaturable SAM in Eq. (9)). A SAM saturability is necessary for DS stabilization [174]. Equality in Eq. (15) corresponds to the DS stability border where  $\sigma = 0$  (see Eq. (9)). The asymptotic  $E \rightarrow \infty$  corresponds to a *perfectly energy-scalable* DS or to a phenomenon of *dissipative soliton resonance* (DSR) [37,44,165–167,175–183], which is sufficiently robust, exists in different SAM environments and even within the anomalous GDD range [178,182]. Important property of DSR is that the DS energy  $E$  can be scaled without loss of stability by plain scaling of laser average power or/and its length  $L$  [44,180,184]. The chirp scales with length as well. As a result, the DS peak power and spectrum width tend to a constant for fixed parameters of Eq. (9) (i.e. fixed  $\alpha, \beta_2, \sigma$  and  $\kappa$ ) and the energy scaling is provided by DS stretching in time domain.

This ideology of energy scaling by the pulse stretching goes back to the so-called wave-breaking-free or stretched pulse fibre lasers where the propagation within the anomalous-dispersion fibre sectors alternates with the propagation under normal GDD action [96,101,185–187]. As a result of pulse stretching, the non-linear effects in such systems are reduced, which allows increasing an energy and suppressing a noise. As an alternative approach, one can exclude an anomalous GDD at all and to realize a so-called *similariton* regime, when a pulse accumulates an extremely large chirp and, thereby, an energy [33,103,188]. However, a self-similar regime is not soliton-like one in nature; therefore, we will focus on the all-normal-dispersion fibre lasers (ANDi) which produces DSs possessing a high stability within a broad range of laser parameters [97,101,185,186,189,190]. In Figure 8, the energy-scalable DS lasers are sub-divided into three main types: (1) *all-fibre*, (2) *fibre with a free-space sector* and (2) LMA including rod and photonic-crystal fibre PCF.

The advantage of the *first type* of lasers is their integrity, which does not require an operational alignment, includes potentially compressing and delivering sections, and provides environment insensitivity and easy integrability with fibre-amplifier cascades [200]. The last advantage is especially attractive because it allows a direct seeding of DS into chirp-pulse amplifier

without preliminary pulse stretching. The *second type* of the DS fibre lasers can be considered as a testbed for development of the first type. No wonder that the results achieved here are more impressive (Figure 8). At last, the *third type* of the DS fibre lasers is most akin to the thin-disk solid-state ones with simultaneous advantage of the broad gainbands. Such lasers provide the DS energy scalability by scaling of laser beam area in combination with the scaling of laser period and average power. Nevertheless, one has to keep in mind that both LMA and PCF technologies have some disadvantages (see above) which make them similar to solid-state lasers.



**Figure 8.** Experimental realizations of DSs in fibre lasers (Refs. [7-9,82,83,100,102,105-107,109-111,122,124,125,130,132,135,138,139,140,141,187,191-209,232]; Ref. [111] corresponds to a DS in the anomalous GDD region).

The diversity of the results obtained (Figure 8) needs a comprehension from a *unified viewpoint*; therefore, let's survey briefly some theoretical aspects relevant to the DS fibre lasers. There is vast literature regarding the theory of DSs. Some preliminary systematization can be found in [44]. However, it is necessary at first to declare the stumbling block of this theory: *absence of a unified viewpoint*. There exist unbroken walls between the circles of scientific community exploiting and exploring the DS concept: walls between the solid-state and fibre laser representations of the theory, condensed-matter one, numerical and analytical approaches, etc. Briefly and conditionally, the relevant theoretical approaches can be divided into (1) numerical, (2) exact analytical and (3) approximated analytical. The last includes the models based on (3.1) perturbative, (3.2) adiabatic models (AM) as well as those based on (3.3) phase-space truncation (i.e. variational approximation (VA) and method of moments (MM)).

As was emphasized repeatedly, both linear and non-linear dissipations are crucial for the DS formation. The simplest and most studied models for such a type of phenomena are based on the different versions of CNGLE (e.g. Eq. (9)).

Extensive *numerical study* of DSs of the cubic-quintic CNGLE has been carried out by N. N. Akhmediev with co-authors [35–37,42,157,166,175,176,178,182]. The simulations have allowed finding the DS stability regions for some two-dimensional projections of CNGLE parametrical space. The summarizing description of the results obtained is presented in [44]. Most impressive results are: (i) parametric space of DS has a reduced dimensionality resulting, in particular, in the appearance of DSR; (ii) DSR remains in a model with lumped evolution that is typical for the most of fibre lasers; (iii) DSR and, correspondingly, DS exist within the anomalous GDD region as well. However, the main shortcomings of the numerical approaches are: the parametrical space under consideration is not physically relevant, and the true dimensionality of DS parametric space is not identified. It is clear that the only advanced and self-consistent *analytical* theory of DS would provide, in particular, a true representation of DS parametric space and DSR conditions.

As was mentioned above, the evolution equations describing DSs are not-integrable. The efforts based on the algebraic techniques [62,213,214] and aimed to finding the generalized DS solutions of CNGLE were not successful to date. Nevertheless, few exact partial DS-solutions are known. For instance, *sole known exact analytical DS-solution* of cubic-quintic CNGLE is [110,166,176,182,189,211,212]:


$$A = \sqrt{\frac{A_0}{\cosh(t/\tau) + B}} \exp\left[\frac{i\psi}{2} \ln(\cosh(t/\tau) + B) + i\phi z\right], \quad (16)$$

where  $A_0$ ,  $B$ ,  $\tau$ ,  $\psi$  and  $\phi$  are real constants [189]. This solution belongs to a fixed-point solution class, which means that it exists only if some constraints are imposed on the cubic-quintic CNGLE parameters. Solution (16) provides with important insights into properties of DSs. In particular, the systematical classification of DS spectra (truncated concave, convex, Lorentzian and structured spectra) and DS temporal profiles (from *sech*-shaped  $A(t)$  to tabletop one) is possible in the framework of analytical approach. The transition to a DSR-regime reveals itself in the ‘time-spectral’ duality shown in Figure 9 [141,189]. The sense of this ‘duality’ will be explained below from the point of view of adiabatic theory of DSs.

The crucial shortcoming of the approach based on few exact DS solutions of evolution equations is that the strict restrictions are imposed on the equation parameters. As a result, the DS cannot be traced within a broad multidimensional parametric range and the picture obtained is rather sporadic and is of interest only in the close relation with the numerical results and experiment. Some additional information can be obtained on the basis of perturbation theory which provides with a quite accurate approximation for a low-energy DS [215–217].


Most powerful approaches to the theory of DSs have been developed in the framework of *approximated techniques* (for review see [44]): AM [165,167,217–221], VA [77,177,222–224,225]





**Figure 9.** ‘Time-spectral duality’ for an energy-scalable DS: bell-like time-profile (blue curve on the left picture) transforms into tabletop one with truncated edges (right) and, inversely, truncated tabletop spectrum (red curve on the left picture) transforms into bell-like one (right) with the DS energy growth.

and MM [175,210,222,226]. The most impressive results obtained are: (i) physically relevant representation of DS parametric space was revealed (it is a so-called *master diagram*, see [44] for review and Figure 10); (ii) such a representation allows understanding the structural properties of DS and its energy-scaling laws (i.e. DSR conditions) for different mode-locking techniques; (iii) DS dynamics and an issue of optimal arrangement of laser elements providing the maximum DS stability and energy have been explored [224,225,227,228]; (iv) vectorial extension of VA concerning a vector DS (VDS) was endeavoured [229].



**Figure 10.** DS master diagrams for the cubic-quintic CNGLE (black) and the CNGLE with a SAM defined by Eq. (12) (red) (in the last case  $\kappa=1/\gamma P_s$ ). DSR ranges correspond to a so-called *positive branch* of DS [44,167], which has a highest DS energy-scalability and stability [221]. The dashed curve corresponds to a DS stability border obtained from numerical simulations of cubic-quintic CNGLE taking into account a quantum noise [243]. Dot blue curve shows the DS border under effect of SRS [260]. The parametrical space shown is the *physically relevant* parametrical space of DS. Points correspond to different scenarios of DS destabilization (see text).

Both AM and VA demonstrate two-dimensional representation of DS parametric space in the form of master diagram. Dimensionality can grow with complication of CNGLE non-linearity when SPM becomes saturable so that the cubic non-linear term in Eq. (9) has to be replaced by  $[(\kappa - \kappa\zeta |A|^2) - i(\gamma + \chi |A|^2)] |A|^2 A$  [217]. This effect can appear in a fibre laser with NPR (e.g. see [179] where such a completely cubic-quintic CNGLE is connected with the NPR mode-locking technique). In this case, DS soliton exists in both normal and anomalous GDD regions [175,182,233].

The master diagram is a manifold of isogains (i.e. curves with  $\sigma = \text{const} \leq 0$ ). Figure 10 demonstrates the zero-isogains ( $\sigma = 0$ ) corresponding to the DS stability limit (upper curves) as well as the borders between ‘energy-scalable’ and ‘energy-non-scalable’ branches of DS (lower curves, see [44,167] for a formal definition<sup>3</sup>). Figure 10 demonstrates that the saturation of SAM (so-called reverse saturable absorber provided, for instance, by NPR or graphene [141]; black curves) enhances the DS stability in comparison with an unsaturable SAM (red curves). Since  $\lim_{C \rightarrow \text{const}} E = \infty$  ( $\text{const} \in$  the DSR range and its maximum value are of  $2/3$ ) for a saturable SAM (cubic-quintic CNGLE, black curves in Figure 10). Such a property of isogain curves corresponds to the DSR phenomenon. Since the stability threshold is defined by the condition of  $C \equiv 2\alpha\gamma / \beta_2\kappa \leq 2/3$ , one may conclude that the broadening of spectral filter band (or gainband) enhances stability against multi-pulsing ( $\alpha \propto 1/\delta\Omega^2$ , see above) [107,108]. Simultaneously, SPM has to be balanced by GDD ( $C \propto \gamma / \beta_2$ ) that, in combination with normalization of energy, gives the energy-scaling law  $E \propto L$  along a DSR curve. VA predicts [230]:

$$E \propto \beta_2 / \sqrt{\alpha} \tag{17}$$

which agrees with experimental observations of linear growth of DS energy with bandwidth [107,108] as well with a rule  $E \propto L$  since  $\beta_2 \propto L$  in the frameworks of distributed CNGLE.

In the case of unsaturable SAM corresponding to SESAM, some nanotube and graphene absorbers, Kerr-lensing, etc. (see Eq. (12)), the energy scaling requires scaling of the control parameter  $C$ . In this case, the asymptotic *energy scaling law* for  $E\kappa^2 / \gamma\sqrt{\alpha} \gg 1$  becomes [44]:

$$E \propto \beta_2 / C\sqrt{\alpha}. \tag{18}$$

The spectral properties of DS are described clearly in the frameworks of AM [44,167]. In the simplest case of cubic-quintic CNGLE, the DS spectrum  $\tilde{A}(\omega)$  in the limit of  $|\psi| \gg 1$  is a Lorentzian profile which has a characteristic width  $\Omega_L$  and is truncated at frequencies  $\pm\Delta$  [44,167,218]:

$$\tilde{A}(\omega) \propto \frac{H(\Delta^2 - \omega^2)}{\Omega_L^2 + \omega^2}, \tag{19}$$

<sup>3</sup> Energy-non-scalable branch has two distinguishing characteristics: it turns into solution of Eq. (9) with  $\zeta, \chi \rightarrow 0$  (‘Schrödinger limit’ [218]) and is unstable in absence of dynamic gain saturation, i.e. if  $\sigma$  is not energy-dependent [221].

where H is a Heaviside function. The DS energy is

$$E \propto \arctan(\Delta/\Omega_L)/\Omega_L, \quad (20)$$

and

$$\begin{aligned} \gamma|A(t)|^2 &= q - \frac{\beta_2}{2} \left( \frac{d\phi(t)}{dt} \right)^2, \\ q &= \gamma P_0 = \frac{\beta_2}{2} \Delta^2 = \frac{3\gamma}{2\zeta} \left( 1 - \frac{C}{2} \right), \\ \frac{\beta_2}{2} \Omega_L^2 &= \frac{\gamma}{\zeta} (1+C) - \frac{5}{3} \gamma P_0, \end{aligned} \quad (21)$$

Here, we trace the zero-isogain  $\sigma = 0$ . The DS time-profile is defined by an implicit expression:

$$t = \tau \left[ \operatorname{arctanh} \left( \frac{d\phi/dt}{\Delta} \right) + \frac{\Delta}{\Omega_L} \arctan \left( \frac{d\phi/dt}{\Delta} \right) \right], \quad (22)$$

with the DS width of  $\tau \propto \Delta^{-1}(\Delta^2 + \Omega_L^2)^{-1}$ . Now, there are the following limiting cases:

$$\lim_{C \rightarrow 2} \begin{cases} E \rightarrow 0 \\ P_0 \rightarrow 0 \\ \Delta \rightarrow 0 \\ \Omega_L \rightarrow \text{const} \\ \tau \rightarrow 0 \\ \gamma|A|^2 = q - \beta\Delta^2 \tanh^2(t/\tau) \end{cases} \quad (23)$$

It is clear that in this 'low-energy' sector the DS time-profile is bell-like and its spectrum has tabletop form ( $\Omega_L \gg \Delta$ ). In the DSR limit, one has:

$$\lim_{C \rightarrow 2} \begin{cases} E \rightarrow \infty \\ P_0 \rightarrow 1/\zeta \\ \Delta \rightarrow \sqrt{\gamma/\beta\zeta} \\ \Omega_L \rightarrow 0 \\ \tau \rightarrow \text{const} \\ \gamma|A|^2 = q - \beta\Omega_L^2 \tan^2(t\Omega_L/\tau\Delta) \end{cases} \quad (24)$$

that is, a DS in the DSR sector has a flattop temporal profile and a Lorentzian spectrum ( $\Omega_L \ll \Delta$ ). Eqs. (24) demonstrate that asymptotical growth of DS energy leads to a spectral condensation ( $\Omega_L \rightarrow 0$ ) without a parallel temporal thermalization ( $\tau \not\rightarrow \infty$ ), which means an inevitable destabilization of a plain energy-scalability [180]. This conclusion does not mean a participial impossibility of DS energy-scaling in the frameworks of cubic-quintic CNGLE model. For instance, a saturable SPM allows DSs with tabletop profiles and  $|d\phi/dt| \rightarrow \infty$  on the pulse edges. Such a DS possesses enhanced energy scalability and was observed experimentally [231].

#### 4. DS spectrum and stability

As was explained, the dual balances in frequency domain:

$$\begin{cases} \beta_2 \Delta^2 \leftrightarrow \gamma P_0 \\ \alpha \Delta^2 \leftrightarrow \kappa P_0 \end{cases} \quad (25)$$

are formative for DS existence and stabilization. No wonder that the spectrum of DS is benchmark of its inherent properties.





Figure 11. Perturbed DS spectrum [241].

Prior to consider the aspects of interweaving of spectral and stability properties of DSs, one has to point to a possibility of multi-wavelength multi-pulsing DSs provided by DS robustness.

As was demonstrated in [234] theoretically, the multi-DSs compounds in a mode-locked laser can be stabilized at multiple frequencies. Experimentally, such multi-frequency DS compounds can be realized by birefringence filters with a periodical (interference-like) dependence of transmission on wavelength under conditions of sufficiently broad gainband and powerful pump [235–239]<sup>4</sup>.




**Figure 12.** Exploding DS corresponding to parameters of point A in Figure 10. Left: contour-plot of instant power, right: 3D-graph of instant power in dependence on local time  $t$  and propagation distance  $z$  (arbitrary units) [243].


As was demonstrated in previous section, DS has non-trivial internal structure due to energy fluxes inside it. The elements of this structure (*internal modes*) can be excited that causes pulsating or chaotic dynamics of DS with preservation of its temporal and spectral localization [241]. The spectral envelope acquires a shape of ‘glass with boiling water’ (Figure 11). Appearance of such perturbations is understandable in frameworks of the DS perturbation theory in spectral domain [242]. One has to note that such perturbations take a place inside the DS stability region where  $\sigma < 0$  (below the corresponding upper curves in Figure 10). Above the stability boarder (‘no DS’ region in Figure 10), there are three main destabilization scenarios [243]. For small energies and in the vicinity of stability border (point A in Figure 10), DS is *exploding* (Figure 12), which means its aperiodic disappearance with excitation of continuous waves and subsequent DS recreation [37,244–248]. With the energy growth (point B in Figure 10), the *rogue* DSs develop (Figure 13) [37,249–253]. Such a regime can be interpreted as DS structural chaotization, that is, generation of multiple DSs with strong interactions causing extreme dynamics.

For sufficiently large energies in the vicinity of stability border (point C in Figure 10), the typical destabilization scenario is the generation of multiple DSs (Figure 14). The source of this destabilization is the growth of spectral dissipation caused by DS spectral broadening with approaching to stability border so that the DS splitting becomes more energy advantageous [161]. Moreover, the DS splitting can be enhanced by its phase inhomogeneity because the gain (energy in-flow) is maximum at the points of stationary phase  $d\phi/dt = 0$  [254]. Such a splitting

<sup>4</sup> A multi-porting configuration of a DS laser supports even simultaneous generation of conventional and dissipative wavelength-separated solitons [240].



**Figure 13.** Rogue DSs corresponding to parameters of point *B* in Figure 10. Left: contour-plot of instant power, right: 3D-graph of instant power in dependence on local time  $t$  and propagation distance  $z$  (arbitrary units) [243].




**Figure 14.** DS molecule corresponding to parameters of point *C* in Figure 10. Left: contour-plot of instant power, right: 3D-graph of instant power in dependence on local time  $t$  and propagation distance  $z$  (arbitrary units) [243].

can result in an extreme dynamics like solitonic turbulence (Figure 13) [255] or regular multi-DS complexes (Figure 14, so-called DS molecules) which can have non-extreme internal dynamics (soliton gas) and interact with a background (soliton liquid) [256-259]. If the dynamic gain saturation (see [44]) contributes, the multi-DS complexes can evolve slowly into a set of equidistant DSs with repetition rate multiple of laser one (harmonic mode-locking) [158,159].

The numerical simulations of cubic-quintic CNGLE with taking into account a quantum noise validated the fact of inconsistency of spectral condensation and absence of temporal thermo-lization that breaks the DS energy scalability (see previous section) [243]. As a result, the DS stability region breaks abruptly with energy growth (dashed curve in Figure 10) and multitude of turbulent scenarios of DSs evolution develops (Figure 15) [34,243].


Serious limitations on power and energy scalability of DSs in fibre lasers arise from stimulated Raman scattering (SRS) [2,109]. The stability border of DS under action of SRS is shown in Figure 10 by dot blue curve (DS is stable on the left of this curve) [260]. As was found, SRS enhances the tendency to multi-pulsing with energy growth caused by enhancement of



**Figure 15.** DS turbulent regimes (contour-plot of instant power, arbitrary units) corresponding to the parameters of points  $D$ ,  $E$ ,  $F$ ,  $G$  and  $H$  in Figure 10 [243].

spectral dissipation due to SRS [260]. Simultaneously, generation of anti-Stokes radiation causes chaotization of DS dynamics and irregular modulation of DS temporal and spectral profiles [261] (Figure 16). DS profile remains localized, but it is strongly cut by colliding dark and grey soliton-like structures [34].

As was shown, the DS dynamics can be regularized by formation of *dissipative Raman soliton* (DRS). DRS can exist in the form of DS which is Stokes-shifted due to self-Raman scattering (Figure 17, *a*) [260] or as bound DS–DRS complex (Figure 17, *b*) [262]. In the last case, stabilization is achieved by feedback, i.e. reinjection of Stokes signal through a delay line [263]. In the absence of a feedback, the Raman pulse is noisy [264] (see dash line spectrum in Figure 16, *b*).



**Figure 16.** Wigner function of turbulent DS in the presence of SRS.




Figure 17. (a) Wigner function of single DRS [260] and (b) spectrum of bound DS-DRS complex [262].

## 5. Vector DSs


As was pointed above, SOP can play leading role in a fibre laser dynamics. In particular, it can contribute to mode-locking or/and spectral filtering. However, diapason of polarization phenomena in a DS fibre laser spreads essentially broader. As was found, intrinsic fibre birefringence (Figure 2) can lead to DS splitting into two independent SOPs [78]. This phenomenon is used to realize the NPR mode-locking mechanism where a DS SOP evolves (or remains locked) as a whole during propagation [265-269]. The polarization dynamics can be fast ( $\leq T$ ) or slow ( $\gg T$ ) and vary from regular (with possible period multiplication or harmonic mode-locking) to chaotic one [270,271]. There are evidences of ultrafast SOP evolution when SOP changes across a DS profile [272].

The specific multiple pulse instability of vector dissipative solitons (VDSs) leads to generation of the bound states of DSs with different SOPs (*vector soliton molecules*) which are locked by a non-linear coupling [273,274] or by a group-velocity locking produced by spectrum shift between DSs with different SOP [275]. As was shown experimentally (Figure 18) [276], the dynamics of VDS molecules can be highly non-trivial and demonstrate both fast and slow periodic switching between fixed SOPs as well as SOP procession, which is especially interesting for fibre laser telecommunications based on polarization multiplexing.

The important breakthrough in the recent theory of VDSs is the demonstration of insufficiency of approaches based on the coupled CNGLEs (like (10)) for adequate description of DS polarization dynamics. It was demonstrated that an active medium polarizability contributes to DS dynamics substantially [277]. As was shown, the SOP-sensitive interaction between DS and a slowly relaxing active medium with taking into account the birefringence of fibre laser elements and light-induced anisotropy caused by elliptically polarised pump field change the SOP at a long time scale that results in fast and slowly evolving SOPs of VDSs (Figure 19).


The non-trivial contribution of active medium kinetics and polarizability with taking into account the pump SOP and SPM demonstrates a complex dynamics including spiral attractors






**Figure 18.** Polarization dynamics of VDS molecules in an Er-fibre laser mode-locked by carbon nanotubes [276]. Top row demonstrates the evolution of Stokes parameters. The bottom row reproduces this evolution on the Poincaré sphere (each point corresponds to SOP after one laser round-trip).

and dynamic chaos (Figure 20) [278]. One may assume that such a non-trivial polarization dynamics is of great importance for DS energy scaling, in particular, due to vector nature of SRS [279]. These topics remain unexplored to date.



**Figure 19.** Polarization dynamics of VDS with taking into account an Er-fibre polarizability [277]. Both fast and slow SOP dynamics exist in dependence on fiber laser birefringence strength.



**Figure 20.** Dynamics of active fibre inversion components, which are polarization-sensitive ( $n_{12}$  and  $n_{22}$ ) and non-sensitive ( $n_0$ ), with the corresponding slow SOP evolution of VDS [278].

## 6. Conclusion

The recent progress in development of ultrafast fibre lasers and advances in exploring of DS are interrelated. DSs allowed scoring a great success in ultrashort pulse energy scalability that is defined by unprecedented stability and robustness of DS. At this moment, it is possible to achieve over-MW peak powers for sub-100 fs pulses directly from a fibre laser at over-MHz repetition rates. New spectral diapasons became reachable owing to development of mid-IR active fibres and using the frequency-conversion directly in a laser. Development of new mode-locking techniques, especially based on using of SESAMs, graphene and another quantum-sized structure allowed improving a laser stability, integrity and environment insensitivity. A great advance has been achieved in the theory of DSs. New powerful analytical techniques based on extensive numerical simulations and experimental advances extended understanding of the DS fundamental properties and revealed new prospects in improvement of characteristics of ultrafast fibre lasers. Based on achieved results, one may outline some unresolved problems. As was found, there are stability limits for a DS energy scaling imposed by optical turbulence and SRS. Deeper insight into the nature of these phenomena could allow to overcome these limits without substantial complication of laser setup. Simultaneously, control of intra-laser spectral conversion is a direct way to broadening of spectral range. Then, the dynamics and properties of VDSs remain scantily explored. Recent studies demonstrated a multitude of polarization phenomena, which cannot be grasped in frameworks of existing models. In particular, polarizability and kinetics of an active fibre in combination with birefringence of a laser in a whole can contribute non-trivially to a laser dynamics. As an additional aspect of further development, one may point at the development of new mode-locking techniques, which could improve DS stability and integrity of a fibre laser, decrease

pulse width and extend a diapason of pulse repetition rates. At last, one has to remember that a fibre laser is an ideal playground for study of complex non-linear phenomena and, undoubtedly, new bridges between different fields of science will be built with a further progress of ultrafast fibre lasers.

## Acknowledgements

This work was supported by FP7-PEOPLE-2012-IAPP (project GRIFFON, No. 324391).

## Author details

Vladimir L. Kalashnikov\* and Sergey V. Sergeyev

\*Address all correspondence to: v.kalashnikov@aston.ac.uk

AIPT, Aston University, B 7ET, Birmingham, UK

## References

- [1] Fermann ME, Hartl I. Ultrafast fiber laser technology. *IEEE J Sel Topics in Quantum Electron* 2009;15(1):191–206. DOI: 10.1109/JSTQE.2008.2010246.
- [2] Fermann ME, Galvanauskas A, Sucha G, Harter D. Fiber-lasers for ultrafast optics. *Appl Physics B* 1997;65:259–75.
- [3] Fermann ME, Hartl I. Ultrafast fiber lasers. *Nature Photonics* 2013;7:868–74. DOI: 10.1038/NPHOTON.2013.280.
- [4] Limpert J, Röser F, Schreiber Th, Tünnermann A. High-power ultrafast fiber laser systems. *IEEE J Sel Top Quantum Electron* 2006;12(2):233–44. DOI: 10.1109/JSTQE.2006.872729.
- [5] Lamb WE. Theory of an optical laser. *Phys Rev* 1964;134:A1429–50.
- [6] Siegman AE. *Lasers*. Sausalito: University Science Book; 1986. 1283 p.
- [7] Lefrançois S, Kieu K, Deng Y, Kafka JD, Wise FW. Scaling of dissipative soliton fiber lasers to megawatt peak powers by use of large-area photonic crystal fiber. *Opt Lett* 2010;35(10):1569–71.

- [8] Baumgartl M, Lecaplain C, Hideur A, Limpert J, Tünnermann C. 66 W average power from a microjoule-class sub-100 fs fiber oscillator. *Opt Lett* 2012;37(10):1640–2. DOI: 10.1364/OL.37.001640.
- [9] Chong A, Renninger WH, Wise FW. Route to the minimum pulse duration in normal-dispersion fiber lasers. *Opt Lett* 2008;33(22):2638–40. DOI: 10.1364/OL.33.002638.
- [10] Sucha G. Overview of industrial and medical applications of ultrashort pulse lasers. In: Fermann ME, Galvanauskas A, Sucha G. (Eds.) *Ultrafast Lasers: Technology and Applications*. New York: Marcel Dekker, Inc.; 2003. pp. 323–358.
- [11] Xu C, Wise FW. Recent advances in fibre lasers for nonlinear microscopy. *Nature Photonics* 2013;7:875–82. DOI: 10.1038/NPHOTON.2013.284.
- [12] Müller M, Squier J. Nonlinear microscopy with ultrashort pulse lasers. In: Fermann ME, Galvanauskas A, Sucha G. (Eds.) *Ultrafast Lasers: Technology and Applications*. New York: Marcel Dekker, Inc.; 2003. pp. 661–97.
- [13] Kalashnikov VL, Sorokin E. Soliton absorption spectroscopy. *Phys Rev A* 2010;81:033840. DOI: 10.1103/PhysRevA.81.033840.
- [14] Kalashnikov VL, Sorokin E, Sorokina IT. Chirped dissipative soliton absorption spectroscopy. *Opt Express* 2011;19(18):17480–92.
- [15] Kurtz RM, Sarayba MA, Juhasz T. Ultrafast lasers in ophthalmology. In: Fermann ME, Galvanauskas A, Sucha G. (Eds.) *Ultrafast Lasers: Technology and Applications*. New York: Marcel Dekker, Inc.; 2003. pp. 745–65.
- [16] Clowes J. Next generation light sources for biomedical applications. *Optic Photonics* 2011;3(1):36–8.
- [17] Fujimoto JG, Brezinski M, Drexler W, Hartl I, Kärtner F, Li X, Morgner U. Optical coherence tomography. In: Fermann ME, Galvanauskas A, Sucha G. (Eds.) *Ultrafast Lasers: Technology and Applications*. New York: Marcel Dekker, Inc.; 2003. pp. 699–743.
- [18] Drexler W, Fujimoto JG. (Eds.) *Optical Coherence Tomography*. Berlin: Springer-Verlag; 2008. 1346 p.
- [19] Lanin AA, Fedotov IV, Sidorov-Biryukov DA, Doronina-Amitonova LV, Ivashkina OI, Zots MA, Sun C-K, Ilday FÖ, Fedotov AB, Anokhin KV, Zheltikov AM. Air-guided photonic-crystal-fiber pulse-compression delivery of multimewatt femtosecond laser output for nonlinear-optical imaging and neurosurgery. *Appl Phys Lett* 2012;100:101104. DOI: 10.1063/1.3681777.
- [20] Osellame R, Cerullo G, Ramponi R. (Eds.) *Femtosecond Laser Micromachining*. Heidelberg: Springer; 2012. 483 p.

- [21] Gattass RR, Mazur E. Femtosecond laser micromachining in transparent materials. *Nat Photon*. 2008;2:219–25. DOI: 10.1038/nphoton.2008.47.
- [22] Nakazawa M. Ultrahigh bit rate communication system. In: Fermann ME, Galvanuskas A, Sucha G. (Eds.) *Ultrafast Lasers: Technology and Applications*. New York: Marcel Dekker, Inc.; 2003. pp. 611–660.
- [23] Udem Th, Holzwarth R, Hänsch TW. Optical frequency metrology. *Nature* 2002;416:233–7. DOI: 10.1038/416233a.
- [24] Liu Y, Tschuch S, Rudenko A, Dürr M, Siegel M, Morgner U, Moshhammer R, Ullrich J. Strong-field double ionization of Ar below the recollision threshold. *Phzs Rev Lett* 2008;101:053001. DOI: 10.1103/PhysRevLett.101.053001.
- [25] Sciaini G, Miller RJD. Femtosecond electron diffraction: heralding the era of atomically resolved dynamics. *Rep Prog Phys* 2011;74:096101. DOI: 10.1088/0034-4885/74/9/096101.
- [26] Südmeyer T, Marchese SV, Hashimoto S, Baer CRE, Gingras G, Witzel B, Keller U. Femtosecond laser oscillators for high-field science. *Nat Photon* 2008;2:599–604. DOI: 10.1038/nphoton.2008.194.
- [27] Krausz F, Ivanov M. Attosecond physics. *Rev Mod Phys* 2009;81(1):163–234. DOI: 10.1103/RevModPhys.81.163
- [28] Pfeifer T, Spielmann C, Gerber G. Femtosecond x-ray science. *Rep Prog Phys* 2006;69(2):443–505. DOI: 10.1088/0034-4885/69/2/R04.
- [29] Mourou GA, Tajima T, Bulanov SV. Optics in the relativistic regime. *Rev Mod Phys* 2006;78(2):309–71. DOI: 10.1103/RevModPhys.78.309.
- [30] Martinez A, Sun Z. Nanotube and graphene saturable absorbers for fibre lasers. *Nat Photon* 2013;7:842–5.
- [31] Lecaplain C, Grelu Ph, Soto-Crespo JM, Akhmediev N. Dissipative rogue waves generated by chaotic bunching in a mode-locked laser. *Phys Rev Lett* 2012;108:233901.
- [32] Dudley JM, Dias F, Erkintalo M, Genty G. Instabilities, breathers and rogue waves in optics. *Nat Photon* 2014;8:755–64.
- [33] Dudley JM, Finot Ch, Richardson DJ, Millot G. Self-similarity in ultrafast nonlinear optics. *Nat Phys* 2007;3:597–603.
- [34] Turitsyna EG, Smirnov SV, Sugavanam S, Tarasov N, Shu X, Babin SA, Podivilov EV, Churkin DV, Falkovich G, Turitsyn SK. The laminar–turbulent transition in a fibre laser. *Nat Photon* 2013;7:783–6. DOI: 10.1038/NPHOTON.2013.246.
- [35] Akhmediev NN, Ankiewicz A. *Solitons: Nonlinear Pulses and Beams*. London: Chapman & Hall; 1997.

- [36] Akhmediev NN, Ankiewicz A. (Eds.) *Dissipative Solitons*. Berlin: Springer-Verlag; 2005.
- [37] Grelu Ph, Akhmediev N. Dissipative solitons for mode-locked lasers. *Nat Photon* 2012;6:84–92. DOI: 10.1038/nphoton.2011.345
- [38] Faccio D, Belgiorno F, Cacciatori S, Gorini V, Liberati S, Moschella U. (Eds.) *Analogue Gravity Phenomenology. Analogue Spacetimes and Horizons, from Theory to Experiment*. Heidelberg: Springer; 2013. p. 438. DOI: 10.1007/978-3-319-00266-8.
- [39] Kevrekidis PG, Frantzeskakis DJ, Carretero-González R. (Eds.) *Emergent Nonlinear Phenomena in Bose-Einstein Condensates*. Berlin: Springer-Verlag; 2008.
- [40] Yang Y. *Solitons in Field Theory and Nonlinear Analysis*. New York: Springer; 2001.
- [41] Abdulaev FK, Konotop VV. (Eds.) *Nonlinear Waves: Classical and Quantum Aspects*. Dordrecht: Kluwer Academic Pub.; 2004.
- [42] Akhmediev NN, Ankiewicz A. (Eds.) *Dissipative Solitons: From Optics to Biology and Medicine*. Berlin: Springer-Verlag; 2008.
- [43] Naruse M, (Ed.) *Nanophotonic Information Physics*. Berlin: Springer-Verlag; 2014.
- [44] Kalashnikov VL. Chirped-pulse oscillators: route to the energy-scalable femtosecond pulses. In: Al-Khursan A. (Ed.) *Solid State Laser*. InTech, 2008; pp. 145–184. DOI: 10.5772/37415.
- [45] Baer CRE, Heckl OH, Saraceno CJ, Schriber C, Kränkel C, Südmeyer T, Keller U. Frontiers in passively mode-locked high-power thin-disk laser oscillators. *Opt Express* 2012;20:7054–65. DOI: 10.1364/OE.20.007054.
- [46] Saraceno CJ, Emaury F, Heckl OH, Baer CRE, Hoffmann M, Schriber C, Golling M, Südmeyer Th, Keller U. 275 W average output power from a femtosecond thin disk oscillator operated in a vacuum environment. *Opt Express* 2012;20:23535–41. DOI: 10.1364/OE.20.023535.
- [47] Zhang J, Brons J, Lilienfein N, Fedulova E, Pervak V, Bauer D, Sutter D, Wei Zh, Apolonski A, Pronin O, Krausz F. 260-megahertz, megawatt-level thin-disk oscillator. *Opt Lett* 2015;40:1627–30. DOI: 10.1364/OL.40.001627.
- [48] Brons J, Pervak V, Fedulova E, Bauer D, Sutter D, Kalashnikov V, Apolonskiy A, Pronin O, Krausz F. Energy scaling of Kerr-lens mode-locked thin-disk oscillators. *Opt Lett* 2014;39:6442–5. DOI: 10.1364/OL.39.006442.
- [49] Naumov S, Fernandez A, Graf R, Dombi P, Krausz F, Apolonski A. Approaching the microjoule frontier with femtosecond laser oscillators. *New J Phys* 2005;7:216. DOI: 10.1088/1367-2630/7/1/216.

- [50] Fedulova E, Fritsch K, Brons J, Pronin O, Amotchkina T, Trubetskov M, Krausz F, Pervak V. Highly-dispersive mirrors reach new levels of dispersion. *Opt Express* 2015;23:13788–93. DOI: 10.1364/OE.23.013788.
- [51] Richardson DJ, Nilsson J, Clarkson WA. High power fiber lasers: current status and future prospects. *J Opt Soc Am B* 2010;27:B63–B92.
- [52] Sraceno CJ, Heckl OH, Baer CRE, Schriber C, Golling M, Beil K, Kränkel ST, Huber G, Keller U. Sub-100 femtosecond pulses from a SESAM modelocked thin disk laser. *Appl Phys B* 2012;106:559–62. DOI: 10.1007/s00340-012-4900-5.
- [53] Zhang J, Brons J, Seidel M, Pervak V, Kalashnikov V, Wei Z, Apolonski A, Krausz F, Pronin O. 49-fs Yb:YAG thin-disk oscillator with distributed Kerr-lens mode-locking. In: CLEO/Europe-EQEC Scientific Programme; 21–25 June; Munich, Germany. 2015. p. 166 (PD-A.1 WED).
- [54] Sorokin E, Kalashnikov VL, Naumov S, Teipel J, Warken F, Giessen H, Sorokina IT. Intra- and extra-cavity spectral broadening and continuum generation at 1.5  $\mu\text{m}$  using compact low energy femtosecond Cr:YAG laser. *Applied Phys B* 2003;77(2–3): 197–204.
- [55] Joly NY, Nold J, Chang W, Hölzer P, Nazarkin A, Wong GKL, Biancalana F, Russel P. St J. Bright spatially coherent wavelength-tunable deep-UV laser source using an Ar-filled photonic crystal fiber. *Phys Rev Lett* 2011;106:203901.
- [56] Jackson SD. Towards high-power mid-infrared emission from a fiber laser. *Nat Photon* 2012;28:423–31. DOI: 10.1038/NPHOTON.2012.149.
- [57] Acerbrón JA, Bonilla LL, Vicente CJP, Ritort F, Spigler R. The Kuramoto model: a simple paradigm for synchronization phenomena. *Rev Mod Phys* 2005;77:137–85.
- [58] Pikovsky A, Rosenblum M, Kurths J. Synchronization: a universal concept in nonlinear sciences. Cambridge: Cambridge University Press; 2001.
- [59] Kuizenga DJ, Siegman AE. FM and AM mode locking of the homogeneous laser - Part I: Theory. *IEEE J Quantum Electron* 1970;6(11):694–708. DOI: 10.1109/JQE.1970.1076343.
- [60] Haus HA. Mode-locking of lasers. *IEEE J Sel Topic Quantum Electron* 2000;6(6):1173–85.
- [61] Haus HA. Short pulse generation. In: Duling IN, III (Ed.) *Compact Sources of Ultra-short Pulses*. Cambridge: Cambridge University Press; 1995. pp. 1–56.
- [62] Kalashnikov VL. Mathematical Ultrashort-Pulse Laser Physics [Internet]. 15/09/2000 [Updated: 29/03/2002]. Available from: <http://lanl.arxiv.org/abs/physics/0009056v3>.
- [63] Kuizenga DI, Siegman AE. Modulator frequency detuning effects in the FM mode-locked laser. *IEEE J Quantum Electron* 1970;QE-6:803–8.

- [64] Akhmanov SA, Vysloukh VA, Chirkin AS. *Optics of Femtosecond Laser Pulses*. New York: AIP; 1992.
- [65] Haus HA, Silberberg Y. Laser mode locking with addition of nonlinear index. *IEEE J Quantum Electron* 1986;QE-22(2):325–31.
- [66] Kalashnikov VL, Poloyko IG, Mikhaylov VP. Generation of ultrashort pulses in lasers with external frequency modulation. *Quantum Electron* 1998;28(3):264–8.
- [67] Kolner BH. Space-time duality and the theory of temporal imaging. *IEEE J Quantum Electron* 1994;QE-30(8):1951–63.
- [68] van Howe J, Xu Ch. Ultrafast optical signal processing based upon space-time dualities. *IEEE J Lightwave Technology* 2006;24(7):2649–62. DOI: 10.1109/JLT.2006.875229.
- [69] Salem R, Foster MA, Gaeta AL. Application of space-time duality to ultrahigh-speed optical signal processing. *Adv Optics Photon* 2013;5:274–317. DOI: 10.1364/AOP.5.000274.
- [70] Agrawal GP. *Nonlinear Fiber Optics*. Third Edition ed. San Diego: AP; 2001. 466 p.
- [71] Newell AC. *Solitons in Mathematics and Physics*. Philadelphia: SIAM; 1985. 246 p.
- [72] Lai Y, Haus HA. Quantum theory of solitons in optical fibers. I. Time-dependent Hartree approximation. *Phys Rev A* 1989;40:844–53.
- [73] Lai Y, Haus HA. Quantum theory of solitons in optical fibers. II. Exact solution. *Phys Rev A* 1989;40:854–66.
- [74] Yoon B, Negele JW. Time-dependent approximation for a one-dimensional system of bosons with attractive  $\delta$ -function interactions. *Phys Rev A* 1977;16:1451.
- [75] Haus HA. Theory of mode-locking with a fast saturable absorber. *J Appl Phys* 1975;46:3049–58.
- [76] Ablowitz MJ, Clarkson PA. *Solitons, Nonlinear Evolution Equations and Inverse Scattering*. Cambridge: Cambridge University Press; 1991. 516 p.
- [77] Turitsyn SK, Bale BG, Fedoruk MP. Dispersion-managed solitons in fibre systems and lasers. *Phys Rep* 2012;521:135–203. DOI: 10.1016/j.physrep.2012.09.004.
- [78] Menyuk CR. Nonlinear pulse propagation in birefringent optical fibers. *IEEE J Quantum Electron* 1987;23(2):174–6.
- [79] Menyuk GR. Pulse propagation in an elliptically birefringent Kerr medium. *IEEE J Quantum Electron* 1989;25(12):2674–82.
- [80] Hasegawa A. *Optical Solitons in Fibers*. Berlin: Springer-Verlag; 1990. 79 p.



- [81] Ding E, Renninger WH, Wise FW, Grelu Ph, Shlizerman E, Kutz JN. High-energy passive mode-locking of fiber lasers. *Int J Optics* 2012;2012(ID354156):1–17. DOI: 10.1155/2012/354156.
- [82] Wang R, Dai Y, Yan L, Wu J, Xu K, Li Y, Lin J. Dissipative soliton in actively mode-locked fiber laser. *Optics Express* 2012;20(6):6406–11.
- [83] Koliada NA, Nyushkov BN, Ivanenko AV, Kobtsev SM, Harper P, Turitsyn SK, Denisov VI, Pivtsov VS. Generation of dissipative solitons in an actively mode-locked ultralong fibre laser. *Quantum Electron* 2013;43(2):95–8. DOI: 10.1070/QE2013v043n02ABEH015041.
- [84] Wang R, Dai Y, Yin F, Xu K, Li J, Lin J. Linear dissipative soliton in an anomalous-dispersion fiber laser. *Optics Express* 2014;22(24):29314–20. DOI: 10.1364/OE.22.029314.
- [85] Haus HA, Fujimoto JG, Ippen EP. Analytic theory of additive pulse and Kerr lens mode locking. *IEEE J. Quantum Electron* 1992;28(10):2086–96.
- [86] Fermann ME, Andrejco MJ, Silberberg Y, Stock ML. Passive mode locking by using nonlinear polarization evolution in a polarization-maintaining erbium-doped fiber. *Optics Lett* 1993;18(11):894–6. DOI: 10.1364/OL.18.000894.
- [87] Hofer M, Ober MH, Haberl F, Fermann ME. Characterization of ultrashort pulse formation in passively mode-locked fiber lasers. *IEEE J Quantum Electron* 1992;28(3)DOI:720–8.
- [88] Fermann ME. Nonlinear polarization evolution in passively modelocked fiber lasers. In: Duling IN, III (Ed.) *Compact sources of ultrashort pulses*. Cambridge: Cambridge University Press; 1995. pp. 179–207.
- [89] Winful HG. Polarization instabilities in birefringent nonlinear media: application to fiber-optic devices. *Optics Lett* 1986;11(1):33–5.
- [90] Kalashnikov VL, Kalosha VP, Mikhailov VP. Self-mode locking of continuous-wave solid-state lasers with a nonlinear Kerr polarization modulator. *J Opt Soc Am B* 1993;10:1443–6.
- [91] Ding E, Kutz JN. Operating regimes, split-step modeling, and the Haus master mode-locking model. *J Opt Soc Am B* 2009;26(12):2290–300.
- [92] Ding E, Shlizerman E, Kutz JN. Generalized master equation for high-energy passive mode-locking: the sinusoidal Ginzburg-Landau equation. *IEEE J Quantum Electron* 2011;47(5):705–14.
- [93] Komarov A, Leblond H, Sanchez F. Multistability and hysteresis phenomena in passively mode-locked fiber lasers. *Phys Rev A* 2005;71(5):053809.

- [94] Komarov A, Leblond H, Sanchez F. Quantic complex Ginzburg-Landau model for ring fiber lasers. *Phys Rev E* 2005;72(2):025604.
- [95] Matsas VJ, Newson TP, Richardson DJ, Payne DN. Selfstarting passively mode-locked fibre ring soliton laser exploiting nonlinear polarization rotation. *Electron Lett* 1992;28(15):1391–3.
- [96] Tamura K, Ippen EP, Haus HA, Nelson IE. 77-fs pulse generation from a stretched-pulse mode-locked all-fiber ring laser. *Optics Lett* 1993;18(13):1080–2.
- [97] Chong A, Buckley J, Renninger W, Wise F. All-normal-dispersion femtosecond fiber laser. *Optics Express* 2006;14(21):10096–100.
- [98] Zhao LM, Tang DY, Wu J. Gain-guided soliton in a positive group-dispersion fiber laser. *Optics Lett* 2006;31(12):1788–90.
- [99] Cabasse A, Ortac B, Martel G, Hideur A, Limpert J. Dissipative solitons in a passively mode-locked Er-doped fiber with strong normal dispersion. *Optics Express* 2008;16(23):19323–9.
- [100] Kieu K, Renninger WH, Chong A, Wise FW. Sub-100 fs pulses at watt-level powers from a dissipative-soliton fiber laser. *Optics Lett* 2009;34(5):593–5.
- [101] Ruehl A, Wandt D, Morgner U, Kracht D. Normal dispersive ultrafast fiber oscillators. *IEEE J Sel Topic Quantum Electron* 2009;15(1):170–81.
- [102] Wu X, Tang DY, Zhang H, Zhao LM. Dissipative soliton resonance in an all-normal-dispersion erbium-doped fiber laser. *Optics Express* 2009;17(7):5580–4.
- [103] Renninger WH, Chong A, Wise F. Self-similar evolution in an all-normal-dispersion laser. *Phys Rev A* 2010;82:021805(R).
- [104] Zhao L, Tang D, Wu X, Zhang H. Dissipative soliton generation in Yb-fiber laser with an invisible intracavity bandpass filter. *Optics Lett* 2010;35(16):2756–8.
- [105] Chichkov NB, Hausmann K, Wandt D, Morgner U, Neumann J, Kracht D. High-power dissipative solitons from an all-normal dispersion erbium fiber oscillator. *Optics Lett* 2010;35(16):2807–9.
- [106] Baumgartl M, Ortac B, Lecaplain C, Hideur A, Limpert J, Tuenermann A. Sub-80 fs dissipative soliton large-mode-area fiber laser. *Optics Lett* 2010;35(13):2311–3.
- [107] Lecaplain C, Baumgartl M, Schreiber T, Hideur A. On the mode-locking mechanism of a dissipative-soliton fiber laser. *Optics Express* 2011;19(27):26742–51.
- [108] Zhang Z, Dai G. All-normal-dispersion dissipative soliton Ytterbium fiber laser without dispersion compensation and additional filter. *IEEE Photon J* 2011;3(6):1023–9. DOI: 10.1109/JPHOT.2011.2170057.

- [109] Kharenko DS, Podivilov EV, Apolonski AA, Babin SA. 20 nJ 200 fs all-fiber-highly chirped dissipative soliton oscillator. *Optics Lett* 2012;37(19):4104–6.
- [110] Li X, Wang Y, Zhao W, Liu X, Wang Y, Tsang YH, Zhang W, Hu X, Yang Zh, Gao C, Li Ch, Shen D. All-fiber dissipative solitons evolution in a compact passively Yb-doped mode-locked fiber laser. *J Lightwave Tech* 2012;30(15):2502–7.
- [111] Duan L, Liu X, Mao D, Wang L, Wang G. Experimental observation of dissipative soliton resonance in an anomalous-dispersion fiber laser. *Optics Express* 2012;20(1):265–70.
- [112] Doran NJ, Wood D. Nonlinear-optical loop mirror. *Optics Lett* 1988;13(1):56–8.
- [113] Ippen EP, Haus HA, Liu LY. Additive pulse modelocking. *J Opt Soc Am B* 1989;6:1736–45.
- [114] Duling IN III, Dennis ML. Modelocking of all-fiber lasers. In: Duling IN III, (Eds.) *Compact Sources of Ultrashort Pulses*. Cambridge: Cambridge University Press; 1995. pp. 140–178.
- [115] Mark J, Liu LY, Hall KL, Haus HA, Ippen EP. Femtosecond pulse generation in a laser with a nonlinear external resonator. *Optics Lett* 1989;14(1):48–50.
- [116] Kalashnikov VL, Kalosha VP, Mikhailov VP, Poloyko IG. Multi-frequency continuous wave solid-state laser. *Optics Commun* 1995;116(4–6):383–8.
- [117] Kalashnikov VL, Kalosha VP, Mikhailov VP, Poloyko IG, Demchuk MI. Efficient self-mode locking of continuous-wave solid-state lasers with resonant nonlinearity in an additional cavity. *Optics Commun* 1994;109:119–25.
- [118] Kalashnikov VL, Kalosha VP, Mikhailov VP, Poloyko IG, Demchuk MI. Self-mode-locking of cw solid-state lasers with a nonlinear antiresonant ring. *Quantum Electron* 1994;24(1):35–9.
- [119] Richardson DJ, Laming RI, Payne DN, Matsas V, Phillips MW. Selfstarting, passively modelocked Erbium fibre laser based on amplifying Sagnac switch. *Electronics Lett* 1991;27(6):542–3.
- [120] Nicholson JW, Andrejco M. A polarization maintaining, dispersion managed, femtosecond figure-eight fiber laser. *Optics Express* 2006;14(18):8160–7.
- [121] Yun L, Liu X, Mao D. Observation of dual-wavelength dissipative solitons in a figure-eight erbium-doped fiber laser. *Optics Express* 2012;20(19):20992–7.
- [122] Wang S-K, Ning Q-Y, Luo A-P, Lin A-B, Luo Z-C, Xu W-C. Dissipative soliton resonance in a passively mode-locked figure-eight fiber laser. *Optics Express* 2013;21(2):2402–7.

- [123] Zhao LM, Bartnik AC, Tai QQ, Wise FW. Generation of 8 nJ pulses from a dissipative-soliton fiber laser with a nonlinear optical loop mirror. *Optics Lett* 2013;38(11):1942–4.
- [124] Lin H, Guo Ch, Ruan Sh, Yang J. Dissipative soliton resonance in an all-normal-dispersion Yb-doped figure-eight fibre laser with tunable output. *Laser Phys Lett* 2014;11:085102.
- [125] Xu Y, Song Y, Du G, Yan P, Guo Ch, Zheng G, Ruan Sh. Dissipative soliton resonance in a wavelength-tunable Thulium-doped fiber laser with net-normal dispersion. *IEEE Photonics J* 2015;7(3):1502007. DOI: 10.1109/JPHOT.2015.2424855.
- [126] Keller U. Recent developments in compact ultrafast lasers. *Nature* 2003;424(14):831–8.
- [127] Keller U. Semiconductor Nonlinearities for Solid-State Laser Modelocking and Q-switching. In: Garmire E, Kost A. (Eds.) *Nonlinear Optics in Semiconductors II (Semiconductors and Semimetals, Vol. 59)*. San Diego: AP; 1999. pp. 211–86.
- [128] Keller U, Weingarten KJ, Kaertner FX, Kopf D, Braun B, Jung ID, Fluck R, Hoenninger C, Matuschek N, Aus der Au J. Semiconductor saturable absorber mirrors (SESAM's) for femtosecond to nanosecond pulse generation in solid-state lasers. *IEEE J Sel Top Quantum Electron* 1996;2(3):435–53.
- [129] Kaertner F. Lecture Notes: Introduction to Ultrafast Optics, Chapter 8 [Internet]. 2005. Available from: <http://ocw.mit.edu/courses/electrical-engineering-and-computer-science/6-977-ultrafast-optics-spring-2005/lecture-notes/chapter8.pdf>.
- [130] Chong A, Renninger WH, Wise FW. Environmentally stable all-normal-dispersion femtosecond fiber laser. *Optics Lett* 2008;33(10):1071–3. DOI: 10.1364/OL.33.001071.
- [131] Cabasse A, Martel G, Oudar JL. High power dissipative soliton in an Erbium-doped fiber laser mode-locked with a high modulation depth saturable absorber mirror. *Optics Express* 2009;17(12):9537–42.
- [132] Tang M, Wang H, Becheker R, Oudar J-L, Gaponov D, Godin T, Hideur A. High-energy dissipative solitons generation from a large normal dispersion Er-fiber laser. *Optics Lett* 2015;40(7):1414–7.
- [133] Gumenyuk R, Vartianen I, Tuovinen H, Okhotnikov OG. Dissipative dispersion-managed soliton 2mm thulium/holmium fiber laser. *Optics Lett* 2011;36(5):609–11.
- [134] Lecourt J-B, Duterte C, Narbonneau F, Kinet D, Hernandez Y, Giannone D. All-normal dispersion, all-fibered PM laser mode-locked by SESAM. *Optics Express* 2012;20(11):11918–23. DOI: 10.1364/OE.20.011918.
- [135] Jiang K, Ouyang C, Wu K, Wong JH. High-energy dissipative soliton with MHz repetition rate from an all-fiber passively mode-locked laser. *Optics Commun* 2012;285(9):2422–5. DOI: 10.1016/j.optcom.2012.01.033.

- [136] Poloyko IG, Kalashnikov VL. Semiconductor saturable absorber mirrors as mode-locking device for femtosecond lasers: nonlinear Fabry-Perot resonator approach. *Optics Commun* 1999;168:167–75.
- [137] Sun Z, Hasan T, Ferrari AC. Ultrafast lasers mode-locked by nanotubes and graphene. *Physica E* 2012;44:1082–91. DOI: 10.1016/j.physe.2012.01.012.
- [138] Zhang H, Tang D, Knize RJ, Zhao L, Bao Q, Loh KP. Graphene mode locked, wavelength tunable, dissipative soliton fiber laser. *Appl Phys Lett* 2010;96:111112.
- [139] Zhao LM, Tang DY, Zhang H, Wu X, Bao Q, Loh KP. Dissipative soliton operation of an Ytterbium-doped fiber laser mode locked with atomic multilayer graphene. *Optics Lett* 2010;35(21):3622–4.
- [140] Cui YD, Liu XM, Zeng C. Conventional and dissipative solitons in a CFBG-based fiber laser mode-locked with a graphene-nanotube mixture. *Laser Phys Lett* 2014;11:055106.
- [141] Cheng Zh, Li H, Shi H, Ren J, Yang Q-H, Wang P. Dissipative soliton resonance and reverse saturable absorption in graphene oxide mode-locked all-normal-dispersion Yb-doped fiber laser. *Optics Express* 2015;23(6):7000–6. DOI: 10.1364/OE.23.007000.
- [142] Liu X, Cui Y, Han D, Yao X, Sun Zh. Distributed ultrafast laser. *Sci Rep* 2014;5:9101. DOI: 10.1038/srep9101.
- [143] Im JH, Choi SY, Rotermund F, Yeom D-I. All-fiber Er-doped dissipative soliton laser based on evanescent field interaction with carbon nanotube saturable absorber. *Optics Express* 2010;18(21):22141–6.
- [144] Du J, Wang Q, Jiang G, Xu Ch, Zhao Ch, Xiang Y, Chen Y, Wen Sh, Zhang H. Ytterbium-doped fiber passively mode locked by few-layer molybdenum disulfide ( $\text{MoS}_2$ ) saturable absorber functioned with enhanced field interaction. *Sci Rep* 2014;4:6346. DOI: 10.1038/srep06346.
- [145] Paschotta R, Keller U. Passive mode locking with slow saturable absorbers. *Appl Phys B* 2001;73:653–62. DOI: 10.1007/s003400100726.
- [146] Haus HA, Silberberg Y. Theory of mode locking of a laser diode with a multiple-quantum-well structure. *J Opt Soc Am B* 1985;2(7):1237–43.
- [147] Kaertner F. Lecture Notes: Introduction to Ultrafast Optics. Chapter 7 [Internet]. 2005. Available from: <http://ocw.mit.edu/courses/electrical-engineering-and-computer-science/6-977-ultrafast-optics-spring-2005/lecture-notes/chapter7.pdf>.
- [148] Lam C-K, Malomed BA, Chow KW, Wai PKA. Spatial solitons supported by localized gain in nonlinear optical waveguides. *Eur Phys J Special Topics* 2009;173:233–43. DOI: 10.1140/epjst/e2009-01076-8.

- [149] Sakaguchi H, Malomed BA. Stable two-dimensional solitons supported by radially inhomogeneous self-focusing nonlinearity. *Optics Lett* 2012;37(6):1035–7. DOI: 10.1364/OL.37.001035.
- [150] Borovkova OV, Kartashov YV, Vysloukh VA, Lobanov VE, Malomed BA, Torner L. Solitons supported by spatially inhomogeneous nonlinear losses. *Optics Express* 2012;20(3):2657–67. DOI: 10.1364/OE.20.002657.
- [151] Kartashov YV, Konotop VV, Vysloukh VA. Two-dimensional dissipative solitons supported by localized gain. *Optics Lett* 2011;36(1):82–4. DOI: 10.1364/OL.36.000082.
- [152] Kalosha VP, Chen L, Bao X. Feasibility of Kerr-lens mode locking in fiber lasers. In: Vallée R, Piché M, Mascher P, Cheben P, Côté D, LaRochelle S, Schriemer HP, Albert J, Ozaki T. (Eds.) *Proc SPIE 7099, Photonics North 2008; 2–4 June 2008; Montréal, Canada*. Bellingham: SPIE; 2008. p. 70990S. DOI: 10.1117/12.807415.
- [153] Kalashnikov V, Apolonski A. Simulation of a Kerr fiber laser. In: *Proc. Meeting on Russian Fiber Lasers; 27–30 March; Novosibirsk, Russia*. 2012. pp. 113–114.
- [154] Shaw JK. *Mathematical Principles of Optical Fiber communications*. Philadelphia: SIAM; 2004. p. 93.
- [155] Kaup DJ. Exact quantization of the nonlinear Schroedinger equation. *J Math Phys* 1975;16:2036–41.
- [156] Thacker HB, Wilkinson D. Inverse scattering transform as an operator method in quantum field theory. *Phys Rev D* 1979;19:3660–5.
- [157] Akhmediev NN, Ankiewicz A. Solitons around us: integrable, Hamiltonian and dissipative systems. In: Porsezian K, Kuriakose VC. (Eds.) *Optical Solitons: Theoretical and Experimental Challenges*. Berlin: Springer-Verlag; 2002. pp. 105–126.
- [158] Kuszelewicz R, Barbay S, Tissoni G, Almuneau G. Editorial on dissipative optical solitons. *Eur Phys JD* 2010;59:1–2. DOI: 10.1140/epjd/e2010-00167-7.
- [159] Martinez OE, Fork RL, Gordon JP. Theory of passively mode-locked lasers for the case of a nonlinear complex-propagation coefficient. *J Opt Soc Am B* 1985;2(5):753–60. DOI: 10.1364/JOSAB.2.000753.
- [160] Haus HA, Fujimoto JG, Ippen EP. Structures for additive pulse mode locking. *J Opt Soc Am B* 1991;8(10):2068–76.
- [161] Kalashnikov VL, Sorokin E, Sorokina IT. Multipulse operation and limits of the Kerr-lens mode locking stability. *IEEE J Quantum Electron* 2003;39(2):323–36.
- [162] Proctor B, Westwig E, Wise F. Characterization of a Kerr-lens mode-locked Ti:sapphire laser with positive group-velocity dispersion. *Optics Lett* 1993;18(19):1654–6. DOI: 10.1364/OL.18.001654.

- [163] Chen S, Liu Y, Mysyrowicz A. Unusual stability of one-parameter family of dissipative solitons due to spectral filtering and nonlinearity saturation. *Phys Rev A* 2010;81:061806(R).
- [164] Kalashnikov VL. Chirped-pulse oscillators: route to the energy-scalable femtosecond pulses. In: Al-Khursan AH. (Ed.) *Solid State Laser*. InTech; 2012. pp. 145–184. DOI: 10.5772/37415.
- [165] Kalashnikov VL, Apolonski A. Chirped-pulse oscillators: a unified standpoint. *Phys Rev A* 2009;79:043829.
- [166] Akhmediev N, Soto-Crespo JM, Grelu Ph. Roadmap to ultra-short record high-energy pulses out of laser oscillators. *Phys Lett A* 2008;372:3124–8. DOI: 10.1016/j.physleta.2008.01.027.
- [167] Kalashnikov VL. Chirped dissipative solitons. In: Babichev LF, Kuvshinov VI. (Eds.) *Nonlinear Dynamics and Applications*. Minsk: 2010. pp. 58–67.
- [168] Brons J, Pervak V, Fedulova E, Bauer D, Sutter D, Kalashnikov V, Apolonskiy A, Proinin O, Krausz F. Energy scaling of Kerr-lens mode-locked thin-disk oscillators. *Optics Lett* 2014;39(22):6442–5. DOI: 10.1364/OL.39.006442.
- [169] Hu M-L, Wang Ch-L, Tian Zh, Xing Q-R, Chai L, Wang Ch-Y. Environmentally stable, high pulse energy Yb-doped large-mode-area photonic crystal fiber laser operating in the soliton-like regime. *IEEE Photon Technol Lett* 2008;20(13):1088–90. DOI: 10.1109/LPT.2008.924300.
- [170] Ramachandran S, Fini JM, Mermelstein M, Nicholson JW, Ghalmi S, Yan MF. Ultra-large effective-area, higher-order mode fibers: a new strategy for high-power lasers. *Laser Photon Rev* 2008;2:429–48. DOI: 10.1002/lpor.200810016.
- [171] Vukovic N, Healy N, Peacock AC. Guiding properties of large mode area silicon microstructured fibers: a route to effective single mode operation. *J Opt Soc Am B* 2011;28(6):1529–33. DOI: 10.1364/JOSAB.28.001529.
- [172] Jansen F, Stutzki F, Otto H-J, Eidam T, Liem A, Jauregui C, Limpert J, Tünnermann A. Thermally induced waveguide changes in active fibers. *Optics Express* 2012;20(4):3997–4008. DOI: 10.1364/OE.20.003997.
- [173] Kalashnikov VL, Sorokin E. Dissipative Raman soliton. *Optics Express* 2014;22(24):30118–26. DOI: 10.1364/OE.22.030118.
- [174] Chernykh AI, Turitsyn SK. Soliton and collapse regimes of pulse generation in passively mode-locking laser systems. *Optics Lett* 1995;20(4):398–400. DOI: 10.1364/OL.20.000398.
- [175] Chang W, Ankiewicz A, Soto-Crespo JM, Akhmediev N. Dissipative soliton resonances. *Phys Rev A* 2008;78:023830. DOI: 10.1103/PhysRevA.78.023830.

- [176] Chang W, Ankiewicz A, Soto-Crespo JM, Akhmediev N. Dissipative soliton resonances in laser models with parameter management. *J Opt Soc Am B* 2008;25(12):1972–7.
- [177] Kalashnikov VL, Apolonski A. Energy scalability of mode-locked oscillators: a completely analytical approach to analysis. *Optics Express* 2010;18(25):25757–70.
- [178] Grelu Ph, Chang W, Ankiewicz A, Soto-Crespo JM, Akhmediev N. Dissipative soliton resonance as a guideline for high-energy pulse laser oscillators. *J Opt Soc Am B* 2010;27(11):2336–41.
- [179] Ding E, Grelu Ph, Kutz JN. Dissipative soliton resonance in a passively mode-locked fiber laser. *Optics Lett* 2011;36(7):1146–8.
- [180] Kharenko DS, Shtyrina OV, Yarutkina IA, Podivilov EP, Fedoruk MP, Babin SA. Generation and scaling of highly-chirped dissipative solitons in an Yb-doped fiber laser. *Laser Phys Lett* 2012;9(9):662–8. DOI: 10.7452/Japl.201210060.
- [181] Cheng Zh, Li H, Wang P. Simulation of generation of dissipative soliton, dissipative soliton resonance and noise-like pulse in Yb-doped mode-locked fiber lasers. *Optics Express* 2015;23(5):5972–81. DOI: 10.1364/OE.23.005972.
- [182] Chang W, Soto-Crespo JM, Ankiewicz A, Akhmediev N. Dissipative soliton resonances in the anomalous dispersion regime. *Phys Rev A* 2009;79:033840. DOI: 10.1103/PhysRevA.79.033840.
- [183] Zh-Ch, Ning Q-Y, Mo H-L, Cui H, Liu J, Wu L-J, Luo A-P, Xu W-Ch. Vector dissipative soliton resonance in a fiber laser. *Optics Express* 2013;21(8):109910204. DOI: 10.1364/OE.21.010199.
- [184] Smirnov SV, Kobtsev SM, Kukarin SV, Turitsyn SK. Mode-locked fibre lasers with high-energy pulses. In: Jakubczak K. (Ed.) *Laser Systems for Applications*. InTech; 2011. pp. 39–58.
- [185] Wise FW, Chong A, Renninger WH. High-energy femtosecond fiber lasers based on pulse propagation at normal dispersion. *Laser Photon Rev* 2008;2(1–2):58–73. DOI: 10.1002/lpor.200710041.
- [186] Renninger WH, Chong A, Wise FW. Pulse shaping and evolution in normal-dispersion mode-locked fiber lasers. *IEEE J Sel Top Quantum Electron* 2012;18(1):389–98. DOI: 10.1109/JSTQE.2011.2157462.
- [187] Nelson LE, Fleischer SB, Lenz G, Ippen EP. Efficient frequency doubling of a femtosecond fiber laser. *Optics Lett* 1996;21(21):1759–61. DOI: 10.1364/OL.21.001759.
- [188] Oktem B, Ülgüdür C, Ilday FÖ. Soliton–similariton fibre laser. *Nat Photon* 2010;4:307–11. DOI: 10.1038/nphoton.2010.33.
- [189] Renninger WH, Chong A, Wise FW. Dissipative solitons in normal-dispersion fiber lasers. *Phys Rev A* 2008;77:023814. DOI: 10.1103/PhysRevA.77.023814.



- [190] Chong A, Renninger WH, Wise FW. Properties of normal-dispersion femtosecond fiber lasers. *J Opt Soc Am B* 2008;25(2):140–8.
- [191] Schultz M, Karow H, Prochnow O, Wandt D, Morgner U, Kracht D. *Optics Express* 2008;16(24):19562–7. DOI: 10.1364/OE.16.019562.
- [192] Im JH, Choi SY, Rotermund F, Yeom D-I. All-fiber Er-doped dissipative soliton laser based on evanescent field interaction with carbon nanotube saturable absorber. *Optics Express* 2010;18(21):22141–6. DOI: 10.1364/OE.18.022141.
- [193] Kieu K, Wise FW. All-fiber normal-dispersion femtosecond laser. *Optics Express* 2008;16(15):11453–8. DOI: 10.1364/OE.16.011453.
- [194] Yang H, Wang A, Zhang Zh. Efficient femtosecond pulse generation in an all-normal-dispersion Yb: fiber ring laser at 605 MHz repetition rate. *Optics Lett* 2012;37(5):954–6. DOI: 10.1364/OL.37.000954.
- [195] Chichkov NB, Hausmann K, Wandt D, Morgner U, Neumann J, Kracht D. 50 fs pulses from an all-normal dispersion erbium fiber oscillator. *Optics Lett* 2010;35(18):3081–3. DOI: 10.1364/OL.35.003081.
- [196] Ruehl A, Kuhn V, Wandt D, Kracht D. Normal dispersion erbium-doped fiber laser with pulse energies above 10 nJ. *Optics Express* 2008;16(5):3130–5. DOI: 10.1364/OE.16.003130.
- [197] Lhermite J, Machinet G, Lecaplain C, Boulet J, Traynor N, Hideur A, Cormier E. High-energy femtosecond fiber laser at 976 nm. *Optics Express* 2010;35(20):3459–61. DOI: 10.1364/OL.35.003459.
- [198] Buckley J, Chong A, Zhou Sh, Renninger W, Wise FW. Stabilization of high-energy femtosecond ytterbium fiber lasers by use of a frequency filter. *J Opt Soc Am B* 2007;24(8):1803–6. DOI: 10.1364/JOSAB.24.001803.
- [199] Buckley JR, Wise FW, Ilday FÖ, Sosnowski T. Femtosecond fiber lasers with pulse energies above 10 nJ. *Optics Lett* 2005;30(14):1888–90. DOI: 10.1364/OL.30.001888.
- [200] Renninger WH, Chong A, Wise FW. Giant-chirp oscillators for short-pulse fiber amplifiers. *Optics Lett* 2008;33(24):3025–7. DOI: 10.1364/OL.33.003025.
- [201] Chong A, Renninger WH, Wise FW. All-normal-dispersion femtosecond fiber laser with pulse energy above 20 nJ. *Optics Lett* 2007;32(16):2408–10. DOI: 10.1364/OL.32.002408.
- [202] Ortaç B, Lecaplain C, Hideur A, Schreiber T, Limpert J, Tünnermann A. Passively mode-locked single-polarization microstructure fiber laser. *Optics Express* 2008;16(3):2122–8. DOI: 10.1364/OE.16.002122.

- [203] Lefrancois S, Sosnowski ThS, Liu Ch-H, Galvanauskas A, Wise FW. Energy scaling of mode-locked fiber lasers with chirally-coupled core fiber. *Optics Express* 2011;19(4): 3464–70. DOI: 10.1364/OE.19.003464.
- [204] Lecaplain C, Ortaç B, Hideur A. High-energy femtosecond pulses from a dissipative soliton fiber laser. *Optics Lett* 2009;34(23):3731–3. DOI: 10.1364/OL.34.003731.
- [205] Lecaplain C, Chédot C, Hideur A, Ortaç B, Limpert J. High-power all-normal-dispersion femtosecond pulse generation from a Yb-doped large-mode-area microstructure fiber laser. *Optics Lett* 2007;32(18):2738–40. DOI: 10.1364/OL.32.002738.
- [206] Ortaç B, Schmidt O, Schreiber T, Limpert J, Tünnermann A, Hideur A. High-energy femtosecond Yb-doped dispersion compensation free fiber laser. *Optics Express* 2007;15(17):10725–32. DOI: 10.1364/OE.15.010725.
- [207] Lhermite J, Lecaplain C, Machinet G, Royon R, Hideur A, Cormier E. Mode-locked 0.5  $\mu$ J fiber laser at 976 nm. *Optics Lett* 2011;36(19):3819–21. DOI: 10.1364/OL.36.003819.
- [208] Lecaplain C, Ortaç B, Machinet G, Boulet J, Baumgartl M, Schreiber T, Cormier E, Hideur A. High-energy femtosecond photonic crystal fiber laser. *Optics Lett* 2010;35(19):3156–8. DOI: 10.1364/OL.35.003156.
- [209] Ortaç B, Baumgartl M, Limpert J, Tünnermann A. Approaching microjoule-level pulse energy with mode-locked femtosecond fiber lasers. *Optics Lett* 2009;34(10): 1585–7. DOI: 10.1364/OL.34.001585.
- [210] Ankiewicz A, Devine N, Akhmediev N, Soto-Crespo JM. Dissipative solitons and antisolitons. *Phys Lett A* 2007;370:454–8. DOI: 10.1016/j.physleta.2007.06.001.
- [211] van Saarloos W, Hohenberg PC. Fronts, pulses, sources and sinks in generalized complex Ginzburg-Landau equations. *Physica D* 1992;56:303–67.
- [212] Soto-Crespo JM, Akhmediev NN, Afanasjev VV, Wabnitz S. Pulse solutions of the cubic-quintic complex Ginzburg-Landau equation in the case of normal dispersion. *Phys Rev E* 55;4:4783–96.
- [213] Conte R. (Ed.) *The Painlevé Property: One Century Later*. New York: Springer-Verlag; 1999. 810 p.
- [214] Greco AM. (Ed.) *Direct and Inverse Methods in Nonlinear Evolution Equations*. Berlin: Springer; 2003. 282 p.
- [215] Kivshar YS, Malomed BA. Dynamics of solitons in nearly integrable systems. *Rev Mod Phys* 1989;61(4):763. DOI: <http://dx.doi.org/10.1103/RevModPhys.61.763>.
- [216] Malomed BA, Nepomnyashchy AA. Kinks and solitons in the generalized Ginzburg-Landau equation. *Phys Rev A* 1990;42:6009.

- [217] Kalashnikov VL. Chirped dissipative solitons of the complex cubic-quintic nonlinear Ginzburg-Landau equation. *Phys Rev E* 2009;80:046606.
- [218] Podivilov E, Kalashnikov VL. Heavily-chirped solitary pulses in the normal dispersion region: new solutions of the cubic-quintic complex Ginzburg-Landau equation. *JETP Lett* 2005;82(8):467–71.
- [219] Kalashnikov VL, Podivilov E, Chernykh A, Apolonski A. Chirped-pulse oscillators: theory and experiment. *Appl Phys B* 2006;83(4):503–10.
- [220] Ablowitz MJ, Horikis ThP. Solitons in normally dispersive mode-locked lasers. *Phys Rev A* 2009;79:063845.
- [221] Kharenko DS, Shtyrina OV, Yarutkina IA, Podivilov EV, Fedoruk MP, Babin SA. Highly chirped dissipative solitons as a one-parameter family of stable solutions of the cubic-quintic Ginzburg-Landau equation. *J Opt Soc Am B* 2011;28(10):2314–9.
- [222] Malomed BA. Variational methods in nonlinear fiber optics and related fields. In: Wolf E. (Ed.) *Progress in Optics*, Vol. 43. North-Holland: Elsevier; 2002. pp. 71–193.
- [223] Ankiewicz A, Akhmediev N, Devine N. Dissipative solitons with a Lagrangian approach. *Optical Fiber Technol* 2007;13(2):91–7.
- [224] Bale BG, Boscolo S, Kutz JN, Turitsyn SK. Intracavity dynamics in high-power mode-locked fiber lasers. *Phys Rev A* 2010;81:033828.
- [225] Bale BG, Kutz JN. Variational method for mode-locked lasers. *J Opt Soc Am B* 2008;25(7):1193–202. DOI: 10.1364/JOSAB.25.001193.
- [226] Tsoy EN, Ankiewicz A, Akhmediev N. Dynamical models for dissipative localized waves of the complex Ginzburg-Landau equation. *Phys Rev E* 2006;73:036621.
- [227] Bale BG, Kutz JN, Chong A, Renninger WH, Wise FW. Spectral filtering for high-energy mode-locking in normal dispersion fiber lasers. *J Opt Soc Am B* 2008;25(10):1763–70. DOI: 10.1364/JOSAB.25.001763.
- [228] Bale BG, Boscolo S, Turitsyn SK. Dissipative dispersion-managed solitons in mode-locked lasers. *Optics Lett* 2009;34(21):3286–8. DOI: 10.1364/OL.34.003286.
- [229] Ding E, Kutz JN. Stability analysis of the mode-locking dynamics in a laser cavity with a passive polarizer. *J Opt Soc Am B* 2009;26(7):1400–11. DOI: 10.1364/JOSAB.26.001400.
- [230] Kalashnikov VL. Dissipative soliton energy scaling. *Phys. Rev. A*. Forthcoming.
- [231] Liu X. Pulse evolution without wave breaking in a strongly dissipative dispersive laser system. *Phys Rev A* 2010;81(5):053819.

- [232] Shen X, Li W, Zeng H. Polarized dissipative solitons in all-polarization-maintained fiber laser with long-term stable self-started mode-locking. *Appl Phys Lett* 2014;105:101109.
- [233] Chen Sh, Liu Y, Mysyrowicz A. Unusual stability of one-parameter family of dissipative solitons due to spectral filtering and nonlinearity saturation. *Phys Rev Lett* 1997;79:4047.
- [234] Farnum ED, Kutz JN. Multifrequency mode-locked lasers. *J Opt Soc Am B* 2008;25(6):1002–10.
- [235] Zhang H, Tang DY, Wu X, Zhao LM. Multi-wavelength dissipative soliton operation of an erbium-doped fiber laser. *Optics Express* 2009;17(15):12692–7.
- [236] Yun L, Liu X, Mao D. Observation of dual-wavelength dissipative solitons in a figure-eight erbium-doped fiber laser. *Optics Express* 2012;20(19):20992–7.
- [237] Zhang ZX, Xu Z, Zhang L. Tunable and switching dual-wavelength dissipative soliton generation in an all-normal-dispersion Yb-doped fiber laser with birefringence fiber filter. *Optics Express* 2012;20(24):26736–42.
- [238] Xu ZW, Zhang ZX. All-normal-dispersion multi-wavelength dissipative soliton Yb-doped fiber laser. *Laser Phys Lett* 2013;10:085105.
- [239] Huang S, Wang Y, Yan P, Zhao J, Li H, Lin R. Tunable and switchable multi-wavelength dissipative soliton generation in a graphene oxide mode-locked Yb-doped fiber laser. *Optics Express* 2014;22(10):11417–26.
- [240] Mao D, Liu X, Han D, Lu H. Compact all-fiber laser delivering conventional and dissipative solitons. *Optics Lett* 2013;38(16):3190–3.
- [241] Kalashnikov VL, Chernykh A. Spectral anomalies and stability of chirped-pulse oscillators. *Phys Rev A* 2007;75:033820.
- [242] Kalashnikov VL. Dissipative solitons: perturbations and chaos formation. In: Skiadas CH, Dimotikalis I, Skiadas C. (Eds.) *Chaos Theory: Modeling, Simulation and Applications*. Singapore: Worlds Scientific Publishing; 2011. pp. 199–206.
- [243] Kalashnikov VL. Dissipative solitons in presence of quantum noise. *Chaotic Model Simulat* 2014;(1):29–37.
- [244] Soto-Crespo JM, Akhmediev N, Ankiewicz A. Pulsating, creeping, and erupting solitons in dissipative systems. *Phys Rev Lett* 2000;85(14):2937–40.
- [245] Soto-Crespo JM, Akhmediev N. Exploding soliton and front solutions of the complex cubic–quintic Ginzburg–Landau equation. *Math Computers Simulat* 2005;69(5–6):526–36. DOI: 10.1016/j.matcom.2005.03.006.
- [246] Cundiff ST, Soto-Crespo JM, Akhmediev N. Experimental evidence for soliton explosions. *Phys Rev Lett* 2002;88(7):073903.

- [247] Crtes C, Descalzi O, Brand HR. Exploding dissipative solitons in the cubic-quintic complex Ginzburg-Landau equation in one and two spatial dimensions. *Eur Phys J Special Topics* 2014;223:2145–59. DOI: 10.1140/epjst/e2014-02255-2.
- [248] Crtes C, Descalzi O, Brand HR. Noise can induce explosions for dissipative solitons. *Phys Rev E* 2012;85(015205(R)). DOI: 10.1103/PhysRevE.85.015205.
- [249] Arecchi FT, Bortolozzo U, Montina A, Residori S. Granularity and inhomogeneity are the joint generators of optical rogue waves. *Phys Rev Lett* 2011;106:153901.
- [250] Onorato M, Residori S, Bortolozzo U, Montina A, Arecchi FT. Rogue waves and their generating mechanisms in different physical contexts. *Phys Rep* 2013;528:47–89.
- [251] Solli DR, Ropers C, Koonath P, Jalali B. Optical rogue waves. *Nature* 2007;450:1054–7.
- [252] Soto-Crespo JM, Grelu Ph, Akhmediev N. Dissipative rogue waves: Extreme pulses generated by passively mode-locked lasers. *Phys Rev E* 2011;84(1):016604.
- [253] Zavyalov A, Egorov O, Iliev R, Lederer F. Rogue waves in mode-locked fiber lasers. *Phys Rev A* 2012;85:013828.
- [254] Komarov A, Sanchez F. Structural dissipative solitons in passive mode-locked fiber lasers. *Phys Rev E* 2008;77:066201.
- [255] Zakharov V, Dias F, Pushkarev A. One-dimensional wave turbulence. *Phys Rep* 2004;398:1–65.
- [256] Yun L, Han D. Bound state of dissipative solitons in a nanotube-mode-locked fiber laser. *Optics Commun* 2014;313:70–3.
- [257] Amrani F, Haboucha A, Salhi M, Leblond H, Komarov A, Sanchez F. Dissipative solitons compounds in a fiber laser. Analogy with the states of the matter. *Appl Phys B* 2010;99:107–14. DOI: 10.1007/s00340-009-3774-7.
- [258] Kalashnikov VL. Dissipative solitons: structural chaos and chaos of destruction. *Chaotic Model Simulat* 2011;(1):51–9.
- [259] Zhang L, Pan Zh, Zhuo Zh, Wang Y. Three multiple-pulse operation states of an all-normal-dispersion dissipative soliton fiber laser. *Int J Optics* 2014;(169379). DOI: 10.1155/2014/169379.
- [260] Kalashnikov VL, Sorokin E. Dissipative Raman solitons. *Optics Express* 2014;22(24):30118–126. DOI: 10.1364/OE.22.030118.
- [261] Kalashnikov VL. Chaotic dissipative Raman solitons. *Chaotic Model Simulat* 2014;(4):403–10.
- [262] Babin SA, Podivilov EV, Kharenko DS, Bednyakova AE, Fedoruk MP, Kalashnikov VL, Apolonski A. Multicolour nonlinearly bound chirped dissipative solitons. *Nature Commun* 2014;5:4653. DOI: 10.1038/ncomms5653.

- [263] Bednyakova AE, Babin SA, Kharenko DS, Podivilov EV, Fedoruk MP, Kalashnikov VL, Apolonski A. Evolution of dissipative solitons in a fiber laser oscillator in the presence of strong Raman scattering. *Optics Express* 2013;21(18):29556–64. DOI: 10.1364/OE.21.02056.
- [264] Kharenko DS, Bednyakova AE, Podivilov EV, Fedoruk MP, Apolonski A, Babin SA. Feedback-controlled Raman dissipative solitons in a fiber laser. *Optics Express* 2015;23(2):1857–62. DOI: 10.1364/OE.23.001857.
- [265] Haus JW, Shaulov G, Kuzin EA, Sanchez-Mondragon J. Vector soliton fiber lasers. *Optics Lett* 1999;24(6):376–8.
- [266] Barad Y, Silberberg Y. Polarization evolution and polarization instability of solitons in a birefringent optical fiber. *Phys Rev Lett* 1997;78(17):3290–3.
- [267] Lei T, Tu Ch, Lu F, Deng Y, Li E. Numerical study on self-similar pulses in mode-locking fiber laser by coupled Ginzburg-Landau equation model. *Optics Express* 2009;17(2):585–91.
- [268] Cundiff ST, Collings BC, Akhmediev NN, Soto-Crespo JM, Bergman K, Knox WH. Observation of polarization-locked vector solitons in an optical fiber. *Phys Rev Lett* 1999;82(20):3988–91.
- [269] Akhmediev N, Buryak A, Soto-Crespo JM. Elliptically polarised solitons in birefringent optical fibers. *Optics Commun* 1994;112(5–6):278–82. DOI: 10.1016/0030-4018(94)90631-9.
- [270] Wu J, Tang DY, Zhao LM, Chan CC. Soliton polarization dynamics in fiber lasers passively mode-locked by the nonlinear polarization rotation technique. *Phys Rev E* 2006;74:046605. DOI: 10.1103/PhysRevE.74.046605.
- [271] Zhang H, Tang DY, Zhao LM, Wu X, Tam HY. Dissipative vector solitons in a dispersion-managed cavity fiber laser with net positive cavity dispersion. *Optics Express* 2009;17(2):455–60.
- [272] Kong L, Xiao X, Yang Ch. Polarization dynamics in dissipative soliton fiber lasers mode-locked by nonlinear polarization rotation. *Optics Express* 2011;19(19):18339–44.
- [273] Mesentsev VK, Turitsyn SK. Stability of vector solitons in optical fibers. *Optics Lett* 1992;17(21):1497–9.
- [274] Zhang H, Tang D, Zhao L, Bao Q, Loh KP. Vector dissipative solitons in graphene mode locked fiber lasers. *Optics Commun* 2010;283:3334–8. DOI: 10.1016/j.optcom.2010.04.064.
- [275] Luo Zh-Ch, Ning Q-Y, Mo H-L, Cui H, Liu L, Wu L-J, Luo A-P, Xu W-Ch. Vector dissipative soliton resonance in a fiber laser. *Optics Express* 2013;21(8):10199–204. DOI: 10.1364/OE.21.010199.

- [276] Tsaturian V, Sergeyev SV, Mou Ch, Rozhin A, Mikhailov V, Rabin B, Westbrook PS, Turitsyn SK. Polarization dynamics of vector soliton molecules in mode locked fibre laser. *Sci Rep* 2013;3:3154. DOI: 10.1038/srep03154.
- [277] Sergeyev SV. Fast and slowly evolving vector solitons in mode-locked fibre lasers. *Philos Transac Royal Soc A* 2014;372:20140006. DOI: 10.1098/rsta.2014.0006.
- [278] Sergeyev SV, Mou Ch, Turitsyna EG, Rozhin A, Turitsyn SK, Blow K. Spiral attractor created by vector solitons. *Light Sci Applic* 2014;3:e131. DOI: 10.1038/lssa.2014.12.
- [279] Lin Q, Agrawal GP. Vector theory of stimulated Raman scattering and its application to fiber-based Raman amplifiers. *J Opt Soc Am B* 2003;20(8):1616–31.