



**Noémia Rita Alves  
Fernandes**

**Fragmentos de tRNA com capacidade de  
silenciamento em peixe zebra**

**Assessing silencing ability of tRNA derived  
fragments in zebrafish**



## **DECLARAÇÃO**

Declaro que este relatório é integralmente da minha autoria, estando devidamente referenciadas as fontes e obras consultadas, bem como identificadas de modo claro as citações dessas obras. Não contém, por isso, qualquer tipo de plágio quer de textos publicados, qualquer que seja o meio dessa publicação, incluindo meios eletrônicos, quer de trabalhos acadêmicos.





**Noémia Rita Alves  
Fernandes**

**Fragmentos de tRNA com capacidade de  
silenciamento em peixe zebra**

**Assessing silencing ability of tRNA derived  
fragments in zebrafish**

Dissertação apresentada à Universidade de Aveiro para cumprimento dos requisitos necessários à obtenção do grau de Mestre em Biologia Molecular e Celular, realizada sob a orientação científica da Doutora Ana Raquel Santos Calhã Mano Soares, Investigadora de Pós-doutoramento do Departamento de Biologia da Universidade de Aveiro

Apoio financeiro da FCT e do FSE no âmbito do III Quadro Comunitário de Apoio



## **o júri**

presidente

**Prof. Doutora Maria de Lourdes Gomes Pereira**  
Professora Associada com agregação da Universidade de Aveiro

arguente

**Doutora Joana Carvalho**  
Investigadora de Pós-doutoramento no Institute of Molecular Pathology and Immunology of the University of Porto (IPATIMUP)

orientadora

**Doutora Ana Raquel Soares**  
Investigadora de Pós-doutoramento do Departamento de Biologia da Universidade de Aveiro





## **agradecimentos**

A realização desta dissertação apenas foi possível devido ao contributo de várias pessoas a quem não posso deixar de expressar o meu sincero reconhecimento e gratidão:

À Doutora Ana Soares, minha orientadora, pela competência científica, acompanhamento e disponibilidade reveladas ao longo deste ano de trabalho, assim como pelas críticas, correções e sugestões relevantes feitas durante a orientação.

Ao Professor Doutor Manuel Santos pela oportunidade de desenvolver este trabalho no seu laboratório, pelo seu entusiasmo contagiante por esta área científica e por todos os conhecimentos e conselhos partilhados.

A todos os colegas de laboratório por toda a ajuda disponibilizada, em especial à Violeta pela partilha dos seus conhecimentos sobre a manutenção e cuidados com os peixe-zebra.

Ao Tomás, pela compreensão, paciência e carinho que sempre me dedicou e por me fazer acreditar em mim e nas minhas capacidades nos momentos mais difíceis.

A todos os meus amigos, em especial à Inês, à Corine e à Raquel pela amizade e carinho incondicionais, pelo apoio, o incentivo e a coragem e por estarem presentes nos momentos bons e menos bons.

À minha família, em especial aos meus pais pela oportunidade, pelo incentivo e pelo carinho e compreensão imprescindíveis ao longo de todo este tempo que sempre demonstraram. E ao meu irmão Pedro que com a sua personalidade peculiar consegue animar-me em qualquer momento.

Por fim quero agradecer à Universidade de Aveiro, em especial ao Departamento de Biologia pela disponibilização das instalações e de todas as condições necessárias à realização deste projeto.



## palavras-chave

Pequenos RNAs não-codificantes, Fragmentos de tRNA, Capacidade de silenciamento, peixe zebra

## resumo

Desde a sua descoberta que os pequenos RNAs não codificantes (pRNanc) têm sido implicados em diversos mecanismos de regulação em organismos eucarióticos, nomeadamente no silenciamento génico pós-transcricional, na regulação da cromatina e no desenvolvimento das células germinativas. Apesar dos microRNAs (miRNAs) serem a classe de pRNanc mais estudada, o desenvolvimento de novas tecnologias de sequenciação permitiu a descoberta de novas classes de pRNanc.

Recentemente foram identificados novos pRNanc que derivam de tRNAs (fragmentos de tRNA) cujas vias de produção e funções ainda não estão bem definidas. Nos últimos anos, o laboratório de RNA da Universidade de Aveiro identificou fragmentos derivados de tRNAs (tRFs) em peixe zebra. A abundância de dois destes fragmentos - tRF\_3 e tRF\_4, que derivam da porção 5' de tRNAs maduros – e o facto de ambos serem conservados em vertebrados, processados pela Dicer e apresentarem alguma capacidade de associação com proteínas Argonautas, sugere que estas moléculas representam uma nova classe de RNAs reguladores que evoluíram cedo nos vertebrados e que podem estar envolvidos em mecanismos de regulação pós-transcricional de forma análoga aos miRNAs.

Tendo esta informação em consideração, nesta tese estudou-se a capacidade de silenciamento tanto do tRF\_3 como do tRF\_4. Para isso foram realizadas experiências com um sistema repórter duplo que permite a análise da regulação do gene alvo por moléculas endógenas. Este ensaio revelou que ambos os fragmentos têm capacidade de silenciamento, embora apenas o tRF\_4 tenha uma região cuja destabilização inibe a sua capacidade de silenciamento, de forma semelhante aos miRNAs. Foi também realizada a previsão computacional de genes alvo para o tRF\_4, sendo que estes se relacionam com o padrão de expressão do tRF\_4. Esta tese demonstra que os tRFs em peixe zebra possuem capacidade de silenciamento e estudos futuros são necessários para determinar experimentalmente os seus alvos moleculares.



**keywords**

Small non-coding RNAs, tRNA-derived fragments, Silencing ability, Zebrafish

**abstract**

Since their discovery, small non-coding RNAs (sncRNAs) have been implicated in several eukaryotic regulatory mechanisms, such as post-transcriptional gene silencing, chromatin regulation and germline development. Although miRNAs constitute the most studied class of sncRNAs, the advent of high-throughput sequencing technologies allowed the discovery of new classes of sncRNA molecules.

Recently, novel sncRNAs derived from tRNAs (tRNA-derived fragments) have been identified, whose biogenesis and functions are not yet well defined. In the last years the RNA laboratory of the Aveiro University has also identified tRNA derived fragments (tRFs) in zebrafish. The abundance of two of these tRFs, namely tRF\_3 and tRF\_4, which derive from the tRNA 5'-portion – and the fact that both are conserved in vertebrates, processed by Dicer and exhibit some ability to associate with Argonaut proteins, suggest that these fragments represent a novel class of regulatory RNAs that have evolved early in vertebrates and may be involved in ancient mechanisms of genome regulation.

Taking this information into account, in this thesis the silencing ability of both tRF\_3 and tRF\_4 was studied. In order to do that, experiments were performed with a dual reporter system, which allows for the analysis of target regulation by the endogenous molecule. This assay revealed that both fragments have silencing ability, although only tRF\_4 has a region whose destabilization inhibits its silencing ability, similarly to miRNAs. Gene target computational predictions for tRF\_4 were also performed. The results obtained are related with tRF\_4 expression pattern. This thesis shows that tRFs in zebrafish have silencing ability and further studies are required in order to determine, experimentally, their molecular targets.



## Contents

<b>List of abbreviations.....</b>	<b>I</b>
<b>List of Figures .....</b>	<b>III</b>
<b>List of Tables.....</b>	<b>V</b>
<b>1. State of the art.....</b>	<b>1</b>
1.1 The RNA .....	3
1.2 The RNA interference .....	4
1.3 A novel class of small RNAs derived from tRNAs .....	5
1.3.1 The tRNA.....	5
1.3.2 The tRNA fragments.....	7
1.3.2.1 tRNA fragments biogenesis.....	8
1.3.2.2 Biological functions.....	12
1.4 Zebrafish and gene silencing.....	15
1.4.1 sncRNAs in Zebrafish.....	17
1.5 Objectives.....	18
<b>2. Materials and Methods .....</b>	<b>19</b>
2.1. Zebrafish development and tissue tRF profiling.....	21
2.2 Reporter Assay .....	21
2.2.1 Primer Annealing .....	22
2.2.2 Ligation .....	22
2.2.3 Bacterial Transformation .....	23
2.2.4 Plasmid Extraction and Linearization.....	23
2.2.5 Zebrafish maintenance .....	24
2.2.6 Zebrafish Embryos Microinjection .....	24
2.3 Gene Target Prediction.....	24
2.3.1 Duplex Injections .....	24
2.3.2 Alcian Blue Assay .....	25
2.3.3 RNA extraction .....	25
2.3.4 cDNA preparation.....	26
2.3.5 sec23b and tub1a RT-qPCR.....	26
<b>3. Results.....</b>	<b>27</b>
3.1 tRF_3 and tRF_4 profiling .....	29

3.2 tRF_3 and tRF_4 have Silencing Ability .....	31
3.3 Gene Target Prediction.....	35
3.3.1 Alcian blue staining .....	38
<b>4. Discussion .....</b>	<b>43</b>
<b>5. Conclusion .....</b>	<b>49</b>
<b>6. Bibliography.....</b>	<b>53</b>



## **List of abbreviations**

aaRS - Aminoacyl-tRNA synthetases

Ago - Argonaute

AMP - Adenosine monophosphate

ATP - Adenosine tri-phosphate

bp - Base pair

cDNA - Complementary DNA

DFRS - Dual fluorescent reporter sensor

DGCR8 - DiGeorge syndrome critical region gene 8

DNA - Deoxyribonucleic acid

dpf - Days post fertilization

dsRBD - Double-stranded RNA-binding

eIF4A - Eukaryotic initiation factor-4A

eIF4E - Eukaryotic initiation factor-4E

eIF4G - Eukaryotic initiation factor-4G

FW - Forward

GFP - Green fluorescent protein

hpf – Hours post fertilization

hrs - Hours

lincRNA - Long intergenic non-coding RNAs

min - Minutes

miRISC - miRNA-induced silencing complex

miRNA - microRNA

MO - Morpholino

mpf - Months post fertilization

mRFP - Red fluorescent protein

mRNA - messenger RNA

ncRNA - non-coding RNA

ng – Nanograms

nt - Nucleotides

PCR - Polimerase chain reaction

piRNA - Piwi-interacting RNA

Piwi - P-element induced wimpy testis  
pL - Picoliter  
rasiRNA - Repeat associated small interfering RNA  
Rev - Reverse  
RISC - RNA-induced silencing complex  
RNA - ribonucleic acid  
RNAi – interference RNA  
rpm - rotations per minute  
rRNA - ribosomal RNA  
RT-PCR - reverse transcription-polymerase chain reaction  
scaRNA - Small Cajal body-specific RNAs  
sec - seconds  
siRNA - small interfering RNA  
snoRNA – small nucleolar RNA  
snRNA - small nuclear RNA  
TALENs - Transcription Activator-Like Effector Nucleases  
TILLING - Targeting Induced Local Lesions in Genome  
tiRNA - tRNA-derived stress-induced fragments  
TRBP - HIV-1 TAR RNA binding protein  
tRNA - transfer RNA  
tRNA<sup>f</sup>/tRF – tRNA-derived fragments  
UTR - un-translated region  
YB-1 - Y box binding protein 1  
ZFN - Zinc Finger Nucleases  
μg - microgram  
μL - microliter  
μM - micromolar

## List of Figures

**Figure 1.** The RNA content of a cell

**Figure 2.** tRNA synthesis pathway

**Figure 3.** tRNA aminoacylation

**Figure 4.** tRFs classes

**Figure 5.** tRNA halves biogenesis

**Figure 6.** miRNA Biogenesis

**Figure 7.** Small tRFs biogenesis

**Figure 8.** 24hpf zebrafish embryo and organ formation

**Figure 9.** Transient assays for unveil miRNA functions in Zebrafish

**Figure 10.** Primers inserted in DFRS plasmid

**Figure 11.** Plasmid construction

**Figure 12 .** tRF sequences and their possible mature tRNAs

**Figure 13.** tRF\_3 abundance increases during development

**Figure 14.** tRF\_3 is tissue specific

**Figure 15.** tRF\_4 expression increases during development

**Figure 16.** tRF\_4 expression is tissue specific

**Figure 17.** tRf\_3 has silencing ability

**Figure 18.** tRf\_4 has silencing ability

**Figure 19.** tRF\_3 do not have a seed region

**Figure 20.** tRF\_4 5'-portion is crucial for silencing

**Figure 21.** Protein processing in endoplasmic reticulum in *Danio rerio*

**Figure 22.** 5 day-post-fertilization zebrafish larvae skeleton stained for cartilage with Alcian Blue

**Figure 23.** tRF\_3 over-expression does not affect cartilage development

**Figure 24.** tRF\_4 over-expression affects cartilage development in zebrafish

**Figure 25.** sec23b does not alter its expression in the presence of tRF\_4.



## **List of Tables**

**Table 1.** Putative gene targets for tRF\_4



# **1. State of the art**






## **1. State of the art**

### **1.1 The RNA**

The RNA was first extracted in 1960, by James Darnell and colleagues (Scherrer, 2003), and since then, many RNA molecules have been described. The RNA is a single-stranded nucleotide chain, which makes it more flexible and capable of making a wide variety of complex 3D molecular shapes. Each nucleotide is comprised of a phosphate group and a ribose sugar which is bond to one of the four bases, adenine (A), cytosine (C), guanine (G) and uracil (U). Also, like some proteins, RNA can have catalytic functions, known as ribozymes (Griffiths, 2004; Pierce, 2010).

RNA molecules are divided into several classes according to their function, size and location in the cell. There are two main types of RNA, namely coding and non-coding RNA. Messenger RNA (mRNAs) constitutes the group of coding RNAs. These molecules are translated into proteins and their size varies with the size of the gene being transcribed. Despite the fact that the mRNA represents 60% of the total transcription of the cell it rarely makes up more than 4% of the total RNA due to its degradation soon after synthesis, being a short-lived molecule (Brown, 2002; Liu, *et al.*, 2006; Arraiano and Fialho, 2007). On the other hand, the non-coding RNAs (ncRNAs) do not encode proteins (Liu, *et al.*, 2006) and account for the remaining 96% of the total RNA (Brown, 2002). The most abundant non-coding RNA is ribosomal RNA (rRNA), which makes up 80% of the total cell RNA. These molecules are important components of the ribosomes, the structures on which protein synthesis takes place, and are found in the cytoplasm (Brown, 2002). The other 16% of RNA contains the transfer RNA (tRNA) and small non-coding RNAs (Brown, 2002). Transfer RNA length ranges from 70 to 90nt and can be found in the cytoplasm and in mitochondria (Elliott and Lodomery, 2011) and plays an important role in protein synthesis, as it carries the correct aminoacid to the ribosome on which translation occurs. Small non-coding RNAs include small nuclear RNA (snRNA), interference RNA (RNAi), long intergenic non-coding RNAs (lincRNA) and anti-sense RNA (Figure 1) (Brown, 2002; Smith, 2009). These RNAs play key roles in several cellular processes and have regulatory functions, being involved in several biological processes, namely development, differentiation, immune response, infection and cancer (Arraiano and Fialho, 2007).



**Figure 1.** The RNA content of a cell (Adapted from Smith, 2009). In a human cell 4% of the RNA (mRNA) encodes proteins. The other 96% contribute for the translation process like rRNA and tRNA or have regulatory functions like RNAi.

## 1.2 The RNA interference

The RNA interference (RNAi) was one of the most remarkable discoveries of the last 20 years. For the first time, it was clear that RNA molecules not only have important roles in translation, but also in regulation of gene expression in almost all eukaryotic organisms (Nilsen, 2007). The RNAi includes any post-transcriptional gene regulation event in which a small sized single-stranded RNA (20 to 30nt) forms a RNA-induced silencing complex (RISC) with an Argonaute protein (Ago), regulating, therefore, the expression of a cognate mRNA either by repressing translation or degradation. In this complex, the role of the small RNA is to guide it to the proper targets by complementary base-pairing interactions, while the Ago proteins function as effector molecules (Nilsen, 2007; Kelly and Hurlstone, 2011; Kang, 2011). Some Ago proteins have catalytic activity and can specifically cleave the mRNA, but others confer the RISC complex the ability to induce DNA methylation, chromatin structure alteration and repression of mRNA translation (Nilsen, 2007).

There are several types of small non-coding RNAs, namely repeated-associated small-interfering RNA (rasiRNA), piwi-interacting RNA (piRNA), small interfering RNA (siRNA) and microRNA (miRNA), being the last two the most studied classes (Lee, *et al.*, 2009) (Figure 1). All of the above have gene expression regulatory functions. Both rasiRNAs and piRNAs are thought to be involved in germline silencing of repeat transcripts, in *Drosophila* and mammals, possibly by chromatin modification (Matera, *et al.*, 2007 and Elliott and Ladomery, 2011). piRNAs are also responsible for cleaving RNAs made by transposable elements and for regulating the embryonic development and

telomere protections (Huang, *et al.*, 2013). siRNAs cleave RNAs derived from viruses, retroelements and repeat sequences. They also regulate long-term gene expression by directing cytosine methylation through RNA directed DNA methylation (Matera, *et al.*, 2007 and Huang, *et al.*, 2013). miRNAs, which are the most well-known class, regulate fundamental biological processes like proliferation, metabolism, embryogenesis, aging, and cell death, by targeting and silencing a huge number of human mRNA sequences (Elliott and Ladomery, 2011 and Huang, *et al.*, 2013).

The advances in high throughput sequencing technologies allowed the discovery and characterization of an increasing number of small ncRNAs and prevented many to be discarded and considered as merely transcriptional “noise”. These small ncRNAs have turned out to be a hotspot in the life sciences, as it is becoming more and more evident that they have key regulatory functions (Huang, *et al.*, 2013). The new sequencing technologies have, among others, unveiled a novel class of non-coding RNAs derived from tRNAs, the tRNA derived fragments (Lee, *et al.*, 2009).

### **1.3 A novel class of small RNAs derived from tRNAs**


#### **1.3.1 The tRNA**

The transfer RNA (tRNA) is the bridge that links the genetic code and the polypeptide sequence, in other words it is the molecular adaptor that associates the mRNA codon to an amino acid (Elliott and Ladomery, 2011). This molecule is usually 70 to 90 nt long and its secondary structure is described as a cloverleaf as it folds into four base paired stems. These structures are called the D loop, T loop, anticodon loop and the accept stem. The anticodon loop contains the anticodon, which recognizes its correspondent codon in the mRNA. The acceptor stem consists in the base paired between the 5' and the 3' ends, having the 3' end a CCA residue which is the location for the amino acid binding. The tRNA has an extensive base modification which allows its tertiary L-shape structure (Elliott and Ladomery, 2011; Sobala and Hutvagner, 2011).


In summary, the synthesis of the transfer RNA comprises three steps: cleavage/trimming, modification and aminoacylation (Figure 2).

Transfer RNA from eukaryotic organisms is synthesized by RNA polymerase III (Alberts, *et al.*, 2008). The tRNA precursor (pre-tRNA) chain is usually synthesized with additional nucleotides in one or both ends, the 5' leader and the 3' trailer. These extra sequences are then removed by the endonuclease RNase P, which recognizes the global L-shape tRNA structure and can specifically hydrolyze the phosphodiester bond, liberating the 5'-end of the molecule leaving a 5'-phosphate group (Krebs, *et al.*, 2011; Elliott and Lodomery, 2011). The 3'-end is first cleaved in the middle by RNase E and RNase III and then trimmed by RNase II, RNase BN, RNase PH and RNaseZ. This last enzyme cleaves the phosphodiester bond leaving a free 3'-OH end for the CCA residues addition by tRNA

nucleotidyltransferase (Elliott and Lodomery, 2011). After, the tRNA suffers modifications in specific bases. The modifications are performed by specific tRNA-modifying enzymes. Usually 15 to 20% of the tRNA nucleotides are modified. These modifications are important for the tertiary structure formation and, when occurring in the anticodon, it creates a phenomenon known as wobble, in which a particular anticodon can recognize more than one codon (Krebs, *et al.*, 2011; Elliott and Lodomery, 2011). Once all these steps have taken place, the tRNA can be charged by the aminoacyl-tRNA synthetases in the 3'-CCA terminal (Figure 3) (Krebs, *et al.*, 2011).



**Figure 2.** tRNA synthesis pathway (Adapted from (Nobel Media AB, 2013). tRNA genes transcription by RNA-Pol III produce a pre-tRNA which contains 5' and 3' extensions. RNase P and RNase Z cleave 5' and 3' end respectively. Then tRNA nucleotidyltransferases create modifications in specific nucleotides and aminoacyl-tRNA synthetases add a CCA triplet at the 3' end.



**Figure 3.** tRNA aminoacylation (Maxwell, 2012). First, a specific amino acid and an ATP molecule bind to the aminoacyl-tRNA synthetases (aaRS). The amino acid is activated by binding to the AMP and a phosphate is released. Then, the correct tRNA binds to the synthetase and the amino acid is covalently attached to it, releasing the AMP. Finally the charged tRNA is released.

### 1.3.2 The tRNA fragments

In 1971, it was observed that, when infected by bacteriophage T4, *E. coli* would produce specific leucine tRNA fragments, one with 48 nt and the other one with 39 nt corresponding to the 5' end and to the 3' end, respectively (Yudelevich, 1971).


Until recently, little importance was given to this finding, as it was thought that these molecules were random degradation products of tRNA, and, therefore, junk. Using protocols usually applied to miRNA profiling, the reads coming from tRNAs often constitute a minor fraction of the total library (1,5% to 10%) and were therefore, ignored from a deeper analysis as sequencing artefacts or degradation products of the mature tRNAs. However, comparative analysis between new and existing sequencing data revealed that tRNA fragments are conserved across distantly related species and correspond to a particular domain of tRNAs, drawing researcher's attention for their possible cellular importance. Also, several deep sequencing, genetic screens, transcriptional, and molecular analyses studies showed that these small RNAs, are associated with several biological processes, such as development and stress response (Martens-Uzunova, *et al.*, 2013).

These newly identified tRNA fragments seem to have different sizes and origins and can be specific to a tissue or a developmental stage (Cole, *et al.*, 2009; Haussecker, *et al.*, 2010; Saikia, *et al.*, 2012).

Since tRNA fragments were discovered, their nomenclature has been highly inconsistent. Several researchers have been using names like tRNA halves, tRNA-derived

halves, tRNA-derived RNA fragments (tRNAf), tRNA-derived stress-induced fragments (tiRNAs) or tRNA-derived small RNAs to refer to similar entities (Martens-Uzunova, *et al.*, 2013). Recently, two different nomenclatures have been proposed, one based on the tRNA fragments size and another on the part of the tRNA or pre-tRNA from which the tRNA fragments are derived (Sobala and Hutvagner, 2011).

Therefore, tRNAfs can be divided into two major groups: the tRNA halves, which include tRNA fragments that range from 30 to 35nt and are produced in response to stress conditions, by cleavage near the anticodon; and the small tRNA fragments (tRFs) (Figure 4). This group contains tRFs with approximately 20nt and can be divided in three sub-classes: 5'tRFs, 3'CCA tRFs and 3'U tRFs. The first class includes all the tRFs derived from the 5' end of the tRNA and cleaved near its D-loop. The second class consists in all tRFs derived from the 3' end and cleaved near the tRNA T-loop. The third class of tRFs is composed by those derived from the 3' end of the pre-tRNA.




**Figure 4.** tRFs classes (Garcia-Silva, *et al.*, 2012). 3'U tRF is represented in yellow and derives from the immature tRNA. Green and blue segments correspond to 5'tRF and 3'CCA tRF respectively and derived from the mature tRNA.

Most of these tRFs end in a series of U residues, generated by RNA polymerase III (Sobala and Hutvagner, 2011; Martens-Uzunova, *et al.*, 2013).

### 1.3.2.1 tRNA fragments biogenesis

The tRNA halves, or the tRNA fragments-derived from cleavage within the anticodon, have been associated with response to stress conditions (Thompson and Parker, 2009; Fu, *et al.*, 2009; Wang, *et al.*, 2012). As stated before, tRNAs are a key component in the translation machinery, however they also have roles in cell proliferation and stress response. For example, during amino acid starvation, tRNA uncharged can act as signaling molecules and activate Gcn2 kinase, which instead phosphorylates eIF2 $\alpha$  (Thompson and Parker, 2009; Saikia, *et al.*, 2012). Stress-induced tRNA cleavage occurs through specific enzymes secreted from stressed cells such as Rny1 in yeast and angiogenin in mammals (Figure 5) (Fu, *et al.*, 2009; Saikia, *et al.*, 2012). Both of these ribonucleases belong to families, RNase T2 and RNase A, respectively. These enzyme families have little substrate

specificity, pointing to possible cleavage of other RNAs during stress. Wang, et al. noticed that the nucleotides before the major cleavage were Gs and Cs, which reduces the randomness theory (Wang, *et al.*, 2012). Rny1 and angiogenin are secreted ribonucleases. Rny1 is secreted in the vacuole and released to the cytoplasm during oxidative stress, where



**Figure 5.** tRNA halves biogenesis (Garcia-Silva, *et al.*, 2012)


it can access the cytosolic RNAs (Thompson and Parker, 2009). Angiogenin is most located in the nucleus or the nucleolus, but is also found in the cytoplasm, bound to its inhibitor RNH1. Stress conditions promote the dissociation of angiogenin from RNH1 or the ribonuclease release from the nucleus (Thompson and Parker, 2009).

tRNA cleavage may inhibit specific steps of translation, leading to its global decrease or produce tRNA halves that can target specific mRNAs, which in conjunction with tRNA processing enzymes or Argonaute proteins, inhibit their translation or leads to their degradation (Thompson and Parker, 2009).

Small tRNA fragments have been often compared with miRNAs. Their features, namely size and association with Ago proteins and even part of their biogenesis are very similar to those of miRNAs (Pederson, 2010).

Very briefly miRNAs, which in general are present in the genome as clusters, are processed from precursor transcripts (pri-miRNAs), which are mostly transcribed by RNA polymerase II from independent genes (Ghildiyal and Zamore, 2009; Carthew and Sontheimer, 2009; Krol, *et al.*, 2010). The pri-miRNA usually has a long imperfect stem and, like most mRNAs, has a 5'-cap structure and a polyadenylated tail (Hussain, 2012). After being transcribed, the pri-miRNA is cleaved in the nucleus by the Microprocessor Drosha-DGCR8 (Pasha in *Drosophila* and *C. elegans*) into a 60-70 nt precursor hairpin, the pre-miRNA, which contains in its 3' end 2 nt overhang and a 5'-phosphate group (Ghildiyal and Zamore, 2009; Krol, *et al.*, 2010). Then a nuclear export protein, Exportin-5 in a RanGTP-dependent manner, exports the pre-miRNA through the nuclear pore (Ghildiyal and Zamore, 2009).

Once in the cytoplasm, Dicer and its dsRBD assistant protein TRBP (or Loqs in flies) cleave the pre-miRNA. This connection to Dicer, can only be possible because, unlike Drosha, it possesses a Paz domain that interacts with the 2 nt overhanging in 3'UTR, left by Drosha (Ghildiyal and Zamore, 2009; Carthew and Sontheimer, 2009). Dicer cleaves off the loop and originates an approximately 20 nt double stranded RNA, a miRNA/miRNA\* duplex, with two nucleotides overhanging at each 3' end. (Ghildiyal and Zamore, 2009; Winter, *et al.*, 2009; Krol, *et al.*, 2010). Dicer and TRBP then dissociates from the duplex and an helicase unwinds and separates both strands (Winter, *et al.*, 2009). Usually only one strand of the miRNA/miRNA\* is incorporated into a miRNA-induced silencing complex (miRISC) being the other strand degraded. The criteria for choosing one strand over the other is probably thermodynamic (Ghildiyal and Zamore, 2009; Carthew and Sontheimer, 2009; Krol, *et al.*, 2010) (Figure 6). miRISC then targets the homologous transcripts by base pairing interactions and silences the target mRNA. The crucial feature for target is excellent complementarity between the 3' end of the target mRNA and the miRNA 5' end, also called “seed” region. This “seed” region starts at position 2 in the guide sequence, up to the 6th or 8th nucleotide. The complementarity between this two regions must be perfect in order to have an effective silencing by the miRNA (Elliott and Ladomery, 2011).



**Figure 6.** miRNA Biogenesis (Winter, *et al.*, 2009). RNA polymerase II transcribes miRNA genes, generating a primary miRNA transcript with hairpin structures. Microprocessor Drosha-DGCR8 then cleaves the pri-miRNA and the resulting hairpin, pre-miRNA, is exported from the nucleus to the cytoplasm by Exportin-5-Ran-GTP. Once in the cytoplasm Dicer-TRBP cleaves the the pre-miRNA into miRNA duplexes. The duplexes are unwound and the functional strand as well as Dicer are loaded into the RNA-induced silencing complex (miRISC). The mature miRNA can then guide the RISC complex to the 3'UTR of a mRNA causing its cleavage or translation repression or deadenylation.




The tRFs are originated from tRNA gene transcripts which may be cleaved by Dicer, ELAC2 or RNase Z, giving rise to 5'tRFs, 3'CCA tRFs and 3'U tRFs. 5'tRFs biogenesis can be Dicer-dependent (Cole, *et al.*, 2009) in mammals or Dicer-independent as it was reported in *Schizosaccharomyces pombe* (Buhler, *et al.*, 2008). The differences in their size (19nt and 23nt, respectively), support the idea that there must be different mechanisms/enzymes for each organisms (Cole, *et al.*, 2009; Sobala and Hutvagner, 2011). Computational studies also point that some 5'tRFs can actually acquire an hairpin structure, like the one observed for miRNAs, which may point towards a similar mechanism of action or function (Cole, *et al.*, 2009 and unpublished data). 3'CCA tRFs are cleaved by dicer after CCA addition to the tRNA. In fact, it was reported that 5'tRFs studied in *Tetrahymena* carry a 5' monophosphate, which is typical of Dicer cleavage and differs from the 5-hydroxyls described on longer half-tRNAs generated by starvation-induced cleavage (Couvillion, *et al.*, 2010). Moreover, Lee *et al.*, found that the preferential cleavage site for these fragments was located in AU sites in prostate cancer cells (Lee, *et al.*, 2009). The 3'U tRFs are generated from pre-tRNA, which still did not lost the polyuridine tract. These fragments are cleaved by RNase Z (Sobala and Hutvagner, 2011) or, ELAC2, a cytoplasm enzyme (Lee *et al.*, 2009). In both studies 3'U tRFs are only found in the cytoplasm, leading to speculate that these transcripts may be cleaved in the nucleus and rapidly exported to the cytoplasm (Haussecker, *et al.*, 2010).

After cleavage both tRNA halves and tRFs are thought to bind to Ago proteins. Until now investigators have found that these molecules associate more efficiently with Ago 3 and Ago4 (Haussecker, *et al.*, 2010). In mammals it is known that only Ago2 has cleavage activity and Ago1, 3 and 4 do not have this activity, therefore their function has not been totally elucidated yet. It was also seen that, in *Tetrahymena*, 3'CCA tRFs associate with a PIWI protein, Twi12, which does not have cleavage activity (Couvillion, *et al.*, 2010).

Although both miRNAs and tRFs are generated by Dicer and associate with Ago proteins, they differ from each other in several aspects. Mammalian miRNAs are not methylated in the 3' terminal ribose. Yet, Cole *et al.*, saw that tRFs-derived from tRNA<sup>Gln</sup> have 3' terminal modification. Still they cannot tell if the modification occurs after or prior Dicer processing. Also, miRNAs form a stable complex, when incorporated with Argonautes, so when Dicer is knocked down there is no significant change in miRNA steady-state level.

However, when knocking down Dicer, Cole *et al.*, noticed smaller fractions of tRFs than in the minimal RISC (AGO2 and tRF), which suggests that either the tRFs associate inefficiently with Ago proteins, or they do not associate with the proteins at all, or the tRFs are incorporated into non-Argonaute complexes, which have a faster turnover rate. This inefficiency in RISC incorporation can be due to the 3' terminal modification (Figure 7) (Cole, *et al.*, 2009).



**Figure 7.** Small tRFs biogenesis (Sobala and Hutvagner, 2011) 3'U tRFs can be generated from pre-tRNA by RNase Z, either in the cytoplasm or in the nucleus. After processed by RNase P and Z and added the CCA-terminal, the tRNA can be permanently exported to the cytoplasm and processed by Dicer. Mature aminoacylated tRNAs may enter the small RNA pathway via Dicer, which produces the 5'tRFs and the 3' CCA tRFs. White triangle - RNase Z cleavage site; Black triangle - RNase P cleavage site; Red arrows - premature nuclear export.

### 1.3.2.2 Biological functions

The tRNA derived fragments have long been regarded as random byproducts of tRNA biogenesis or degradation. Yet, the discovery of the small tRFs, which biogenesis is dependent on Dicer automatically raised the question of whether or not these fragments have a biological function (Cole, *et al.*, 2009; Gebetsberger, *et al.*, 2012). Several lines of work suggested that tRFs are indeed novel biological entities. First of all, tRFs biogenesis seems to be regulated and the tRFs have approximately the same size distribution. In fact the cleavage sites showed to be conserved and specific enzymes are required to cleave the

tRNA. (Lee, *et al.*, 2009; Cole, *et al.*, 2009; Sobala and Hutvagner, 2011; Martens-Uzunova, *et al.*, 2013). Furthermore the results show that tRFs are produced after intron excision and even after 5'-CCA-3' addition (Martens-Uzunova, *et al.*, 2013). Also tRNA halves biogenesis seems to be also controlled by RNA methylation and induced by specific stress (Martens-Uzunova, *et al.*, 2013) or cell proliferation (Wang, *et al.*, 2012). Secondly, until now only tRFs corresponding to the 5' end and the 3' end of the tRNA were reported, and no tRF matching the anticodon loop was stated in mammals (Sobala and Hutvagner, 2011). Pederson, stated that this feature might be selected by evolution, and as the anticodon is optimally configured for base-pairing, it did not have any selective advantage on a regulation path being, therefore, excluded from the sequence (Pederson, 2010). Third, when performing deep sequencing it is clear that the sequencing number of different tRFs does not correlate to those of the tRNA gene copies or with tRNA gene expression or codon usage distribution. Moreover, when RNA degradation increases there is not an increase in the fragments abundance (Lee, *et al.*, 2009; Sobala and Hutvagner, 2011; Martens-Uzunova, *et al.*, 2013). At last, there has been evidence that tRFs can associate with Ago proteins and some can actually have trans-silencing activity (Lee, *et al.*, 2009; Sobala and Hutvagner, 2011).

These evidences indicate that the tRNA-derived fragments can be novel biological entities. Although, their roles are not totally clarified yet, there are already several lines suggesting that tRNA-derived fragments participate in important biological processes like stress response and gene expression regulation.

Haussecker, *et al.*, focused their work on tRFs (3'U and 3'CCA) ability to associate with Ago proteins and their propensity to have an effect on luciferase reporters. They found that both 3'U and 3'CCA bind more effectively to Ago 3 and Ago4 than with Ago1 and Ago2 and that 3'CCA and that the latter has a moderate effect on the reporter while 3'U does not (Haussecker, *et al.*, 2010).

Lee, *et al.*, also studied 3'U but focus their attention on their possible function. The authors studied in particular tRF-1001, and by knocking down this tRF with a siRNA, they saw a decrease in cell proliferation (Lee, *et al.*, 2009).

Using HeLa cells, Cole *et al.* conjecture that tRNAs may compete with pre-miRNA for Dicer cleavage leading to an unbalanced miRNA homeostasis. An increase in tRNA levels leads to cell transformation maybe because Dicer would be forced to accept them as

a substrate and thus altering miRNA homeostasis and function. They also verified that 5'tRFs do not associate with Ago2, but Ago1. Ago1 has been implicated in transcription gene silencing by miRNAs and siRNAs that target promoter regions and promotes heterochromatin formation. This suggests that tRFs may also be involved in transcriptional gene silencing pathway (Cole, *et al.*, 2009). In another study, using deep sequencing and computational comparison, it was showed that there are 3'CCA tRFs that do not undergo Dicer processing, but can Associate with Ago2 and guide down regulation *in vitro* (Li, *et al.*, 2012). Wang *et al.* also demonstrated that a 5'tRF derived tRNA<sup>glu</sup> (CTC) has trans-silencing activity. This tRF is located in the cytoplasm and the mechanism by which silencing is carried out is different from miRNA/siRNA pathway, as the binding region to the target RNA is positioned in the 3'end and not in the 5'end (Wang, *et al.*, 2012).

Recently, several works from Paul Anderson group have shown that tiRNAs (tRNA-derived stress-induced fragments) can inhibit translation. They have demonstrated that Angiogenin is responsible for producing tiRNAs in stress conditions and U2OS cells transfection with 5'-tiRNAs could induce phospho-eIF2 $\alpha$ -independent translational arrest, and promote stress granules assembly. The tiRNAs efficiency was higher when possessing a 5'-monophosphate in the 5'-end. Ivanov *et al.*, found that tiRNAs can efficiently inhibit translation of uncapped mRNAs. 5'-tiRNAs can directly or indirectly bind eIF4G, eIF4A, or to the complex eIF4G/ eIF4A and inhibit translation. It can also bind to the eIF4F, favouring eIF4E:4E-BP complex formation over eIF4F. These tiRNAs also interact with YB-1, whose function is to regulate transcription and translation. The cold shock domain of this protein can displace eIF4E from m<sup>7</sup>G cap and the carboxyl terminus interferes with eIF4G binding. This suggests that tiRNAs and YB-1 cooperate to eIF4G/A binding and consequently translation initiation and that this protein is require for stress granule assembly by the tiRNAs (Yamasaki, *et al.*, 2009; Emara, *et al.*, 2010; Ivanov, *et al.*, 2011).

Sobala and Hutvagner, also found that 5'tRFs can inhibit protein translation in human cells, by interfering with the elongation process. Using HeLa cells, they studied a highly-express 5'tRF (Gln) and find that it can inhibit reporter genes translation *in vitro* and *in vivo*. Unlike miRNA/siRNA, they noticed that this tRF did not need complementary sites, instead it recognizes a "GG" dinucleotide sequence, which appears to be conserved in 5'tRFs at position 17-8 or 18-19, depending on the parent tRNA. They observed that the 5'tRF cellular fraction can associate actively with polysomes, interfering with the


elongation process or with other important translation processes like tRNA-charge specificity (Sobala and Hutvagner, 2013).

Using 454 DNA pyrosequencing technologies to sncRNA discovery in zebrafish, the laboratory of RNA in the university of Aveiro unveiled 11 tRFs, which align with the tRNAs in the 3'end (6tRFs), the 5' end (4tRFs) and the pre-tRNA (1tRF) (unpublished data). Some of these tRFs showed to be more abundant in adult tissues like muscle, fins and skin, than in developmental stages. Three of the tRFs identified are homologous to other tRFs identified in human cells or tissues. tRF\_3 is similar to tRF-5002 found in prostate cancer cell lines and also to another tRNA<sup>glu</sup> fragment in human fetus hepatic tissue and it was recently described in human cells after a RSV infection. tRNAf\_2 was also found in human fetus hepatic tissue and derives from tRNA<sup>val</sup>, tRNAf\_11 is identical to cand15 and derives from tRNA<sup>arg</sup>. These results show that these tRFs are likely conserved in vertebrates and therefore they must have a biological function. *In vitro* biogenesis assays performed with two of the most expressed tRNA derived fragments found in zebrafish - tRF\_3 and tRF\_4, showed that they were cleaved by Dicer and may fold into a similar structure to that of pre-miRNAs (unpublished data). Also, both tRF\_3 and tRF\_4 have an uridine at the 5' end, which is a miRNA characteristic and bind, although weakly, to Ago1 (unpublished data), suggesting that these fragments can act like miRNAs and be involved in gene silencing.

#### **1.4 Zebrafish and gene silencing**

*Danio rerio*, also known as zebrafish, is a small size tropical water fish, which inhabits Asia. For the past 20 years this organism has become one of the favourite *in vivo* model systems for studying developmental processes and human diseases (Santoriello and Zon, 2012; Kardash, 2013). Recently sequencing studies unveiled that zebrafish has 26,000 protein coding genes and by direct comparison to the human genome it was possible to conclude that 71,4% of the human genes have at least one zebrafish orthologue. On the other hand, 69% of zebrafish genes have at least one orthologue in humans (Santoriello and Zon, 2012; Chitramutu, 2013; Howe, *et al.*, 2013; Kardash, 2013). Also, it was possible to see that 82% (2,601 out of 3,176) of the genes related to human diseases can be related to a zebrafish orthologue (Howe, *et al.*, 2013). Besides the great genetic similarity, zebrafish has many advantages for which it is considered an attractive model in research.

Zebrafish has a high fecundity and a single female can lay 200-300 eggs. The embryos develop outside the body and are transparent, which allows following organogenesis. The development is rapid and the major organs are formed within 24hours after fertilization (Figure 8). Furthermore, its raise and maintenance is easy and inexpensive when compared to other vertebrate models (Santoriello and Zon, 2012; Chitramutu, 2013).




**Figure 8.** 24hpf zebrafish embryo and organ formation. (Weigta, *et al.*, 2010)H – Head; E - Eye; S/O - sacculi/otoliths;T – Tail; TT – Tail Tip

At first, zebrafish was used in forward genetics, in which chemical and insertional mutation, namely the DNA transposon system known as Sleeping Beauty, has

led to the identification of the causative genes for a given phenotype, by positional cloning or through candidate gene approach (Santoriello and Zon, 2012; Chitramutu, 2013). Recently, reverse genetic techniques have been developed and successfully applied to zebrafish. These techniques include antisense morpholino (MO) oligonucleotide mediated gene knockdown technology, Targeting Induced Local Lesions In Genome (TILLING), Zinc Finger Nucleases (ZFN), Transcription Activator-Like Effector Nucleases (TALENs), Tol2 mediated Transgenesis, GAL4-UAS System, Tol2-mediated Gal4/UAS, Cre/Lox system and a tamoxifen-inducible Cre/lox method (Chitramuthu, 2013).

Microinjection is the most frequently used method when working with zebrafish embryos. This is an efficient and rapid vehicle for adding either mRNA, morpholino or RNA duplex into hundreds of embryos per hour (Figure 9) (Rosen, *et al.*, 2009, Fjose and Zhao, 2010). Microinjection of fertilized eggs, easily obtained by mating adult fish, provides the opportunity to study directly their functions throughout zebrafish development. Injection of chemically synthesized miRNA mimics allows determining the effect of an over-expression of a specific miRNA during development and, in some cases, to detect variations in predicted mRNA targets. Also the validation of these targets and their binding sites can be achieved by co-injection of a reporter gene, which consists in mRNA of GFP and the miRNA binding site in its 3'UTR (Fjose and Zhao, 2010).



**Figure 9.** Transient assays for unveil miRNA functions in Zebrafish (Adapted from: Fjose and Zhao, 2010). **A)** Gain-or-loss of function experiments can be performed by injection of miRNA or miRNA inhibitors in embryos. The effects can be analyzed by assessing the phenotype or the expression levels of the mRNA. **(B–D)** Illustrates the effects on the mRNA levels by injection of either a mature miRNA duplex or MOs. **B)** miRNA duplex injection leads to a decreased level of the target mRNA and therefore the protein encoded by it. **C)** MOs injection blocks miRNA from binding its target site, leading to an increase in protein level. **D)** Mature miRNA and GFP reporter mRNA co-injections can be used to validate predicted binding sites of a specific mRNAs.

### 1.4.1 sncRNAs in Zebrafish

According to miRBase, 346 microRNAs have already been unveiled in zebrafish (miRBase, 2013). Recently, Wei et al, using Next-Generation sequencing technology unveiled the expression pattern of 198 known miRNAs within 122 different miRNA families in zebrafish embryos. The same study showed that 225 of the 56,311 reads derived mostly from 3' and 5' ends of mature tRNAs. These fragments were different and had different cleavage-sites from those detected in angiogenin-dependent fragments found in stressed cells. 3' tRNA reads size did not fluctuate much (18nt), unlike the 5' which were distributed more widely and lead to the conclusion that they might be under different selection mechanisms or have different functions. Also, it was possible to notice that tRFs abundance increases dramatically from 24hpf, being more abundant those derived from the 5'end (Wei, *et al.*, 2012).

As stated before, 454 DNA pyrosequencing technologies applied to sncRNA discovery in zebrafish by our laboratory unveiled 11 tRFs in this organism. From those, tRF\_3 and tRF\_4 were chosen to study in this thesis, as they were the most expressed ones. tRF\_3 is a ~30nt fragment which derives from the 5'end of a mature tRNA through Dicer cleavage. This fragment is conserved in zebrafish, humans and mouse and appears to bind to Ago1. Preliminary studies also indicate that its expression is tissue dependent. tRF\_4 is a smaller fragment, with ~19nt. This is not a “common” tRF, as it does not start at the 5'end of the mature tRNA, but near its D-loop. Its biogenesis is Dicer dependent and it can also bind to Ago1 protein. tRF\_4 expression varies according to the tissue and in some tissues its

expression is higher than its corresponding tRNA. As the tRF\_3, tRF\_4 is conserved in zebrafish, humans and mouse.

Zebrafish has been used to a widely range of molecular studies. Although, until now, studies on the function of tRNA-derived fragments are scarce.

### **1.5 Objectives**

The aim of this work was to elucidate the function of two tRFs - tRF\_3 and tRF\_4 - previously described in zebrafish, by our lab. The main objective was to test if these tRNA-derived fragments have silencing ability, similarly to miRNAs and investigate putative targets that can be silenced by them.



## **2. Materials and Methods**



## 2. Materials and Methods

### 2.1. Zebrafish development and tissue tRF profiling

tRF profiling in zebrafish was performed by Northern blot. Northern blot is a technique that allows detecting specific RNAs by using a radioactive hybridization probe. For profiling of tRNA fragments during development samples from 24hpf, 48hpf, 72hpf, 10dpf, 1mpf and 2mpf were used. For zebrafish tissue profiling samples from gut, muscle, eyes, brain, fins, bones, skin and gills were used. After RNA extraction, 30µg of total RNA were fractionated on a denaturing polyacrylamide gel (10%). The RNA was transferred to an Hybond-N membrane (GE Healthcare) for 35min in a semidry transfer system followed by UV-crosslinked. Complementary antisense oligonucleotides to tRF\_3 and tRF\_4 were used as hybridization probes after being labeled with [<sup>32</sup>P]-ATP and T4 polynucleotide kinase (Takara). Membrane pre-hybridization lasted for 4hrs in hybridization buffer (5× Denhardt's solution, 1% SDS and 6,6x SSPE) at 64°C (tRF\_3) and 40°C (tRF\_4). The probes were added and incubated with the membranes over night at the previous temperatures. The membranes were washed twice with washing solution (2xSSPE and 0,1%SDS) at room temperature and twice at 57°C for 3min each and then exposed to a phosphor screen (Biorad®) overnight followed by scanning with a Molecular Imager® FX (Biorad), equipped with Quantity One FX software.

The probes used were the following:

tRF\_3: 5'-GCCGAATCCTAACCCTAGACCACCAGGGA-3'

tRF\_4: 5'-GCGAGAATCATACCCCTA-3'

### 2.2 Reporter Assay

In this work a double reporter system, provided by Huttner's lab, was used in order to verify the tRFs silencing ability (Tonelli, *et al*, 2006). The DFERS plasmid (pDSV2-EGFP-mRFP) has two fluorescent proteins, GFP (green fluorescent protein) and RFP (red fluorescent protein), which are controlled by the same promoter. GFP identifies the tissues expressing the plasmid and the RFP, which contains a 3'UTR cassette complementary to the tRF of interest, functions as a silencing sensor. Two different assays were performed. First, a total complementary sequence to the tRF\_3 and tRF\_4 was used in order to see whether they have silencing ability or not. Then, mutations were made in specific locations in order to study the tRFs functional domains.

### 2.2.1 Primer Annealing

In order to have an efficient ligation of the insert (tRF and its complementary sequence) to the DFRS plasmid, the primers must be phosphorylated. This is accomplished by using T4 PNK enzyme. A mix with 3µL of the sense and anti-sense oligo at 100µM, 3 µL of 10x PNK buffer, 2µL of 10µM ATP, 2 µL of T4 PNK and 17µL of milliQ H<sub>2</sub>O was prepared, for a total of 30µL of solution. The mix was then incubated at 37°C for 90min. Then 4µL NaCl 0,5M was added, incubated at 95°C for 2min and cool to the room temperature (~2hrs). The mix was then stored at 4°C until use. The primers used were the following:

Ctrl

FW: 5' TCGAGTGCAGTTTCGAACCTACATAACTGGATATCCTGACGTTTCGAACCTACATAACTT 3'

Rev: 5' CTAGAAGTTATGTAAGTTTCGAACGTCAGGATATCCAGTTATGTAAGTTTCGAACGTCAC - 3'

tRF\_3

Fw: 5' TCGAGGCCGAATCCTAACCCTAGACCACCAGGGAGGATATCCGCCGAATCCTAACCCTAGACCACCAGGGAT3'

Rev: 5' CTAGATCCCTGGTGGTCTAGTGGTTAGGATTCGGCGGATATCCTCCCTGGTGGTCTAGTGGTTAGGATTCGGCC3'

tRF\_3\_Mut3

Fw: 5' TCGAGGTTGCCCTCCTAACCCTAGACCACCAGGGAGGATATCCGTTGCCCTCCTAACCCTAGACCACCAGGGAT3'

Rev: 5' CTAGATCCCTGGTGGTCTAGTGGTTAGGAGGCAACGGATATCCTCCCTGGTGGTCTAGTGGTTAGGAGGCAACC3'

tRF\_3\_Mut5

Fw: 5' TCGAGGCCGAATCCTAACCCTAGACCCTCGAAAGGATATCCGCCGAATCCTAACCCTAGACCCTCGAAAT3'

Rev: 5' CTAGATTTCGAGTGGTCTAGTGGTTAGGATTCGGCGGATATCCTTTTCGAGTGGTCTAGTGGTTAGGATTCGGCC3'

tRF\_4

FW: 5' TCGAGGCGAGAATCATACCCCTAGGATATCCGCGAGAATCATACCCCTAT3'

Rev: 5' CTAGATAGGGGTATGATTCTCGCGGATATCCTAGGGGTATGATTCTCGCC3'

tRF\_4\_Mut3

FW: 5' TCGAGGTAAAGATCATACCCCTAGGATATCCGTAAAGATCATACCCCTAT3'

Rev: 5' CTAGATAGGGGTATGATCTTTACGGATATCCTAGGGGTATGATCTTTACC3'

tRF\_4\_Mut5

FW: 5' TCGAGGCGAGAATCATATTCACAGGATATCCGCGAGAATCATATTCACAT3'

Rev: 5' CTAGATGTGAATATGATTCTCGCGGATATCCTGTGAATATGATTCTCGCC3'

**Figure 10.** Primers inserted in DFRS plasmid. The primers are composed by two sequences (black) separated by EcoRV cleavage site (orange). Restriction sites are shown in blue and mutated nucleotides in red.

Forward and reverse primers both from tRF\_3 and tRF\_4 were ordered from Sigma and the tRF\_3\_Mut3, tRF\_3\_Mut5, tRF\_4\_Mut3, tRF4\_Mut5 were ordered from Eurofins.

### 2.2.2 Ligation

The ligation of the insert to the plasmid was performed following the protocol described below. Briefly, three mixes with different ratios of plasmid (concentration 10ng/µL) and insert (concentration 10ng/µL) were prepared: 1:0, 1:3 and 1:5, respectively. 2µL of T4 DNA ligase 5x buffer and 1µL of T4 DNA ligase were added to each mix (ligation reaction) in a total volume of 20 µL. All ligation reactions were incubated at 16°C overnight (Figure 11).



**Figure 11.** Plasmid construction. The DFRS plasmid contains GFP (green) and a mRFP (red) sequences. Both are controlled by similar promoters (SV40). mRFP 3'UTR, where all the sequences were inserted, is shown in blue. PA is the polyadenilation site.

### 2.2.3 Bacterial Transformation

Transformation is the process by which cells uptake a foreign DNA present in its surroundings. In this case the cell is *E.coli* and the DNA is the DFRS plasmid. 20  $\mu$ L of plasmid DNA was added to 200  $\mu$ L competent cells, incubated on ice for 30 min, heat shocked at 42°C for 90 sec, placed on ice and to it, 800  $\mu$ L SOC media was added. The cultures were then covered and incubated with shaking at 37 °C for 1 hour. Positive transformants were selected by plating on LB agar medium containing ampicilin and growing overnight at 37 °C.

### 2.2.4 Plasmid Extraction and Linearization

Plasmid extraction was performed with GeneJET™ Plasmid Miniprep Kit from Fermentas, following the manufacturers' protocol and quantified by nanodrop. The plasmid was sequenced by stabvida.

For a most effective expression in zebrafish, the plasmid has to be injected in its linear form. The linearization was performed using 2 $\mu$ L of 10x FD buffer, 2 $\mu$ L of plasmid DNA (up to 12 $\mu$ g), 1 $\mu$ L of FastDigest BglII and miliQ H<sub>2</sub>O up to a final volume of 20  $\mu$ L. The mix was then incubated for 40 minutes at 37°C. The final digestion product was cleared using QIAquick® Nucleotide Removal Kit from QIAGEN, following the manufactures' protocol. Briefly, 100  $\mu$ L of PNI buffer was added and the mixture was applied to the QIAquick spin column followed by centrifugation (6000rpm, 1min). 750  $\mu$ L of PE buffer was added and the mixture was centrifuge first at 6000rpm, 1min and then

13,000 rpm, 1min to remove residual ethanol residues from PE buffer. 30 µL of MiliQ water were applied to the centre of the QIAquick membrane and after 2 min the tubes were centrifuged (13rpm, 1min).

### **2.2.5 Zebrafish maintenance**

In this study wild-type AB zebrafish embryos were used. The embryos were collected and kept at 28°C under standard laboratory conditions. Wild-type AB zebrafish were maintained in CESAM laboratory zebrafish facility in Aveiro University. The study was approved by the DGV (Direção-Geral de Veterinária).

### **2.2.6 Zebrafish Embryos Microinjection**

One cell zebrafish eggs were microinjected with a solution containing 20ng/µL of the reporter plasmid. The solution was made with Phenol red and 0,9M KCl.

One-cell stage eggs were injected with approximately 1000pL of the solution. Only wild type AB zebrafish strain was used in this study and the embryos were collected and kept at 28°C in the CESAM's zebrafish facility of Aveiro University.

They were analyzed at 24hpf by fluorescent microscopy.

## **2.3 Gene Target Prediction**

3'UTRs from zebrafish mRNAs were acquired from Biomart (<http://www.biomart.org>) and blasted against tRF\_4 sequence. Those which showed perfect match between nucleotides 2 and 7 (seed region) and no more than 5 mismatches were confirmed thermodynamically by RNAhybrid and considered for further analysis.

### **2.3.1 Duplex Injections**

tRF\_3 and tRF\_4 RNA duplexes, a duplex containing a scrambled sequence were obtained as siRNAs, the first two from IDT and the last one from SIGMA:

#### **tRF\_3 duplex:**

sense: 5'-UCCCUGGUGGUCUAGUGGUUAGGAUUCGGC-3'

antisense: 5'-GCCGAAUCCUAACCACUAGACCACCAGGGA-3'

#### **tRF\_4 duplex:**

sense 5'-UAGGGGUAUGAUUCUCGC-3'

antisense 5'-GCGAGAAUCAUACCCCUA-3'

#### **Scrambled duplex:**

sense 5'-GUGUAACACGUCUAUACGCCCA-3';

antisense 5'-GGCGUAUAGACGUGUUACAC [dT][dT]-3';

The siRNAs were injected into one-cell fertilized embryos at 5  $\mu$ M concentration in TE Buffer (pH=7,5). Approximately 1000pL of the RNA duplex were injected.

### **2.3.2 Alcian Blue Assay**

Alcian blue assay stains cartilage and bones allowing its visualization and analysis.

Five days post fertilization (dpf) zebrafish embryos were fixed for 5hrs in 4% (p/v) phosphate-buffered formaldehyde and maintained in methanol 100% at -20°C until use. The embryos were washed 2 times in phosphate-buffered saline with 0.1% Tween-20 (PBT). Then they were bleached in 30% hydrogen peroxide until the eyes became sufficiently translucent (~2hrs). Next the embryos were rinsed twice with 1 ml of PBT and transferred into 0,5ml of an Alcian blue solution (1% concentrated hydrochloric acid, 70% ethanol, 0.1% Alcian blue), overnight. The specimens were rinsed 3 times with 1 ml acidic ethanol (5% concentrated hydrochloric acid, 70% ethanol, HCl-EtOH) and incubated for 20 min in 1ml of the same solution. Re-hydration was performed in 1ml HCl-EtOH/H<sub>2</sub>O<sub>d</sub> series: 75% / 25% HCl-EtOH/ H<sub>2</sub>O<sub>d</sub>; 50/50 HCl-EtOH/ H<sub>2</sub>O<sub>d</sub>; 25/75 HCl-EtOH/ H<sub>2</sub>O<sub>d</sub> and 100% H<sub>2</sub>O<sub>d</sub>. Finally the H<sub>2</sub>O<sub>d</sub> was removed and the embryos were stored in 1ml KOH. The embryos were visualized in a dissecting microscope (Nikon SMZ1500).

### **2.3.3 RNA extraction**

The RNA was extracted using TRIZOL® Reagent, from approximately 100 injected embryos (24hpf), and following the manufacturer's protocol. For cDNA preparation the samples were treated with DNase I (Invitrogen), according to the protocol provided by Invitrogen, with minor changes. Briefly, 500ng of total RNA were mixed with 1 $\mu$ L DNase I and DNaseI buffer incubated for 40 min at 37°C. 1 $\mu$ L of EDTA is added in order to inhibit the reaction and the mix is incubated for 10 min at 65°C. Two RNA extractions were performed, for the first 100  $\mu$ L phenol:chlorophorm:iso-amyl was used followed by centrifugation (12,000rpm for 15minutues at 4°C ), for the second 1volume of chlorophorm was added to the supernatant and the samples were centrifuge (12,000rpm for 15minutues at 4°C). The supernatant was removed, the pellet washed with 500 $\mu$ L of 75% ethanol and centrifuged again at 7,500rpm for 5minutes at 4°C. The pellet was dried, resuspended in 30 $\mu$ L of miliQ H<sub>2</sub>O and quantified by nanodrop.

### **2.3.4 cDNA preparation**

cDNA is DNA obtained from mRNA template in a reaction catalysed by the enzymes reverse transcriptase. This technique can be combined with PCR (RT-PCR) allowing detection of gene expression.

cDNA was prepared using SuperScript®II RT (Invitrogen) from 500ng of total RNA. In the first step 1µL of Oligo d(T) 12-18, 1µL of dNTPs mix (10µM) and H<sub>2</sub>O up to a total volume of 12µL were added. The mix was incubated at 65°C for 5 minutes and then cooled down for another 5 minutes. Next, 4µL of 5xFirst Strand buffer, 2µL of 0,1 µM DTT and 1µL of RNase out were added. The solution was mixed and incubated at 42°C for 2 minutes. One uL of SSII RT was added and the samples were incubated for 15 minutes at 70°C. Then 0,5µL of RNase H was added and after incubated for 20 minutes at 37°C.

### **2.3.5 sec23b and tub1a RT-qPCR**

In order to quantify sec23b transcripts a quantitative real time PCR was performed. qPCR was carried out by using the synthesized cDNA and SYBR® Green detection reagent, Universal qPCR Primer provided in the kit, and the forward primer for sec23b and tub1a (the endogenous control) detection (the same used for RT-PCR). It was performed in the ABI Prism 7500 Real time PCR System (Applied Biosystems) and the reactions were incubated in a 96-well optical plate. The cycling began with template denaturation at 95°C for 2min, followed by 40 cycles at 95°C for 30 sec and 60°C for 1 min. The samples and the no-template controls were carried out in triplicate and run at the same time.

The primers used were the following:

**Sec23b\_fw**–5'-TGGTGGGATCGGCCATGTG-3';

**Tub1a\_fw**–5'-GGCATCAACTACCAGCCTCC-3';



## **3. Results**



### 3. Results

#### 3.1 tRF\_3 and tRF\_4 profiling


The tRF\_3 is a 30nt non-coding RNA derived from the 5' end of a mature tRNA (Figure 12). Previous work from our lab has shown that this fragment requires Dicer for its biogenesis and that its expression is conserved in humans, mouse and in zebrafish. The tRF\_4 derives from the 5' end of a mature tRNA and it is 19nt long (Figure 12). This fragment is rather peculiar as it does not start near the D-loop and not in the 5' end. Its biogenesis is also dependent on Dicer and it derives from tRNAs that can acquire pre-miRNA-like conformations. Given all these evidences and the similarities between this fragment and miRNA/siRNA molecules, it is possible to assume that tRF\_4 is not a random product of tRNA degradation and may have analogous biological functions in the cell.

```
tRF_3      Type: Glu      Anticodon: CTC
TCCCTGGTGGTCTAGTGGTtAGGATTcGGcGCTCTcACCGCCGCAGCCCGGGTTCGATTCCCGGTCAGAGAA
TCCCTGGTGGTCTAGTGGTtAGGATTcGGc


tRF_4      Type: Pro     Anticodon: CGG
GGCTTGTTGGTCTAGGGGTATGATTCTCGCTTAGGGTGTGAGAGGtCCCGGGTTCAAATCCTGGACGAGCCC
TAGGGGTATGATTCTCGC
```

**Figure 12.** tRF sequences and their possible mature tRNAs. tRFs are shown in orange and the mature tRNA in black. Sequences are written in 5' to 3' direction. Glu – Glutamine; Pro-Proline

Despite of being a tRNA halve, previous results shown that the tRF\_3 expression is not related to stress conditions in zebrafish and it is different from tissue to tissue. In this study the expression of this fragment was analysed both during zebrafish development and in its different tissues in adulthood. It was possible to see that its expression increases during its development (Figure 13), being more prominent in adult tissues such as gut, skin, bone and fins (Figure 14).




**Figure 13.** tRF\_3 abundance increases during development. tRF\_3 development profiling was performed by northern blot, using RNA samples from different stages of development (24hpf, 48hpf, 72hpf, 10dpf, 1mpf and 2mpf).




**Figure 14.** tRF\_3 is tissue specific. tRF\_3 tissue profiling was obtained by northern blot using RNA samples from different tissues from adult fish.

Like the tRF\_3, tRF\_4 has also shown to be differently expressed from tissue to tissue. Its profiling revealed that it increases during development, especially after 10 days post fertilization (Figure 15) and it is more abundant in certain tissues, namely eyes, gills, bone, gut and skin (Figure 16). It was also possible to see that in these tissues, the abundance of the fragment is higher than the mature tRNA, which confirmed the previous results obtained in the lab (Figure 16).



**Figure 15.** tRF\_4 expression increases during development. Development profiling of tRF\_4 was performed using RNA samples from different stages of development (24hpf, 48hpf, 72hpf, 10dpf, 1mpf and 2mpf), by northern blot.




**Figure 16.** tRF\_4 expression is tissue specific. tRF\_4 profiling was obtained by northern blot using RNA samples from different tissues from adult fish.

### 3.2 tRF\_3 and tRF\_4 have Silencing Ability


In order to study the silencing ability of the highly expressed tRF\_3 and tRF\_4 in zebrafish, a dual-fluorescence GFP-Reporter/mRFP-Sensor plasmid (Dual fluorescence reporter system – DFRS) was injected in one cell-stage zebrafish embryos. The zebrafish was chosen as a model to this work, as the fragments in this study had already been reported in this organism and are conserved among vertebrates (unpublished data). The DFRS plasmid contains both GFP and mRFP under the control of identical constitutive promoters, the SV40 promoter. As the sensor-based strategy relies on the silencing of a transcript, the GFP-reporter is used to identify the cells that are actually expressing the plasmid, and the mRFP-sensor contains a 3'untranslated region with a tandem cassette, which is complementary to the tRF of interest (Tonelli, *et al*, 2006). Therefore, only the embryos expressing GFP, and hence the DFRS plasmid were analysed and the expression

of the injected plasmid was visible in the zebrafish embryo muscle. For each condition three types of pictures were taken, one showing only GFP signal, another showing RFP signal and a third containing both GFP and RFP signals (Merge).

Using the DRFS plasmid, with the reverse complementary sequence to the tRFs inserted in the mRFP 3'UTR, it was evident that tRF\_3 and tRF\_4 have silencing ability, as there was a decrease in the expression of mRFP signal when compared with the control. For both tRF\_3 and tRF\_4, the majority of the embryos injected with the plasmid containing the GFP and the RFP, it was visible that the RFP signal was absent, although the GFP was still detectable (Figures 17 and 18 respectively). This means that the plasmid was being expressed in the zebrafish embryo muscles cells, but due to their ability for silencing by establishing a bound with a complementary sequence, the RFP signal was not detected. However, it was also noticed that the silencing ability for the tRF\_4 fragment is stronger than in tRF\_3. This may be due to the higher expression of tRF\_4 compared to tRF\_3 in these tissues (Figure 16). The tRF\_3 and tRF\_4 have, therefore, ability to recognize a complementary sequence as a target site and suppress its expression.



**Figure 17.** tRF\_3 has silencing ability. A, B, C – Embryo injected with 20ng/μl of DRFS control plasmid, showing GFP, mRFP and merge signal respectively. D, E, F - Embryo injected with 20ng/μl of DRFS plasmid containing the reverse complementary sequence to the tRF\_3, showing GFP, mRFP and merge signal respectively.




**Figure 18.** tRF\_4 has silencing ability. A, B, C – Embryo injected with 20ng/ $\mu$ l of DFRS control plasmid, showing GFP, mRFP and merge signal respectively. D, E, F - Embryo injected with 20ng/ $\mu$ l of DFRS plasmid containing the reverse complementary sequence to the tRF\_4, showing GFP, mRFP and merge signal respectively

### 3.2.1 Functional domains

As stated before, miRNAs usually use their 5' end, more specifically, the region between nucleotide 2 and 8, to target and bind to an mRNA target. This region, also known as “seed”, must be totally complementary to the target mRNA, in order to accomplish an efficient silencing. As the fragments in study, specially tRF\_4, have similar features to a miRNA, mutations were inserted in the complementary reverse sequence (present in the DFRS plasmid) in order to affect binding to the 3' portion or the 5' portion of the tRFs, and, therefore, study whether or not they would also need a “seed” region for an effective silencing ability, similarly to miRNAs.

Neither mutations introduced in the tRF\_3 reporter affected considerably tRF\_3 silencing ability. In fact, there was no total recovery of fluorescence after the injection of both reporters. Therefore, it was not evident that a portion of this fragment was more critical for the silencing than the other (Figure 19).


It is clear that tRF\_3 can recognize a complementary sequence as a target site and therefore has silencing ability. However, no region is crucial for the silencing ability of this molecule, indicating that there may be other feature that is responsible for it.



**Figure 19.** tRF\_3 do not have a seed region. A,B,C – Embryos injected with the tRF\_3 reverse complementary sequence plasmid showing GFP. mRFP and merge respectively. D,E,F – Embryos injected with a plasmid containing a sequence with mutations affecting 3'-portion complementarity. D,E,F – Embryos injected with a plasmid containing a sequence with mutations affecting 5'-portion complementarity.

For the tRF\_4 it was possible to see fluorescence recovery in embryos injected with the tRF\_4\_Mut5 DFRS plasmid, which means that, when the complementarity affected is in the 5'-portion of the tRF\_4, it cannot recognize its target and the silencing ability is compromised. On the other hand, the embryos injected with the tRF\_4\_Mut3 DFRS plasmid, did not display the red fluorescence. In this case, the complementarity was only being affected in the fragment's 3'-portion, leaving the 5'-portion free for targeting and silencing the RFP transcript. Thereby, it is possible to conclude that 5' portion was essential for target recognition and silencing ability, as when this region was affected by the mutation, the tRF\_4 was not able to recognize the tRF\_4\_Mut5 sequence, not even the regions that kept the complementary (Figure 20).





**Figure 20.** tRF\_4 5-portion is crucial for silencing. A,B,C – Embryos injected with the tRF\_4 reverse complementary sequence plasmid showing GFP, mRFP and merge respectively. D,E,F – Embryos injected with a plasmid containing a sequence with mutations affecting 3'-portion complementarity to the tRF\_4. G,H,I– Embryos injected with a plasmid containing a sequence with mutations affecting 5'-portion complementarity to the tRF\_4.

This new tRF\_4 feature as well as its size, its Dicer-dependent biogenesis and its ability to bind Ago proteins, reinforces the theory that this fragment, for resembling a miRNA, can act as such, entering a similar pathway and, therefore may have similar functions. miRNA molecules are well known for its ability for targeting and silencing, by binding to the target mRNA using their 5'-portion (seed region). In fact, when the seed region is affected, for example by a mutation, the miRNA cannot recognize its target and loses its silencing ability. In the tRF\_4 case, the mutations inserted in the reverse complementary sequence (tRF4\_Mut5), affected the 5'-region and inhibited tRF\_4 to recognize its target, similarly to miRNA molecules.

### 3.3 Gene Target Prediction

Since the tRF\_4 has miRNA-like features, it is possible to assume that this fragment will be capable of silencing genes. Therefore, bioinformatics gene target

prediction was performed. These predictions were performed taking into account that tRF\_4 silencing ability is dependent on its 5' portion and no more than 5 mismatches were allowed in the remaining sequence. Interestingly, the best gene target predictions correspond to genes that are expressed in bone and muscular tissues, which are tissues where tRF\_4 is highly expressed as shown previously. These genes were Sec23 homolog B (*S.cerevisiae*) and parvin, alpha a.

Sec23b: 5' GTTTGAATGATACCCCTG 3'

Parvin, alpha a: 5' CTGTTAATTATACCCCTA 3'


**Table 1.** Putative gene targets for tRF\_4. Mismatches are shown in red

Putative Target Gene	N. of mismatches	RNAhybrid predictions	Minimum free energy (kcal/mol)
Sec23 homolog B ( <i>S.cerevisiae</i> )	5	<pre> target5' A   UUU   G           G 3'               GUG   GAAU AUACCCCU               CGC   CUUA UAUGGGGA tRF_4 3' U           G           5'           </pre>	-25,3
Parvin, alpha a	5	<pre> target5' A   UUU   G           G 3'               GUG   GAAU AUACCCCU               CGC   CUUA UAUGGGGA tRF_4 3' U           G           5'           </pre>	-23,8

Sec23b is located in the zebrafish's chromosome 6 and produces a 766aa protein (NCBIa, 2013; UniProtKB, 2013). This gene is conserved in human (20p11.23), chimpanzee (chromosome20), Rhesus monkey (chromosome 10), dog (chromosome 24), cow (chromosome 13), mouse (chromosome 2), chicken (chromosome 3) (NCBIa, 2013).

Sec23B is part of the coat protein II complex (COPII), along with other proteins like Sec23A, Sec24A, Sec24B, Sec24C and Sec24D. This complex is involved in protein traffic between endoplasmic reticulum and Golgi complex. Sec23 proteins activate Sar1GTPase, which recruits COPII to the ER membrane. Sec24 proteins are involved in cargo sorting (Figure 21) (Melville, Montero-Balaguer, Levic, Bradley, Smith, & Hatzopoulos, 2011). One of the proteins transported by endoplasmic reticulum–Golgi intermediate compartment (ERGIC), in a COPII-dependent manner, is protocollagen, which is an important component for cartilage development (Lang, *et al*, 2006).

Morpholino injections in zebrafish showed that *sec23b* knockdown leads to a complete loss of ventral pharyngeal skeleton, under developed neurocranium and absence of ear capsules. Deficits in neurocranium growth were also observed in *Sec23a* and *Sec23b* experiments, leading to the conclusion that *Sec23* genes are crucial for embryonic neurocranium morphogenesis and cartilage development (Lang, *et al.*, 2006). *Sec23b* has been reported to be involved in congenital dyserythropoietic anemia II (CDAII), as it plays a role in erythrocytes development and differentiation (Schwarz, *et al.*, 2009).




**Figure 21.** Protein processing in endoplasmic reticulum in *Danio rerio* (KEGG, 2010). New synthesized peptides enter the endoplasmic reticulum through the *sec61* pore and are glycosylated and folded. Properly folded proteins are then carried to the Golgi complex in transport vesicles. 60-90nm COPII (*Sec23*-*Sec24* and *Sec13*-*Sec31* heterodimeres) vesicles transport small globular proteins.

Parvin alpha is located in the chromosome 25 in zebrafish and is conserved in human (11p15.3), chimpanzee (chromosome 11), Rhesus monkey (chromosome 14), dog (chromosome 21), cow (chromosome 15), mouse (chromosome 7), rat (1q33), and chicken (chromosome 5) (NCBIb, 2013). This gene encode for 372aa actin-binding protein, which is a member of the parvin family. These proteins are present on focal contacts as they contain calponin homology domains that bind to the actin filaments. Parvin alpha play a role in the integrin-linked kinase signaling complex and is involved in cell adhesion,

motility and survival. It is also involved in sprouting angiogenesis, outflow tract septum morphogenesis and smooth muscle cell chemotaxis (NCBIc, 2013).

### 3.3.1 Alcian blue staining


Alcian blue solution allows the visualization of cartilage patterns in larvae and adult zebrafish as it dyes proteoglycan components of the extracellular matrix associated with chondrocytes. This assay may be performed in 54hpf embryos, but the full pattern of early cranial cartilage will only be visible at 72hpf (Javidan and Schilling, 2009). In a 5dpf zebrafish, stained with alcian blue it is possible to distinguish in a ventral view the ethmoid plate (eth), which is formed by the medial fusion of the two trabeculae. In the mandibular arch (first pharyngeal arch) is located the Meckel's cartilage (mk) and the quadrate (qu) and in the hyoid arch (second pharyngeal arch) is possible to see the basihyal (bh), the ceratohyal (ch) and the hyosymplectic (hys). The gill arch derivatives are the basibranchials (bb) and the ceratobranchials (cb), which is composed by five arches. In a lateral view besides the ethmoid, the Meckel's cartilage and the ceratohyal it is also possible to see the pterygoid process of the quadrate (pty), the auditory capsule (aud) (figure 22) (Neuhauss, et al., 1996)



**Figure 22.** 5 day-post-fertilization zebrafish larvae skeleton stained for cartilage with Alcian Blue (Neuhauss, et al., 1996). A. Ventral view B. Lateral view

Alcian blue cartilage staining assay was performed for embryos injected with scrambled\_duplex, tRF\_3\_duplex, tRF\_4\_duplex. Duplex injection was performed in order to study the over expression of the tRFs. These duplexes may target and silence different mRNAs, and if the target is sec23b the embryos will show phenotypes similar to sec23b knockdown. The concentration 8 $\mu$  was chosen in order to avoid unspecific phenotypes


derived from duplex toxicity. It has been reported, that for higher concentrations, embryos show truncated tails, loss of eye and brain structures, enlarged heart cavities and growth retardation (Gruber, *et al.*, 2005). In embryos injected with tRF\_3\_duplex no alteration was visible in cartilage development. Both the mandibular and the hyoid arches were well developed as well as the five ceratobranchial arches. In addition, the head, eyes and mouth appeared to have the right shape and size (figure 23). The lack of phenotype in fishes injected with this duplex allowed concluding that this tRF is not involved in cartilage and bone related genes regulation, which was expected as this tRF is not expected to silence genes that affect these pathways. Nevertheless we cannot exclude that this tRF may be involved in other regulatory pathways not yet explored.



**Figure 23.** tRF\_3 over-expression does not affect cartilage development. 5dpf zebrafish. Zebrafish 1A and 1B were injected with a 2 $\mu$ m solution containing a siRNA scramble sequence which does not have any target in zebrafish. Zebrafish 2A and 2B were injected with a 8 $\mu$ m solution containing a siRNA tRF 3 sequence. A. Lateral view; B. Ventral view.


On the other hand, embryos injected with tRF\_4\_duplex showed significant alterations. It was possible to see that the first and the second arches were severely disformed, wider than the one seen in the scrambled embryos. Also, the first arch is positioned closest to the mouth than normal (Figure 20.2B arrows), which may affect its opening. Furthermore, the ceratobranchial arches present an unusual shape and the ethmoid plate is underdeveloped. It was also possible to see that the head was smaller and the eyes size and shape indicated underdevelopment (Figure 24.2B arrows). These phenotypes

were similar to those observed by Lang and colleagues in *sec23b* knock down after a morpholino microinjection.



**Figure 24.** tRF\_4 over-expression affects cartilage development in zebrafish. 5dpf zebrafish. Zebrafish 1A and 1B were injected with a 2µm solution containing a siRNA scramble sequence. Zebrafish 2A and 2B were injected with a 8µm solution containing a siRNA\_tRF\_4 sequence. A. Lateral view; B. Ventral view.

Given that the previous results indicated that *sec23b* could be affected by duplex injections, its mRNA levels were analyzed, by RT-qPCR, in order to investigate whether or not these changes could result from a decrease in *sec23b* expression. Embryos injected with scrambled duplex were used as a control between samples and the internal control used in this assay was tubulin  $\alpha$ . There was not any significant alteration between the controls and the embryos injected with the siRNA\_tRF\_4 (Figure 26).



**Figure 25.** *sec23b* does not alter its expression in the presence of *tRF\_4*. Scrambled was used as an external control and *tub1a* as an internal control. Both scrambled and *tRF\_4* were normalized using *tub1a*. *tRF\_4* is a mean of 2 replicates.

Despite the previous evidences (gene target prediction and alcian blue assay) point to the regulation of *sec23b* by *tRF\_4*, there was no significant difference in its mRNA abundance when *tRF\_4* was over-expressed. However, it was clear that this tRF is capable of binding and silencing a target gene by using its 5'-portion and that it interferes with cartilage and bone development. As the tRFs silencing pathway is not well understood yet, it is possible that the *tRF\_4* can silence *sec23b* in other way rather than mRNA degradation.





## **4. Discussion**



#### 4. Discussion

In this work two specific tRFs found in Zebrafish were studied. Both of them derived from the 5' portion of different tRNAs. tRF\_4 is derived from a proline tRNA and tRF\_3 from a glutamine tRNA and have different sizes, 19nt and 31nt respectively.

Regarding their expression, it was possible to observe that their production varies according to the developmental stages and tissues. tRF\_3 expression dramatically increases from 1 month to 2 months, and in adulthood is highly expressed in gut, skin, bone and fins, while in brain its abundance is low. On the other hand, during development, tRF\_4 expression increases substantially at 10dpf. This fragment is extremely abundant in eyes, gills, bone, gut and skin but little expression was noticed in fins. These results may indicate that tRF expression is controlled and may play a role, during development, in tissue differentiation, and, in adulthood, in keeping tissue identity.

As for their silencing ability, this work clearly shows that both of the fragments can silence a target mRNA.

Concerning tRF\_3, it was possible to conclude that it has silencing ability, as it was able to recognize and target its reverse complementary sequence inhibiting mRFP expression. However, it did not appear to need any functional domain for silencing. This fragment was also studied by Wang and colleagues using A549 cells after a respiratory syncytial virus infection (Wang, *et al*, 2012). Using a different reporter system (luciferase sensor plasmid), they also saw a decrease in luciferase signal when compared with the control, concluding that tRF5-GluCTC has silencing ability. Yet, they stated that the 3'portion of the fragment was the essential domain for targeting and therefore for its silencing ability (Wang, *et al*, 2012).

The differences between the results may be due to the different organisms and conditions in study. This study was performed using a whole organism (zebrafish embryos) which was not exposed to any kind of stress, while Wang and colleagues performed their experiences in cell lines exposed to a stress (infection). In fact, Wang et al reported that its biogenesis pathway is angiogenin-dependent (Wang, *et al*, 2012), which is not consistent to Dicer-dependent biogenesis (unpublished data) saw in zebrafish. It is possible that the same fragment may arise in different cell conditions, from different biogenesis pathways and have different functions.

Overall, it is possible to conclude that tRF\_3 can target and silence a RNA sequence, yet in a miRNA/siRNA distinct pathway. In fact, Sobala and Hutvagner,

proposed that 5'tRFs, which have a "GG" dinucleotide sequence either in position 17-18, just like the tRF\_3, can, in some cases, need these dinucleotides to repress translation (Sobala and Hutvagner, 2013). Further studies should be performed, namely by mutating "GG" dinucleotides, in order to evaluate whether or not the guanosine residues could play a role in tRF\_3 silencing ability.

The tRF\_4 has not been described yet, by any other investigators. It is known that its size is smaller (~19nt) than tRF\_3, and its biogenesis is Dicer-dependent. Like the tRF\_3 this fragment can also target and silence a specific RNA target. However, it was clear that this fragment behaves like a miRNA. It requires its 5'-portion, also known as "seed" region, to be able to silence gene expression.

Based on the previous results, target predictions were then performed for tRF\_4 and several genes were listed. From those, only *sec23b* and *parvin alpha* were considered in this study. These genes are totally complementary to the tRF\_4 "seed" region, have 5 mismatches and contribute for cartilage and muscle development respectively. Over-expression of tRF\_4 was then performed by injecting a duplex and phenotype alterations were analysed by Alcian blue. This assay showed phenotypes consistent with a *sec23b* under-expression. The ceratobranchial arches presented an unusual shape, being the first arch closest to the mouth than normal. The embryos revealed to be underdeveloped, showing smaller heads and eyes. This phenotype was similar to those seen by Lang and colleagues, when using a *sec23b* morpholino (Lang, *et al*, 2006). Yet, their phenotypes were slightly more severe. These differences may be due to the concentration of duplex used in this work. As the high concentrations of duplexes may induce unspecific phenotypes, a small concentration was injected in zebrafish embryos. Therefore, the phenotypes obtained were less severe.

Although Alcian blue results pointed to *sec23b* loss of function, the PCR results revealed normal *sec23b* transcriptional levels. It is possible that the silencing ability of this fragment may not be through the *sec23b* degradation. tRFs silencing pathway is still unknown and it is possible that silencing may be due to blocking translation itself and not by mRNA degradation.

Furthermore this fragment may silence in other position like the 5'UTR or the coding region or it is also possible that *sec23b* is not the tRF\_4 target. Zebrafish genome is continually actualized in databases and some sequences remain provisional. The

development of new sequencing technologies and continuous studies in this field might reveal new and more accurate sequences that may be suitable gene targets.

In order to clarify whether or not *sec23b* is the tRF\_4 target two different approaches could be made. One by using a GFP reporter plasmid: the GFP 3'UTR could be replaced with the 3'UTR of *sec23b* and see if in this case the GFP signal disappears. The other by using western blot to quantify the amount of *sec23b* protein in fishes over-expressing tRF\_4.



## **5. Conclusion**





## **5. Conclusion**

Although the RNA has been discovered for more than 50 years, there is still a lot to discover in this field. Small non-coding RNAs revolutionized the way scientists viewed gene expression and novel classes, functions and targets are uncovered every day. Recently, new sequence technologies and bioinformatics algorithms helped to identify a novel non-coding RNA class, which was thought to be a degradation product, the tRNA-derived fragments. Their controlled biogenesis, especially the use of specific cleavage enzymes, their specific length and cleavage sites and their conservation among species lead to the conclusion that they were in fact novel biological identities.

Through this work it was possible to conclude that these particular fragments can play a role in gene expression regulation. In fact, tRF\_4 has demonstrated to share several characteristics like size, functional domain and most likely silencing pathway to a miRNA.

Further studies should be performed in order to unveil more tRFs and to understand their exact biogenesis, mechanistic, pathway and function in the organism. Their similarity to miRNAs may allow them to recruit the miRNA machinery, blocking miRNA function and leading to aberrant phenotypes. On the other hand, they may have their own machinery and cooperate with other small non-coding RNAs on controlling important biological processes.



## **6. Bibliography**



## 6. Bibliography

- Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., & Walter, P. (2008). *The molecular Biology of the Cell* .5<sup>th</sup> ed. New York: Garland Science.
- Arraiano, C., and Fialho, A. (2007). *O mundo do RNA - novos desafios e perspectivas futuras* .1<sup>st</sup> ed. Lousã: Lidel.
- Brown, T. (2002). *Genomes 2*. 2<sup>nd</sup> ed. New York: John Wiley & Sons Inc.
- Buhler, M., Spies, N., Bartel, D., and Moazed, D. (2008). TRAMP-mediated RNA surveillance prevents spurious entry of RNAs into the *Schizosaccharomyces pombe* siRNA pathway. *Nat Struct Mol Biol* , 15, 1015–1023.
- Carthew, R., and Sontheimer, E. (2009). Origins and Mechanisms of miRNA and siRNA. *Cell* , pp. 642-655.
- Chitramuthu, B. (2013). Modeling Human Disease and Development in Zebrafish. *Human Genet Embryol* , pp. 1-3.
- Cole, C., Sobala, A., and Lu, C. (2009). Filtering of deep sequencing data reveals the existence of abundant Dicer-dependent small RNAs derived from tRNAs. *RNA* , 15, pp. 2147-2160.
- Couvillion, M., Sachidanandam, R., and Collins, K. (2010). Ravi Sachidanandam Tetrahymena Piwi protein carries tRNA fragment cargo. *Genes & Development* , pp. 2742–2747.
- Elliott, D., and Lodomery, M. (2011). *Molecular Biology of RNA*. 1<sup>st</sup> ed.. Oxford University Press.
- Emara, M., Ivanov, P., Hickman, T., Dawra, N., Tisdale, S., Kedersha, N., *et al.* (2010). Angiogenin-induced tRNA-derived Stress-induced RNAs Promote Stress-induced Stress Granule Assembly. *The Journal of Biological Chemistry* , pp. 10959-10968.
- Fjose, A., and Zhao, X. (2010). Exploring microRNA functions in zebrafish. *New Biotechnology* , pp. 250-255.

- Fu, H., Feng, J., Liu, Q., Sun, F., Tie, Y., Zhu, J., *et al.* (2009). Stress induces tRNA cleavage by angiogenin in mammalian cells. *FEBS Letters* , pp. 437-442.
- Garcia-Silva, M., Cabrera-Cabrera, F., Güida, M., and Cayota, A. (2012). Hints of tRNA-Derived Small RNAs Role in RNA Silencing Mechanisms. *Genes* , pp. 603-614.
- Gebetsberger, J., Zywicki, M., Kunzi, A., and Polacek, N. (2012). tRNA-Derived Fragments Target the Ribosome and Function as Regulatory Non-Coding RNA in *Haloferax volcanii*. *Archaea* , pp. 1-11.
- Ghildiyal, M., and Zamore, P. (2009). Small silencing RNAs: an expanding universe. *Nature reviews. Genetics*, *10*, pp. 94-108.
- Griffiths, A. (2004). *An Introduction to Genetic Analysis* .8<sup>th</sup> ed.. W. H. Freeman.
- Gruber, J., Manninga, H., Tuschl, T., Osborn, M. and Weber, K. (2005). Specific RNAi Mediated Gene Knockdown in Zebrafish Cell Lines. *RNA*, pp. 101-105.
- Haussecker, D., Huang, Y., Lau, A., Parameswaran, P., Fire, A., and Kay, M. (2010). Human tRNA-derived small RNAs in the global regulation of RNA silencing. *RNA* , *16*, pp. 673–695.
- Howe, K., Clark, M., Torroja, C., Torrance, J., Berthelot, C., Muffato, M., *et al.* (2013). The zebrafish reference genome sequence and its relationship to the human genome. *Nature* , pp. 1-6.
- Huang, Y., Zhang, J., Yu, X., Xu, T., Wang, Z., and Cheng, X. (2013). Molecular Functions of Small Regulatory Noncoding RNA. *Biochemistry (Moscow)* , *78*, 221-230.
- Hussain, M. (2012). Micro-RNAs (miRNAs): genomic organization, biogenesis and mode of action. *Cell Tissue Res* , pp. 405-413.
- Ivanov, P., Emara, M., Villen, J., Gygi, S., and Anderson, P. (2011). Angiogenin-Induced tRNA Fragments Inhibit Translation Initiation. *Molecular Cell* , pp. 613-623.
- Javidan, Y., and Schilling, T. (2009). Development of Cartilage and Bone. In M. Westerfield, L. Zon, and W. Detrich, *Essential Zebrafish Methods*. Oxford: Elsevier., pp. 415-436

- Kang, C. (2011). *Gene Therapy - Developments and Future Perspectives* . 1<sup>st</sup> ed.. InTech.
- Kardash, E. (2013). *Labome*. Obtained on 17<sup>th</sup> April 2013, from <http://www.labome.com/method/Current-Methods-in-Zebrafish-Research.html>
- Kelly, A., and Hurlstone, A. F. (2011). The use of RNAi technologies for gene knockdown in zebrafish. *Briefings in functional genomics* , pp. 189-196.
- Krebs, J., Goldstein, E., and Kilpatrick, S. (2011). *Lewin's Genes X* .10<sup>th</sup> ed.. Jones and Bartlett Publishers.
- Krol, J., Loedige, I., and Filipowicz, W. (2010). The widespread regulation of microRNA biogenesis, function and decay. *Nature Reviews, Genetics* , pp. 597-610.
- Laboratoires, K. (27<sup>th</sup> December 2010). *KEGG*. Obtained on 16<sup>th</sup> September 2013, from Kanehisa Laboratories: <http://www.kegg.jp/pathway/dre04141>
- Lang, M., Lapierre, L., Frotscher, M., Goldenring, J., and Knapik, E. (2006). Secretory COPII coat component Sec23a is essential for craniofacial chondrocyte maturation. *Nature Genetics* , 782006, pp. 1198-1203.
- Lee, Y., Shibata, Y., Malhotra, A., and Dutta, A. (2009). A novel class of small RNAs: tRNA-derived RNA fragments (tRFs). *Genes Dev* , 23, pp. 2639–2649.
- Liu, J., Gough, J., and Rost, B. (2006). Distinguishing Protein-Coding from Non-Coding RNA through Support Vector Machines. *PLoS Genetics* , pp. 529-536.
- Martens-Uzunova, E., Olvedy, M., and Jenster, G. (2013). Beyond microRNA - Novel RNAs derived from small non-coding RNA and their implication in cancer. *Cancer Letters* .
- Matera, A., Terns, T., and Terns, M. (2007). Non-coding RNAs: lessons from the small nuclear and small nucleolar RNAs. *Molecular Cell Biology* , 8, 209-220.
- Maxwell, R. (2012). *Biology MOOC Newsletter Archive*. Obtained on 25<sup>th</sup> July 2013, from <http://biomocnews.blogspot.pt/2012/10/daily-newsletter-october-31-2012-codons.html>

Melville, D., Montero-Balaguer, M., Levic, D., Bradley, K., Smith, J., and Hatzopoulos, A. (2011). The feelgood mutation in zebrafish dysregulates COPII-dependent secretion of select extracellular matrix proteins in skeletal morphogenesis. *Disease Models & Mechanisms* , pp. 763-776.

miRBase. (2013). *miRBase*. Obtained on 16<sup>th</sup> November 2013, from [http://www.mirbase.org/cgi-bin/mirna\\_summary.pl?org=dre](http://www.mirbase.org/cgi-bin/mirna_summary.pl?org=dre)

NCBIa. (4<sup>th</sup> August 2013). *NCBI - National Center for Biotechnology Information*. Obtained on 16<sup>th</sup> November 2013, from U.S. National Library of Medicine: <http://www.ncbi.nlm.nih.gov/homologene/74571>

NCBIb. (4<sup>th</sup> August 2013). *NCBI-National Center for Biotechnology Information*. Obtained on 17<sup>th</sup> November 2013, from U.S. National Library of Medicine: <http://www.ncbi.nlm.nih.gov/gene/442922>

NCBIc. (8<sup>th</sup> August 2013). *NCBI-National Center for Biotechnology Information*. Obtained on 17<sup>th</sup> November 2013, from U.S. National Library of Medicine: <http://www.ncbi.nlm.nih.gov/gene/55742>

Neuhauss, S., Solnica-Krezel, L., Schier, A., Zwartkruis, F., Stemple, D., Malicki, J., *et al.* (1996). Mutations affecting craniofacial development in zebrafish. *Development* 123 , pp. 357-367.

Nilsen, T. (2007). RNA 1997-2007: A Remarkable Decade of Discovery. *Molecular Cell* , pp. 715-720.

Nobel Media AB. (2013). *Nobelprize.org*. obtained on 8<sup>th</sup> October 2013, from [http://www.nobelprize.org/educational/medicine/dna/a/translation/trna\\_processing.html](http://www.nobelprize.org/educational/medicine/dna/a/translation/trna_processing.html)

Pederson, T. (2010). Regulatory RNAs derived from transfer RNA? *RNA* , 16, pp. 1865-1869.

Pierce, B. (2010). *Genetics Essentials: Concepts and Connections*. 1<sup>st</sup> ed.. New York: W. H. Freeman and Company.



- Rosen, J., Sweeney, M., and Mably, J. (2009). Microinjection of Zebrafish Embryos to Analyze Gene Function. *J. Vis. Exp.* , pp. 1-5.
- Saikia, M., Krokowski, D., Guan, B., Ivanov, P., Parisien, M., Hu, G., et al. (2012). Genome-wide identification and quantitative analysis of cleaved tRNA fragments induced by cellular stress. *JBC* , pp. 42708-42725.
- Santoriello, C., and Zon, L. (2012). Hooked! Modeling human disease in zabrafish. *J Clin Invest* , pp. 2337-2343.
- Scherrer, K. (2003). Historical review: The discovery of ‘giant’ RNA and RNA processing: 40 years of enigma. *TRENDS in Biochemical Sciences* , pp. 566-571.
- Schwarz, K., Iolascon, A., Verissimo, F., Trede, N., Horsley, W., Chen, W., et al. (2009). Mutations affecting the secretory COPII coat component SEC23B cause congenital dyserythropoietic anemia typeII. *Nature Genétics* , 41, pp. 936-940.
- Smith, T. (2009). *Finch Talk*. obtained on 5<sup>th</sup> Dezembro 2012, from <http://finchtalk.geospiza.com/2009/05/small-rnas-get-smaller.html>
- Sobala, A., and Hutvagner, G. (2013). Small RNA derived from the 5' end of tRNAs can inhibit protein translation in human cells . *RNA Biology* , pp. 525-535.
- Sobala, A., and Hutvagner, G. (2011). Trnsfer RNA-derived fragments:origins, processing, and functions. *John Wiley & Sons,Ltd* , 2, pp. 853-862.
- Thompson, D., and Parker, R. (2009). Stressing out over tRNA Cleavage. *Cell* , pp. 215-219.
- Tonelli, D., Calegari, F., Fei, J., and Nomura, T. (2006). Single-cell detection of microRNAs in developing vertebrate embryos after acute administration of a dual-fluorescence reporter/sensor plasmid. *BioTechniques* , 41, pp. 727-732.
- Wang, Q., Lee, I., Ren, J., Ajay, S., Lee, Y., an Bao, X. (2012). Identification and Functional Characterization of tRNA-derived RNA Fragments (tRFs) in Respiratory Syncytial Virus Infection. *The American Society of Gene & Cell Therapy* , pp. 1-12.

Wei, C., Salichos, L., Wittgrove, C., Rokas, A., and Patton, J. (2012). Transcriptome-wide analysis of small RNA expression in early zebrafish development. *RNA* , pp. 915–929.

Weigta, S., Hueblera, N., Braunbeckb, T., Landenberga, F., and Broscharda, T. (2010). Zebrafish teratogenicity test with metabolic activation (mDarT): Effects of phase I activation of acetaminophen on zebrafish *Danio rerio* embryos. *Toxicology* , pp. Pages 36–49.

Winter, J., Jung, S., Keller, S., Gregory, R., and Diederichs, S. (2009). Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nature Cell Biology* , pp. 228-234.

Yamasaki, S., Ivanov, P., Hu, G., and Anderson, P. (2009). Angiogenin cleaves tRNA and promotes stress-induced translational repression. *J. Cell Biol* , pp. 35-42.

Yudelevich, A. (1971). Specific cleavage of an *Escherichia coli* leucine transfer RNA following bacteriophage T4 infection. *Journal of Molecular Biology* , pp. 25-29.