

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

**4,800**

Open access books available

**122,000**

International authors and editors

**135M**

Downloads

Our authors are among the

**154**

Countries delivered to

**TOP 1%**

most cited scientists

**12.2%**

Contributors from top 500 universities



**WEB OF SCIENCE™**

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# On the Far Side of Telomeres: The Many Roles of Telomerase in the Acquisition and Retention of Cancer Stemness

---

Kerem Terali and Açelya Yilmazer

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/65762>

---

## Abstract

It is well recognised that upregulation/reactivation of telomerase, the telomere-lengthening enzyme, is the *sine qua non* of cellular immortalisation and malignant transformation. But there is also convincing evidence that telomerase stands at a crossroads where several developmental signalling pathways meet and that its upregulation/reactivation has effects beyond maintaining telomere length, such as altering energy metabolism and modulating gene expression. We believe that it is important to realise that, in a pathological context, such extratelomeric effects of telomerase are related to the emergence and persistence of the cancer stem cell (CSC) phenotype. Given the common conception of cancer stemness as a major contributor to therapy resistance and tumour relapse, a more complete annotation of biological mechanisms for its regulation by telomerase will provide the opportunity to develop telomerase-targeted anticancer therapies which kill or differentiate CSCs effectively.

**Keywords:** telomeres, telomerase, TERT, cancer stemness, CSCs, targeted anticancer therapy

---

## 1. Introduction

Telomeres are specialised structures that define the very ends of linear eukaryotic chromosomes and provide for their stability by protecting against degradation or end-to-end fusion. In mammals, telomeres are localised throughout the nucleus and associated with the nuclear matrix. Telomeric DNA of human cells is composed of a long (5 to 15 kb) stretch of the repeating hexanucleotide sequence 5'-TTAGGG-3' on one strand (the G-rich strand) and the complementary 5'-CCCTAA-3' on the other (the C-rich strand). The G-rich strand has a short (35 to

600 nt) single-stranded overhang at its 3' end (the G-overhang) which folds back and base pairs with the C-rich strand, forming a T-loop [1]. In humans, hexameric telomere repeats act as binding sites for various telomere-binding proteins collectively termed the shelterin complex, a dynamic ensemble of interactions that allows the cell to distinguish between natural chromosome ends and DNA double-strand breaks, preventing the cell's DNA damage response (DDR) from improper activation [2]. Telomeres undergo progressive shortening with each cell division as a result of incomplete lagging strand synthesis, less widely known end-processing events, and oxidative damage [3, 4]. This so-called telomere erosion operates as a kind of mitotic clock that determines ageing of the whole organism and suppresses malignant transformation of its constituent cells. The biological function of telomeres is heavily regulated and relies on both a minimal length of telomeric DNA and the proper functioning of the associated shelterin complex. A unique enzyme termed telomerase assists in replicating linear chromosomes through *de novo* synthesis of telomeric repeats, thereby counteracting the progressive telomere erosion that would otherwise occur in its partial or complete absence. In addition to its role in telomere length homeostasis, telomerase also performs telomere length-independent functions such as modulation of gene expression. In a pathological context, telomerase's new talents are intimately related to tumour development and progression to metastatic disease. This chapter summarises the newly discovered extracurricular activities of telomerase and describe how these are involved in regulating cancer stemness, the stem-like component of human tumours.

## 2. Telomerase and the cancer connection

Telomerase is a conserved RNA-dependent DNA polymerase canonically responsible for the maintenance of telomere length above a critical threshold. Human telomerase is primarily localised in the nucleus, as deducible from its role in telomere biology, but it can also be found in other cellular compartments such as the cytosol and mitochondria [5]. Telomerase is ribonucleoprotein in nature and consists minimally and essentially of a protein catalytic subunit (telomerase reverse transcriptase, TERT) and a large RNA subunit (telomerase RNA, TER). Active human telomerase has a bilobal architecture where one TERT subunit and one TER subunit participate in the formation of each lobe and a hinge region connects the two lobes [6]. This conformationally flexible, dimeric structure of the human enzyme undoubtedly has profound functional implications with respect to the catalytic cycle. Firstly, during the synthesis of telomeric DNA by telomerase, the 3' end of the G-overhang is positioned in the active site of TERT and aligned by base pairing with the 3' end of the RNA template in TER. Secondly, TERT catalyses the addition of deoxyribonucleotides to the chromosome substrate through reverse transcribing TER into hexameric telomere repeats until the 5' end of the RNA template is reached. Lastly, telomerase translocates and realigns with the newly synthesised 3' end of the chromosome substrate to restart the catalytic cycle [7, 8]. In spite of the fact that TERT and TER are the two subunits that provide the catalytic core of telomerase, there are several other molecules that associate with telomerase and are involved in its biogenesis, trafficking, recruitment, and activation. Some of the most well-known telomerase-associated

proteins include the nucleolar protein dyskerin [9], the three other nucleolar proteins NOP10, NHP2 and GAR1 [10], the two AAA+ ATPases pontin and reptin [11], and the WD40-repeat protein TCAB1 [12]. It should be noted that not all cells necessarily rely upon telomerase to maintain telomere length. Some telomerase-negative immortalised cell lines and tumours are able to elongate their telomeres by the much rarer alternative lengthening of telomeres (ALT) pathway. In contrast to telomerase, which utilises an RNA template to *de novo* synthesise telomeric repeats, the ALT pathway utilises a DNA template for DNA copying in an inter- or intramolecular recombination event [13].

Cancer is usually an age-related genetic disease, manifesting only when normal cells develop genomic instability over a reasonable period of time and acquire unlimited replicative potential that leads to the generation of macroscopic tumours. Telomerase upregulation/reactivation is observed in at least 85% of advanced human tumours, strongly suggesting a crucial role during human tumour pathogenesis [14, 15]. The most widely accepted multistep model of general tumourigenesis for explaining the part played by telomerase in telomere maintenance and cellular immortalisation is provided in section 3.2. Besides being found in primary tumours, telomerase activity is also detected in circulating tumour cells in, for instance, breast [16], ovarian [17] and prostate [18] cancers. Telomerase is upregulated/reactivated in premalignant cells by five common mechanisms: (i) increased transcriptional activation of *TERT* and/or *TER*; (ii) loss of transcriptional repressors of *TERT*; (iii) mutations in the *TERT* gene promoter/enhancer region (which result in the transactivation of this gene); (iv) several kinases (which phosphorylate and thus enhance the activity of *TERT*); and (v) gain of copy number of *TERT* and/or *TER* [13]. Somatic mutations in the *TERT* gene promoter region are frequent events in cancers of the bladder, central nervous system, skin (melanoma) and thyroid (follicular cell-derived) [19]. Two mutually exclusive and highly recurrent *TERT* promoter mutations are C250T and C228T [20, 21]. Although both mutations create a similar binding motif for E-twenty-six (ETS) transcription factors, they are functionally distinct in such a way that the the C250T *TERT* promoter but not the C228T *TERT* promoter additionally requires non-canonical NF- $\kappa$ B signalling in order to be transcriptionally driven [22]. Collectively, these findings highlight the contribution of *TERT* promoter mutations and non-canonical NF- $\kappa$ B signalling to tumourigenesis and decipher a fundamental mechanism for the reactivation of *TERT* in various tumours.

### 3. Cancer stemness

Cancer cells within a single tumour often exist in distinct phenotypic states which differ in functional attributes. This so-called intratumoural heterogeneity originates from a myriad of cell types recruited to the tumour as well as from genetic, epigenetic and metabolic differences amongst the cancer cells themselves and may result in variable or unpredictable responses to treatment [23]. Postulated to be the driving force behind tumour maintenance, hypermalignant stem-like cells called cancer stem cells (CSCs) represent a unique dimension of intratumoural heterogeneity. This often-small subpopulation of cancer cells is thought to play pivotal roles

in tumour initiation and progression, spreading, therapy resistance, and recurrence, all of which lead to poor prognosis.


### 3.1. Definitions and measurements

CSCs have critical implications for nearly, if not quite, all types of cancers, including leukaemias [24–26], lymphomas [27], melanomas [28, 29], sarcomas [30], and various carcinomas such as brain [31], skin [32], head and neck [33], lung [34, 35], liver [36], gastric [37], colorectal [38, 39], bladder [40], pancreatic [41, 42], prostate [43], breast [44] and ovarian [45] cancers. The functions assumed by CSCs in human cancers are diverse, ranging from sustaining tumour growth and dissemination to treatment failure and tumour relapse. The three clear-cut features that contribute to the aforementioned functions of CSCs, aka cancer stemness traits, are: (i) their unrestricted ability to self-renew; (ii) their aberrant ability to differentiate into mixed populations of tumour cells; and (iii) their high ability to transition from a proliferative to a quiescent state. In spite of the fact that these operational characteristics are, to a large extent, shared by both CSCs and physiological stem cells, CSCs are the ones that are known to be related to several malignant phenotypes, including induction of invasion and metastasis and resistance to apoptosis. In addition, CSCs are distinguished from bulk tumour cells by their capacity to form nonadherent spheres when cultured in stem cell media, their propensity to found fresh tumours when transplanted into severe combined immunodeficient mice and their expression of a selected repertoire of stem cell-surface markers [46]. Therefore, it is important here to realise that cancer stemness, no matter how it is measured, stresses the ways in which CSCs differ from bulk cancer cells as well as from physiological stem cells. The cancer stemness phenomenon is of considerable clinical importance and significance because it prognosticates that successful anticancer therapy must involve strategies that will eradicate CSCs, as these cells are able to dominate any residual tumour cells that survive conventional anticancer therapies.

### 3.2. Determinants and signatures

In the past, the cancer stemness model was widely seen as a static one which suggested a stable CSC population and a hierarchical organisation of cell division and differentiation. In recent times, however, the cancer stemness model has evolved into a dynamic one where CSCs are rather accepted as a functional subpopulation of cancer cells and can also be formed by the process of dedifferentiation from mature cancer cells under proper environmental conditions [47]. Accumulating data have revealed that cancer stemness is governed by genetic changes (such as oncogene activation and oncosuppressor gene inactivation), epigenetic changes (such as miRNA targeting and promoter DNA hypomethylation/hypermethylation) and metabolic changes (such as the shift to aerobic glycolysis) concomitant with changes in the tumour microenvironment, especially the CSC niche (Figure 1). These changes are a prerequisite for the oncogenic transformation and cellular (nuclear and metabolic) reprogramming of non-CSCs to CSCs and precipitate a spectrum of drastic cellular consequences, including overproduction of certain oncoproteins, disruption of certain oncosuppressor proteins, upregulation/reactivation of telomerase, reactivation of the epithelial-to-mesenchymal transition (EMT)

programme, modulation of energy metabolism, stimulation of a number of embryonic/ oncogenic signalling pathways, and differential expression of several microRNAs (miRNAs).



**Figure 1. How do CSCs arise?** The joint impact of genetic, epigenetic, metabolic and microenvironmental factors is believed to determine the conversion of non-CSCs (which could be normal stem cells, mature cancer cells, or others) to CSCs. This process somehow is a dynamic one and CSCs are a functional and not merely territorial subpopulation of cancer cells.

Disruption of diverse oncosuppressor proteins with antiproliferative, prodifferentiative and/or proapoptotic effects accounts for an early molecular event accompanying the emergence of cancer stemness traits. p53, pRB, PTEN, and p16<sup>INK4A</sup> are by far among the most commonly inactivated oncosuppressor proteins in advanced human tumours [48]. Their inactivation permits premalignant cells to avoid replicative senescence, apoptosis, or both, thereby continuing to divide and accumulating further tumourigenic alterations like genomic (chromosomal) instability that follows telomere erosion [49]. The subsequent upregulation/reactivation of telomerase compensates for telomere erosion (which would otherwise trigger entry of cells into a period of crisis with massive cell death), suppressing genomic (chromosomal) instability and allowing premalignant cells to proliferate for a virtually infinite number of cell divisions. Additionally, and surprisingly, there is accumulating evidence that telomerase upregulation/reactivation provides susceptible cells with cancer stemness traits. The many functions of telomerase in the development and maintenance of cancer stemness will be addressed in more detail in the next sections of this chapter.

A further means by which non-CSCs acquire cancer stemness traits is through the EMT process. Physiologically, EMT causes cells to change from a stationary epithelial to a motile mesenchymal morphology, thereby allowing for wound healing, tissue regeneration and organ fibrosis in adults and cell migration and tissue remodelling in developing embryos [50]. In the context of epithelium-derived carcinoma, however, the reactivation of the EMT programme contributes to the evolution of primary tumours towards increasingly aggressive phenotypes. The complex molecular, cellular and morphological alterations linked to pathological EMT are generally mediated by the joint action of the signals from the tumour microenvironment that induce the EMT programme (for example, TGF- $\beta$  signalling, inflammatory cytokines, and hypoxia), the transcription factors that coordinate the EMT programme (for example, SNAI-1, SNAI-2, ZEB-1, ZEB-2, TWIST-1, and TWIST-2), and the effector proteins that execute the EMT programme (for example, low levels of E-cadherin and high levels of vimentin, N-cadherin, fibronectin, CD44, and MMPs) [51]. Such cooperation between the cell-extrinsic signals and the cell-intrinsic regulators is fully important and primarily responsible for endowing epithelial tumour cells with CSC-like properties, ranging from cell motility to invasiveness to cell survival, which are indispensable to metastatic dissemination from the primary tumour site, secondary tumour growth at a distant site, and resistance to therapy, respectively. Although CSCs originating from bulk tumour cells within epithelium-derived carcinomas achieve their final stemness state possibly *via* EMT, the degree to which they resemble or depart from CSCs originating from adult stem cells has yet to be fully explored.

Metabolic reprogramming is also an obvious mechanism of intervening and redirecting the cell fate of differentiated (normal or non-CSC tumour) cells. Traditionally, energy metabolism was widely accepted as a passive process that generated ATP and building blocks to meet the demands of the specialised cell types of the body in response to extra- and/or intracellular signals. Today, however, the modulation of energy metabolism and build-up of oncogenic metabolites are viewed as the harbingers of cancer stemness [52]. Typically, cancer cells are dependent more on glycolysis for energy production, even in the presence of sufficient oxygen to support oxidative phosphorylation. This phenomenon of aerobic glycolysis is commonly referred to as the Warburg effect and appears to fulfil the requirement of proliferating cancer cells to rapidly yield ATP and to provide anabolic substrates (such as amino acids, nucleotides, and phospholipids) for their daughter cells [53]. Besides, increased lactate generation during aerobic glycolysis provokes the acidification of the tumour microenvironment, ultimately giving rise to motile, invasive/metastatic and drug-resistant cells [54]. In agreement with this, a recent report confirmed and substantiated the need for a metabolic switch to glycolysis in the emergence of EMT-driven CSC-like characteristics in basal-like breast cancer cells [55]. Another report utilising nasopharyngeal carcinoma as a model system established that behaviourally-selected and accordingly-assayed CSCs, as distinct from their differentiated progenies, exhibit a metabolic shift from oxidative phosphorylation to glycolysis for ATP supply [56]. Nevertheless, contradictory evidence on the metabolic profile of CSCs has also been presented; two independent research groups reported that the bioenergetic and biosynthetic demands of quiescent/slow-cycling CSCs are likely to be met by oxidative phosphorylation, not by glycolysis [57, 58].

CSCs maintain their prolonged residence in the stemness state through diverting and co-opting elegant signalling pathways that are normally active during embryonic development. The embryonic/oncogenic signalling pathways operating in CSCs include Notch, Hedgehog (HH), Wnt/ $\beta$ -catenin, cytokine receptor-mediated JAK/STAT, TNF- $\alpha$  receptor-mediated NF- $\kappa$ B, growth factor receptor (receptor tyrosine kinase)-mediated PI3K/AKT/mTOR, TGF- $\beta$ /BMP receptor-mediated SMAD, and Hippo-YAP/TAZ [59–61]. Sustained activation of and crosstalk between these cascades ultimately enhance the expression of cell-surface proteins (for example, CD133, CD44, integrins, and CXCR4), prosurvival proteins (for example, BCL-2, BCL-XL, MCL-1, survivin, and MIC-1), induced pluripotency-associated transcription factors (for example, BMI-1, OCT-3/4, SOX-2, MYC, and NANOG), EMT-associated proteins (for example, MMPs, vimentin, N-cadherin, SNAIL, TWIST, and ZEB), glycolysis-associated proteins (for example, GLUTs and glycolytic enzymes), treatment resistance-associated proteins (for example, GSH, ALDH1, ABCB1, ABCC1, ABCG2, CHK-1, and CHK-2), proangiogenic factors (for example, VEGF and COX-2), and proinflammatory cytokines (for example, IL-6 and TNF- $\alpha$ ) [60, 62, 63].

Lastly, several miRNAs have been observed to support the emergence of cancer stemness traits through targeting signalling elements and gene groups implicated in CSC biology. miRNAs are a fast-growing class of short (19 to 22 nt), noncoding, regulatory RNA molecules that customarily bind to the 3'-untranslated region (3'-UTR) of their target transcripts to induce translational repression, degradation, or destabilisation. Although miRNAs generally help regulate the transitions between different stages of development, they are also linked with tumourigenesis. As such, CSC-specific miRNA expression profiles may be useful for prognostic purposes. Those miRNAs that are highly expressed in CSCs of a specific tumour are termed oncomiRs; those that are excluded from CSCs of the same tumour are known as tumour-suppressor miRs. Breast CSCs were the first CSCs in which differential expression of miRNAs was demonstrated [64].

#### **4. Regulation of cancer stemness by telomerase**

Most adult somatic cells do not or only transiently express telomerase and undergo telomere shortening with every cell division until the cell eventually dies. Most tumour cells, including CSCs, however, display high levels of telomerase activity and possess the ability to continually regenerate their telomeres [65]. As a result, telomerase upregulation/reactivation serves as an important mechanism for CSCs to attain indefinite (or at least extremely long) replicative lifespans. In fact, in reality, telomerase undertakes roles that significantly diverge from its normal role in elongating telomeres, as suggested by contemporary research on manipulation of telomerase expression and/or function in cells representing potential targets for oncogenic transformation and cellular (nuclear and metabolic) reprogramming. Central to the extratelomeric roles of telomerase (particularly of TERT) is its interaction with key downstream components of the main embryonic/oncogenic signalling pathways or with other macromolecules (such as DNA and transcription factors) by which gene expression is regulated. The presence of a few to several hundred copies of TERT, which are not assembled into telomerase,




in human immortalised cell lines reasonably provides a molecular basis for the formidable power of TERT as a transcriptional cofactor in oncogenic transformation and cellular (nuclear and metabolic) reprogramming, irrespectively of its TER-dependent DNA polymerase activity [66]. A very recent systemic review of the literature by us disclosed that most of the non-canonical responsibilities of telomerase identified so far strongly relate to the control of cancer stemness traits [67]. Telomerase/TERT-controlled aspects of the CSC phenotype involve proliferation, survival, therapy resistance, induced pluripotency, motility, glycolytic metabolism, and niche establishment and integrity (Figure 2). Equally strikingly, there seems to be a positive feedback loop between a number of gene products targeting TERT and TERT expression itself, plausibly amplifying the effects of central oncogenes and oncogenic pathways associated with the generation and/or maintenance of cancer stemness traits in a cell-autonomous manner. Although some of the observed cell-intrinsic/microenvironmental changes may require a catalytically active enzyme, there are several examples of oncogenic alterations brought about by catalytically inactive telomerase, as in the case of alternatively spliced (AS) TERT variants. To date, as many as twenty different AS TERT variants have been identified [68]. These variants tend to occur more frequently in cancer cells than in normal cells, indicating that they may be evolutionarily favoured in the context of pathology.

#### 4.1. Stimulation of CSC proliferation

Given their role in the expansion of a tumour cell population, CSCs must display extensive proliferative capacity. Cell proliferation is both a matter of progression through the cell cycle and an issue necessitating cell growth (biosynthesis). There is a wealth of information in the literature on the promotive role of telomerase, independent of its telomere-elongating function, in cell proliferation. In an early study of the association between telomerase and cell proliferation, telomerase was shown to support the proliferation of human mammary epithelial cells through elevated EGFR signalling (even though it is quite ambiguous whether this effect is telomere length-independent or not) [69]. Moreover, TERT confers CSC characteristics to glioma cells by inducing EGFR expression, disconnectedly from its role in telomere biology [70]. Interestingly, telomerase upregulation was found to be closely linked to EGFR expression in actively proliferating normal human epithelial cells [71]. These observations imply the existence of a feed-forward loop that involves telomerase/TERT and EGFR. A plausible mechanism linking the EGFR–telomerase axis to cancer is that aberrant EGFR signalling may render CSCs less dependent on exogenous mitogens/growth factors and reinforce the persistent expression of telomerase in CSCs, thus playing a critical role in tumour development and progression.

Expanding these findings, one research group demonstrated that TERT promotes the proliferation of mammalian tissue progenitor cells *via* transcriptional control of a MYC- and Wnt-related developmental program [72]. To be more precise, TERT physically occupies the promoters of Wnt/ $\beta$ -catenin target genes, including those encoding cyclin D1 and MYC [73]. Cyclin D1 is a cell cycle control protein with oncogenic potential and has both enzymatic and nonenzymatic activities which are of great significance in tumour cells [74]. An additional molecular component involved in cyclin D1 expression in proliferating cells is nucleolar



**Figure 2. Emergence and persistence of cancer stemness by telomerase/TERT (figure adapted from [67] with permission).** Clearly and unmistakably, telomerase/TERT is a powerful regulator of many aspects of the CSC phenotype, including: (a) proliferation, survival, therapy resistance; and (b) induced pluripotency, motility, glycolytic metabolism, niche establishment and integrity. This multifaceted ribonucleoprotein complex exerts its telomere-independent tumour-promoting effects partly by diverting and co-opting developmental signalling pathways and modulating gene expression. A cross symbol denotes an inhibition (blockage). A dashed arrow indicates that a given cancer stemness trait is not a direct consequence of the process shown in the preceding box, but of the inhibition of that process. AG, angiogenesis; DDR, DNA damage response; dsDNA, double-stranded DNA; dsRNA, double-stranded RNA; GSH, glutathione; INF, inflammation; mtDNA, mitochondrial DNA; nDNA, nuclear DNA; rDNA, ribosomal DNA; RMRP, RNA component of mitochondrial RNA-processing endoribonuclease; ROS, reactive oxygen species; siRNAs, small interfering RNAs

antigen p120 (NOL1), as suggested by the results of a very recent study [75]. In this study, telomerase was found to interact, in a TER-dependent fashion, with NOL1 and activate the transcription of the gene coding for cyclin D1. The relationship between TERT and cyclin D1 expression was corroborated also by other scientists [76–79]. Further support for TERT involvement in the stimulation of Wnt signalling-mediated cell proliferation was evidenced by an independent approach employing the  $\Delta 4-13$  AS variant of human TERT that is devoid of reverse transcriptase activity [80]. In this approach, ectopic expression and small interfering RNA (siRNA)-mediated knockdown of the  $\Delta 4-13$  AS variant in ALT cells, transformed telomerase-positive cell lines and telomerase-negative normal cells unquestionably proved that the proliferative effect of TERT is not coupled to telomerase activity. Because  $\beta$ -catenin is known to modulate TERT expression in stem cells and tumour cells [81], it is tempting to speculate that telomerase and Wnt/ $\beta$ -catenin signalling may act together in a positive feedback circuit to actively encourage the proliferation of CSCs.

Intriguingly, one group failed to find evidence that TERT promotes Wnt signalling in human breast cancer cells, indicating that TERT's effect on Wnt signalling is possibly context- and cell type-dependent [82]. Their findings are in the same direction as those from a former study on telomerase-null mouse models, where TERT loss-of-function in a physiological setting was reported to have no evident effects on Wnt signalling [83]. An alternative mechanism of action of telomerase on cell proliferation, as deduced by reverse genetics in human mammary epithelial cells, is that TERT-induced cell proliferation may result primarily from decreased levels of the RNA component of mitochondrial RNA-processing endoribonuclease (RMRP), not from increased Wnt signalling [78]. TERT and RMRP form a definite ribonucleoprotein complex that exhibits RNA-dependent RNA polymerase activity and, using RMRP as a template, produces double-stranded RNAs that can be later processed into siRNAs in a Dicer-dependent fashion [84]. siRNAs regulate gene expression at the posttranscriptional level as well as at the level of chromatin structure; therefore, it is reasonable to question whether their mutations or altered expression correlate with human cancers.

Aside from activating Wnt signalling and regulating gene expression (in the presence of RMRP), telomerase also stimulates ribosomal biogenesis through increased Pol I-directed ribosomal DNA transcription, exerting a positive influence on cell cycle and proliferation dynamics [85]. This may ultimately improve the protein synthesis capacity of CSCs for unrestrained growth. The molecular mechanism behind telomerase-induced ribosomal biogenesis was investigated in a very recent report where a MYC-driven oncogenesis model was proposed [79].

#### **4.2. Promotion of CSC survival**

Apoptosis is a form of cell death induced by miscellaneous stimuli and mediated by a subset of cysteine proteases termed caspases. A cancer cell's ability to evade death signals, thus preventing self-destruction by the activation of an apoptotic programme, is regarded as one of the hallmarks of cancer [86]. Several pieces of information suggest that telomerase exerts antiapoptotic effects in cancer cells through telomere-independent mechanisms. In the case of CSCs, telomerase-mediated inhibition of apoptosis may contribute to the enhanced and

continued survival of these cells in tumours. In keeping with its cytoprotective function, TERT was revealed to inhibit cell death by blocking the death receptor-initiated (or extrinsic) apoptotic pathway in acute promyelocytic cells [87]. Similarly, yet in a mechanistically different way, TERT was also shown to block the mitochondrion-initiated (or intrinsic) apoptotic pathway in colon and cervical carcinoma cell lines [88]. A more thorough probing of the molecular mechanism behind telomerase-mediated suppression of the intrinsic apoptotic pathway found that TERT overexpression upregulates the expression of the antiapoptotic mitochondrial protein BCL-2, downregulates the expression of some proapoptotic mitochondrial proteins (for example, BAX) and reduces the activation of some caspases (for example, caspase-9) in ovarian surface epithelial cells [89]. Given the capacity of BCL-2 to increase telomerase activity in human colorectal and cervical carcinoma cell lines [90], it is conceivable that telomerase and BCL-2 are engaged in a positive feedback loop that impedes apoptosis. In this connection, it is well to add that the introduction of a constitutively expressed TERT construct into colon carcinoma and Burkitt's lymphoma cell lines was, independently of telomerase activity, associated with the reversion of a transcriptional programme coordinated by p53, a potent and common activator of both the intrinsic and extrinsic apoptotic pathways [91].

Apart from suppressing mitochondrion-initiated cell death, overexpression of TERT was also found to suppress, in a telomere-independent manner, endoplasmic reticulum (ER) stress-induced cell death in murine primary neural cells and human cancer cell lines [92]. ER stress arises as a result of perturbations in ER function and elicits the unfolded protein response (UPR), a conserved signal transduction pathway for dealing with misfolded proteins. When the UPR-induced mechanisms fail to alleviate ER stress, both the intrinsic and extrinsic apoptotic pathways may become activated [93]. Reciprocally, specific activation of ER stress was demonstrated to upregulate TERT expression in a breast cancer cell line [94]. It therefore seems reasonable to suggest that TERT and ER stress are involved in a dynamic interplay supporting CSC survival in abnormal metabolic conditions such as glucose starvation.

### **4.3. Induction of pluripotency**

Restoration of the molecular circuitry that forms the necessary base of pluripotency in embryonic stem cells (ESCs) strongly correlates with the gaining and retention of cancer stemness. In ESCs, this circuitry is made up of special transcription factors and function as a repressor of differentiation. Takahashi and Yamanaka were the first to demonstrate that a cocktail of four transcription factors (namely OCT-3/4, SOX-2, MYC, and KLF4) are necessary and sufficient for nuclear reprogramming into an ESC-like state [95]. In CSCs, the so-called Yamanaka factors, besides driving the induction of pluripotency, are additionally involved in inhibiting apoptosis [96]. A valued piece of work documented that TERT forms a ternary complex with the nucleolar GTP-binding protein NS/GNL3L and the chromatin remodelling factor BRG1 and that the resulting NS/GNL3L TERT BRG1 complex is required for NS/GNL3L-induced upregulation of the nuclear reprogramming factors OCT-3/4, MYC, and KLF-4 [97]. The likely part played by TERT/telomerase in contributing to the pluripotent character of CSCs is also congruous with the later finding that siRNA-mediat-

ed hTERT depletion in gastric CSCs downregulates the induced pluripotency-associated transcription factor OCT-4 [98].

#### 4.4. Increase of CSC motility and invasiveness

Migrating CSCs and EMT-phenotypic cells have the ability to disseminate from their primary site and are thus present in the invasive front of tumours. The initial evidence for telomerase/TERT participation in cell migration came from experiments measuring the *in vitro* migration rate of telomerase-positive progenitor cells and cancer cell lines [99, 100]. Later experiments aiming at uncovering the molecular mechanism behind this positive trend showed that telomerase reconstitution boosts cell migration through the activation of Rho family members and the SDF-1–CXCR4 axis [101]. With respect to CSC motility, expression of the chemokine receptor CXCR4 may enable CSCs to migrate along a gradient of the ligand SDF-1 and thus help facilitate their spread. Therapeutic strategies intended to interfere with the SDF-1–CXCR4 axis can possibly have useful clinical relevance and application in the prevention of metastatic disease.

Differentiated epithelial cells that have undergone EMT may as well exhibit augmented motility and invasiveness leading to metastasis. The ternary complex containing TERT, BRG1, and NS or GNL3L (see section 4.3) acts in a telomere-independent mode to activate the EMT programme *via* NS/GNL3L-induced upregulation of vimentin, SNAI, and TWIST, three of the mesenchymal cell markers, in genetically defined cancer cells [97]. TERT additionally stimulates EMT in gastric cancer cells through directly regulating the expression of Wnt/ $\beta$ -catenin target genes like those coding for vimentin and SNAI-1 [98]. Equally important is the fact that TERT, in a telomere-independent manner, regulates the expression of several MMP family members, such as MMP-9, *via* the NF- $\kappa$ B pathway [102]. MMPs are the key mediators promoting extracellular matrix (ECM) degradation and remodelling, both of which pave the way for EMT and subsequent metastasis. The indirect involvement of TERT in dissemination was also highlighted by a separate set of data which documented that changes in the motility and invasiveness of malignant cells are likely to result from the TERT-induced upregulation of the metastasis-implicated proteins RhoC and MMP-9 [103]. Interestingly, MMP-9 silencing was shown to downregulate TERT expression *via* ITGB1-mediated FAK signalling in glioma xenograft cells [104]. It is worth mentioning here that a very recent report found that ITGB1 itself is regulated by TERT and that TERT may promote the invasion and metastasis of gastric cancer cells by enhancing ITGB1 protein levels [105]. Collectively, these findings reinforce the notion that there is an indirect, metastasis-favouring interaction between TERT, MMP-9 and ITGB1 in cancer cells. Another study discovered that TERT upregulates the levels of MAC2BP, a metastasis-related secreted ECM glycoprotein, in gastric cancer cells [106]. MAC2BP is believed to support metastasis through interacting with galectins and altering cell–cell and cell–matrix adhesion properties [107].

Another contribution to knowledge came from a very recent report in which TERT was found to stimulate the expression of oncomiRs, including miR-21, in human leukaemia and HeLa cell lines [108]. Extant research identifies miR-21 as being among the most frequently upregulated miRNAs in epithelial cell-derived solid tumours [109] and also as having a

decisive role in the conservation of CSC phenotype *via* the AKT and ERK1/2 signalling pathways targeting PTEN [110]. The centrality of miR-21 to cancer stemness was confirmed in a recent study on the antisense oligonucleotide-mediated inhibition of miR-21 in two different anaplastic thyroid carcinoma (ATC) cell lines, where the knockdown of miR-21 disturbed the stemness state of ATC cells, as assessed by a decreased expression of the genes encoding OCT-4 and ABCG2 [111].

#### 4.5. Modulation of energy metabolism

Apparently, genetic, epigenetic and microenvironmental changes that regulate the transition to a CSC-like state cannot occur without the presence of a favourable metabotype. In general, stimulation of aerobic glycolysis promotes metabolic reprogramming, whereas inhibition of glycolytic enzymes impairs metabolic reprogramming. In harmony with the concept that metabolism is involved in the control of cancer stemness, a microarray-based gene expression profiling study elucidated that ribozyme-mediated targeting of telomerase in murine melanoma cells downregulates the expression of more than a few glycolytic pathway genes such as those coding for phosphofructokinase and aldolase C [112]. Additionally, a very recent report showed that siRNA-mediated knockdown of TERT in human lymphoma cells lowers the expression of MYC-regulated target genes such as those coding for the glycolytic enzymes lactate dehydrogenase, hexokinase 2, and pyruvate kinase M2 isoform [79]. Due to the fact that MYC is a well-established oncogenic transcription factor activating TERT expression [113], a feed-forward mechanism for the rewiring of glucose metabolism in CSCs is likely to prevail between TERT and MYC.

#### 4.6. Contribution to therapy resistance

CSCs are notorious for their resistance to existing cancer treatment regimens, including radiotherapy and chemotherapy. Both cell-intrinsic and microenvironmental factors appear to contribute to the emergence of therapy resistance in CSCs. Radiotherapy works by directing ionising radiation toward tumours to induce the generation of reactive oxygen species (ROS) which react with and cause damaging of DNA. By the same token, a number of chemotherapeutic agents such as platinum-based antitumour drugs are known to bind to and cause crosslinking of DNA. In this regard, implementation by CSCs of fast and efficient DNA repair mechanisms as well as potent antioxidant/scavenger systems may prove vital to circumvent the deleterious effects of irradiation and several classes of antitumour compounds.

Growing evidence points to a role for telomerase in modulating DDR and contributing to DNA repair. In a prior report, TERT was proposed to, independently of its effect on telomere length, set in motion a transcriptional programme leading to enlarged ribonucleotide (NTP) pools, enhanced DNA repair, and increased chromosomal stability [114]. A circumstantial investigation into the enhanced DNA repair capability of telomerase-expressing cells suggested that TERT/telomerase increases DNA end-joining repair and accelerates nucleotide excision repair through recruiting proteinaceous factors to sites where DNA damage is occurring [115]. The aforementioned findings are consistent with a newer report which showed that TERT expression affords a means of protecting human transformed cells against double-stranded DNA-

damaging drugs and increases their endurance to chromosomal instability [116]. Specifically expressing TERT mutants lacking catalytic activity in ALT cells, the authors of the same report reached the conclusion that the observed cytoprotective effect of telomerase is distinct from its function in telomere biosynthesis. In a separate study providing evidence for an epigenetic component to telomerase-induced treatment resistance, stable short hairpin RNA (shRNA)-mediated suppression of TERT expression was demonstrated to, in a telomere length-independent way, diminish the response of human fibroblasts to DNA double strand breaks, most likely through a mechanism altering the overall state (that is to say, configuration) of chromatin [117].

Telomerase upregulation/reactivation also seems to be involved in counteracting oxidative stress-induced intracellular injury that often follows therapy, as evident from several experimental studies. The initial study examining the extratelomeric function of telomerase under oxidative stress found that mitochondrially-localised TERT decreases cellular peroxide levels and mitochondrial superoxide production, increases mitochondrial membrane potential and protects mitochondrial DNA from oxidative damage in human lung fibroblasts [118]. The observation that telomerase provides resistance to oxidative stress was validated and extended in an ensuing study where TERT was shown to bind to mitochondrial DNA and accordingly protect it and its function against damage [119]. An alternative explanation for telomerase-induced resistance to oxidative stress came from a more recent study in which TERT overexpression in cancer cells was demonstrated to alleviate basal ROS levels and intracellular ROS production through potentiating the effects of endogenous antioxidants or free radical scavengers such that the proportion of reduced to oxidised glutathione (GSH/GSSG) is increased and peroxiredoxin is replenished in the interior of the cell [120]. Apart from serving to keep mitochondrial DNA damage-free, mitochondrially-localised telomerase also guards nuclear DNA against oxidative attack through decreasing mitochondrial ROS production [121]. Finally, it is appropriate to mention that the  $\beta$ -deletion variant, a catalytically defective AS variant of TERT, localises to both mitochondria and the nucleus and, distinct from the canonical role of TERT in telomere extension, protects three basal breast cancer cell lines from cisplatin-induced apoptosis, endowing breast tumours with chemotherapy resistance [122].

#### **4.7. Establishment and integrity of the CSC niche**

The tumour microenvironment (TME) is an umbrella term that encompasses all cellular and non-cellular components surrounding a tumour. These components include tumour-adjacent stromal cells (for example, endothelial cells and fibroblasts), diverse effectors of the immune system (for example, lymphocytes and mesenchymal stem cells), ECM elements, proteases, and networks of cytokines, growth factors and other soluble factors. Specifically, both the immediate TME (cell–cell and cell–matrix connections) and the extended TME (for example, vascular bed) are thought to be implicated in tumour progression. The TME is also capable of creating a niche for CSCs, in which they remain in an undifferentiated state until stimulated to differentiate into non-CSC tumour cells and form tumour bulk. Modulation of gene expression and metabolism by telomerase in CSCs may recondition the CSC niche in favour of the hypermalignant (that is to say, highly metastatic, therapy-resistant) nature of these cells.

It is in this regard that a recent study demonstrated that telomerase binds to p65 and localises to promoters of NF- $\kappa$ B target genes, such as those encoding IL-6, TNF- $\alpha$ , and IL-8, proinflammatory cytokines that are the critical triggers of inflammatory responses [123]. Inflammation is considered an enabling characteristic of cancer for the reason that it supplies bioactive molecules (for example, growth factors and EMT-inducing ligands) to the TME and primes cells to release ROS and other chemicals that drive the mutagenesis, and hence genetic evolution, of nearby tumour cells toward hypermalignancy [86]. Since NF- $\kappa$ B is known to transcriptionally upregulate telomerase levels [124], this finding implies that a positive feedback loop between telomerase and NF- $\kappa$ B may explain the grounds for the coexistence of chronic inflammation and sustained telomerase activity in neoplastic lesions.

Furthermore, it was reported that TERT activates the transcription of VEGF, an endothelial mitogen and master orchestrator of angiogenesis, independently of telomerase activity in HeLa cells [125]. Further dissection of the underlying regulatory mechanism led to the conclusion that TERT upregulates VEGF expression through its interaction with the specificity protein 1 (Sp1) transcription factor [126]. Angiogenesis is the process of growth or formation of fresh blood vessels from the pre-existing vasculature, and its induction is widely considered an essential attribute of tumour growth as well as metastasis as solid tumours larger than 1 cm<sup>3</sup> have to develop their own blood supply to circumvent necrotic cell death. Given the prior discovery that VEGF stimulates the production of TERT [127], it may be that there is a positive feedback circuit between TERT and VEGF. This regulation may account for the combined and continuing contribution of these two proteins to the maintenance of CSCs in solid tumours. Moreover, siRNA-mediated knockdown of TERT was shown to downregulate the prostaglandin-synthesising enzyme COX-2 in pancreatic cancer cells [128]. COX-2, like VEGF, is a proangiogenic factor that has the potential to establish a selective niche favourable to the preservation of CSCs. Subsequent studies showed that COX-2 stimulates the expression of TERT in cervical cancer cells [129]. Collectively, these data indicate that a feed-forward regulation, which could be important in carcinoma growth and progression, occurs between TERT and COX-2.


## 5. Prospects for telomerase-targeted anticancer therapies

There is little doubt that telomerase is not only an effector in human tumour pathogenesis but also a regulator of essentially all aspects of malignant behaviour, including spreading to secondary sites. This being the case, specifically and sensitively measuring telomerase activity in clinical samples represents both an early diagnostic marker and a negative prognostic factor for patients with malignant disease. The obvious capacity of telomerase to directly regulate an ever-expanding number of tumour-promoting genes and pathways renders this gifted ribonucleoprotein particle an attractive and almost universal target for human cancers. Given the integral role played by cancer stemness in disease recurrence, selectively targeting and eradicating CSCs with low toxicity to somatic cells and with minimal side effects holds the promise of a full cure for cancer. Besides being pursued with vigour in the hope of improving patient outcomes, successfully suppressing cancer stemness may also provide the ultimate



evidence for the CSC concept. As summarised in this chapter, telomerase contributes to carcinogenesis most likely through the emergence and persistence of conspicuous CSC qualities. Accordingly, telomerase inhibition in CSCs is predicted to: (i) shrink telomeres; (ii) restrain anchorage-independent growth and inhibit proliferation by cell cycle arrest; (iii) induce CSC death; (iv) induce CSC differentiation; (v) inhibit CSC migration and reverse the EMT programme; (vi) deteriorate glucose metabolism; (vii) enhance radio- and chemosensitivity; and (viii) disrupt the CSC niche. Serious telomere shrinkage is assumed to be a long-term effect of telomerase inhibition in CSCs so the tumour mass will continue to expand for a time after treatment until its constituent cells enter crisis and begin to die in large numbers. The rest of the aforementioned effects, however, are likely to occur after short-term exposure of CSCs to telomerase inhibitors, inducing relatively rapid initial responses to treatment. Natural telomerase inhibitors (phytochemicals) and small-molecule telomerase inhibitors, antisense oligonucleotides and chemically modified nucleic acids, immunotherapeutic agents, and telomerase-directed gene therapy are promising treatment options and may play a larger role in the near future [130]. Imetelstat (GRN163L), which was designed by Geron Corporation in 2003, is the first telomerase inhibitor to advance to clinical development. It is a lipid-conjugated 13-mer (5'-TAGGGTTAGACAA-3') antisense oligonucleotide that is complementary to and binds with high affinity to TER, thereby directly inhibiting telomerase activity and interfering with telomere length homeostasis. It is perhaps safe here to assume that Imetelstat impairs the regulatory role of telomerase in CSC biology not only through telomere shortening but also through negatively influencing its telomere length-independent tumour-promoting functions. In support of this, short-term (72-hour) Imetelstat exposure was shown to promote the differentiation and inhibit the colony-forming ability of multiple myeloma CSCs through a telomere length-independent mechanism [131]. Similarly, *in vitro* Imetelstat treatment was found to deplete breast and pancreatic CSCs, as measured by the reduced proportion of ALDH-positive and CSC-surface marker-expressing cells, through a mechanism of action independent of telomere shortening [132]. Although Imetelstat is known to form thermodynamically stable and sequence-specific duplexes with TER, the possibility that even less thermodynamically stable tetraplexes of Imetelstat may bind to and interfere with some other, yet to be identified, proteins (particularly those that interact with telomerase) should not be excluded [133]. There also exists a possibility that telomerase inhibitors like Imetelstat may be coupled with conventional therapies such as surgical (debulking) therapy, radiotherapy, and chemotherapy, all of which have their own weaknesses and inadequacies. Such combination therapy is predicted to result in rapid and durable clinical responses in broad tumour types (Figure 3). As shown by the sources provided earlier in this chapter and elsewhere in the literature, the principal signalling pathways governing CSC biology operate in physiological stem cells as well. This complicates telomerase inhibition therapy because of the risk of telomerase inhibitors exerting an adverse influence on the size of the physiological stem cell pool and/or on the integrity of the physiological stem cell niche. The notion that physiological stem cells only transiently express telomerase and have relatively long telomeres [65], however, means that there is likely to be a narrow but safe therapeutic window where only CSCs will be depleted by telomerase inhibitors and normal stem cells will remain unaffected. Furthermore, rational approaches that disrupt the interactions of telomerase with important downstream components of embryonic/oncogenic signalling pathways (Wnt/ $\beta$ -catenin and NF- $\kappa$ B being the most prominent of all so far) may be conceived and executed as therapeutic tactics to

specifically target and eliminate CSCs. As far as targeting protein–protein interactions is concerned, a future challenge is to add to existing pathways or identify new ones where telomerase, independently of its role in preventing telomere loss, intervenes and contributes to CSC phenotype. The essential nature of telomerase in the promotion and maintenance of cancer stemness provides a logical basis for believing that CSCs will not develop resistance to any of the aforementioned telomerase-focused therapeutics, in contrast to other targeted anticancer therapies whose targets are likely to be compensated for by functionally equivalent gene products and signalling pathways.



**Figure 3. Combination therapy for predicting rapid and durable clinical responses in broad tumours.** Here, telomerase-focused therapy is proposed as a strategy to effectively target CSCs. Combining telomerase-focused therapy with conventional therapy may provide scope for the elimination of bulk tumour cells while preventing recurrence by simultaneously eradicating the stem-like component of the tumour. Combination therapy may also prevent non-CSC tumour cells from acquiring cancer stemness traits *via*, for example, EMT. A bar-headed solid line denotes a strong inhibitory (negative) effect. A bar-headed dashed line denotes a weak inhibitory (negative) effect.

## 6. Abbreviations

ABC, ATP-binding cassette; Akt, protein kinase B; ALDH1, aldehyde dehydrogenase 1; BAX, BCL-2-associated protein X; BCL, B-cell lymphoma family protein; BMI-1, B lymphoma Mo-MLV insertion region 1 homolog; BMP, bone morphogenetic protein; BRG1, Brahma-related gene 1; CD, cluster of differentiation; CHK, checkpoint kinase; COX-2, cyclooxygenase-2;

CXCR4, C–X–C chemokine receptor type 4; EGFR, epidermal growth factor receptor; ERK1/2, extracellular signal-related kinase 1/2; FAK, focal adhesion kinase; GLUTs, glucose transporters; GNL3L, guanine nucleotide-binding protein-like 3-like; IL, interleukin; ITGB1, integrin beta-1; JAK, Janus kinase; KLF-4, Kruppel-like factor-4; MAC2BP, Mac-2-binding protein; MCL-1, myeloid cell leukaemia-1; MIC-1, macrophage inhibitory cytokine-1; MMPs, matrix metalloproteinases; MYC, v-myc avian myelocytomatosis viral oncoprotein homolog; mTOR, mammalian target of rapamycin; NANOG, Nanog homeobox transcription factor; NF- $\kappa$ B, nuclear factor- $\kappa$ B; NS, nucleostemin; OCT-3/4, octamer-binding transcription factor-3/4; PI3K, phosphoinositide 3-kinase; PTEN, phosphatase and tensin homolog; SDF-1, stromal cell-derived factor-1; SMAD, small mother against decapentaplegic homolog; SNAI, snail family zinc-finger transcription factor; SOX-2, SRY (sex determining region Y)-box 2; STAT, signal transducer and activator of transcription; TAZ, Tafazzin; TGF- $\beta$ , transforming growth factor- $\beta$ ; TNF- $\alpha$ , tumour necrosis factor- $\alpha$ ; TWIST, twist family bHLH (basic helix-loop-helix) transcription factor; VEGF, vascular endothelial growth factor; Wnt, Wingless ligand; YAP, Yes-associated protein; ZEB, zinc-finger E-box-binding homeobox family protein

## Author details

Kerem Terali<sup>1\*</sup> and Açelya Yilmazer<sup>2</sup>

\*Address all correspondence to: kerem.terali@neu.edu.tr

1 Department of Medical Biochemistry, Faculty of Medicine, Near East University, Nicosia, North Cyprus, via Mersin, Turkey

2 Department of Biomedical Engineering, Faculty of Engineering, Ankara University, Tandogan, Ankara, Turkey

## References

- [1] Samassekou O, Gadji M, Drouin R, Yan J. Sizing the ends: normal length of human telomeres. *Ann Anat.* 2010;192:284–91.
- [2] Palm W, de Lange T. How shelterin protects mammalian telomeres. *Annu Rev Genet.* 2008;42:301–34.
- [3] de Lange T. Mammalian telomeres. In: de Lange T, Lundblad V, Blackburn E, editors. *Telomeres*. 2nd ed. NY: Cold Spring Harbor Laboratory Press; 2006.
- [4] von Zglinicki T. Oxidative stress shortens telomeres. *Trends Biochem Sci.* 2002;27:339–44.

- [5] Saretzki G. Extra-telomeric functions of human telomerase: cancer, mitochondria and oxidative stress. *Curr Pharm Des.* 2014;20:6386–403.
- [6] Sauerwald A, Sandin S, Cristofari G, Scheres SHW, Lingner J, Rhodes D. Structure of active dimeric human telomerase. *Nat Struct Mol Biol.* 2013;20:454–60.
- [7] Blackburn EH, Collins K. Telomerase: an RNP enzyme synthesizes DNA. *Cold Spring Harb Perspect Biol.* 2011;3:a003558.
- [8] Pfeiffer V, Lingner J. Replication of telomeres and the regulation of telomerase. *Cold Spring Harb Perspect Biol.* 2013;5:a010405.
- [9] Mitchell JR, Wood E, Collins K. A telomerase component is defective in the human disease dyskeratosis congenita. *Nature.* 1999;402:551–5.
- [10] Fu D, Collins K. Purification of human telomerase complexes identifies factors involved in telomerase biogenesis and telomere length regulation. *Mol Cell.* 2007;28:773–85.
- [11] Venteicher AS, Meng Z, Mason PJ, Veenstra TD, Artandi SE. Identification of ATPases pontin and reptin as telomerase components essential for holoenzyme assembly. *Cell.* 2008;132:945–57.
- [12] Venteicher AS, Abreu EB, Meng Z, McCann KE, Terns RM, Veenstra TD, Terns MP, Artandi SE. A human telomerase holoenzyme protein required for Cajal body localization and telomere synthesis. *Science.* 2009;323:644–648.
- [13] Reddel RR. Telomere maintenance mechanisms in human cancer: clinical implications. *Curr Pharm Des.* 2014;20:6361–74.
- [14] Kim NW, Piatyszek MA, Prowse KR, Harley CB, West MD, Ho PL, Coviello GM, Wright WE, Weinrich SL, Shay JW. Specific association of human telomerase activity with immortal cells and cancer. *Science.* 1994;266:2011–5.
- [15] Shay JW, Bacchetti S. A survey of telomerase activity in human cancer. *Eur J Cancer.* 1997;33:787–91.
- [16] Soria JC, Gauthier LR, Raymond E, Granotier C, Morat L, Armand JP, Boussin FD, Sabatier L. Molecular detection of telomerase-positive circulating epithelial cells in metastatic breast cancer patients. *Clin Cancer Res.* 1999;5:971–5.
- [17] Sapi E, Okpokwasili NI, Rutherford T. Detection of telomerase-positive circulating epithelial cells in ovarian cancer patients. *Cancer Detect Prev.* 2002;26:158–67.
- [18] Goldkorn A, Ely B, Tangen CM, Tai YC, Xu T, Li H, Twardowski P, Veldhuizen PJ, Agarwal N, Carducci MA, et al. Circulating tumor cell telomerase activity as a prognostic marker for overall survival in SWOG 0421: A phase III metastatic castration resistant prostate cancer trial. *Int J Cancer.* 2015;136:1856–62.

- [19] Vinagre J, Almeida A, Pópulo H, Batista R, Lyra J, Pinto V, Coelho R, Celestino R, Prazeres H, Lima L, et al. Frequency of TERT promoter mutations in human cancers. *Nat Commun.* 2013;4:2185.
- [20] Huang FW, Hodis E, Xu MJ, Kryukov GV, Chin L, Garraway LA. Highly recurrent TERT promoter mutations in human melanoma. *Science.* 2013;339:957–9.
- [21] Horn S, Figl A, Rachakonda PS, Fischer C, Sucker A, Gast A, Kadel S, Moll I, Nagore E, Hemminki K, et al. TERT promoter mutations in familial and sporadic melanoma. *Science.* 2013;339:959–61.
- [22] Li Y, Zhou QL, Sun W, Chandrasekharan P, Cheng HS, Ying Z, Lakshmanan M, Raju A, Tenen DG, Cheng SY, et al. Non-canonical NF-kappaB signalling and ETS1/2 cooperatively drive C250T mutant TERT promoter activation. *Nat Cell Biol.* 2015;17:1327–38.
- [23] Kreso A, Dick JE. Evolution of the cancer stem cell model. *Cell Stem Cell.* 2014;14:275–91.
- [24] Lapidot T, Sirard C, Vormoor J, Murdoch B, Hoang T, Caceres-Cortes J, Minden M, Paterson B, Caligiuri MA, Dick JE. A cell initiating human acute leukaemia after transplantation into SCID mice. *Nature.* 1994;367:645–8.
- [25] Bonnet D, Dick JE. Human acute myeloid leukemia is organized as a hierarchy that originates from a primitive hematopoietic cell. *Nat Med.* 1997;3:730–7.
- [26] Graham SM, Jørgensen HG, Allan E, Pearson C, Alcorn MJ, Richmond L, Holyoake TL. Primitive, quiescent, Philadelphia-positive stem cells from patients with chronic myeloid leukemia are insensitive to STI571 in vitro. *Blood.* 2002;99:319–25.
- [27] Chen Z, Ayala P, Wang M, Fayad L, Katz RL, Romaguera J, Caraway N, Neelapu SS, Kwak LW, Simmons PJ, McCarty N. Prospective isolation of clonogenic mantle cell lymphoma-initiating cells. *Stem Cell Res.* 2010;5:212–25.
- [28] Fang D, Nguyen TK, Leishear K, Finko R, Kulp AN, Hotz S, Van Belle PA, Xu X, Elder DE, Herlyn M. A tumorigenic subpopulation with stem cell properties in melanomas. *Cancer Res.* 2005;65:9328–37.
- [29] Schatton T, Murphy GF, Frank NY, Yamaura K, Waaga-Gasser AM, Gasser M, Zhan Q, Jordan S, Duncan LM, Weishaupt C, Fuhlbrigge RC, Kupper TS, Sayegh MH, Frank MH. Identification of cells initiating human melanomas. *Nature.* 2008;451:345–9.
- [30] Suvà ML, Riggi N, Stehle JC, Baumer K, Tercier S, Joseph JM, Suvà D, Clément V, Provero P, Cironi L, Osterheld MC, Guillou L, Stamenkovic I. Identification of cancer stem cells in Ewing's sarcoma. *Cancer Res.* 2009;69:1776–81.
- [31] Singh SK, Hawkins C, Clarke ID, Squire JA, Bayani J, Hide T, Henkelman RM, Cusimano MD, Dirks PB. Identification of human brain tumour initiating cells. *Nature.* 2004;432:396–401.

- [32] Malanchi I, Peinado H, Kassen D, Hussenet T, Metzger D, Chambon P, Huber M, Hohl D, Cano A, Birchmeier W, Huelsken J. Cutaneous cancer stem cell maintenance is dependent on beta-catenin signalling. *Nature*. 2008;452:650–3.
- [33] Prince ME, Sivanandan R, Kaczorowski A, Wolf GT, Kaplan MJ, Dalerba P, Weissman IL, Clarke MF, Ailles LE. Identification of a subpopulation of cells with cancer stem cell properties in head and neck squamous cell carcinoma. *Proc Natl Acad Sci USA*. 2007;104:973–8.
- [34] Kim CF, Jackson EL, Woolfenden AE, Lawrence S, Babar I, Vogel S, Crowley D, Bronson RT, Jacks T. Identification of bronchioalveolar stem cells in normal lung and lung cancer. *Cell*. 2005;121:823–35.
- [35] Eramo A, Lotti F, Sette G, Piloizzi E, Biffoni M, Di Virgilio A, Conticello C, Ruco L, Peschle C, De Maria R. Identification and expansion of the tumorigenic lung cancer stem cell population. *Cell Death Differ*. 2008;15:504–14.
- [36] Ma S, Chan KW, Hu L, Lee TK, Wo JY, Ng IO, Zheng BJ, Guan XY. Identification and characterization of tumorigenic liver cancer stem/progenitor cells. *Gastroenterology*. 2007;132:2542–56.
- [37] Takaishi S, Okumura T, Tu S, Wang SS, Shibata W, Vigneshwaran R, Gordon SA, Shimada Y, Wang TC. Identification of gastric cancer stem cells using the cell surface marker CD44. *Stem Cells*. 2009;27:1006–20.
- [38] O'Brien CA, Pollett A, Gallinger S, Dick JE. A human colon cancer cell capable of initiating tumour growth in immunodeficient mice. *Nature*. 2007;445:106–10.
- [39] Ricci-Vitiani L, Lombardi DG, Piloizzi E, Biffoni M, Todaro M, Peschle C, De Maria R. Identification and expansion of human colon-cancer-initiating cells. *Nature*. 2007;445:111–5.
- [40] Chan KS, Espinosa I, Chao M, Wong D, Ailles L, Diehn M, Gill H, Presti J Jr, Chang HY, van de Rijn M, Shortliffe L, Weissman IL. Identification, molecular characterization, clinical prognosis, and therapeutic targeting of human bladder tumor-initiating cells. *Proc Natl Acad Sci USA* 2009;106:14016–21.
- [41] Li C, Heidt DG, Dalerba P, Burant CF, Zhang L, Adsay V, Wicha M, Clarke MF, Simeone DM. Identification of pancreatic cancer stem cells. *Cancer Res*. 2007;67:1030–7.
- [42] Hermann PC, Huber SL, Herrler T, Aicher A, Ellwart JW, Guba M, Bruns CJ, Heeschen C. Distinct populations of cancer stem cells determine tumor growth and metastatic activity in human pancreatic cancer. *Cell Stem Cell*. 2007;1:313–23.
- [43] Collins AT, Berry PA, Hyde C, Stower MJ, Maitland NJ. Prospective identification of tumorigenic prostate cancer stem cells. *Cancer Res*. 2005;65:10946–51.
- [44] Al-Hajj M, Wicha MS, Benito-Hernandez A, Morrison SJ, Clarke MF. Prospective identification of tumorigenic breast cancer cells. *Proc Natl Acad Sci USA*. 2003;100:3983–8.

- [45] Zhang S, Balch C, Chan MW, Lai HC, Matei D, Schilder JM, Yan PS, Huang TH, Nephew KP. Identification and characterization of ovarian cancer-initiating cells from primary human tumors. *Cancer Res.* 2008;68:4311–20.
- [46] Valent P, Bonnet D, De MR, Lapidot T, Copland M, Melo JV, Chomienne C, Ishikawa F, Schuringa JJ, Stassi G, Huntly B, Herrmann H, Soulier J, Roesch A, Schuurhuis GJ, Wohrer S, Arock M, Zuber J, Cerny-Reiterer S, Johnsen HE, Andreeff M, Eaves C. Cancer stem cell definitions and terminology: the devil is in the details. *Nat Rev Cancer.* 2012;11:767–75.
- [47] Islam F, Qiao B, Smith RA, Gopalan V, Lam AK. Cancer stem cell: fundamental experimental pathological concepts and updates. *Exp Mol Pathol.* 2015;98:184–91.
- [48] Mimeault M, Batra SK. Recent insights into the molecular mechanisms involved in aging and the malignant transformation of adult stem/progenitor cells and their therapeutic implications. *Ageing Res Rev.* 2009;8:94–112.
- [49] Feldser DM, Hackett JA, Greider CW. Telomere dysfunction and the initiation of genome instability. *Nat Rev Cancer.* 2003;3:623–7.
- [50] Thiery JP, Acloque H, Huang RY, Nieto MA. Epithelial-mesenchymal transitions in development and disease. *Cell.* 2009;139:871–890.
- [51] Tsai JH, Yang J. Epithelial–mesenchymal plasticity in carcinoma metastasis. *Genes Dev.* 2013;27:2192–206.
- [52] Menendez JA, Alarcón T. Metabostemness: a new cancer hallmark. *Front Oncol.* 2014;4:262.
- [53] Lunt SY, Vander Heiden MG. Aerobic glycolysis: meeting the metabolic requirements of cell proliferation. *Annu Rev Cell Dev Biol.* 2011;27:441–64.
- [54] Abaza M, Luqmani YA. The influence of pH and hypoxia on tumor metastasis. *Expert Rev Anticancer Ther.* 2013;13:1229–42.
- [55] Dong C, Yuan T, Wu Y, Wang Y, Fan TW, Miriyala S, Lin Y, Yao J, Shi J, Kang T, Lorkiewicz P, St Clair D, Hung MC, Evers BM, Zhou BP. Loss of FBP1 by Snail-mediated repression provides metabolic advantages in basal-like breast cancer. *Cancer Cell.* 2013;23:316–31.
- [56] Shen YA, Wang CY, Hsieh YT, Chen YJ, Wei YH. Metabolic reprogramming orchestrates cancer stem cell properties in nasopharyngeal carcinoma. *Cell Cycle.* 2015;14:86–98.
- [57] Vlashi E, Lagadec C, Vergnes L, Matsutani T, Masui K, Poulou M. Metabolic state of glioma stem cells and nontumorigenic cells. *Proc Natl Acad Sci USA.* 2011;108:16062–7.
- [58] Lagadinou ED, Sach A, Callahan K, Rossi RM, Neering SJ, Minhajuddin M, Ashton JM, Pei S, Grose V, O'Dwyer KM, Liesveld JL, Brookes PS, Becker MW, Jordan CT. BCL-2 inhibition targets oxidative phosphorylation and selectively eradicates quiescent human leukemia stem cells. *Cell Stem Cell.* 2013;12:329–41.

- [59] Pattabiraman DR, Weinberg RA. Tackling the cancer stem cells—what challenges do they pose? *Nat Rev Drug Discov*. 2014;13:497–512.
- [60] Mimeault M, Batra SK. Altered gene products involved in the malignant reprogramming of cancer stem/progenitor cells and multitargeted therapies. *Mol Aspects Med*. 2014;39:3–32.
- [61] Ajani JA, Song S, Hochster HS, Steinberg IB. Cancer stem cells: the promise and the potential. *Semin Oncol*. 2015;42(Suppl 1):S3–17.
- [62] Mimeault M, Batra SK. Molecular biomarkers of cancer stem/progenitor cells associated with progression, metastases, and treatment resistance of aggressive cancers. *Cancer Epidemiol Biomarkers Prev*. 2014;23:234–54.
- [63] Kim JK, Jeon HY, Kim H. The molecular mechanisms underlying the therapeutic resistance of cancer stem cells. *Arch Pharm Res*. 2015;38:389–401.
- [64] Yu F, Yao H, Zhu P, Zhang X, Pan Q, Gong C, Huang Y, Hu X, Su F, Lieberman J, et al. let-7 regulates self renewal and tumorigenicity of breast cancer cells. *Cell*. 2007;131:1109–23.
- [65] Shay JW, Wright WE. Telomeres and telomerase in normal and cancer stem cells. *FEBS Lett*. 2010;584:3819–25.
- [66] Xi L, Cech TR. Inventory of telomerase components in human cells reveals multiple subpopulations of hTR and hTERT. *Nucleic Acids Res*. 2014;42:8565–77.
- [67] Terali K, Yilmazer A. New surprises from an old favourite: the emergence of telomerase as a key player in the regulation of cancer stemness. *Biochimie*. 2016;121:170–78.
- [68] Bollmann FM. Physiological and pathological significance of human telomerase reverse transcriptase splice variants. *Biochimie*. 2013;95:1965–70.
- [69] Smith LL, Collier HA, Roberts JM. Telomerase modulates expression of growth-controlling genes and enhances cell proliferation. *Nat Cell Biol*. 2003;5:474–9.
- [70] Beck S, Jin X, Sohn YW, Kim JK, Kim SH, Yin J, Pian X, Kim SC, Nam DH, Choi YJ, Kim H. Telomerase activity-independent function of TERT allows glioma cells to attain cancer stem cell characteristics by inducing EGFR expression. *Mol Cells*. 2011;31:9–15.
- [71] Kunimura C, Kikuchi K, Ahmed N, Shimizu A, Yasumoto S. Telomerase activity in a specific cell subset co-expressing integrin beta1/EGFR but not p75NGFR/bcl2/integrin beta4 in normal human epithelial cells. *Oncogene*. 1998;17:187–97.
- [72] Choi J, Southworth LK, Sarin KY, Venteicher AS, Ma W, Chang W, Cheung P, Jun S, Artandi MK, Shah N, Kim SK, Artandi SE. TERT promotes epithelial proliferation through transcriptional control of a Myc- and Wnt-related developmental program. *PLoS Genet*. 2008;4:e10.
- [73] Park JI, Venteicher AS, Hong JY, Choi J, Jun S, Shkreli M, Chang W, Meng Z, Cheung P, Ji H, McLaughlin M, Veenstra TD, Nusse R, McCrea PD, Artandi SE. Telomerase



- modulates Wnt signalling by association with target gene chromatin. *Nature*. 2009;460:66–72.
- [74] Musgrove EA, Caldon CE, Barraclough J, Stone A, Sutherland RL. Cyclin D as a therapeutic target in cancer. *Nat Rev Cancer*. 2011;11:558–72.
- [75] Hong J, Lee JH, Chung IK. Telomerase activates transcription of cyclin D1 gene through an interaction with NOL1. *J Cell Sci*. 2016;129:1566–79.
- [76] Jagadeesh S, Banerjee PP. Telomerase reverse transcriptase regulates the expression of a key cell cycle regulator, cyclin D1. *Biochem Biophys Res Commun*. 2006;347:774–80.
- [77] Li J, Huang X, Xie X, Wang J, Duan M. Human telomerase reverse transcriptase regulates cyclin D1 and G1/S phase transition in laryngeal squamous carcinoma. *Acta Otolaryngol*. 2011;131:546–51.
- [78] Mukherjee S, Firpo EJ, Wang Y, Roberts JM. Separation of telomerase functions by reverse genetics. *Proc Natl Acad Sci USA*. 2011;108:E1363–71.
- [79] Koh CM, Khattar E, Leow SC, Liu CY, Muller J, Ang WX, Li Y, Franzoso G, Li S, Guccione E, Tergaonkar V. Telomerase regulates MYC-driven oncogenesis independent of its reverse transcriptase activity. *J Clin Invest*. 2015;125:2109–22.
- [80] Hrdličková R, Nehyba J, Bose HR Jr. Alternatively spliced telomerase reverse transcriptase variants lacking telomerase activity stimulate cell proliferation. *Mol Cell Biol*. 2012;32:4283–96.
- [81] Hoffmeyer K, Raggioli A, Rudloff S, Anton R, Hierholzer A, Del Valle I, Hein K, Vogt R, Kemler R. Wnt/beta-catenin signaling regulates telomerase in stem cells and cancer cells. *Science*. 2012;336:1549–54.
- [82] Listerman I, Gazzaniga FS, Blackburn EH. An investigation of the effects of the core protein telomerase reverse transcriptase on Wnt signaling in breast cancer cells. *Mol Cell Biol*. 2014; 34:280–9.
- [83] Strong MA, Vidal-Cardenas SL, Karim B, Yu H, Guo N, Greider CW. Phenotypes in mTERT<sup>+/−</sup> and mTERT<sup>−/−</sup> mice are due to short telomeres, not telomere-independent functions of telomerase reverse transcriptase. *Mol. Cell Biol*. 2011;31:2369–79.
- [84] Maida Y, Yasukawa M, Furuuchi M, Lassmann T, Possemato R, Okamoto N, Kasim V, Hayashizaki Y, Hahn WC, Masutomi K. An RNA-dependent RNA polymerase formed by TERT and the RMRP RNA. *Nature*. 2009;461:230–5.
- [85] Gonzalez OG, Assfalg R, Koch S, Schelling A, Meena JK, Kraus J, Lechel A, Katz SF, Benes V, Scharffetter-Kochanek K, Kestler HA, Gunes C, Iben S. Telomerase stimulates ribosomal DNA transcription under hyperproliferative conditions. *Nat Commun*. 2014;5:4599.
- [86] Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144:646–74.

- [87] Dudognon C, Pendino F, Hillion J, Saumet A, Lanotte M, Segal-Bendirdjian E. Death receptor signaling regulatory function for telomerase: hTERT abolishes TRAIL-induced apoptosis, independently of telomere maintenance. *Oncogene*. 2004;23:7469–74.
- [88] Del Bufalo D, Rizzo A, Trisciuglio D, Cardinali G, Torrisi MR, Zangemeister-Wittke U, Zupi G, Biroccio A. Involvement of hTERT in apoptosis induced by interference with Bcl-2 expression and function. *Cell Death Differ*. 2005;12:1429–38.
- [89] Bermudez Y, Erasso D, Johnson N, Matte AD, Lowell N, Kruk P. Telomerase confers resistance to caspase-mediated apoptosis. *Clin Intervent Aging*. 2006;1:155–67.
- [90] Mandal M, Kumar R. Bcl-2 modulates telomerase activity. *J Biol Chem*. 1997;272:14183–7.
- [91] Rahman R, Latonen L, Wiman KG. hTERT antagonizes p53-induced apoptosis independently of telomerase activity. *Oncogene*. 2005;24:1320–7.
- [92] Zhou J, Mao B, Zhou Q, Ding D, Wang M, Guo P, Gao Y, Shay JW, Yuan Z, Cong YS. Endoplasmic reticulum stress activates telomerase. *Aging Cell*. 2014;13:197–200.
- [93] Sano R, Reed JC. ER stress-induced cell death mechanisms. *Biochim Biophys Acta*. 2013;1833:3460–70.
- [94] Hosoi T, Inoue Y, Nakatsu K, Matsushima N, Kiyose N, Shimamoto A, Tahara H, Ozawa K. TERT attenuated ER stress-induced cell death. *Biochem Biophys Res Commun*. 2014;447:378–82.
- [95] Takahashi K, Yamanaka S. Induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. *Cell*. 2006;126:663–76.
- [96] Liu A, Yu X, Liu S. Pluripotency transcription factors and cancer stem cells: small genes make a big difference. *Chin J Cancer*. 2013;32:483–487.
- [97] Okamoto N, Yasukawa M, Nguyen C, Kasim V, Maida Y, Possemato R, Shibata T, Ligon KL, Fukami K, Hahn WC, Masutomi K. Maintenance of tumor initiating cells of defined genetic composition by nucleostemin. *Proc Natl Acad Sci USA*. 2011;108:20388–93.
- [98] Liu Z, Li Q, Li K, Chen L, Li W, Hou M, Liu T, Yang J, Lindvall C, Bjorkholm M, Jia J, Xu D. Telomerase reverse transcriptase promotes epithelial-mesenchymal transition and stem cell-like traits in cancer cells. *Oncogene*. 2013;32:4203–13.
- [99] Sato N, Maehara N, Mizumoto K, Nagai E, Yasoshima T, Hirata K, Tanaka M. Telomerase activity of cultured human pancreatic carcinoma cell lines correlates with their potential for migration and invasion. *Cancer*. 2001;91:496–504.
- [100] Murasawa S, Uevadot J, Silver M, Isner JM, Losordo DW, Asahara T. Constitutive human telomerase reverse transcriptase expression enhances regenerative properties of endothelial progenitor cells. *Circulation*. 2002;106:1133–9.

- [101] Qu Y, Mao M, Li X, Zhang L, Huang X, Yang C, Zhao F, Xiong Y, Mu D. Enhanced migration and CXCR4 over-expression in fibroblasts with telomerase reconstitution. *Mol Cell Biochem.* 2008;313:45–52.
- [102] Ding D, Xi P, Zhou J, Wang M, Cong YS. Human telomerase reverse transcriptase regulates MMP expression independently of telomerase activity via NF-kappaB-dependent transcription. *FASEB J.* 2013;27:4375–83.
- [103] Chen P, Peng J, Huang L, Li W, Wang W, Cui Z, Han H, Gong L, Xiang D, Qiao S, Yu X, Wei Y, Ma L, Li N, Zhu J, Leng X. Overexpression of human telomerase reverse transcriptase promotes the motility and invasiveness of HepG2 cells in vitro. *Oncology Reports.* 2013;30:1157–64.
- [104] Ponnala S, Chetty C, Veeravalli KK, Dinh DH, Klopfenstein JD, Rao JS. MMP-9 silencing regulates hTERT expression via beta1 integrin-mediated FAK signaling and induces senescence in glioma xenograft cells. *Cell Signal.* 2011;23:2065–75.
- [105] He B, Xiao YF, Tang B, Wu YY, Hu CJ, Xie R, Yang X, Yu ST, Dong H, Zhao XY, Li JL, Yang SM. hTERT mediates gastric cancer metastasis partially through the indirect targeting of ITGB1 by microRNA-29a. *Sci Rep.* 2016;6:21955.
- [106] Park YP, Choi SC, Kim JH, Song EY, Kim JW, Yoon DY, Yeom YI, Lim JS, Kim JW, Paik SG, Lee HG. Up-regulation of Mac-2 binding protein by hTERT in gastric cancer. *Int J Cancer.* 2007;120:813–20.
- [107] Grassadonia A, Tinari N, Iurisci I, Piccolo E, Cumashi A, Innominato P, D'Egidio M, Natoli C, Piantelli M, Iacobelli S. 90K (Mac-2 BP) and galectins in tumor progression and metastasis. *Glycoconj J.* 2004;19:551–6.
- [108] Lassmann T, Maida Y, Tomaru Y, Yasukawa M, Ando Y, Kojima M, Kasim V, Simon C, Daub CO, Carninci P, Hayashizaki Y, Masutomi K. Telomerase reverse transcriptase regulates microRNAs. *Int J Mol Sci.* 2015;16:1192–208.
- [109] Volinia S, Calin GA, Liu CG, Ambs S, Cimmino A, Petrocca F, Visone R, Iorio M, Roldo C, Ferracin M, Prueitt RL, Yanaihara N, Lanza G, Scarpa A, Vecchione A, Negrini M, Harris CC, Croce CM. A microRNA expression signature of human solid tumors defines cancer gene targets. *Proc Nat Acad Sci USA.* 2006;103:2257–61.
- [110] Han M, Liu M, Wang Y, Chen X, Xu J, Sun Y, Zhao L, Qu H, Fan Y, Wu C. Antagonism of miR-21 reverses epithelial–mesenchymal transition and cancer stem cell phenotype through AKT/ERK1/2 inactivation by targeting PTEN. *PLoS One.* 2012;7:e39520.
- [111] Haghpanah V, Fallah P, Tavakoli R, Naderi M, Samimi H, Soleimani M, Larijani B. Antisense-miR-21 enhances differentiation/apoptosis and reduces cancer stemness state on anaplastic thyroid cancer. *Tumor Biol.* 2016;37:1299–308.
- [112] Bagheri S, Nosrati M, Li S, Fong S, Torabian S, Rangel J, Moore DH, Federman S, Laposa RR, Baehner FL, Sagebiel RW, Cleaver JE, Haqq C, Debs RJ, Blackburn EH, Kashani-Sabet M. Genes and pathways downstream of telomerase in melanoma metastasis. *Proc Natl Acad Sci USA.* 2006;103:11306–11.

- [113] Wu KJ, Grandori C, Amacker M, Simon-Vermot N, Polack A, Lingner J, Dalla-Favera R. Direct activation of TERT transcription by c-MYC. *Nat Genet.* 1999;21:220–4.
- [114] Sharma GG, Gupta A, Wang H, Scherthan H, Dhar S, Gandhi V, Iliakis G, Shay JW, Young CSH, Pandita TK. hTERT associates with human telomeres and enhances genomic stability and DNA repair. *Oncogene.* 2003;22:131–46.
- [115] Shin KH, Kang MK, Dicterow E, Kameta A, Baluda MA, Park NH. Introduction of human telomerase reverse transcriptase to normal human fibroblasts enhances DNA repair capacity. *Clin Cancer Res.* 2004;10:2551–60.
- [116] Fleisig HB, Hukezalie KR, Thompson CAH, Au-Yeung TTT, Ludlow AT, Xhao CR, Wong JMY. Telomerase reverse transcriptase expression protects transformed human cells against DNA-damaging agents, and increases tolerance to chromosomal instability. *Oncogene.* 2015;doi:10.1038/onc.2015.75.
- [117] Masutomi K, Possemato R, Wong JM, Currier JL, Tothova Z, Manola JB, Ganesan S, Lansdorp PM, Collins K, Hahn WC. The telomerase reverse transcriptase regulates chromatin state and DNA damage responses. *Proc Natl Acad Sci USA.* 2005;102:8222–27.
- [118] Ahmed S, Passos JF, Birket MJ, Beckmann T, Brings S, Peters H, Birch-Machin MA, von Zglinicki T, Saretzki G. Telomerase does not counteract telomere shortening but protects mitochondrial function under oxidative stress. *J Cell Sci.* 2008;121:1046–53.
- [119] Haendeler J, Dröse S, Büchner N, Jakob S, Altschmied J, Goy C, Spyridopoulos I, Zeiher AM, Brandt U, Dimmeler S. Mitochondrial telomerase reverse transcriptase binds to and protects mitochondrial DNA and function from damage. *Arterioscler Thromb Vasc Biol.* 2009;29:929–35.
- [120] Indran I, Hande P, Pervaiz S. hTERT overexpression alleviates intracellular ROS production, improves mitochondrial function, and inhibits ROS-mediated apoptosis in cancer cells. *Cancer Res.* 2011;71:266–76.
- [121] Singhapol C, Pal D, Czapiewski R, Porika M, Nelson G, Saretzki GC. Mitochondrial telomerase protects cancer cells from nuclear DNA damage and apoptosis. *PLoS One.* 2013;8:e52989.
- [122] Listerman I, Sun J, Gazzaniga FS, Lukas JL, Blackburn EH. The major reverse transcriptase-incompetent splice variant of the human telomerase protein inhibits telomerase activity but protects from apoptosis. *Cancer Res.* 2013;73:2817–28.
- [123] Ghosh A, Saginc G, Leow SC, Khattar E, Shin EM, Yan TD, Wong M, Zhang Z, Li G, Sung WK, Zhou J, Chng WJ, Li S, Liu E, Tergaonkar V. Telomerase directly regulates NF-kappaB-dependent transcription. *Nat Cell Biol.* 2012;14:1270–81.
- [124] Yin L, Hubbard AK, Giardina C. NF-kappa B regulates transcription of the mouse telomerase catalytic subunit. *J Biol Chem.* 2000;275:36,671–5.

- [125] Zhou L, Zheng D, Wang M, Cong YS. Telomerase reverse transcriptase activates the expression of vascular endothelial growth factor independent of telomerase activity. *Biochem Biophys Res Commun.* 2009;386:739–43.
- [126] Liu N, Ding D, Hao W, Yang F, Wu X, Wang M, Xu X, Ju Z, Liu JP, Song Z, Shay JW, Guo Y, Cong YS. hTERT promotes tumor angiogenesis by activating VEGF via interactions with the Sp1 transcription factor. *Nuc Acids Res.* 2016 (in press)
- [127] Bermudez Y, Yang H, Saunders BO, Cheng JQ, Nicosia SV, Kruk PA. VEGF- and LPA-induced telomerase in human ovarian cancer cells is Sp1-dependent. *Gynecol Oncol.* 2007;106:526–37.
- [128] Zhong YQ, Xia ZS, Fu YR, Zhu ZH. Knockdown of hTERT by siRNA suppresses growth of Capan-2 human pancreatic cancer cell via the inhibition of expressions of Bcl-2 and COX-2. *J Dig Dis.* 2010;11:176–84.
- [129] Liu L, Liu C, Lou F, Zhang G, Wang X, Fan Y, Yan K, Wang K, Xu Z, Hu S, Björkholm M, Xu D. Activation of telomerase by seminal plasma in malignant and normal cervical epithelial cells. *J Pathol.* 2011;225:203–11.
- [130] Jäger K, Walter M. Therapeutic targeting of telomerase. *Genes.* 2016;7:39.
- [131] Brennan SK, Wang Q, Tressler R, Harley C, Go N, Bassett E, Huff CA, Jones RJ, Matsui W. Telomerase inhibition targets clonogenic multiple myeloma cells through telomere length-dependent and independent mechanisms. *PLoS One.* 2010;5:e12487.
- [132] Joseph I, Tressler R, Bassett E, Harley C, Buseman CM, Pattamatta P, Wright WE, Shay JW, Go NF. The telomerase inhibitor imetelstat depletes cancer stem cells in breast and pancreatic cancer cell lines. *Cancer Res.* 2010;70:9494–504.
- [133] Gryaznov SM. Oligonucleotide n3'→p5' phosphoramidates and thio-phosphoramidates as potential therapeutic agents. *Chem Biodivers.* 2010;7:477–93.