

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Gene-Environment Interactions: The Case of Asbestosis

Vita Dolzan, Metoda Dodic-Fikfak and
Alenka Franko

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/66120>

Abstract

It is becoming evident that both environmental/lifestyle and genetic factors may influence the development of many diseases. This chapter highlights the importance of considering gene-environment interactions, which is shown on the example of our studies into asbestosis, one of the most frequent asbestos-related diseases. Asbestos fibres induce generation of reactive oxygen and nitric species (ROS and RNS), and it is generally accepted that ROS and RNS are involved in the pathogenesis of asbestos-related diseases. Human tissues contain specific enzymes that metabolise ROS and RNS, such as superoxide dismutases (SODs), catalase (CAT), glutathione-S-transferases (GSTs) and inducible nitric oxide synthase (iNOS). As these enzymes are encoded by polymorphic genes, genetic variability in an individual's capacity to detoxify these reactive species may modify the risk for disease. Our previous studies into asbestosis showed that the associations between the risk of asbestosis and *MnSOD* Ala-9Val polymorphism and between asbestosis and *iNOS* genotypes were modified by *CAT* -262C>T polymorphism. A strong interaction was also found between smoking (lifestyle factor) and *GSTM1*-null polymorphism, between smoking and *iNOS* (CCTTT)_n polymorphism and between cumulative asbestos exposure (environmental factor) and *iNOS* (CCTTT)_n polymorphism. The findings of our studies and other studies indicate that in addition to environmental and/or occupational exposure to different hazards and lifestyle factors, genetic factors as well as the interactions between different genotypes, between genotypes and lifestyle factors and between genotypes and environmental/occupational exposure to hazards may also have an important role on the development of diseases and should be further investigated.

Keywords: asbestosis, exposure, gene-environment interactions, gene-gene interactions

1. Introduction

It is becoming evident that both environmental and genetic factors may influence the development of many diseases [1–7]. It is therefore important to consider gene-environment interactions when studying diseases related to exposure to different hazards and lifestyle factors.

Environmental and lifestyle factors have been investigated in many epidemiological studies using self-reported information obtained by questionnaires, interviews, records or measurements of exposure. However, very few epidemiological studies included the information on genetic risk factors. Similarly, many studies investigating genetic factors obtained little information on environmental factors and lifestyle. Genetic predisposition can be presumed from family history, from phenotypic characteristics (e.g. metabolic capacity) or, most importantly, from an analysis of deoxyribonucleic acid (DNA) sequence [8].

The research into gene-environment interactions requires the information on both environmental/lifestyle and genetic factors [7, 8]. Primary candidates for gene-environment interaction studies have been mostly genes coding for xenobiotic-metabolising enzymes [3]. Genetic variability in these genes may lead to interindividual differences in the capacity for xenobiotic metabolism, thus modifying an individual's susceptibility to the development of disease [3].

The approach to the analysis of gene-environment interactions is presented using the example of our study into asbestosis, which is one of the most frequent asbestos-related diseases. According to the model of causation, asbestos exposure, genetic factors and possibly also unknown causes have a crucial role in the occurrence of asbestosis [9]. Although asbestos-related diseases are among the most extensively studied occupational diseases, and the causal relationship between asbestos exposure and asbestosis has been well proved [10–14], relatively little has been known about the genetic factors that might modify an individual's susceptibility to the development of this disease [6, 15–17].

2. Asbestos exposure

Asbestos is a commercial name for a group of fibrous silicates with certain toxic properties, such as the ability to produce inflammation, fibrous scarring and cancer [18–20]. Based on their physical and chemical structures, asbestos fibres can be classified into two major groups: chrysotile and amphiboles [20–25].

Occupational exposure to asbestos occurs in asbestos mining, production and milling of asbestos fibres; in asbestos cement industry; in construction; in machine and insulation product industry; in ship building or repair; in car industry; in production of brakes and clutches; in car, bus, lorry, railway carriage and aeroplane repair; in asphalt mixing; in disposal of asbestos waste and materials; in brickworks; in textile industry and in other industries and activities [20, 22, 26–28].

Local population can be exposed to asbestos mostly in the neighbourhood of factories where asbestos is produced or used (exposure to polluted air, water and food). The source of environmental asbestos exposure may also be asbestos cement sheets, asbestos insulators and other asbestos-containing products. Asbestos fibres may also be found in water which flows through asbestos cement pipes, especially if they have been damaged. Workers exposed to asbestos may bring asbestos home to the family members on clothing or hair [26–28].

3. Asbestos-related diseases

Asbestos exposure has been associated with the development of asbestosis; pleural diseases, such as pleural plaques, diffuse pleural thickening and pleural effusion and several types of cancer: lung cancer, diffuse malignant mesothelioma of the pleura and peritoneum, cancer of the larynx, cancer of the ovary as well as the cancers of the buccal mucosa, the pharynx, the gastrointestinal tract and the kidney [11, 12, 16, 25, 29–41].

4. Clinical presentation of asbestosis

Asbestosis is an interstitial pulmonary process that develops into diffuse pulmonary fibrosis after a long latency period [42, 43]. The disease continues to progress even after the cessation of exposure, and the process is irreversible. One of the earliest symptoms may be dyspnoea, which is manifested at first only after strenuous exertion, but subsequently with less and less exertion, and eventually it appears even at rest. Another non-specific symptom and usually late manifestation of the disease is irritating and dry, usually non-productive cough, sometimes associated with chest pain [42, 44]. Pulmonary function changes are characterised mostly by a restrictive impairment [27, 28, 42–44]. Later, obstructive airway impairment may also occur [27, 28]. On chest radiographs, small irregular opacities appear initially in the lower lung fields that may enlarge with more advanced disease and involve also middle lung fields [27, 42–44]. Characteristic features of asbestosis on high-resolution computed tomography (HRCT) include fibrotic intralobular interstitial thickening and interlobular septal thickening, sub-pleural lines and opacities, parenchymal bands, ground-glass opacities and, in more severe disease, variable honeycombing [27].

5. Reactive oxygen and nitric species: the link between asbestos exposure and the development of asbestosis

The pathogenesis of asbestosis is still poorly understood. The findings of studies on cell cultures and animal models indicate that reactive oxygen and nitric species (ROS and RNS) are involved in the pathogenesis of this disease [23, 30, 45–55]. The most important reactive metabolites in the pathogenesis of asbestos-related lung diseases are superoxide anion (O_2^-),

hydrogen peroxide (H_2O_2), hydroxyl radical (OH^\cdot) and nitric oxide (NO) [46, 48, 56, 57]. Asbestos may stimulate the production of ROS in two different ways. The first mechanism involves redox-active iron (Fe^{2+} , Fe^{3+}) in asbestos that catalyses the formation of OH^\cdot , whereas the second mechanism involves the production of ROS by alveolar macrophages during the phagocytosis of asbestos fibres [58–60]. Reactive oxygen species in lungs may lead to the production of cytotoxic and potentially genotoxic electrophilic compounds [46].

It has also been suggested that asbestos fibres may upregulate the activity of inducible nitric oxide synthase (iNOS), thus inducing the production of NO by alveolar macrophages and pulmonary epithelial cells [51, 61–64]. Because NO is a free radical, it reacts readily with other reactive oxygen metabolites (as, for instance, O_2^-), leading to the formation of toxic metabolites, most importantly peroxynitrite [65–69]. Nitric oxide may play a role in the initiation and progression of asbestosis [51, 64, 70, 71]. However, the data presented by Dörger et al. [72] indicate that iNOS-derived NO plays a dual role in acute asbestos-induced lung injury and that although iNOS deficiency resulted in an exacerbated inflammatory response, it improved oxidant-promoted lung tissue damage.

Reactive oxygen species and RNS can damage all types of biomolecule, including lipids, proteins and deoxyribonucleic acid (DNA). Complex defence mechanisms, including enzymes, proteins and antioxidants, are involved in the prevention of cell damage [73, 74].

6. Enzymes involved in the detoxification of reactive oxygen and nitric species

Human tissues contain specific enzyme systems to detoxify ROS and RNS. Superoxide dismutases (SODs) and catalase (CAT) together with glutathione peroxidases represent an important line of the primary antioxidant enzyme defence system against ROS. Superoxide dismutases catalyse the dismutation of O_2^- to H_2O_2 and oxygen (O_2), whereas CAT subsequently catalyses the conversion of H_2O_2 to water (H_2O) and O_2 [48, 75–82]. Three distinct SOD isoenzymes have been identified in mammals: a cytosolic copper-zinc SOD (CuZnSOD or SOD1) localised in cytoplasmic compartment with copper (Cu) and zinc (Zn) in the catalytic centre, manganese SOD (MnSOD or SOD2) that is localised in mitochondria and uses manganese (Mn) as a cofactor and extracellular SOD (ECSOD or SOD3) that also contains Cu and Zn in the catalytic centre and is located in the extracellular space [74, 82, 83].

Another important family of enzymes involved in the detoxification of xenobiotics and electrophiles produced by ROS and RNS is glutathione S-transferases (GSTs) [84–87]. They catalyse the conjugation of reduced glutathione to different electrophiles [88]. These conjugation reactions mostly result in less reactive products [89]; however, in some cases, the products are more reactive and consequently more harmful than the parent compound [90, 91]. Seven classes of cytosolic GST isoenzymes have been recognised in mammals (Alpha, Mu, Pi, Sigma, Theta, Omega, Zeta) [84–86, 91, 92]. The major GST enzyme in the human lung is GSTP1, which belongs to the Pi class [90, 91, 93], while GSTM1 (Mu class) and GSTT1 (Theta class) were most frequently investigated [90, 91].

7. Genetic variability of metabolic enzymes

Genetic polymorphisms are the most common cause for genetic variability of detoxification and antioxidative enzymes [15–17, 80, 91, 94–99].

The most common functional single nucleotide polymorphism (SNP) of the *MnSOD* gene is C to T substitution (c.201C>T, rs4880), which results in alanine (Ala) to valine (Val) amino acid change at position –9 of the mitochondrial targeting sequence (MnSOD p.Ala-9Val) [96, 97, 100]. It has been suggested that this SNP alters the secondary structure of the protein and hence may affect the efficiency of transport of the MnSOD into the mitochondria, where it would be biologically available [96, 97].

ECSOD is secreted into extracellular space where it binds lung matrix components and inhibits their fragmentation in response to oxidative stress [101, 102]. In the *ECSOD* gene, a C to G substitution (c.896C>G, rs1799895) leads to amino acid change from arginine (Arg) to glycine (Gly) at position 213 (p.Arg213Gly) [89, 100, 103–105]. This polymorphism causes an 8- to 15-fold increase in the concentration of plasma ECSOD levels due to impaired binding to the extracellular matrix [103, 104].

The most common functional SNP of the catalase gene (*CAT*) consists of a C to T substitution at position –262 in the promoter region (*CAT* c.–262C>T) and has a substantial impact on the basal expression as well as the *CAT* levels in red blood cell [80]. The findings of later studies indicated lower *CAT* activity in subjects with the –262TT genotype than those with the CT and CC genotypes [106–111].

Regarding GSTs, the most common polymorphism of the *GSTM1* and *GSTT1* genes in most of the populations is null polymorphism due to homozygous deletion (null genotype) of these genes, which result in the absence of the *GSTM1* and *GSTT1* enzyme activity [17, 91]. *GSTM1*-null genotype has been associated with an increased risk of asbestosis in some studies [16, 86], while this association has not been proved in the others [15, 17]. No association has been found between *GSTT1* deletion polymorphism and asbestosis in the studies published so far [17, 86]. As for the *GSTP1* gene, two common single nucleotide polymorphisms in the coding sequence were reported to result in amino acid substitution that may lead to reduced conjugating activity of the enzyme [91, 98, 112, 113]. The first polymorphism is characterised by adenine (A) to guanine (G) transition of nucleotide 313 in exon 5 (c.A313G), which causes an isoleucine (Ile) to valine (Val) substitution at position 105 of the *GSTP1* enzyme (p.Ile105Val), resulting in three possible genotypes: 105 Ile/Ile, 105 Ile/Val or 105 Val/Val. The second polymorphism involves the cytosine (C) to thymine (T) transition at nucleotide 341 in exon 6 (c.C341T), which results in alanine (Ala) to Val substitution at position 114 of the *GSTP1* enzyme (p.Ala114Val). Regarding codon 114, three genotypes are also possible: 114 Ala/Ala, 114 Ala/Val or 114 Val/Val [91, 98]. Based on the presence of the polymorphisms in both codons 105 and 114, *GSTP1* genotypes can be combined into groups with a presumed high, intermediate or low conjugation capacity of the enzyme.

The human *iNOS* gene is also known to be polymorphic. Several types of polymorphisms have been identified in the promoter region of the *iNOS* gene [99, 114]. The CCTTT pentanucleotide

tandem repeat polymorphisms have been associated with the transcriptional promoter activity, which has been shown to increase with the CCTTT repeat number. Based on that, alleles with 11 or fewer CCTTT repeats are usually defined as short alleles (S) and the ones with 12 or more repeats as long alleles (L). Accordingly, the subjects can have SS, SL or LL genotype [115].

8. Gene-environment interactions and asbestosis

We are presenting the example of an approach to gene-environment interaction research by summarising and building on the results of our studies that aimed to investigate the influence of interactions between different genotypes (*MnSOD*, *ECSOD*, *CAT*, *GSTM1*, *GSTT1*, *GSTP*, *iNOS*), between genotypes and smoking and between genotypes and cumulative asbestos exposure on the risk of developing asbestosis [6, 14, 116–119].

A nested case-control study included 262 cases with asbestosis and 265 controls with no asbestos-related disease. All the subjects included in the study were employed in the asbestos cement manufacturing plant of Salonit Anhovo, Slovenia, and occupationally exposed to asbestos. Data on smoking were obtained from all subjects using a standardised questionnaire [25, 120] and checked during the interview. The data on the cumulative asbestos exposure, expressed in fibres/cm³-years [intensity in fibres per cm³ of air multiplied by time of exposure expressed in years], were available for all the subjects from the previous study [25]. The diagnosis of asbestosis or 'no asbestos-related disease' was based on the Helsinki Criteria for Diagnosis and Attribution of Asbestos Diseases [121] and on the American Thoracic Society recommendations [122]. Each case was confirmed by an interdisciplinary group of experts (consisting of an occupational physician, a radiologist and a pulmonologist) of the State Board for Recognition of Occupational Asbestos Diseases at the Clinical Institute of Occupational Medicine. Capillary blood samples from the finger tips of all cases and controls have been collected on FTA Mini Cards (Whatman Bioscience) for the isolation of deoxyribonucleic acid (DNA) and genotyping. All the genetic analyses were performed using PCR-based approaches as previously described [6, 14, 116–119].

Before testing interactions, the associations between outcome [in our case asbestosis] and individual variables were assessed using univariate logistic regression analysis. As expected, asbestosis was associated with cumulative asbestos exposure, whereas no association was found with smoking (OR = 0.98, 95% CI = 0.69–1.39 for ever versus never smoking) [14]. Analysing the association between asbestosis and individual genotypes, an important association was observed between asbestosis and *MnSOD* genotype (OR = 1.50, 95% CI = 1.01–2.24 for -9Ala/Ala versus combined Ala/Val and Val/Val genotypes) [118]. Only non-significantly elevated risk of asbestosis was observed for the *ECSOD* and *CAT* genotypes (OR = 1.63, 95% CI = 0.62–4.27 for *ECSOD* 213Arg/Gly versus the Arg/Arg genotype and OR = 1.36, 95% CI = 0.70–2.62 for *CAT* -262 TT compared to combined CT and CC genotypes, respectively) [117, 118]. Regarding GSTs, no association was found between asbestosis and *GSTM1*-null genotype (OR = 1.01, 95% CI = 0.71–1.43), while the presence of *GSTT1*-null genotype showed

a protective effect for this disease (OR = 0.61, 95% CI = 0.40–0.94) [14]. On the other hand, *GSTP1* genotype coding for an enzyme with a high conjugation capacity versus genotypes resulting in an intermediate or low enzyme activity significantly increases the risk of developing asbestosis (OR = 1.49, 95% CI 1.06–2.10) [116]. A slightly elevated risk of asbestosis was also found for the *iNOS* LL genotype compared to the combined SL and SS genotypes (OR = 1.20, 95% CI = 0.85–1.69) [119]. Based on the above-mentioned results, it could be suggested that the genotypes may increase, decrease or have no effect on the risk of disease, in our case asbestosis.

Univariate modelling was followed by multivariate analysis and interactions as the genes usually do not act independently, but may interact. To test the interactions, simple categorical models based on stratification were constructed first, followed by logistic regression models using dummy variables. The analysis showed that the association between asbestosis and *MnSOD* Ala-9Val genotypes was modified strongly by *CAT* -262 C>T genotypes. An increased risk of developing asbestosis was observed for the combined *MnSOD* -9Ala/Val and Val/Val genotypes compared to the Ala/Ala genotype only among those subjects who also had *CAT* -262TT genotype, suggesting an interaction, which was further confirmed by logistic regression analysis using dummy variables (OR = 4.49, 95% CI = 1.08–18.61) [6]. Considering that both MnSOD and *CAT* constitute a part of the primary defence system against ROS and catalyse the consecutive reactions in the detoxification of ROS [48, 74, 80, 82], this interaction could be considered as logical and biologically plausible. Similarly, the association between asbestosis and *iNOS* (CCTTT)_n genotypes was also modified by *CAT* -262 C>T genotypes, where a higher asbestosis risk for the *iNOS* LL genotype versus the combined SL and SS genotypes was observed only among those who had *CAT* -262 TT genotype (OR = 4.78, 95% CI = 1.15–19.81) [6]. Taking into account that reactions between ROS and NO have been proposed to potentiate the cytotoxic and mutagenic effect of asbestos fibres [48, 51, 64, 71] and based on the assumption that NO produced by the catalytic activity of iNOS can function as a protective agent against toxic effects of H₂O₂ [123], which is detoxified by *CAT* [48, 74, 80, 82], and vice versa that H₂O₂ decreases the cytotoxicity of NO [124], this interaction could also be considered as biologically plausible [6].

Next, interactions between different genotypes and an important lifestyle factor—in our case smoking—have been tested. We observed that the *GSTM1*-null polymorphism did modify the association between smoking and asbestosis, although there was no independent association between either *GSTM1*-null polymorphism or smoking and asbestosis risk (OR = 2.67, 95% CI = 1.31–5.46) [6]. We can explain this modifying effect with the observation that both asbestos and smoking increase the production of ROS [46, 125, 126], which are known to be involved in the pathogenesis of asbestosis [23, 30, 46, 48–50]. It has been suggested that cigarette smoke and asbestos increase DNA damage and ROS production in pulmonary cells synergistically [125–127]. In line with these reports and considering the role of *GSTM1* in the defence against ROS [84–87], this observation could also be considered as biologically plausible [6]. Similarly, the association between smoking and asbestosis was modified by *iNOS* (CCTTT)_n polymorphism (OR = 2.00, 95% CI = 0.99–4.03) [6]. Knowing that cigarette smoke is the largest source of NO that humans are exposed to and can also increase the expression and activity of iNOS [128, 129] and based on the suggestion that asbestos fibres may upregulate the activity

of iNOS and thus the production of NO, which is thought to play an important role in the initiation and progression of asbestosis [51, 70], this interaction could also be physiologically explained [6].

Finally, we present an example of the interaction between genotypes and environmental exposure, in our case occupational exposure to asbestos. In order to assess the interactions between the genotypes and occupational cumulative asbestos exposure, we have first constructed simple categorical models that included cumulative asbestos exposure categorised as follows: ≤ 11.23 fibres/cm³-years and >11.23 fibres/cm³-years (11.23 fibres/cm³-years was the average cumulative asbestos exposure for the controls). In our analysis, we have observed that the association between asbestosis and cumulative asbestos exposure was modified by the iNOS (CCTTT)_n genotypes (OR = 5.74; 95% CI = 3.30–9.99) [6].

9. Conclusions

The findings of our studies suggest that in addition to environmental and/or occupational exposure to different hazards and lifestyle factors, the genetic factors and the interactions between different genotypes, between genotypes and lifestyle factors and between genotypes and environmental/occupational exposure to hazards may have an important influence on the development of diseases and should be further investigated [6, 130–133]. In agreement with our observations, an increasing number of molecular epidemiological studies support the importance of investigating not only genetic predisposition but also gene-gene and gene-environment interactions when assessing the risk of developing diseases [134–136]. Novel high-throughput technologies may also allow the investigation of interactions between exposure to hazards and epigenetic changes in disease risk assessment [137].

Acknowledgement

This work was financially supported by the Slovenian Research Agency [grants L3-3648 and P1-0170].

Author details

Vita Dolzan¹, Metoda Dodic-Fikfak² and Alenka Franko^{2*}

*Address all correspondence to: alenka.franko@siol.net

1 Pharmacogenetics Laboratory, Institute of Biochemistry, Faculty of Medicine, University of Ljubljana, Slovenia

2 Clinical Institute of Occupational Medicine, University Medical Centre, Ljubljana, Slovenia

References

- [1] Greenland S, Rothman KJ. Concepts of interaction. In: Rothman KJ, Greenland S, editors. *Modern epidemiology* Philadelphia, PA: Lippincott-Raven, 1998. pp. 329–342.
- [2] Furberg AH, Ambrosone CB. Molecular epidemiology, biomarkers and cancer prevention. *Trends Mol Med*. 2001;7:517–521. PubMed PMID: 11689338.
- [3] Mucci LA, Wedren S, Tamimi RM, Trichopoulos D, Adami HO. The role of gene-environment interaction in the aetiology of human cancer: examples from cancers of the large bowel, lung and breast. *J Intern Med*. 2001;249:477–493. PubMed PMID: 11422654.
- [4] Boks MP, Schipper M, Schubart CD, Sommer IE, Kahn RS, Ophoff RA. Investigating gene environment interaction in complex diseases: increasing power by selective sampling for environmental exposure. *Int J Epidemiol*. 2007;36:1363–1369. PubMed PMID: 17971387.
- [5] Khoury MJ, Millikan R, Gwinn M. Genetic and molecular epidemiology. In: Rothman KJ, Greenland S, Lash TL editors. *Modern Epidemiology* Philadelphia, Baltimore, New York, London, Buenos Aires, Hong Kong, Sydney, Tokyo: Lippincott-Raven, 2008. pp. 564–597.
- [6] Franko A, Dolžan V, Arnerić N, Dodič-Fikfak M. The influence of gene-gene and gene-environment interactions on the risk of asbestosis. *Biomed Res Int*. 2013;2013:405743. PubMed PMID: 23984360; doi: 10.1155/2013/405743.
- [7] Lu M, Lee HS, Hadley D, Huang JZ, Qian X. Logistic principal component analysis for rare variants in gene-environment interaction analysis. *IEEE/ACM Trans Comput Biol Bioinform*. 2014;1:1020–1028. PubMed PMID: 26357039; doi: 10.1109/TCBB.2014.2322371.
- [8] Hunter DJ. Gene-environment interactions in human diseases. *Nat Rev Genet*. 2005;6:287–298. PubMed PMID: 15803198.
- [9] Rothman KJ, Greenland S, Poole C, Lash TL. Causation and cause inference. In: Rothman KJ, Greenland S, Lash TL editors. *Modern Epidemiology* Philadelphia, Baltimore, New York, London, Buenos Aires, Hong Kong, Sydney, Tokyo: Lippincott-Raven, 2008. pp. 5–22.
- [10] Mossman BT, Gee JB. Asbestos-related diseases. *N Engl J Med*. 1989;320:1721–1730. PubMed PMID: 2659987.
- [11] Dement JM, Brown DP, Okun A. Follow-up study of chrysotile asbestos textile workers: cohort mortality and case-control analyses. *Am J Ind Med*. 1994;26:431–447. PubMed PMID: 7810543.

- [12] Dement JM, Brown DP. Lung cancer mortality among asbestos textile workers: a review and update. *Ann Occup Hyg.* 1994;38:525–532. PubMed PMID: 7978974.
- [13] Jakobsson K, Strömberg U, Albin M, Welinder H, Hagmar L. Radiological changes in asbestos cement workers. *Occup Environ Med.* 1995;52:20–27. PubMed PMID: 7697136.
- [14] Franko A, Dodič-Fikfak M, Arnerić N, Dolžan V. Glutathione S- transferases GSTM1 and GSTT1 polymorphisms and asbestosis. *J Occup Environ Med.* 2007;49:667–671. PubMed PMID: 17563610.
- [15] Jakobsson K, Rannug A, Alexandrie AK, Rylander L, Albin M, Hagmar L. Genetic polymorphism for glutathione-S-transferase mu in asbestos cement workers. *Occup Environ Med.* 1994;51:812–816. PubMed PMID: 7849864.
- [16] Smith CM, Kelsey KT, Wiencke JK, Leyden K, Levin S, Christiani DC. Inherited glutathione-S-transferase deficiency is a risk factor for pulmonary asbestosis. *Cancer Epidemiol Biomarkers Prev.* 1994;3:471–477. PubMed PMID: 8000297.
- [17] Hirvonen A, Saarikoski ST, Linnainmaa K, Koskinen K, Husgafvel-Pursiainen H, Mattson K, Vainio H. Glutathione S-transferase and N-acetyltransferase genotypes and asbestos-associated pulmonary disorders. *J Natl Cancer Inst.* 1996;88:1853–6185. PubMed PMID: 8961976.
- [18] Becklake MR. Exposure to asbestos and human disease. *N Engl J Med.* 1982; 06:1480–1482. PubMed PMID: 7078593.
- [19] Craighead JE, Mossman BT. The pathogenesis of asbestos-associated diseases. *N Engl J Med.* 1982;306:1446–1455. PubMed PMID: 7043267.
- [20] Wagner GR, Hearl FJ. Mineral dust: asbestos, silica, coal, manufactured fibers. In: Rosenstock L, Cullen MR, Brodtkin CA, Redlich CA, editors. *Textbook of clinical occupational and environmental medicine.* 2nd ed. Philadelphia, Edinburgh, New York, St Louis, Sydney, Toronto: Elsevier Saunders, 2005. pp. 1073–1078.
- [21] International Agency for Research on Cancer [IARC]. IARC Working Group. *Asbestos.* Lyon: IARC, 1972.
- [22] International Agency for Research on Cancer [IARC]. IARC monographs on the valuation of carcinogenic risk of chemicals to man. Some inorganic and organometallic compounds. Lyon: IARC, 1973. pp. 17–47.
- [23] Mossman BT. Mechanism of asbestos carcinogenesis and toxicity. The amphibole hypothesis revised. *Br J Ind Med.* 1993;50:673–676. PubMed PMID: 8398854.
- [24] Piolatto G, Negri E, La Vecchia C, Pira E, Decarli A, Peto J. An update of cancer mortality among chrysotile asbestos miners in Balangero, Northern Italy. *Br J Ind Med.* 1990;47:810–814. PubMed PMID: 2176805.

- [25] Dodič-Fikfak M, Kriebel D, Quinn MM, Eisen EA, Wegman DH. A case control study of lung cancer and exposure to chrysotile and amphibole at a Slovenian asbestos-cement plant. *Ann Occup Hyg.* 2007;51:261–268. PubMed PMID: 17351264.
- [26] Dodič-Fikfak M., Šešok J. National guidelines for asbestos: final report (1.12.1998–31.10.1999). Ljubljana: Inštitut za varovanje zdravja RS, 1999.
- [27] Brodtkin CA, Rosenstock L. Asbestos and asbestos-related pleural disease. In: Rosenstock L, Cullen MR, Brodtkin CA, Redlich CA editors. *Textbook of clinical occupational and environmental medicine*. 2nd ed. Philadelphia, Edinburgh, New York, St Louis, Sydney, Toronto: Elsevier Saunders, 2005. pp. 364–377.
- [28] Rom WN. Asbestosis, pleural fibrosis, and lung cancer. In: Rom WN, Markowitz SB, editors. *Environmental and occupational medicine*. 4th ed. Philadelphia, Baltimore, New York, London, Buenos Aires, Hong Kong, Sydney, Tokyo: Wolters Kluwer, Lippincott Williams & Wilkins, 2007. pp. 298–316.
- [29] Frumkin H, Berlin J. Asbestos exposure and gastrointestinal malignancy review and meta-analysis. *Am J Ind Med.* 1988;14:79–95. PubMed PMID: 3189361.
- [30] Mossman BT, Bignon J, Corn M, Seaton A, Gee JB. Asbestos: scientific developments and implications for public policy. *Science.* 1990;247:294–301. PubMed PMID: 2153315.
- [31] Selikoff IJ, Seidman H. Asbestos-associated deaths among insulation workers in the United States and Canada, 1967–1987. *Ann N Y Acad Sci.* 1991;643:1–14. PubMed PMID: 1809121.
- [32] McDonald JC, Liddell FD, Dufresne A, McDonald AD. The 1891–1920 birth cohort of Quebec chrysotile miners and millers: mortality 1976–88. *Br J Ind Med.* 1993;50:1073–1081. PubMed PMID: 8280638.
- [33] Giaroli C, Belli S, Bruno C, Candela S, Grignoli M, Minisci S, Poletti R, Ricco G, Vecchi G, Venturi G, Ziccardi A, Comba P. Mortality study of asbestos cement workers. *Int Arch Occup Environ Health.* 1994;66:7–11. PubMed PMID: 7927845.
- [34] Hughes JM. Human evidence: lung cancer mortality risk from chrysotile exposure. *Ann Occup Hyg.* 1994;38:555–560. PubMed PMID: 7978978.
- [35] Jakobsson K, Albin M, Hagmar L. Asbestos, cement, and cancer in the right part of the colon. *Occup Environ Med.* 1994;51:95–101. PubMed PMID: 8111470.
- [36] Tarchi M, Orsi D, Comba P, De Santis M, Pirastu R, Battista G, Valani M. Cohort mortality study of rock salt workers in Italy. *Am J Ind Med.* 1994;25:251–256. PubMed PMID: 8147397.
- [37] Straif K, Benbrahim-Tallaa L, Baan R, Grosse Y, Secretan B, El Ghissassi F, Bouvard V, Guha N, Freeman C, Galichet L, Cogliano V. WHO International Agency for Research on Cancer Monograph Working Group. A review of human carcinogens—part C: metals, arsenic, dusts, and fibres. *Lancet Oncol.* 2009;10:453–454. PubMed PMID: 19998521.

- [38] Hughes JM, Weill H. Asbestosis as a precursor of asbestos-related lung cancer: results of a prospective mortality study. *Br J Ind Med*. 1991;48:229–233. PubMed PMID: 2025587.
- [39] Karjalainen A, Pukkala E, Kauppinen T, Partanen T. Incidence of cancer among Finnish patients with asbestos-related pulmonary or pleural fibrosis. *Cancer Causes Control*. 1999;10:51–57. PubMed PMID: 10334642.
- [40] Weiss W. Asbestosis: a marker for the increased risk of lung cancer among workers exposed to asbestos. *Chest*. 1999;115:536–549. PubMed PMID: 10027457.
- [41] Reid A, de Klerk N, Ambrosini GL, Olsen N, Pang SC, Berry G, Musk AW. The effect of asbestosis on lung cancer risk beyond the dose related effect of asbestos alone. *Occup Environ Med*. 2005;62:885–889. PubMed PMID: 16299098.
- [42] Peters GA, Peters BJ. Medical aspects. In: Peters GA, Peters BJ, editors. *Sourcebook on asbestos diseases: medical, legal, and engineering aspects*. New York, London: Garland STPM Press, 1980. pp. B2–B6.
- [43] Speizer FE, Balmes JR. Environmental lung disease. In: Fauci AS, Braunwald E, Kasper DL, Hauser SL, Longo DL, Jameson JL, Loscalzo J, editors. *Harrison's principles of internal medicine*. 17th ed. New York, Chicago, San Francisco, Lisbon, London, Madrid, Mexico City, New Delhi, San Juan, Seoul, Singapore, Sydney, Toronto: McGraw Hill, 2008. pp. 1612–1614.
- [44] Holland JP, Smith DD. Asbestos. In: Greenberg MI, Hamilton RJ, Phillips SD, McCluskey GJ, editors. *Occupational, industrial, and environmental toxicology*. 2nd ed. Philadelphia: Mosby, An Affiliate of Elsevier Science, 2003. pp. 655–658.
- [45] Garcia JG, Griffith DE, Cohen AB, Callahan KS. Alveolar macrophages from patients with asbestos exposure release increased levels of leukotriene B₄. *Am Rev Respir Dis*. 1989;139:1494–1501. PubMed PMID: 254324.
- [46] Kamp DW, Graceffa P, Pryor WA, Weitzman SA. The role of free radicals in asbestos-induced diseases. *Free Radic Biol Med*. 1992;12:293–315. PubMed PMID: 1577332.
- [47] Ding M, Dong Z, Chen F, Pack D, Ma WY, Ye J, Shi X, Castranova V, Vallyathan V. Asbestos induces activator protein-1 transactivation in transgenic mice. *Cancer Res*. 1999;59:1884–1889. PubMed PMID: 10213496.
- [48] Kinnula VL. Oxidant and antioxidant mechanisms of lung disease caused by asbestos fibres. *Eur Respir J* 1999;14:706–16. PubMed PMID: 10543297.
- [49] Li J, Huang B, Shi X, Castranova V, Vallyathan V, Huang C. Involvement of hydrogen peroxide in asbestos-induced NFAT activation. *Mol Cell Biochem*. 2002;234–235:161–168. PubMed PMID: 12162429.

- [50] Xu A, Zhou H, Yu DZ, Hei TK. Mechanisms of the genotoxicity of crocidolite asbestos in mammalian cells: implication from mutation patterns induced by reactive oxygen species. *Environ Health Perspect.* 2002;110:1003–1008. PubMed PMID: 12361925.
- [51] Castranova V. Role of nitric oxide in the progression of pneumoconiosis. *Biochemistry [Mosc].* 2004;69:32–37. PubMed PMID: 14972015.
- [52] Castranova V. Signaling pathways controlling the production of inflammatory mediators in response to crystalline silica exposure: role of reactive oxygen/nitrogen species. *Free Radic Biol Med.* 2004;37:916–925. PubMed PMID: 15336307.
- [53] Fattman CL, Chang LY, Termin TA, Petersen L, Enghild JJ, Oury TD. Enhanced bleomycin-induced pulmonary damage in mice lacking extracellular superoxide dismutase. *Free Radic Biol Med.* 2003;35:763–771. PubMed PMID: 14583340.
- [54] Fattman CL, Tan RJ, Tobolewski JM, Oury TD. Increased sensitivity to asbestos-induced lung injury in mice lacking extracellular superoxide dismutase. *Free Radic Biol Med.* 2006;40:601–6007. PubMed PMID: 16458190.
- [55] Kinnula VL, Hodgson UA, Lakari EK, Tan RJ, Sormunen RT, Soini YM, Kakko SJ, Laitinen TH, Oury TD, Pääkkö PK. Extracellular superoxide dismutase has a highly specific localization in idiopathic pulmonary fibrosis/usual interstitial pneumonia. *Histopathology.* 2006;49:66–74. PubMed PMID: 16842247.
- [56] Weitzman SA, Graceffa P. Asbestos catalyzes hydroxyl and superoxide radical generation from hydrogen peroxide. *Arch Biochem Biophys.* 1984;228:373–376. PubMed PMID: 6320737.
- [57] Halliwell B, Gutteridge JMC. Reactive species as useful biomolecules. In: Halliwell B, Gutteridge JMC, editors. *Free radicals in biology and medicine.* 3rd ed. Oxford, New York: Oxford University Press, 1999. p. 464.
- [58] Kamp DW, Panduri V, Weitzman SA, Chandel N. Asbestos-induced alveolar epithelial cell apoptosis: role of mitochondrial dysfunction caused by iron-derived free radicals. *Mol Cell Biochem.* 2002;234–235:153–160. PubMed PMID: 12162428.
- [59] Panduri V, Weitzman SA, Chandel N, Kamp DW. The mitochondria-regulated death pathway mediates asbestos-induced alveolar epithelial cell apoptosis. *Am J Respir Cell Mol Biol.* 2003;28:241–248. PubMed PMID: 12540492.
- [60] Poser I, Rahman Q, Lohani M, Yadav S, Becker HH, Weiss DG, Schiffmann D, Dopp E. Modulation of genotoxic effects in asbestos-exposed primary human mesothelial cells by radical scavengers, metal chelators and a glutathione precursor. *Mutat Res.* 2004;559:19–27. PubMed PMID: 15066570.
- [61] Thomas G, Ando T, Verma K, Kagan E. Asbestos fibers and interferon-gamma up-regulate nitric oxide production in rat alveolar macrophages. *Am J Respir Cell Mol Biol.* 1994;11:707–715. PubMed PMID: 7524571.

- [62] Quinlan TR, BeruBe KA, Hacker MP, Taatjes DJ, Timblin CR, Goldberg J, Kimberley P, O'Shaughnessy P, Hemenway D, Torino J, Jimenez LA, Mossman BT. Mechanisms of asbestos-induced nitric oxide production by rat alveolar macrophages in inhalation and in vitro models. *Free Radic Biol Med.* 1998;24:778–788. PubMed PMID: 9586808.
- [63] Tanaka S, Choe N, Hemenway DR, Zhu S, Matalon S, Kagan E. Asbestos inhalation induces reactive nitrogen species and nitrotyrosine formation in the lungs and pleura of the rat. *J Clin Invest.* 1998;102:445–454. PubMed PMID: 9664087.
- [64] Aldieri E, Ghigo D, Tomatis M, Prandi L, Fenoglio I, Costamagna C, Pescarmona G, Bosia A, Fubini B. Iron inhibits the nitric oxide synthesis elicited by asbestos in murine macrophages. *Free Radic Biol Med.* 2001;31:412–417. PubMed PMID: 11461780.
- [65] Förstermann U, Kleinert H. Nitric oxide synthase: expression and expressional control of the three isoforms. *Naunyn Schmiedebergs Arch Pharmacol.* 1995;352:351–364. PubMed PMID: 8532063.
- [66] deRojas-Walker T, Tamir S, Ji H, Wishnok JS, Tannenbaum SR. Nitric oxide induces oxidative damage in addition to deamination in macrophage DNA. *Chem Res Toxicol.* 1995;8:473–477. PubMed PMID: 7578935.
- [67] Beckman JS, Koppenol WH. Nitric oxide, superoxide, and peroxynitrite: the good, the bad, and ugly. *Am J Physiol.* 1996;271:C1424–C1437. PubMed PMID: 8944624.
- [68] Wink DA, Mitchell JB. Chemical biology of nitric oxide: insights into regulatory, cytotoxic, and cytoprotective mechanisms of nitric oxide. *Free Radic Biol Med.* 1998;25:434–456. PubMed PMID: 9741580.
- [69] Halliwell B, Gutteridge JMC. The chemistry of free radicals and related "reactive species". In: Halliwell B, Gutteridge JMC, editors. *Free radicals in biology and medicine.* 3rd ed. Oxford, New York: Oxford University Press, 1999. pp. 73–82.
- [70] Chao CC, Park SH, Aust AE. Participation of nitric oxide and iron in the oxidation of DNA in asbestos-treated human lung epithelial cells. *Arch Biochem Biophys.* 1996;326:152–157. PubMed PMID: 8579364.
- [71] Park SH, Aust AE. Participation of iron and nitric oxide in the mutagenicity of asbestos in hgp^{rt-}, gpt⁺ Chinese hamster V79 cells. *Cancer Res.* 1998;58:1144–1148. PubMed PMID: 9515798.
- [72] Dörger M, Allmeling AM, Kiefmann R, Schropp A, Krombach F. Dual role of inducible nitric oxide synthase in acute asbestos-induced lung injury. *Free Radic Biol Med.* 2002;33:491–501. PubMed PMID: 12160931.
- [73] Halliwell B, Gutteridge JMC. Oxidative stress: adaptation, damage, repair and death. In: Halliwell B, Gutteridge JMC, editors. *Free radicals in biology and medicine.* 3rd ed. Oxford, New York: Oxford University Press, 1999. pp. 246–350.

- [74] Halliwell B, Gutteridge JMC. Antioxidant defences. In: Halliwell B, Gutteridge JMC, editors. *Free radicals in biology and medicine*. 3rd ed. Oxford, New York: Oxford University Press, 1999. pp. 105–243.
- [75] McCord JM, Fridovich I. Superoxide dismutase. An enzymic function for erythrocyte hemocuprein (hemocuprein). *J Biol Chem*. 1969;244:6049–6055. PubMed PMID: 5389100.
- [76] Quan F, Korneluk RG, Tropak MB, Gravel RA. Isolation and characterization of the human catalase gene. *Nucleic Acids Res*. 1986;14:5321–5335. PubMed PMID: 3755525.
- [77] Hendrickson DJ, Fisher JH, Jones C, Ho YS. Regional localization of human extracellular superoxide dismutase gene to 4pter-q21. *Genomics*. 1990;8:736–738. PubMed PMID: 2276747.
- [78] Church SL, Grant JW, Meese EU, Trent JM. Sublocalization of the gene encoding manganese superoxide dismutase [MnSOD/SOD2] to 6q25 by fluorescence in situ hybridization and somatic cell hybrid mapping. *Genomics*. 1992;14:823–825. PubMed PMID: 1427917.
- [79] Folz RJ, Abushama AM, Suliman HB. Extracellular superoxide dismutase in the airways of transgenic mice reduces inflammation and attenuates lung toxicity following hyperoxia. *J Clin Invest*. 1999;103:1055–1066. PubMed PMID: 10194479.
- [80] Forsberg L, Lyrenäs L, de Faire U, Morgenstern R. A common functional C-T substitution polymorphism in the promoter region of the human catalase gene influences transcription factor binding, reporter gene transcription and is correlated to blood catalase levels. *Free Radic Biol Med*. 2001;30:500–505. PubMed PMID: 11182520.
- [81] Bowler RP, Nicks M, Warnick K, Crapo JD. Role of extracellular superoxide dismutase in bleomycin-induced pulmonary fibrosis. *Am J Physiol Lung Cell Mol Physiol*. 2002;282:L719-L726. PubMed PMID: 11880297.
- [82] Zelko IN, Mariani TJ, Folz RJ. Superoxide dismutase multigene family: a comparison of the CuZn-SOD (SOD1), Mn-SOD (SOD2), and EC-SOD (SOD3) gene structures, evolution, and expression. *Free Radic Biol Med*. 2002;33:337–349. PubMed PMID: 12126755.
- [83] Carlsson LM, Jonsson J, Edlund T, Marklund SL. Mice lacking extracellular superoxide dismutase are more sensitive to hyperoxia. *Proc Natl Acad Sci U S A*. 1995;92:6264–6268. PubMed PMID: 7603981.
- [84] Singhal SS, Saxena M, Ahmad H, Awasthi S, Haque AK, Awasthi YC. Glutathione S-transferases of human lung: characterization and evaluation of the protective role of the alpha-class isozymes against lipid peroxidation. *Arch Biochem Biophys*. 1992;299:232–241. PubMed PMID: 1444461.
- [85] Hubbard NE, Erickson KL. Role of 5'-lipoygenase metabolites in the activation of peritoneal macrophages for tumoricidal function. *Cell Immuno*. 1995;160:115–122. PubMed PMID: 7842477.

- [86] Kelsey KT, Nelson HH, Wiencke JK, Smith CM, Levin S. The glutathione S-transferase theta and mu deletion polymorphisms in asbestosis. *Am J Ind Med.* 1997;31:274–279. PubMed PMID: 9055949.
- [87] Ketterer B. A bird's eye view of the glutathione transferase field. *Chem Biol Interact.* 2001;138:27–42. PubMed PMID: 11640913.
- [88] Hayes JD, Pulford DJ. The glutathione S-transferase supergene family: regulation of GST and the contribution of the isoenzymes to cancer chemoprotection and drug resistance. *Crit Rev Biochem Mol Biol.* 1995;30:445–600. PubMed PMID: 8770536.
- [89] Timbrell J. Phase 2 reactions. Conjugation. In: Timbrell J, editors. *Principles of biochemical toxicology.* London, Philadelphia: Taylor & Francis, 2003. pp. 95–101.
- [90] Hayes JD, Strange RC. Glutathione S-transferase polymorphisms and their biological consequences. *Pharmacology.* 2000;61:154–166. PubMed PMID: 10971201.
- [91] Hayes JD, Flanagan JU, Jowsey IR. Glutathione transferases. *Annu Rev Pharmacol Toxicol.* 2005;45:51–88. PubMed PMID: 15822171.
- [92] Blackburn AC, Woollatt E, Sutherland GR, Board PG. Characterization and chromosome location of the gene GSTZ1 encoding the human Zeta class glutathione transferase and maleylacetoacetate isomerase. *Cytogenet Cell Genet.* 1998;83:109–114. PubMed PMID: 9925947.
- [93] Anttila S, Hirvonen A, Vainio H, Husgafvel-Pursiainen K, Hayes JD, Ketterer B. Immunohistochemical localization of glutathione S-transferases in human lung. *Cancer Res.* 1993;53:5643–5648. PubMed PMID: 8242618.
- [94] Stücker I, Boffetta P, Antilla S, Benhamou S, Hirvonen A, London S, Taioli E. Lack of interaction between asbestos exposure and glutathione S-transferase M1 and T1 genotypes in lung carcinogenesis. *Cancer Epidemiol Biomarkers Prev.* 2001;10:1253–8. PubMed PMID: 11751442.
- [95] Hirvonen A, Tuimala J, Ollikainen T, Linnainmaa K, Kinnula V. Manganese superoxide dismutase genotypes and asbestos-associated pulmonary disorders. *Cancer Lett.* 2002;178:71–74. PubMed PMID: 11849743.
- [96] Shimoda-Matsubayashi S, Matsumine H, Kobayashi T, Nakagawa-Hattori Y, Shimizu Y, Mizuno Y. Structural dimorphism in the mitochondrial targeting sequence in the human manganese superoxide dismutase gene. A predictive evidence for conformational change to influence mitochondrial transport and a study of allelic association in Parkinson's disease. *Biochem Biophys Res Commun.* 1996;226:561–565. PubMed PMID: 8806673.
- [97] Rosenblum JS, Gilula NB, Lerner RA. On signal sequence polymorphisms and diseases of distribution. *Proc Natl Acad Sci U S A.* 1996;93:4471–3. PubMed PMID: 8633092.
- [98] Ali-Osman F, Akande O, Antoun G, Mao JX, Buolamwini J. Molecular cloning, characterization, and expression in *Escherichia coli* of full-length cDNAs of three human

- glutathione S-transferase P1 gene variants – evidence for differential conjugation capacity of the encoded proteins. *J Biol Chem.* 1997;272:10004–10012. PubMed PMID: 9092542.
- [99] Tatemichi M, Sawa T, Gilibert I, Tazawa H, Katoh T, Ohshima H. Increased risk of intestinal type of gastric adenocarcinoma in Japanese women associated with long forms of CCTTT pentanucleotide repeat in the inducible nitric oxide synthase promoter. *Cancer Lett.* 2005;217:197–202. PubMed PMID: 15617837.
- [100] Kinnula VL, Lehtonen S, Koistinen P, Kakko S, Savolainen M, Kere J, Ollikainen V, Laitinen T. Two functional variants of the superoxide dismutase genes in Finnish families with asthma. *Thorax.* 2004;59:116–119. PubMed PMID: 14760150.
- [101] Gao F, Kinnula VL, Myllärniemi M, Oury TD. Extracellular superoxide dismutase in pulmonary fibrosis. *Antioxid Redox Signal.* 2008;10:343–354. PubMed PMID: 17999630.
- [102] Kliment CR, Tobolewski JM, Manni ML, Tan RJ, Enghild J, Oury TD. Extracellular superoxide dismutase protects against matrix degradation of heparan sulfate in the lung. *Antioxid Redox Signal.* 2008;10:261–268. PubMed PMID: 17961072.
- [103] Folz RJ, Peno-Green L, Crapo JD. Identification of a homozygous missense mutation [Arg to Gly] in the critical binding region of the human EC-SOD gene [SOD3] and its association with dramatically increased serum enzyme levels. *Hum Mol Genet.* 1994;3:2251–2254. PubMed PMID: 7881430.
- [104] Sandström J, Nilsson P, Karlsson K, Marklund SL. 10-fold increase in human plasma extracellular superoxide dismutase content caused by a mutation in heparin-binding domain. *J Biol Chem.* 1994;269:19163–19166. PubMed PMID: 8034674.
- [105] Yamada H, Yamada Y, Adachi T, Goto H, Ogasawara N, Futenma A, Kitano M, Hirano K, Kato K. Molecular analysis of extracellular-superoxide dismutase gene associated with high level in serum. *Jpn J Hum Genet.* 1995;40:177–184. PubMed PMID: 7662997.
- [106] Jiang Z, Akey JM, Shi J, Xiong M, Wang Y, Shen Y, Xu X, Chen H, Wu H, Xiao J, Lu D, Huang W, Jin L. A polymorphism in the promoter region of catalase is associated with blood pressure levels. *Hum Genet.* 2001;109:95–98. PubMed PMID: 11479740.
- [107] Casp CB, She JX, McCormack WT. Genetic association of the catalase gene [CAT] with vitiligo susceptibility. *Pigment Cell Res.* 2002;15:62–66. PubMed PMID: 11837458.
- [108] Nadif R, Mintz M, Jedlicka A, Bertrand JP, Kleeberger SR, Kauffmann F. Association of CAT polymorphisms with catalase activity and exposure to environmental oxidative stimuli. *Free Radic Res.* 2005;39:1345–1350. PubMed PMID: 16298864.
- [109] Zhou XF, Cui J, DeStefano AL, Chazaro I, Farrer LA, Manolis AJ, Gavras H, Baldwin CT. Polymorphisms in the promoter region of catalase gene and essential hypertension. *Dis Markers.* 2005;21:3–7. PubMed PMID: 15735318.
- [110] Ahn J, Nowell S, McCann SE, Yu J, Carter L, Lang NP, Kadlubar FF, Ratnasinghe LD, Ambrosone CB. Associations between catalase phenotype and genotype: modification

- by epidemiologic factors. *Cancer Epidemiol Biomarkers Prev.* 2006;15:1217–1222. PubMed PMID: 16775184.
- [111] Bastaki M, Huen K, Manzanillo P, Chande N, Chen C, Balmes JR, Tager IB, Holland N. Genotype-activity relationship for Mn-superoxide dismutase, glutathione peroxidase 1 and catalase in humans. *Pharmacogenet Genom.* 2006;16:279–286. PubMed PMID: 16538174.
- [112] Johansson AS, Stenberg G, Widersten M, Mannervik B. Structure-activity relationships and thermal stability of human glutathione transferase P1-1 governed by the H-site residue 105. *J Mol Bio.* 1998;278:687–698. PubMed PMID: 9600848.
- [113] Watson MA, Stewart RK, Smith GB, Massey TE, Bell DA. Human glutathione S-transferase P1 polymorphisms: relationship to lung tissue enzyme activity and population frequency distribution. *Carcinogenesis.* 1998;19:275–280. PubMed PMID: 9498276.
- [114] Xu W, Liu L, Emson PC, Harrington CR, Charles IG. Evolution of a homopurine-homopyrimidine pentanucleotide repeat sequence upstream of the human inducible nitric oxide synthase gene. *Gene.* 1997;204:165–170. PubMed PMID: 9434180.
- [115] Warpeha KM, Xu W, Liu L, Charles IG, Patterson CC, Ah-Fat F, Harding S, Hart PM, Chakravarthy U, Hughes AE. Genotyping and functional analysis of a polymorphic [CCTTT]_n repeat of NOS2A in diabetic retinopathy. *FASEB J.* 1999;13:1825–1832. PubMed PMID: 10506586.
- [116] Franko A, Dolžan V, Arnerić N, Dodič-Fikfak M. The influence of genetic polymorphisms of GSTP1 on the development of asbestosis. *J Occup Environ Med.* 2008;50:7–12. PubMed PMID: 18188076; doi: 10.1097/JOM.0b013e31815cbab5.
- [117] Franko A, Dolžan V, Arnerić N, Dodič-Fikfak M. Asbestosis and catalase genetic polymorphism. *Arh Hig Rada Toksikol.* 2008;59:233–240. PubMed PMID: 19064360; doi: 10.2478/10004-1254-59-2008-1907.
- [118] Franko A, Dodič-Fikfak M, Arnerić N, Dolžan V. Manganese and extracellular superoxide dismutase polymorphisms and risk for asbestosis. *J Biomed Biotechnol.* 2009;2009:493083. PubMed PMID: 19636420; doi: 10.1155/2009/493083.
- [119] Franko A, Dodič-Fikfak M, Arnerić N, Dolžan V. Inducible nitric oxide synthase genetic polymorphism and risk of asbestosis. *J Biomed Biotechnol.* 2011;2011:685870. PubMed PMID: 21660141; doi: 10.1155/2011/685870.
- [120] Ferris BG. Epidemiology standardization project [American Thoracic Society]. *Am Rev Respir Dis.* 1978;118:1–120. PubMed PMID: 742764.
- [121] Asbestos, asbestosis, and cancer: the Helsinki criteria for diagnosis and attribution. Consensus report. *Scand J Work Environ Health.* 1997;23:311–316. PubMed PMID: 9322824.

- [122] American Thoracic Society. Diagnosis and initial management of nonmalignant diseases related to asbestos. *Am J Respir Crit Care Med.* 2004;170:691–751. PubMed PMID: 15355871.
- [123] Yoshie Y, Ohshima H. Nitric oxide synergistically enhances DNA strand breakage induced by polyhydroxyaromatic compounds, but inhibits that induced by the Fenton reaction. *Arch Biochem Biophys.* 1997;342:13–21. PubMed PMID: 9185609.
- [124] Haberstroh K, Heigold S, Bauer G. Transformed cell-derived reactive oxygen species support and inhibit nitric oxide-mediated apoptosis induction. *Int J Oncol.* 2002;21:145–151. PubMed PMID: 12063561.
- [125] Jackson JH, Schraufstatter IU, Hyslop PA, Vosbeck K, Sauerheber R, Weitzman SA, Cochrane CG. Role of oxidants in DNA damage. Hydroxyl radical mediates the synergistic DNA damaging effects of asbestos and cigarette smoke. *J Clin Invest.* 1987;80:1090–1095. PubMed PMID: 2821073.
- [126] Valavanidis A, Vlachogianni T, Fiotakis K. Tobacco smoke: involvement of reactive oxygen species and stable free radicals in mechanisms of oxidative damage, carcinogenesis and synergistic effects with other respirable particles. *Int J Environ Res Public Health.* 2009;6:445–462. PubMed PMID: 19440393; doi: 10.3390/ijerph6020445.
- [127] Valavanidis A, Balomenou H, Macropoulou I, Zarodimos I. A study of the synergistic interaction of asbestos fibers with cigarette tar extracts for the generation of hydroxyl radicals in aqueous buffer solution. *Free Radic Biol Med.* 1996;20:853–858. PubMed PMID: 8728034.
- [128] van der Vliet A, Cross CE. Oxidants, nitrosants, and the lung. *Am J Med.* 2000;109:398–421. PubMed PMID: 11020397.
- [129] Hasnis E, Bar-Shai M, Burbea Z, Reznick AZ. Mechanisms underlying cigarette smoke-induced NF-kappaB activation in human lymphocytes: the role of reactive nitrogen species. *J Physiol Pharmacol.* 2007;58:275–287. PubMed PMID: 18204137.
- [130] Figueroa JD, Koutros S, Colt JS, Kogevinas M, Garcia-Closas M, Real FX, Friesen MC, Baris D, Stewart P, Schwenn M, Johnson A, Karagas MR, Armenti KR, Moore LE, Schned A, Lenz P, Prokunina-Olsson L, Banday AR, Paquin A, Ylaya K, Chung JY, Hewitt SM, Nickerson ML, Tardón A, Serra C, Carrato A, García-Closas R, Lloreta J, Malats N, Fraumeni JF Jr, Chanock SJ, Chatterjee N, Rothman N, Silverman DT. Modification of Occupational Exposures on Bladder Cancer Risk by Common Genetic Polymorphisms. *J Natl Cancer Inst.* 2015;107(11). pii: djv223. PubMed PMID: 26374428; doi: 10.1093/jnci/djv223.
- [131] Huo X, Chen D, He Y, Zhu W, Zhou W, Zhang J. Bisphenol-A and female infertility: a possible role of gene-environment interactions. *Int J Environ Res Public Health.* 2015;12:11101–11116. PubMed PMID: 26371021; doi: 10.3390/ijerph120911101.
- [132] Melkonian SC, Daniel CR, Ye Y, Tannir NM, Karam JA, Matin SF, Wood CG, Wu X. Gene-environment interaction of genome-wide association study-identified susceptibility

- loci and meat-cooking mutagens in the etiology of renal cell carcinoma. *Cancer*. 2016;122:108–115. PubMed PMID: 26551148; doi: 10.1002/cncr.29543.
- [133] Reynolds CA, Gatz M, Christensen K, Christiansen L, Dahl Aslan AK, Kaprio J, Korhonen T, Kremen WS, Krueger R, McGue M, Neiderhiser JM, Pedersen NL, IGEMS consortium. Gene-environment interplay in physical, psychological, and cognitive domains in mid to late adulthood: is APOE a variability gene? *Behav Genet*. 2016;46:4–19. PubMed PMID: 26538244; doi: 10.1007/s10519-015-9761-3.
- [134] Tunesi S, Ferrante D, Mirabelli D, Andorno S, Betti M, Fiorito G, Guarrera S, Casalone E, Neri M, Ugolini D, Bonassi S, Matullo G, Dianzani I, Magnani C. Gene-asbestos interaction in malignant pleural mesothelioma susceptibility. *Carcinogenesis*. 2015;36:1129–1135. PubMed PMID: 26139392; doi: 10.1136/oemed-2015-102803.
- [135] Malhotra J, Sartori S, Brennan P, Zaridze D, Szeszenia-Dabrowska N, Świątkowska B, Rudnai P, Lissowska J, Fabianova E, Mates D, Bencko V, Gaborieau V, Stücker I, Foretova L, Janout V, Boffetta P. Effect of occupational exposures on lung cancer susceptibility: a study of gene-environment interaction analysis. *Cancer Epidemiol Biomarkers Prev*. 2015;24:570–579. PubMed PMID: 25583949; doi: 10.1158/1055-9965.
- [136] Liu CY, Stücker I, Chen C, Goodman G, McHugh MK, D'Amelio AM Jr, Etzel CJ, Li S, Lin X, Christiani DC. Genome-wide gene-asbestos exposure interaction association study identifies a common susceptibility variant on 22q13.31 associated with lung cancer risk. *Cancer Epidemiol Biomarkers Prev*. 2015;24:1564–1573. PubMed PMID: 26199339; doi: 10.1158/1055-9965.EPI-15-0021.
- [137] Schulte PA, Whittaker C, Curran CP. Considerations for using genetic and epigenetic information in occupational health risk assessment and standard setting. *J Occup Environ Hyg*. 2015;12 Suppl 1:S69-S81. PubMed PMID: 26583908; doi: 10.1080/15459624.2015.1060323.