

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Salmonella Fimbriae: What is the Clue to Their Hairdo?

Karine Dufresne and France Daigle

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67189>

Abstract

Fimbriae are important virulence factors for *Salmonella* pathogenesis. They mediate adhesion to host cells (including plants), food, stainless steel and much more. The fimbrial systems are organised in gene clusters of four to fifteen genes that code for structural, assembly and regulatory proteins. There are three kinds of fimbriae depending on their mode of assembly. The chaperone/usher (CU) fimbriae use a dedicated chaperone and usher protein to coordinate the subunit biogenesis on the cell surface. The curli fimbriae are assembled by nucleation/precipitation pathway. The type IV fimbria assembly requires a transmembrane apparatus and ATP to energise the reaction. Several fimbriae are conserved among *Salmonella* serovars, while some are present in a limited set or only specific serovars. Expression and regulation of fimbrial genes are not well understood, and most *Salmonella* fimbriae are poorly expressed during in vitro culture, which further complicates research concerning their regulation and role during infection. However, *Salmonella fim* gene cluster, coding for type-1 fimbriae, was widely studied and presents its own set of regulators. Investigating fimbrial distribution, expression and regulation will further elucidate their roles in bacterial pathogenesis and host specificity. Furthermore, fimbriae are important for developing efficient diagnostic tests and antimicrobial strategies against *Salmonella*.

Keywords: fimbriae, adhesion, chaperone/usher, curli, type IV fimbria, fimbriome, *fim*

1. Introduction

Multiple virulence factors are implicated in *Salmonella* pathogenesis. These factors include type 3 secretion systems (T3SS) encoded in *Salmonella* Pathogenicity Islands (SPI)-1 and SPI-2, other SPIs, flagella, capsule, plasmids and adhesion systems [1, 2]. Among those factors, fimbriae represent a major player in pathogenesis and a source of diversity for *Salmonella*

serovars. Fimbriae are the most common adhesion systems and are differentially expressed and found in a specific pattern among each serovar [3, 4].

Historically, the first observation of fimbriae was described in 1901 in *Bacillus anthracis* by Hinterberger and Reitman which hypothesised that the filaments were implicated in nutrients acquisition [5]. Then, in 1949, Anderson suggested that the filaments were artefacts due to sample preparation for electron microscopy [6]. However, many other studies contradicted Anderson and confirmed the presence of non-flagellar appendages on the bacterial surface. In 1950, Houwink and Van Iterson observed the appendages and described them as shorter and more rigid filaments than the flagella from *Escherichia coli* and suggested that the fibres were implicated in attachment to surface [7]. The name fimbria (Latin word for fibres) was suggested in 1955 by Duguid et al. to describe the filamentous structures [7, 8]. The term fimbria is preferable to use to describe non-flagellar filaments than pili, which is used to designate structures implicated in conjugation [9, 10]. In 1966, Duguid et al. classified fimbriae in seven types (types 1–6 and F) according to the morphology and haemagglutination patterns. However, another classification, based on serology, better predicted genetic relatedness of fimbrial antigens. Nowadays, fimbriae are designated by the mode of assembly of the fibril [8].

A specific fimbrial gene cluster (FGC) encodes for the structural, assembly and sometime regulatory proteins required for the production of the filamentous adhesive appendage on the bacterial surface. FGCs are usually composed of four to fifteen genes [10, 11]. An average of 12 FGCs by strains was observed in *S. enterica*. Despite that all *Salmonella* genome harbours multiple FGCs, very few are characterised so far. Most fimbriae are poorly expressed under laboratory conditions, and the functional redundancy complicates their studies [10]. However, fimbriae are implicated during infection and in a variety of other roles, like biofilm formation, seroconversion, haemagglutination, cellular invasion and macrophage interactions [2, 7, 12–16]. In mice model, *S. Typhimurium* fimbriae demonstrate a role in intestinal cells attachment, caecum colonisation and persistence in gut [17–19]. Moreover, fimbriae are important determinants of host adaptation by *Salmonella* [20].

In this chapter, an overview of *Salmonella* fimbriae is presented. First, the three pathways for fimbrial biogenesis (CU, precipitation/nucleation, type IV fimbriae) are described. Second, the distribution of fimbrial genes among *Salmonella* subspecies and serovars is presented. Third, the regulation of fimbrial genes is described and *fim* FGC regulation is detailed. Finally, the use of fimbriae as diagnostic and therapeutic tools is discussed.

2. Fimbrial biogenesis pathways

Three pathways for fimbrial assembly exist in *Salmonella*, the chaperone/usher (CU), the nucleation/precipitation and the type IV pathway [21]. Fimbriae of the CU pathway employ dedicated chaperones and ushers for the fimbrial assembly. The nucleation/precipitation pathway forms an aggregative fibre by precipitation of the subunits in the presence of the nucleator in the extracellular environment. Finally, the type IV fimbrial pathway uses complex machinery

for the fimbriae formation and needs ATP to drive the reaction of assembly. Furthermore, the type IV fimbriae can retract and reverse its assembly [21].

The three pathways produce quite different fimbriae. CU fimbriae have the classic fimbrial shape with the repetition of major subunits emerging from the usher inserted in the outer membrane. The major subunits can be accompanied by minor subunits and/or adhesins [8]. The fimbriae produced by the nucleation/precipitation pathway have an aggregated shape, due to the precipitation of major subunits together. This kind of fimbriae is highly stable and hardly depolymerised [22]. The type IV fimbriae anchor in the inner membrane and are prolonged by the repetition of the major subunit (pilin) through the periplasm and the outer membrane reaching the extracellular medium [23]. Here, the three fimbrial assembly mechanisms will be detailed.

2.1. Chaperone/usher pathway

The CU fimbriae represent the largest and most diversified class of adhesion systems [24, 25]. Multiple CU fimbriae are present in *Salmonella* suggesting a functional redundancy [23, 26]. The assembly is characterised by an interaction between the subunits, a periplasmic chaperone and an outer membrane usher in order to form a mature fibre (**Figure 1**) [27]. Each fimbria produced by this pathway has its own unique and specific chaperone and usher [11]. Usher sequence is a good discrimination tool and is used to subdivide the CU fimbriae into six phylogenetic clades (α , κ , π , σ , γ , β) [10, 26].




Figure 1. Chaperone/usher pathway. The subunit proteins are synthesised in the cytoplasm and translocated through the periplasm via SecYEG implying also SecDF/YajC inner membrane proteins. When the signal peptide is cleaved from the subunit, the chaperone protein complements the missing strand of the subunit in a process called donor strand complementation. The energy from the folding of the subunit is preserved by the chaperone. The chaperone drives the subunit to the usher and exchanges the donor strand. The subunit is then translocated by the usher to the extracellular medium and added to other subunits to form the fibril. IM = inner membrane; OM = outer membrane.

The biogenesis of the CU fimbriae begins with the production of the subunits in the cytoplasm and their export through the inner membrane by the general secretory pathway (GSP) [23, 27, 28]. It consists in a post-translational translocation implying the SecYEG complex and SecDF/YajC proteins. When the pre-protein is produced, it can be targeted directly to the accessory factor SecA or transported to SecA by the general chaperone SecB. Then, SecA catalyses the hydrolysis of ATP to energise the translocation through SecYEG. Use of ATP, in combination with proton-motive force, triggers the transport of the pre-protein to the periplasm. During the translocation across the inner membrane, the N-terminal signal peptide is cleaved by periplasmic peptidases [27, 29]. To prevent early folding of the subunits, the fimbrial chaperone instantly forms a complex with the translocated subunit in the periplasm [30].

Fimbrial chaperone shares conserved structural features with the general periplasmic chaperones [30]. They are formed of two β -sheet domains oriented to produce an L-shaped molecule and together form a β -barrel. Each domain has an immunoglobulin-like fold and is composed of seven primary β -strands [30–32]. Hydrophobic residues are alternated in the seven strands, facing the internal part of the barrel. These residues form the hydrophobic core of the domain that is implicated in the binding of the subunit. The fimbrial chaperones have an extended loop that lies at the extremity of one arm of the L-shaped molecule. This loop contains a conserved motif that is involved in the complex formation between the chaperone and subunits [30]. The subunit and the chaperone have a similar structure, but the subunit is missing the seventh β -strand of the C-terminal extremity [28]. The chaperone transfers the missing β -strand to the subunit to complete its structure: this mechanism is called the donor strand complementation [25]. The chaperone preserves the folding energy of the subunit to drive the last steps of the assembly due to lack of energy source (ATP) in the periplasmic space [33]. The chaperone also prevents premature fimbrial formation in the periplasm and primes the assembly through the usher [30, 34].

Then, the uncapping of the chaperone by the usher exposes the interactive surface of the subunit to the outer membrane usher and assembly of subunits at the surface can occur [33]. The transfer of the subunit from the chaperone to the usher happens very rapidly *in vivo*. In the absence of the usher *in vitro*, only a slow and inefficient assembly was observed. This suggests that the uncapping of the chaperone is important for the efficiency of mature fimbriae assembly [28, 30]. An interaction between the usher and the subunit and also between the usher and the chaperone is required [31]. This triangular interaction is important for the usher to discriminate subunit-loaded from unloaded chaperone [33]. Fimbrial usher forms a ring in the outer membrane with a transient twin-pore of 2–3 nm diameter to allow passage of subunits to the extracellular environment [35]. The usher catalyses fimbrial polymerisation by involving donor strand exchange where the N-terminal sequence of the subunit is replaced by a short sequence of the last subunit in the polymerised fibril with a zip-in-zip-out mechanism [33]. This step is triggered in part by the chaperone required for the strand exchange between the new subunit and the forming fimbria. The quaternary structure of the subunit is achieved when the protein passes through the pore. The final morphology and structure (rigid or flexible), the length (1–3 μm) and width (2–10 nm)

of the fibre of the CU fimbriae depend on the subunits composition and the interactions between subunits [10, 33].

2.2. Nucleation/precipitation pathway

Curli fimbriae were initially discovered in *Escherichia coli* and are very conserved among the *Enterobacteriaceae* family, compared to any other types of FGC. The amyloid fibrils are particularly known for their role in biofilm formation and its recognition by the immune system [36]. The FGC for curli is named *csg* (curli subunit gene) for *E. coli* and *agf* (thin aggregative fimbriae) for *Salmonella*, but the term *csg* is now commonly used for *Salmonella*. Curli formation depends on two divergent operons, *csgBAC* and *csgDEFG*. The *csgBAC* genes encode for CsgA, the major subunit, CsgB, the nucleator, and CsgC, an oxidoreductase of unknown function. The *csgDEFG* genes encode for the transcription regulator of the operon (CsgD) and for the assembly proteins located in the periplasm (CsgE) or in the outer membrane (CsgG and CsgF) [37].

The curli assembly mechanism is characterised by the exportation of the subunits and their precipitation to each other in the presence of a nucleator that fixes the fibril on the bacterial surface. Exportation of curli proteins also uses the GSP to pass through the inner membrane to the periplasm. Then, the CsgA and CsgB proteins are secreted by the lipoprotein CsgG. CsgG is composed of nine anticodon-binding domain-like units that form a 36-stranded β -barrel complex that is inserted in the outer membrane. CsgG forms a pore in the outer membrane that permits the passage of the subunits and the nucleator. CsgG is accompanied by the accessory proteins CsgE and CsgF. CsgE is a specificity factor that forms a nonameric adaptor that binds to CsgG and closes the periplasmic space. The presence of CsgE optimises the uptake of CsgA by CsgG and translocation of CsgA [38]. CsgF helps the nucleation activity of CsgB. It was suggested that CsgF has a role in specific localisation and/or chaperoning of the nucleator, so CsgB will reach its full activity. Moreover, CsgF depends on CsgG and CsgE for its stability [39].

Once at the bacterial surface, the nucleator polymerises the subunits together into thin aggregative fimbriae (fibrils). This process happens only in the extracellular environment and requires the presence of the nucleator CsgB to polymerise CsgA into a filament. CsgA proteins fold into an insoluble cross β -sheet molecules [26]. CsgB anchors the curli fimbriae on the surface of the bacterial cell (**Figure 2**). In *E. coli*, it was observed that CsgB, in addition to its role of nucleator, is also part of the fimbriae with the CsgA subunits. A structurally different fibril made of CsgB subunits can be formed in the absence of CsgA [40]. CsgA and CsgB share 30% of sequence identity and have the same predicted length [37]. In *E. coli*, interbacterial complementation between a nucleator mutant and a subunit mutant is possible. However, in *Salmonella*, this complementation cannot happen, suggesting that the curli fimbriae are different in their nucleation process. However, the interbacterial complementation was observed in *Salmonella* when a lipopolysaccharide O-antigen mutant was used [41]. The nucleation/precipitation pathway is still poorly understood, and research is actually performed on the different aspects of the curli fimbrial formation.




Figure 2. Nucleation/precipitation pathway. The subunit CsgA is synthesised in the cytoplasm and translocated by the GSP. CsgA passes through the periplasm and is translocated in the extracellular medium by CsgG, helped by CsgE. The nucleator CsgB is also translocated by CsgG and supported by CsgF for its stability on the bacterial surface. When CsgA is in the presence of the nucleator in the extracellular environment, the subunits precipitate in an aggregated fibril. CsgC is an oxidoreductase, but its specific role is still undiscovered. IM, inner membrane; OM, outer membrane.

2.3. Type IV fimbriae

Type IV fimbriae are usually from 1 to 5 μm long and are composed of repeated subunits of a single pilin. Type IV fimbria is subdivided into two groups based on homology of the major subunits: type IVa and type IVb fimbriae [26]. The difference between the two types is in the length of the peptide sequence and the mature major pilin sequence. Specific mechanism of assembly of type IVb fimbriae from *Salmonella* has not been characterised yet [42].

Type IV fimbriae pathway has the most complex machinery. They form an apparatus, composed of various proteins, that goes through the inner and outer membranes allowing the anchor of the fibre and energy accessibility for fimbrial assembly. The gene cluster also encodes numerous proteins with diverse functions, as the fibril is not only assembled but also disassembled. Type IV fimbriae are frequently compared to the type II secretion system (T2SS) which possesses similar structure and mechanism of assembly. Type IV fimbriae are implicated in adherence and twitching motility [11].

Type IV fimbriae are present in a variety of organisms including human pathogens such as *Neisseria gonorrhoeae*, *Neisseria meningitidis*, *Pseudomonas aeruginosa* and *Vibrio cholerae*. For *Salmonella*, they are found in *S. bongori*, *S. enterica* serovars Heidelberg, Paratyphi B and Typhi [42]. *S. bongori* type IV fimbria is encoded by the *sbe* operon that remains uncharacterised and is located on a plasmid, as well as in *S. Paratyphi B*, while the type IVb gene cluster is located on the chromosome of *S. Heidelberg* and *S. Typhi* [26].

For *S. Typhi*, the PilS subunits are produced in the cytoplasm and translocated to the periplasm by the GSP. In the periplasm, the N-terminal sequence of PilS is cleaved by PilU, a prepilin pep-

tidase [23]. The mature pilins are then anchored to the inner membrane on platform proteins and linked together into a fibril (**Figure 3**) [26, 43]. The N-terminal domain of the mature subunits is highly hydrophobic, which permits the PilS proteins to group into a helical structure [22, 42]. The pilins are added one by one, but at three sites simultaneously, each corresponding to a strand to form a three-helix bundle [44]. An ATPase inserted in the inner membrane supplies the energy required for the assembly of the type IV fimbriae. The secretin proteins are inserted in the outer membrane and form a channel that permits the passage of the intact pilus through the bacterial surface [26]. These proteins form complexes that are then assembled in a cage-like final structure [44]. Other proteins are also involved in the assembly/disassembly mechanisms, such as another ATPase dedicated for the disassembly of the fimbriae, lipoproteins of the secretin complex (pilotins), inner membrane proteins or gene products involved in peptidoglycan remodelling to permit the passage of the fibril through the periplasm [22, 44, 45]. This assembly pathway is less understood and requires further investigations [44].




Figure 3. Type IV pathway. The pre-pilins are transported and translocated through the inner membrane (IM) to the periplasm by GSP. A peptidase cleaves the signal peptide of the pre-pilin, and the pilin can be assembled on the platform proteins. An ATPase triggers the reaction. The pilins form a three-helix structure that passes through the outer membrane (OM) by a secretin supported by pilotin. The type IV fimbriae can also retract depending on the environmental conditions.

3. *Salmonella* fimbriome

Each fimbrial pathway described above is present in *Salmonella* creating a great element of genetic diversity. CU fimbriae are the most common fimbriae detected in the *Salmonella* genome. Curli (*csg*) is found in all *Salmonella* genome, whereas only a few serovars have the type IV fimbriae. There are 38 unique FGCs identified so far in 111 sequenced genomes from 34 different serovars (**Table 1**) [46, 47]. Each serovar has its own repertoire of FGCs, but there are seven FGCs that are highly conserved in most *Salmonella* strains forming the core of *Salmonella* FGCs. Most of the FGCs are sporadic or found only in a few strains constituting the signature of each serovar.

Fimbriae	CU clade	Prevalence	Distribution	Fimbriae	CU clade	Prevalence	Distribution
<i>bcf</i>	γ 1	Core	Absent in IV	<i>sdj</i>	γ 4	Sporadic	IIIb <i>diarizonae</i>
<i>csg</i>	curli	Core	All <i>Salmonella</i>	<i>sdk/sfi</i>	π	Sporadic	IIIb, VI
<i>fim</i>	γ 1	Core	Absent in <i>bongori</i>	<i>sdl</i>	π	Sporadic	IIIb <i>diarizonae</i>
<i>lpf</i>	γ 1	Conserved	Absent in ID	<i>sef</i>	γ 3	Sporadic	IB, D (pseudo)
<i>mrk</i>	γ 4	Sporadic	Only in Montevideo	<i>sib</i>	β	Sporadic	VI <i>indica</i>
<i>pef</i>	κ	Sporadic	Only in IA, IC and <i>bongori</i>	<i>fae/skf</i>	κ	Sporadic	IB, IE
<i>peg</i>	γ 4	Conserved	IB, IC, IIIa, VI, <i>bongori</i>	<i>ssf</i>	γ 4	Sporadic	II <i>salamae</i>
<i>peh</i>	γ 4	Sporadic	Only in Montevideo	<i>sta</i>	γ 4	Sporadic	ID
<i>pil</i>		Sporadic	Type IV; ID, IE, <i>bongori</i>	<i>stb</i>	γ 4	Core	I, II, IIIb;
<i>saf</i>	γ 3	Conserved	ssp. I	<i>stc</i>	γ 4	Conserved	IA, IB, ID
<i>sba</i>	γ 4	Sporadic	<i>bongori</i>	<i>std</i>	π	Core	II, IIIa, missing in Gallinarum
<i>sbb/sbf</i>	π	Sporadic	<i>bongori</i>	<i>ste</i>	π	Conserved	Missing in IA, IE
<i>sbc/spf</i>	κ	Sporadic	IV, VI, <i>bongori</i>	<i>stf</i>	π	Conserved	Missing in ID, IE
<i>sbs</i>	β	Sporadic	II <i>salamae</i>	<i>stg</i>	γ 1	Sporadic	ID, <i>bongori</i>
<i>sdc/sas</i>	σ	Sporadic	IIIa <i>arizonae</i>	<i>sth</i>	γ 1	Core	Missing IIIa and IIIb
<i>sdd/smf</i>	γ 1	Sporadic	IE, II, IIIa, IV	<i>sti</i>	γ 1	Conserved	Missing in ID
<i>sde</i>	γ 3	Sporadic	Tennessee (IE)	<i>stj</i>	β	Sporadic	IA, IE
<i>sdh</i>	γ 4	Sporadic	IE	<i>stk</i>	γ 4	Sporadic	IE
<i>sdi/sdf</i>	γ 4	Sporadic	IIIb <i>diarizonae</i>	<i>tcf</i>	α	Sporadic	IC, ID, IE

Table 1. *Salmonella* fimbriome.

Each *Salmonella* strain contains 5–14 different CU fimbriae with an average of 12 fimbriae in *S. enterica*. Representatives from all the six phylogenetic clades are present in *Salmonella* (Table 2) [26]. The γ -fimbriae constitute the largest clade with 22 FGCs and include the highly conserved FGCs (*bcf*, *fim* and *sth*) that belong to the clade γ -1. The most diverse clade is γ -4, with the conserved *stb* and *stc* or *peg* (*stc-peg*) and many of the new sporadic FGCs, while the α clade (for alternate CU), also known as class 5 fimbriae, has one FGC, *tcf*, which is found in several serovars. The σ clade also had only one FGC representative, *sdc*, that was only found in *S. enterica* subspecies IIIa (*arizonae*).

The distribution of the 38 FGCs gave a signature for each species, subspecies and serovars (**Table 3**). Seven FGCs, curli and the CU *fim*, *bcf*, *sth*, *stb*, *stc-peg* and *std*, represent the conserved (core) fimbriae of *Salmonella* (positive in more than 90% of strains). The *fim* fimbriae were found in all *S. enterica* strains, only missing in *S. bongori*. The *bcf* cluster was only missing in *S. enterica* ssp. IV (*houtenae*), and the *sth* cluster was only missing in *S. enterica* ssp. IIIa and IIIb. The *stb* cluster was present in *S. enterica* ssp. I, II, IIIb and the *std* cluster was not detected in *S. enterica* serovar Gallinarum, ssp. II, IIIA and *S. bongori*. The FGC *stc* and *peg* had probably emerged from a common ancestor: they belong to the same clade (γ -4) and are inserted at the same position in the genome (between *thiM* and *mrp*); their distribution is mutually exclusive; and either one is present in the majority of *Salmonella* strains.

CU clade	Fimbriae
α	<i>tcf</i>
β	<i>sbs</i> , <i>sib</i> , <i>stj</i>
γ 1	<i>bcf</i> , <i>fim</i> , <i>lpf</i> , <i>sdd/smf</i> , <i>stg</i> , <i>sth</i> , <i>sti</i>
γ 3	<i>saf</i> , <i>sde</i> , <i>sef</i>
γ 4	<i>mrk</i> , <i>peg</i> , <i>peh</i> , <i>sba</i> , <i>sdh</i> , <i>sdi</i> , <i>sdj</i> , <i>ssf</i> , <i>sta</i> , <i>stb</i> , <i>stc</i> , <i>stk</i>
κ	<i>fael/skf</i> , <i>pef</i> , <i>sbc/spf</i>
π	<i>sbb/sbf</i> , <i>sdk</i> , <i>sdl</i> , <i>std</i> , <i>ste</i> , <i>stf</i>
σ	<i>sdc/sas</i>

Table 2. *Salmonella* fimbrial type.

Most cases of salmonellosis in humans are caused by *S. enterica* ssp. I, and many of the sequenced serovars were from ssp. I. Thus, 27 out of the 38 FGCs are found in ssp. I. The ssp. I was divided into five classes using previous phylogenetic analysis [46, 47] (**Table 3**). The class IA contains broad host range serovars involved in gastroenteritis, mainly serovar Typhimurium. The class IB is formed by serovars Dublin, Enteritidis, Pullorum and Gallinarum, all sharing similar O-antigens and FGCs. The class IC contains serovars Choleraesuis and Paratyphi C and class ID contains the human-specific serovars Typhi and Paratyphi A. A separate branch of class IA, including serovars Heidelberg, Virchow and Hadar, that had the highest number of FGCs, as well as serovars Montevideo, Schwarzengrund, Welterveden, Javiana, Kentucky and Tennessee, was commonly isolated in association with edible plants and constitutes the class IE.

In addition to the seven core FGCs, five highly conserved FGCs (*saf*, *ste*, *stf*, *sti* and *lpf*) were associated with *S. enterica* ssp. I (**Table 3**). The *sti*, *lpf* and *stf* clusters are missing in human-specific serovars (class ID). The *ste* cluster is missing in class IA serovars and in some of the class IE serovars. Thus, *S. enterica* ssp. I harbours the core FGCs (*fim*, *bcf*, *sth*, *stb*, *stc-peg* and *std*), the conserved FGCs (*saf*, *ste*, *stf*, *sti* and *lpf*) and some sporadic FGCs unique to each serovar. Many FGCs of *Salmonella* are sporadic and form the unique repertoire in each serovar.

Despite the presence of many FGCs, extensive gene degradation was observed in most of the host-restricted and warm-blooded host-adapted serovars, mainly Gallinarum, Choleraesuis,

Paratyphi A and Typhi. Genome degradation of FGCs may correspond to the loss of genes rendered unnecessary by niche specialisation or by selective pressure in order to diminish antigen presentation at the bacterial surface during systemic disease. Intriguingly, most of FGCs were intact in Paratyphi B.

There are 11 FGCs that are not in *ssp. I*, with only *sbc* and *sdk* that are shared by more than one serovars. The low numbers of FGCs might be specific for cold-blooded animals' colonisation. A conserved signature specific for each subspecies was observed. As more diverse strains will be sequenced, new FGCs probably be discovered.

	Subspecies	Core	Conserved		Accessory	Absent
<i>Salmonella enterica</i>	I. <i>enterica</i>	<i>bcf, csg, fim, sth, stb, std, stc-peg</i>	<i>sti,, saf, ste, stf, lpf</i>	A	<i>pef, stj</i>	<i>ste</i>
				B	<i>fae, sef</i>	
				C	<i>pef, tcf,</i>	
				D	<i>sef, sta, stg, tcf, pil</i>	<i>sti, lpf, stf</i>
				E	<i>fae, mrk, peh, sdd, sde, sdh, stj, stk, pil</i>	<i>lpf, ste, stf</i>
	VI. <i>indica</i>				<i>sbc, sdk, sib</i>	<i>stb</i>
	II. <i>salamae</i>				<i>sdd, ssf, sbs</i>	<i>std</i>
	IV. <i>houtenae</i>				<i>sbc, sdd</i>	<i>bcf, stb</i>
	IIIb. <i>diarizonae</i>				<i>sdi, sdj, sdk, sdl</i>	<i>sth</i>
	IIIa. <i>arizonae</i>				<i>sdc, sdd</i>	<i>sth, stb, std</i>
<i>Salmonella bongori</i>			<i>lpf</i>		<i>sba, sbb, sbc, sbe, stg(sbd)</i>	<i>fim, stb, std</i>

Table 3. Fimbrial distribution.

4. Fimbrial regulation

Salmonella fimbriae are usually not expressed constitutively and rarely expressed under laboratory condition, except for Fim fimbriae, a type-1 fimbria [3]. Fimbriae are important during infection [19, 48, 49], suggesting that their expression is tightly regulated. Little is known about the regulation mechanisms that promote fimbrial expression. In general, fimbrial expression is positively or negatively regulated at the genetic level. Some regulators are unique to a specific fimbriae, like the regulation of curli by CsgD, while others are global, like Dam, H-NS and Lrp (leucine-responsive regulatory protein) [50]. These mechanisms include regulatory proteins, DNA methylation, cyclic di-GMP and small RNAs [50]. In *Salmonella*, a regulation network exists between the virulence factors. Here, we present the regulation of fimbrial genes including the interaction with motility and invasion. Then, we propose an example of regulation of the *fim* FGC expression in *S. Typhimurium*, the most characterised fimbriae of *Salmonella*.

4.1. General regulation of fimbrial genes

Genes implicated in different aspects of virulence including motility, adhesion, invasion of host cells and intestinal persistence are all regulated during infection. It was proposed that there is a temporal hierarchy between the T3SS of SPI-1 (invasion), flagellar and fimbrial genes, where SPI-1 is first activated, followed by flagellar genes and then type-1 fimbrial genes (*fim*). The crosstalk between these systems seems to be critical for bacterial pathogenesis [51]. Each element of virulence is related to a large regulation network that is not completely understood. DNA adenine methylation (Dam) regulates many virulence genes in *Salmonella* [52]: it is required for SPI-1 and *pef* expression, but it also represses many genes, including the *std*, *csg* and flagellar genes [52–54]. It was also shown that fimbrial FGCs are repressed by the Rcs phosphorelay, a sensor of outer membrane stress [55]. Another example of regulation interaction between motility and fimbrial expression was observed by a deletion of *ydiV* in *S. Typhimurium* that results in the derepression of curli fimbriae (*csgAB*), causing an increase in swimming motility and a decrease in swarming [56].

Crosstalk regulation also occurs between the capsule and the type IVb fimbriae in *S. Typhi*. Both virulence factors are encoded on SPI-7 and facilitate invasion of monocytes, suggesting a regulation overlapped. However, the exact regulation elements that act on those two systems are unknown [57].

One of the post-transcriptional regulation mechanisms uses the binding of small RNAs and the Hfq chaperone. In an *hfq* mutant strain, the expression of fimbrial gene *sefA* was activated when most of the other fimbrial subunit genes were repressed in *S. Enteritidis*. Overall, the *hfq* deletion decreased adherence compared to wild-type strain. Thus, Hfq seems to regulate fimbrial expression of most fimbrial genes from *S. Enteritidis* [58]. There is probably more sRNAs regulation of fimbrial gene expression awaiting to be discovered.

Phase variation is a transcriptional mechanism that controls the switch between fimbriated (ON) and afimbriated (OFF) cells within a bacterial population. In *Salmonella*, expression of *lpf* and *pef* was shown to be controlled by phase variation. The regulators of this mechanism are various and depend on the FGCs concerned [54, 59].

The secondary messenger cyclic-di-GMP controls virulence and biofilm formation in *Salmonella* [60]. In *Salmonella*, curli expression was activated by AdrA, a GGDEF-domain protein that increases intracellular level of cyclic-di-GMP [61]. Fimbrial production regulated by the cyclic-di-GMP level was also observed in other species such as *Klebsiella pneumoniae*, *E. coli* and *P. aeruginosa* [62].

In spite of all those known elements of regulation, how *Salmonella* passes from being afimbriated in vitro to a fimbriated form in vivo is still unknown.

4.2. Regulation of *fim* in *S. Typhimurium*

The *fim* FGC codes for six genes (*fimAICDHF*). This cluster is the most studied and one of the most conserved fimbriae of *Salmonella enterica* and was mainly characterised in *S. Typhimurium*. These fimbriae have a binding specificity for mannose residues [63]. The *fim* fimbria of *Salmonella* is not homologue with its homonym from *E. coli*, except for sharing some morphological and

mechanistic features [64, 65]. Regulation and amino acid sequences of fimbrial proteins are divergent between the two species. The transition from afimbriated to fimbriated stage occurs for *fim*, but there is no inversion of the promoter region as observed for *E. coli* phase variation [66]. For *S. Typhimurium*, the major subunit FimA is accompanied by two other subunits, FimI and FimF, and by the adhesin, FimH. The *fimC* and *fimD* genes encode respectively the chaperone and the usher [67]. Ancillary genes, *fimZYW*, and a rare arginine transfer RNA (tRNA) *fimU* and *STM0551*, inserted between *fimY* and *fimW*, directly regulate *fim* expression [62].

Ancillary genes *fimZYW* regulate the expression of *fimA* [50, 68–70]. FimZ, a sensor DNA-binding protein, is the principal positive regulator of *fimA* [71]. FimY upregulates *fimZ* expression by binding to the *fimZ* promoter. FimY and FimZ then form a complex that activates the *fimA* promoter [63, 69]; *fimY* is itself regulated by the arginine tRNA *fimU* [72]. Lrp is another regulator that binds and activates the *fimZ* promoter, probably by antagonising the binding of the global repressor protein H-NS to this promoter region [67, 70, 73]. H-NS has a high affinity for AT-rich DNA region, and *fimZ* gene has an unusual AT-rich sequence [73].

At the opposite, FimW repressed directly *fimA* expression and indirectly by lowering FimZ availability by its degradation caused by FimW binding [74]. It is also suggested that *STM0551*, an EAL domain protein, is a negative regulator of *fim* expression by lowering the c-di-GMP level [62].

FimZ is also used as a regulator relay by two-component system for expression of *hilA*, the principal regulator of SPI-1 (invasion). The two-component system PhoBR induces *fimZ* expression, and PhoPQ activates *hilE*, one of the negative regulators of *hilA*, by a FimYZ-dependant manner. FimZ also downregulates *flhDC*, genes implicated in flagella expression [75]. Flagellar gene *fliZ* also represses *fimZ* on a post-transcriptional manner, reinforcing the fact that there is an alternated expression of flagella (motility) and fimbriae (adhesion) [51] and confirming a regulation network between SPI-1 (invasion) and fimbrial expression [76]. Thus, a combination of factors directly implicated in fimbrial genes regulation can also impact on other virulence systems of *Salmonella* (Figure 4).




Figure 4. *fim* regulation. The *fim* FGC is activated and inhibited by diverse regulators. FimZ, accompanied by FimY, is the principal activator of *fimA*. Lrp and c-di-GMP also activate *fimA* at the promoter level. H-NS and FimW inhibit *fimA* expression by linking its promoter. FimW also reduces *fimA* expression by linking FimZ and decreases the availability of this activator. The gene *fimZ* is activated by FimZ itself and by FimY, but is repressed by FliZ in a post-transcriptional manner. The tRNA *fimU* regulates *fimY*. FimZ downregulates *flhCD*, genes implicated in flagellar expression.

5. Fimbriae as a tool

Salmonella infections are a major concern for public and animal health. Some serovars are host specific, while others are broad-spectrum pathogens and can be transmitted from food-borne animals to humans. On the other hand, animals can develop health problems and will not be suitable for consumption. To prevent those issues, it is critical to develop ways to detect *Salmonella* and protect potential hosts against infection. The importance of fimbriae for detection of *Salmonella* by molecular techniques and for vaccine development is presented in this section [77].

5.1. *Salmonella* detection using fimbrial genes

Salmonella-specific tests were performed since the end of the 1980s and mainly targeted surface antigens. Those tests include agglutination tests and ELISA (enzyme-linked immunosorbent assays) [77–79]. In 1993, Doran et al. presented a DNA-based test that targets *csgA* (*agfA*), offering a faster and more precise test for genus identification [80]. Then, in early 2000s, PCR (polymerisation chain reaction) tests using fimbrial genes, like *sef* or *csgA* (*agfA*), in combination with other virulence genes were developed to differentiate *Salmonella* strains from each other [81]. Different PCR tests (multiplex, nested and direct PCR) were elaborated for detection of *Salmonella*. Several of those tests integrated detection of fimbrial genes (i.e. *staA*, *fimW*) to discriminate between serovars [81–83]. Recently, a loop-mediated isothermal amplification (LAMP) assay was developed to detect *Salmonella* by targeting *bcfD*, a gene that belongs to the core of FGC. In isothermal conditions, the reaction occurs in an hour permitting rapid detection of *Salmonella* [84].

Salmonella-specific tests evolved from detecting antigens, which can be long and expensive to perform, to detecting specific genes in less than an hour by sensitive methods. Fimbrial genes are tools of choice for detection of *Salmonella*. The presence of conserved fimbrial genes allows the discrimination between *Salmonella* and non-*Salmonella* species. On the other hand, the presence of a specific pattern of fimbrial genes enables the discrimination between serovars.

5.2. Vaccines development

As surface structures, fimbriae constitute antigens of choice for the development of vaccines against *Salmonella* [85]. Fimbriae are difficult to study because they are poorly expressed under laboratory conditions and are redundant. The most interesting fimbriae are the ones expressed during infection. Targeting those fimbriae will confer higher chances to be recognised by the immune system in key moments of infection.

More than 20 fimbrial antigens were detected in typhoid fever patient's blood by transcriptomic analysis: SteD, StaACD, BcfDE, SafBC, TcfBCD, StbBC, FimAIDH, StdBC, StgACD and SthA [86]. Antibodies against immunogenic fimbrial proteins TcfB, StbD and CsgEFG were identified in the blood of typhoid fever patients [12]. Immunoreactive antibodies against SthDA and BcfA were found in lymphocytes supernatant (ALS) of patients with typhoid fever [87].

SefA, a protein from the SEF14 fimbriae of *S. Enteritidis*, was used as an antigen associated with liposomes for oral immunisation of chickens [88]. The immunisation of chickens by fimbrial antigens was efficient for IgG and IgA responses and reduced *Salmonella* colonisation. Four weeks after immunisation, the bacterial excretion from the intestinal tract was significantly reduced [88]. The liposome-associated immunisation was also performed with fimbrial antigen from SEF21 and resulted in a similar efficiency [89]. SefD, another antigen from SEF14, was also used to vaccinate animals in a bacterin preparation, a vaccine prepared from inactivated bacteria. This vaccine was efficient to reduce the presence of *Salmonella* from the spleens of hens [90].

As factors implicated in the first stages of infection, fimbriae are an interesting target for vaccine development [91]. Fimbrial antigens are important for the development of new anti-*Salmonella* therapies [85, 86]. However, a better understanding of their expression pattern in vivo is needed to optimise the therapeutic effects of fimbrial-targeted vaccines. Fimbrial antigens may be combined with other immunogenic proteins to increase the immune response [91].

6. Conclusion

Fimbriae are diverse proteinaceous surface structures. They diverge by their assembly mechanisms and result in different filamentous structures with roles in pathogenesis. However, their roles are not completely understood. They were first known for adherence to cells and inert surfaces, but they seem to be implicated in so much more functions during infection.

The multiplicity of adhesion systems is also an enigma. Most of the *Salmonella* serovars possess 12 fimbrial gene clusters. Some fimbriae are specific to certain serovars and may play a role in these bacteria that do not need to be fulfilled in other serovars. At the opposite, there is a core of fimbrial genes that are present in most of the serovars. Fimbriae are one of the keys to understand *Salmonella* pathogenesis. The specific pattern of each serovar, with further investigations on the sporadic fimbriae, may also bring insights into our understanding of *Salmonella* pathogenesis.

Regulation of fimbrial genes is a complex network that is tightly related to invasion and motility. Virulence factors are finely regulated, and a temporal expression hierarchy allows the success of *Salmonella* infection. General regulators are already known to regulate fimbrial genes such as stress sensor Rcs relay or the Hfq factor. Phase variation from a fimbriated to afimbriated status occurs in *Salmonella*. However, this phenomenon is not from a promoter inversion of the Fim cluster, but from the regulation by ancillary genes related to *fim* gene cluster. These ancillary genes are themselves precisely regulated by a variety of regulators known for their role in other bacterial processes.

The actual understanding of fimbrial expression opens a new area on human health prevention. Some conserved fimbrial genes, in combination with other virulence genes, are precious markers for *Salmonella* detection. These tools could permit a faster diagnostic for human patients, but also a rapid detection of contaminated food or infected animals. Fimbrial proteins can serve as good immunogens in vaccine preparation against *Salmonella* infection.

A better understanding of fimbrial expression, production and regulation processes becomes important for prevention of *Salmonella* infection. It will also enlighten the importance of fimbriae in other human pathogens, as fimbrial systems are part of virulence factors in many bacteria.

Author details

Karine Dufresne and France Daigle*

*Address all correspondence to: france.daigle@umontreal.ca

Department of Microbiology, Infectiology and Immunology, Université de Montréal, Montréal, Canada

References

- [1] Daigle F. Typhi genes expressed during infection or involved in pathogenesis. *J Infect Dev Ctries.* 2008;2(6):431–7.
- [2] Sabbagh SC, Forest CG, Lepage C, Leclerc JM, Daigle F. So similar, yet so different: uncovering distinctive features in the genomes of *Salmonella enterica* serovars Typhimurium and Typhi. *FEMS Microbiol Lett.* 2010;305(1):1–13.
- [3] Humphries AD, Raffatellu M, Winter S, Weening EH, Kingsley RA, Droleskey R, et al. The use of flow cytometry to detect expression of subunits encoded by 11 *Salmonella enterica* serotype Typhimurium fimbrial operons. *Mol Microbiol.* 2003;48(5):1357–76.
- [4] Townsend SM, Kramer NE, Edwards R, Baker S, Hamlin N, Simmonds M, et al. *Salmonella enterica* serovar Typhi possesses a unique repertoire of fimbrial gene sequences. *Infect Immun.* 2001;69(5):2894–901.
- [5] Houwink AL, van IW. Electron microscopical observations on bacterial cytology; a study on flagellation. *Biochim Biophys Acta.* 1950;5(1):10–44.
- [6] Anderson TF. On the mechanism of adsorption of bacteriophages on host cells. In: Miles AA, Pirie, N. W., editor. *The nature of the bacterial surface.* Oxford: Oxford University Press; 1949. p. 76–95.
- [7] Duguid JP, Smith IW, Dempster G, Edmunds PN. Non-flagellar filamentous appendages (fimbriae) and haemagglutinating activity in *Bacterium coli*. *J Pathol Bacteriol.* 1955;70(2):335–48.
- [8] Thanassi DG, Nuccio SP, Shu Kin So S, Baumler AJ. Fimbriae: Classification and Biochemistry. *EcoSal Plus.* 2007;2(2):1–27.
- [9] Ottow JC. Ecology, physiology, and genetics of fimbriae and pili. *Annu Rev Microbiol.* 1975;29:79–108.

- [10] Nuccio SP, Baumler AJ. Evolution of the chaperone/usher assembly pathway: fimbrial classification goes Greek. *Microbiol Mol Biol Rev.* 2007;71(4):551–75.
- [11] Mol O, Oudega B. Molecular and structural aspects of fimbriae biosynthesis and assembly in *Escherichia coli*. *FEMS Microbiol Rev.* 1996;19(1):25–52.
- [12] Harris JB, Baresch-Bernal A, Rollins SM, Alam A, LaRocque RC, Bikowski M, et al. Identification of in vivo-induced bacterial protein antigens during human infection with *Salmonella enterica* serovar Typhi. *Infect Immun.* 2006;74(9):5161–8.
- [13] Boddicker JD, Ledebauer NA, Jagnow J, Jones BD, Clegg S. Differential binding to and biofilm formation on HEp-2 cells by *Salmonella enterica* serovar Typhimurium is dependent upon allelic variation in the *fimH* gene of the *fim* gene cluster. *Mol Microbiol.* 2002;45(5):1255–65.
- [14] Edwards RA, Schifferli DM, Maloy SR. A role for *Salmonella* fimbriae in intraperitoneal infections. *Proc Natl Acad Sci U S A.* 2000;97(3):1258–62.
- [15] Althouse C, Patterson S, Fedorka-Cray P, Isaacson RE. Type 1 fimbriae of *Salmonella enterica* serovar Typhimurium bind to enterocytes and contribute to colonization of swine in vivo. *Infect Immun.* 2003;71(11):6446–52.
- [16] Sabbagh SC, Lepage C, McClelland M, Daigle F. Selection of *Salmonella enterica* serovar Typhi genes involved during interaction with human macrophages by screening of a transposon mutant library. *PLoS One.* 2012;7(5):e36643.
- [17] Chessa D, Winter MG, Jakomin M, Baumler AJ. *Salmonella enterica* serotype Typhimurium Std fimbriae bind terminal alpha(1,2)fucose residues in the cecal mucosa. *Mol Microbiol.* 2009;71(4):864–75.
- [18] Humphries A, Deridder S, Baumler AJ. *Salmonella enterica* serotype Typhimurium fimbrial proteins serve as antigens during infection of mice. *Infect Immun.* 2005;73(9):5329–38.
- [19] Weening EH, Barker JD, Laarakker MC, Humphries AD, Tsois RM, Baumler AJ. The *Salmonella enterica* serotype Typhimurium *lpf*, *bcf*, *stb*, *stc*, *std*, and *sth* fimbrial operons are required for intestinal persistence in mice. *Infect Immun.* 2005;73(6):3358–66.
- [20] Yue M, Han X, De Masi L, Zhu C, Ma X, Zhang J, et al. Allelic variation contributes to bacterial host specificity. *Nat Commun.* 2015;6:8754.
- [21] Fronzes R, Remaut H, Waksman G. Architectures and biogenesis of non-flagellar protein appendages in Gram-negative bacteria. *EMBO J.* 2008;27(17):2271–80.
- [22] Smyth CJ, Marron MB, Twohig JM, Smith SG. Fimbrial adhesins: similarities and variations in structure and biogenesis. *FEMS Immunol Med Microbiol.* 1996;16(2):127–39.
- [23] Wagner C, Hensel M. Adhesive mechanisms of *Salmonella enterica*. *Adv Exp Med Biol.* 2011;715:17–34.
- [24] Soto GE, Hultgren SJ. Bacterial adhesins: common themes and variations in architecture and assembly. *J Bacteriol.* 1999;181(4):1059–71.

- [25] Sauer FG, Barnhart M, Choudhury D, Knight SD, Waksman G, Hultgren SJ. Chaperone-assisted pilus assembly and bacterial attachment. *Curr Opin Struct Biol.* 2000;10(5):548–56.
- [26] Porwollik S. *Salmonella: from genome to function.* Norfolk, UK: Caister Academic Press; 2011. xii, 300, A1 p.
- [27] Gerlach RG, Hensel M. Protein secretion systems and adhesins: the molecular armory of Gram-negative pathogens. *Int J Med Microbiol.* 2007;297(6):401–15.
- [28] Thanassi DG, Saulino ET, Hultgren SJ. The chaperone/usher pathway: a major terminal branch of the general secretory pathway. *Curr Opin Microbiol.* 1998;1(2):223–31.
- [29] Dufresne K, Paradis-Bleau C. Biology and assembly of the bacterial envelope. *Adv Exp Med Biol.* 2015;883:41–76.
- [30] Sauer FG, Remaut H, Hultgren SJ, Waksman G. Fiber assembly by the chaperone-usher pathway. *Biochim Biophys Acta.* 2004;1694(1–3):259–67.
- [31] Barnhart MM, Sauer FG, Pinkner JS, Hultgren SJ. Chaperone-subunit-usher interactions required for donor strand exchange during bacterial pilus assembly. *J Bacteriol.* 2003;185(9):2723–30.
- [32] Saulino ET, Bullitt E, Hultgren SJ. Snapshots of usher-mediated protein secretion and ordered pilus assembly. *Proc Natl Acad Sci USA.* 2000;97(16):9240–5.
- [33] Zavialov A, Zav'yalova G, Korpela T, Zav'yalov V. FGL chaperone-assembled fimbrial polyadhesins: anti-immune armament of Gram-negative bacterial pathogens. *FEMS Microbiol Rev.* 2007;31(4):478–514.
- [34] Waksman G, Hultgren SJ. Structural biology of the chaperone-usher pathway of pilus biogenesis. *Nat Rev Microbiol.* 2009;7(11):765–74.
- [35] Li H, Qian L, Chen Z, Thibault D, Liu G, Liu T, et al. The outer membrane usher forms a twin-pore secretion complex. *J Mol Biol.* 2004;344(5):1397–407.
- [36] Zhan R, Han Q, Zhang C, Tian Z, Zhang J. Toll-Like receptor 2 (TLR2) and TLR9 play opposing roles in host innate immunity against *Salmonella enterica* serovar Typhimurium infection. *Infect Immun.* 2015;83(4):1641–9.
- [37] Barnhart MM, Chapman MR. Curli biogenesis and function. *Annu Rev Microbiol.* 2006;60:131–47.
- [38] Goyal P, Krasteva PV, Van Gerven N, Gubellini F, Van den Broeck I, Troupiotis-Tsailaki A, et al. Structural and mechanistic insights into the bacterial amyloid secretion channel CsgG. *Nature.* 2014;516(7530):250–3.
- [39] Nenninger AA, Robinson LS, Hultgren SJ. Localized and efficient curli nucleation requires the chaperone-like amyloid assembly protein CsgF. *Proc Natl Acad Sci USA.* 2009;106(3):900–5.
- [40] Bian Z, Normark S. Nucleator function of CsgB for the assembly of adhesive surface organelles in *Escherichia coli.* *EMBO J.* 1997;16(19):5827–36.

- [41] White AP, Gibson DL, Collinson SK, Banser PA, Kay WW. Extracellular polysaccharides associated with thin aggregative fimbriae of *Salmonella enterica* serovar enteritidis. *J Bacteriol.* 2003;185(18):5398–407.
- [42] Karupppiah V, Collins RF, Thistlethwaite A, Gao Y, Derrick JP. Structure and assembly of an inner membrane platform for initiation of type IV pilus biogenesis. *Proc Natl Acad Sci USA.* 2013;110(48):E4638–47.
- [43] Takhar HK, Kemp K, Kim M, Howell PL, Burrows LL. The platform protein is essential for type IV pilus biogenesis. *J Biol Chem.* 2013;288(14):9721–8.
- [44] Proft T, Baker EN. Pili in Gram-negative and Gram-positive bacteria—structure, assembly and their role in disease. *Cell Mol Life Sci.* 2009;66(4):613–35.
- [45] Jain S, Moscicka KB, Bos MP, Pachulec E, Stuart MC, Keegstra W, et al. Structural characterization of outer membrane components of the type IV pili system in pathogenic *Neisseria*. *PLoS One.* 2011;6(1):e16624.
- [46] Desai PT, Porwollik S, Long F, Cheng P, Wollam A, Bhonagiri-Palsikar V, et al. Evolutionary Genomics of *Salmonella enterica* Subspecies. *MBio.* 2013;4(2):e00579–12.
- [47] Yue M, Rankin SC, Blanchet RT, Nulton JD, Edwards RA, Schifferli DM. Diversification of the *Salmonella* fimbriae: a model of macro- and microevolution. *PLoS One.* 2012;7(6):e38596.
- [48] van der Velden AW, Baumler AJ, Tsois RM, Heffron F. Multiple fimbrial adhesins are required for full virulence of *Salmonella typhimurium* in mice. *Infect Immun.* 1998;66(6):2803–8.
- [49] Clayton DJ, Bowen AJ, Hulme SD, Buckley AM, Deacon VL, Thomson NR, et al. Analysis of the role of 13 major fimbrial subunits in colonisation of the chicken intestines by *Salmonella enterica* serovar Enteritidis reveals a role for a novel locus. *BMC Microbiol.* 2008;8:228.
- [50] Clegg S, Wilson J, Johnson J. More than one way to control hair growth: regulatory mechanisms in enterobacteria that affect fimbriae assembled by the chaperone/usher pathway. *J Bacteriol.* 2011;193(9):2081–8.
- [51] Saini S, Slauch JM, Aldridge PD, Rao CV. Role of cross talk in regulating the dynamic expression of the flagellar *Salmonella* pathogenicity island 1 and type 1 fimbrial genes. *J Bacteriol.* 2010;192(21):5767–77.
- [52] Balbontin R, Rowley G, Pucciarelli MG, Lopez-Garrido J, Wormstone Y, Lucchini S, et al. DNA adenine methylation regulates virulence gene expression in *Salmonella enterica* serovar Typhimurium. *J Bacteriol.* 2006;188(23):8160–8.
- [53] Aya Castaneda Mdel R, Sarnacki SH, Noto Llana M, Lopez Guerra AG, Giacomodonato MN, Cerquetti MC. Dam methylation is required for efficient biofilm production in *Salmonella enterica* serovar Enteritidis. *Int J Food Microbiol.* 2015;193:15–22.

- [54] Nicholson B, Low D. DNA methylation-dependent regulation of *pef* expression in *Salmonella typhimurium*. *Mol Microbiol*. 2000;35(4):728–42.
- [55] Wang Q, Zhao Y, McClelland M, Harshey RM. The RcsCDB signaling system and swarming motility in *Salmonella enterica* serovar typhimurium: dual regulation of flagellar and SPI-2 virulence genes. *J Bacteriol*. 2007;189(23):8447–57.
- [56] Deditius JA, Felgner S, Sporing I, Kuhne C, Frahm M, Rohde M, et al. Characterization of novel factors involved in swimming and swarming motility in *Salmonella enterica* Serovar Typhimurium. *PLoS One*. 2015;10(8):e0135351.
- [57] Lee FK, Morris C, Hackett J. The *Salmonella enterica* serovar Typhi Vi capsule and self-association pili share controls on expression. *FEMS Microbiol Lett*. 2006;261(1):41–6.
- [58] Meng X, Meng X, Zhu C, Wang H, Wang J, Nie J, et al. The RNA chaperone Hfq regulates expression of fimbrial-related genes and virulence of *Salmonella enterica* serovar Enteritidis. *FEMS Microbiol Lett*. 2013;346(2):90–6.
- [59] Norris TL, Kingsley RA, Bumler AJ. Expression and transcriptional control of the *Salmonella typhimurium* *Ipf* fimbrial operon by phase variation. *Mol Microbiol*. 1998;29(1):311–20.
- [60] Ahmad I, Lamprokostopoulou A, Le Guyon S, Streck E, Barthel M, Peters V, et al. Complex c-di-GMP signaling networks mediate transition between virulence properties and biofilm formation in *Salmonella enterica* serovar Typhimurium. *PLoS One*. 2011;6(12):e28351.
- [61] Kader A, Simm R, Gerstel U, Morr M, Romling U. Hierarchical involvement of various GGDEF domain proteins in *rdar* morphotype development of *Salmonella enterica* serovar Typhimurium. *Mol Microbiol*. 2006;60(3):602–16.
- [62] Wang KC, Hsu YH, Huang YN, Yeh KS. A previously uncharacterized gene *stm0551* plays a repressive role in the regulation of type 1 fimbriae in *Salmonella enterica* serotype Typhimurium. *BMC Microbiol*. 2012;12:111.
- [63] Wang KC, Hsu YH, Huang YN, Lin JH, Yeh KS. FimY of *Salmonella enterica* serovar Typhimurium functions as a DNA-binding protein and binds the *fimZ* promoter. *Microbiol Res*. 2014;169(7–8):496–503.
- [64] Humphries AD, Townsend SM, Kingsley RA, Nicholson TL, Tsolis RM, Baumler AJ. Role of fimbriae as antigens and intestinal colonization factors of *Salmonella* serovars. *FEMS Microbiol Lett*. 2001;201(2):121–5.
- [65] Kisiela DI, Chattopadhyay S, Tchesnokova V, Paul S, Weissman SJ, Medenica I, et al. Evolutionary analysis points to divergent physiological roles of type 1 fimbriae in *Salmonella* and *Escherichia coli*. *MBio*. 2013;4(2).
- [66] Clegg S, Hancox LS, Yeh KS. *Salmonella typhimurium* fimbrial phase variation and *fimA* expression. *J Bacteriol*. 1996;178(2):542–5.

- [67] Wang KC, Hsu YH, Huang YN, Chen TH, Lin JH, Hsuan SL, et al. A low-pH medium in vitro or the environment within a macrophage decreases the transcriptional levels of *fimA*, *fimZ* and *lrp* in *Salmonella enterica* serovar Typhimurium. *J Biosci.* 2013;38(3):499–507.
- [68] Swenson DL, Clegg S. Identification of ancillary *fim* genes affecting *fimA* expression in *Salmonella typhimurium*. *J Bacteriol.* 1992;174(23):7697–704.
- [69] Tinker JK, Clegg S. Characterization of FimY as a coactivator of type 1 fimbrial expression in *Salmonella enterica* serovar Typhimurium. *Infect Immun.* 2000;68(6):3305–13.
- [70] Chuang YC, Wang KC, Chen YT, Yang CH, Men SC, Fan CC, et al. Identification of the genetic determinants of *Salmonella enterica* serotype Typhimurium that may regulate the expression of the type 1 fimbriae in response to solid agar and static broth culture conditions. *BMC Microbiol.* 2008;8:126.
- [71] Yeh KS, Tinker JK, Clegg S. FimZ binds the *Salmonella typhimurium fimA* promoter region and may regulate its own expression with FimY. *Microbiol Immunol.* 2002;46(1):1–10.
- [72] Tinker JK, Clegg S. Control of FimY translation and type 1 fimbrial production by the arginine tRNA encoded by *fimU* in *Salmonella enterica* serovar Typhimurium. *Mol Microbiol.* 2001;40(3):757–68.
- [73] McFarland KA, Lucchini S, Hinton JC, Dorman CJ. The leucine-responsive regulatory protein, Lrp, activates transcription of the *fim* operon in *Salmonella enterica* serovar typhimurium via the *fimZ* regulatory gene. *J Bacteriol.* 2008;190(2):602–12.
- [74] Tinker JK, Hancox LS, Clegg S. FimW is a negative regulator affecting type 1 fimbrial expression in *Salmonella enterica* serovar typhimurium. *J Bacteriol.* 2001;183(2):435–42.
- [75] Clegg S, Hughes KT. FimZ is a molecular link between sticking and swimming in *Salmonella enterica* serovar Typhimurium. *J Bacteriol.* 2002;184(4):1209–13.
- [76] Baxter MA, Jones BD. Two-component regulators control *hilA* expression by controlling *fimZ* and *hilE* expression within *Salmonella enterica* serovar Typhimurium. *Infect Immun.* 2015;83(3):978–85.
- [77] Thorns CJ. *Salmonella* fimbriae: novel antigens in the detection and control of *salmonella* infections. *Br Vet J.* 1995;151(6):643–58.
- [78] Feldsine PT, Falbo-Nelson MT, Husted DL. Polyclonal enzyme immunoassay method for detection of motile and non-motile *Salmonella* in foods: comparative study. *J AOAC Int.* 1993;76(3):694–7.
- [79] Kerr S, Ball HJ, Mackie DP, Pollock DA, Finlay DA. Diagnostic application of monoclonal antibodies to outer membrane protein for rapid detection of *Salmonella*. *J Appl Bacteriol.* 1992;72(4):302–8.
- [80] Doran JL, Collinson SK, Burian J, Sarlos G, Todd EC, Munro CK, et al. DNA-based diagnostic tests for *Salmonella* species targeting *agfA*, the structural gene for thin, aggregative fimbriae. *J Clin Microbiol.* 1993;31(9):2263–73.

- [81] Craciunas C, Keul AL, Flonta M, Cristea M. DNA-based diagnostic tests for *Salmonella* strains targeting *hilA*, *agfA*, *spvC* and *sef* genes. *J Environ Manage*. 2012;95 Suppl:S15–8.
- [82] Zhang JY, Dong LW, Ren Q, Wang XZ, Yang Y, Zhou W, et al. Simple and rapid detection of *Salmonella* by direct PCR amplification of gene *fimW*. *Curr Microbiol*. 2014;69(4):429–35.
- [83] Pratap CB, Kumar G, Patel SK, Verma AK, Shukla VK, Kumar K, et al. Targeting of putative fimbrial gene for detection of *S. Typhi* in typhoid fever and chronic typhoid carriers by nested PCR. *J Infect Dev Ctries*. 2013;7(7):520–7.
- [84] Zhuang L, Gong J, Li Q, Zhu C, Yu Y, Dou X, et al. Detection of *Salmonella* spp. by a loop-mediated isothermal amplification (LAMP) method targeting *bcfD* gene. *Lett Appl Microbiol*. 2014;59(6):658–64.
- [85] Ghosh S, Chakraborty K, Nagaraja T, Basak S, Koley H, Dutta S, et al. An adhesion protein of *Salmonella enterica* serovar Typhi is required for pathogenesis and potential target for vaccine development. *Proc Natl Acad Sci USA*. 2011;108(8):3348–53.
- [86] Sheikh A, Charles RC, Sharmeen N, Rollins SM, Harris JB, Bhuiyan MS, et al. In vivo expression of *Salmonella enterica* serotype Typhi genes in the blood of patients with typhoid fever in Bangladesh. *PLoS Negl Trop Dis*. 2011;5(12):e1419.
- [87] Charles RC, Liang L, Khanam F, Sayeed MA, Hung C, Leung DT, et al. Immunoproteomic analysis of antibody in lymphocyte supernatant in patients with typhoid fever in Bangladesh. *Clin Vaccine Immunol*. 2014;21(3):280–5.
- [88] Pang Y, Zhang Y, Wang H, Jin J, Piao J, Piao J, et al. Reduction of *Salmonella* enteritidis number after infections by immunization of liposome-associated recombinant SefA. *Avian Dis*. 2013;57(3):627–33.
- [89] Pang Y, Wang H, Li Z, Piao J, Piao J, Chi Y, et al. Immune response to liposome-associated recombinant SEF21 following oral immunization in chickens. *Avian Dis*. 2012;56(2):347–53.
- [90] Sanchez-Ingunza R, Guard J, Morales CA, Icard AH. Reduction of *Salmonella* Enteritidis in the spleens of hens by bacterins that vary in fimbrial protein SefD. *Foodborne Pathog Dis*. 2015;12(10):836–43.
- [91] Wizemann TM, Adamou JE, Langermann S. Adhesins as targets for vaccine development. *Emerg Infect Dis*. 1999;5(3):395–403.

