

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

## Chemical, Antioxidant, and Cytotoxic Properties of Native Blue Corn Extract

---

Rosa Guzmán-Gerónimo, Edna Alarcón Aparicio,  
Oscar García Barradas, Jose Chávez-Servia and  
Tania Alarcón-Zavaleta

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67574>

---

### Abstract

In recent years, natural products such as dietary phytoconstituents have been the focus of scientific studies for cancer prevention. Among these are polyphenols, which have shown anticancer properties. Pigmented cereals such as blue maize are a rich source of polyphenols such as anthocyanins. Therefore, the aim of this work is to determine the chemical composition and cytotoxic activity of blue maize extract in several cancer cell lines. The total polyphenol content, total anthocyanins, and antioxidant activity of 16 blue corn samples from the Mixteco race were analyzed. From these, the sample with the highest content of polyphenols, anthocyanins, and antioxidant activity was selected and its anthocyanin fraction was isolated using an amberlite column and analyzed by means of HPLC-ESI-MS. The total polyphenol content ranged from 142.8 to 203.2 mg GAE/100g. The total anthocyanin contents varied between 19.02 and 66.92 mg C3G/100g. The antioxidant activity ranged from 18.5 to 27.8  $\mu\text{mol TE/g}$ . The anthocyanin profile showed eight different compounds, mainly acylated anthocyanins. Cytotoxicity of blue corn extract on cancer cell lines was determined at concentrations of 100 and 500  $\mu\text{g/mL}$  using the SRB assay. A cytotoxic effect was mainly observed on SKLU-1 and HTC-15 cell lines.

**Keywords:** blue corn extract, dietary phytoconstituents, anthocyanin profile, cancer cell lines, cytotoxic activity

---

## 1. Introduction

Molecules derived from natural sources, such as plants, marine organisms, and microorganisms, have become important sources of active compounds in the development of drugs for the treatment of human chronic diseases. In recent years, natural products such as dietary phytoconstituents have been the focus of scientific studies for cancer prevention [1]. Epidemiological and preclinical research indicates that dietary compounds possess chemopreventive properties, for example, garlic consumption has been associated with a lower risk of cancer [2–4]. In addition, supplementation of dietary phytochemicals for chemoprevention is gaining increased attention due to their chemical diversity, biological activity, and good availability.

Currently, more than 1000 dietary compounds belonging to different chemical classes have shown potential chemopreventive activities [5]. Among dietary constituents, polyphenols such as anthocyanins have demonstrated to exert many biological activities including anticancer properties [6]. From the chemical standpoint, anthocyanins are phenolic substances that belong to the group of flavonoids derived from the 2-phenylbenzopyrilic cation found in nature in a glycosylated or acylated form [7]. These compounds are particularly abundant in pigmented cereals such as red, purple, and black rice, black sorghum, and red, blue, or purple maize [8–10].

Mexico is the center of origin and biodiversity of maize (*Zea mays* L.). Species have an extensive genetic diversity, with 59 different races described with different shapes and colors ranging from white to yellow, red, purple, and blue [11]. Pigmented maize genotypes are used in the production of tortillas, tamales, atoles, and other traditional Mexican foods. These maize varieties have been the focus of scientific studies because they are a rich source of polyphenols such as anthocyanins. Recent data indicate that blue maize contains monomeric anthocyanins as well as acylated anthocyanins [12, 13].

Even though blue maize is an important part of the Mexican diet, there is little scientific information regarding its anthocyanin profile and anticancer properties. Chemical composition is a factor that must be considered in the selection of blue maize genotypes due to its impact on biological activity, and thus its potential applications for the treatment of disease such as cancer. For this reason, prior to embarking on cancer phytochemical trials, it is important to carry out a preclinical research in order to evaluate the potential application of phytochemicals from blue maize. It is well known that *in vitro* studies examine preliminary efficacy of phytochemicals for cancer prevention or therapy [14].

Given the above, the aim of this work is to evaluate the total content of polyphenols, anthocyanins, and the antioxidant activity of blue corn from the Mixteco race, and to determine its anthocyanin profile and the cytotoxic activity of the anthocyanin fraction in several cancer cell lines.

## 2. Research methods

### 2.1. Plant material

Sixteen samples of blue maize from the Mixteco race (**Figure 1**) were donated by the Interdisciplinary Research Center for Integral Regional Development (CIDIIR as per the



**Figure 1.** Grains of blue maize from Mixteco race.

Spanish acronym) of the National Polytechnic Institute, Oaxaca Unit in Mexico. Maize kernels were grounded and placed in amber bottles for analysis.

## 2.2. Blue corn extracts

Ground blue corn kernels (1:5 p:v) were homogenized for 20 min with ethanol acidified with citric acid 1M (85:15 v:v). This was performed using an ultrasonic homogenizer at a frequency of 20 kHz and 750 W power (Cole-Palmer Instrument Company, VCX-750, USA) with a tip diameter of 13 mm at an amplitude of 25  $\mu$ m with a pulse of 5 s in the 'On' position and 5 s in the 'Off' position. The sample was placed under refrigeration for 24 h and centrifuged at 4000 rpm for 15 min at a temperature of 5°C. The process was repeated twice and the extract was concentrated using a rotary evaporator under vacuum. The conditions of extraction have been included in a patent request, MX/A/20131011202.

## 2.3. Total phenolic content

For analytical purposes, total polyphenols were evaluated using the colorimetric method previously described by Folin-Ciocalteu and modified by Singleton and Rossi [15]. In this study,

0.2 mL of the extract was mixed with 3.0 mL of distilled water and 0.2 mL of Folin-Ciocalteu reagent. Next, a calcium carbonate saturated solution of 0.75 mL was added. Then the mixture was incubated for 60 min at 37°C in darkness; absorbance was read at 750 nm. This measurement was compared to a standard curve prepared with a gallic acid solution (20–120 mg/L) (Sigma Chemical). The total phenolic content was expressed as milligram equivalents of gallic acid/100 g of fresh weight (mg GAE/100g).

#### **2.4. Total monomeric anthocyanin content**

Monomeric anthocyanin content was evaluated using the differential pH method [16]. Absorbance was measured in a UV-VIS spectrophotometer (Perkin Elmer, Inc., Shelton, CT, USA). For the analysis, samples were diluted in potassium chloride buffer (pH 1.0) and sodium acetate buffer (pH 4.5). The difference in absorption at 510 and 700 nm was determined in buffers at pH 1.0 and 4.5. The monomeric anthocyanin content was expressed as cyanidin 3-glucoside mg/100g.

#### **2.5. Antioxidant activity**

The antioxidant assay was performed according to the DPPH (2,2-Diphenyl-1-picrylhydrazyl) method [17]. Trolox was used to make a calibration curve (100–800  $\mu\text{mol}$ ). About 2.9 mL of DPPH solution was mixed with 0.1 mL of blue corn extract, and then kept in the dark for 1 h. The sample was incubated for 30 min and its absorbance was read at 517 nm. The result was expressed as  $\mu\text{mol eq. trolox/g}$  of sample.

#### **2.6. Isolation and chromatographic analysis of anthocyanins**

For the isolation of anthocyanins, a column was packed with amberlite XAD-7 preconditioned with 5% acetic acid [18]. Then, 1 mL of the blue corn concentrated extract was placed into the column and eluted with acidified ethanol (5% acetic acid). The eluate was then concentrated to dryness using a Buchi rotary evaporator (Heidolph Digital Laborota pump 4011) coupled to Pimo Vacum Buchi V-700. Anthocyanins were analyzed by HPLC-ESI-MS. The HPLC system was coupled to a Brüker MicrOTOF II spectrometer. The column was C-18 ZORBAX eclipse plus column with 100 mm  $\times$  2.1 mm, 3.5  $\mu\text{m}$ . The isocratic elution was done with a mix of methanol:water (2:8 v:v). Mass spectra analysis was carried out in negative ion mode, scan range: 50–3000 amu, capillary voltage 3.8 kV, dry gas flow at 4.0 L/min.

#### **2.7. Cell culture and assay for cytotoxic activity**

Prostate cancer cell lines (PC-3), neoplastic myelogenous leukemic cell lines (K-562), human colon cancer cell lines (HCT-15), human breast cancer cell lines (MCF-7), and lung cancer cell lines were provided by the National Cancer Institute (NCI), USA, and the Center of HIV/AIDS Services Center in Mexico City. Cytotoxicity of blue corn extract on tumor cells was determined at different concentrations (50, 100, and 500  $\mu\text{g/mL}$ ), using the protein-binding dye sulforhodamine B (SRB) assay in microculture to determine cell growth [19]. Cell lines were cultured in RPMI-1640 (Sigma Chemical Co., Ltd., St. Louis, MO, USA), supplemented

with 10% of fetal bovine serum purchased from Invitrogen Corporation, 2 mM L-glutamine, 100 IU/mL, penicillin G, 100 mg/mL streptomycin sulfate, and 0.25 mg/mL amphotericin B (Gibco). They were maintained at 37°C in a 5% CO<sub>2</sub> atmosphere and 95% humidity. For the assay, the following suspensions were prepared: 5 × 10<sup>4</sup> cell/mL (K-562, MCF-7), 7.5 × 10<sup>4</sup> cell/mL (PC-3), and 10 × 10<sup>4</sup> cell/mL (HCT-15, SKLU-1); 100 µL of these cells in suspension were seeded in 96-well micro-titer plates and incubated in order to achieve cell attachment to the plates. After 24 h of incubation, 100 µL of each test compound and positive substances (Cisplatin) were added to each well. After 48 h of incubation, adherent cell cultures were fixed 'in situ' by adding 50 mL of cold 50% (wt/vol) trichloroacetic acid (TCA) and incubated for 60 min. at 4°C. The supernatant was discarded and the plates were washed three times with water and then air-dried. Cultures fixed with TCA were stained for 30 min. with 100 mL of 0.4% SRB solution. Protein-bound dye was extracted with 10 µmol of unbuffered tris base and the optical densities were measured by an Ultra Microplate Reader (Elx808, BIO-TEK Instruments, Inc.) with a test wavelength of 515 nm.

### 3. Results and discussion

#### 3.1. Total content of polyphenols, monomeric anthocyanins, and antioxidant activity

The first aim of this research is to evaluate the total content of polyphenols, anthocyanins, and the antioxidant activity of blue corn extracts. Ethanol acidified with citric acid was used in the preparation of the extracts, since organic acids decrease the decomposition of anthocyanins during the following concentration step [20].

The total polyphenol content was observed in the range of 143–203 mg equivalent of gallic acid/100 g sample (**Table 1**), while the concentration of monomeric anthocyanins varied from 21 to 69 mg cyanidin-3-glucoside/100 g sample. In this study, the total polyphenol and anthocyanin levels were lower than values previously reported for American and Mexican blue corn [21]. Antioxidant activity evaluated with the DPPH method showed values between 18.5 and 26.8 µmol/100 g.

Results for the total content of polyphenols, monomeric anthocyanins, and antioxidant activity were plotted in a polygons graph in order to identify the sample with the best characteristics. **Figure 2** shows that sample CIIDIR-125 had the largest content of anthocyanins and antioxidant activity; therefore, it was selected to undergo the anthocyanin profile analysis and biological tests.

#### 3.2. Anthocyanin profile of blue corn extract

**Figure 3** shows the profile of anthocyanins isolated from blue corn using amberlite XAD-resin. The MS data analysis for blue corn anthocyanins is summarized in **Table 2**. It shows ions at  $m/z = 287$  and 271, suggesting that anthocyanins are derived mainly from cyanidin and pelargonidin. Eight anthocyanins were identified such as: cyanidin-3-(3'',6''-dimalonyl-glucoside), pelargonidin-3-glucoside dimalonate, pelargonidin-3-(sinapoyl glucoside)-5-glucoside,

Sample	Total polyphenols <sup>1</sup>	Monomeric anthocyanins <sup>2</sup>	Antioxidant activity <sup>3</sup>
CIIDIR-02	154.4 <sup>f,g,h,i</sup>	32.5 <sup>de2</sup>	18.5 <sup>e1</sup>
CIIDIR-12	176.7 <sup>c,d</sup>	48.5 <sup>b</sup>	21.2 <sup>d,e</sup>
CIIDIR-54	158.4 <sup>f,g</sup>	31.2 <sup>e</sup>	18.6 <sup>e</sup>
CIIDIR-107	173.3 <sup>d,e</sup>	53.4 <sup>b</sup>	26.7 <sup>a,b,c</sup>
CIIDIR-112	142.8 <sup>i</sup>	30.7 <sup>e</sup>	18.5 <sup>e</sup>
CIIDIR-125	203.2 <sup>a</sup>	66.9 <sup>a</sup>	24.4 <sup>ab,c,d</sup>
CIIDIR-129	164.3 <sup>e,f</sup>	47.5 <sup>b,c</sup>	23.6 <sup>b,c,d</sup>
CIIDIR-131	173.4 <sup>d,e</sup>	53.4 <sup>b</sup>	27.8 <sup>a</sup>
CIIDIR-167	187.1 <sup>b,c</sup>	32.3 <sup>e</sup>	24.4 <sup>a,b,c,d</sup>
CIIDIR-172	147.4 <sup>g,h,i</sup>	28.6 <sup>ef</sup>	22.7 <sup>c,d</sup>
CIIDIR-179	162.1 <sup>f</sup>	30.7 <sup>e</sup>	21.5 <sup>d,e</sup>
CIIDIR-184	146.5 <sup>ij</sup>	21.4 <sup>f</sup>	20.6 <sup>d,e</sup>
CIIDIR-185	192.9 <sup>a,b</sup>	35.1 <sup>de</sup>	26.8 <sup>a,b</sup>
CIIDIR-189	157.7 <sup>f,g,h</sup>	40.3 <sup>cd</sup>	22.8 <sup>c,d</sup>
CIIDIR-190	148.1 <sup>g,h,i</sup>	33.1 <sup>de</sup>	23.5 <sup>b,c,d</sup>
CIIDIR-197	175.3 <sup>d</sup>	35 <sup>de</sup>	26.7 <sup>a,b,c</sup>

Samples with the same letters are not significant statistically ( $p < 0.05$ ).

<sup>1</sup>mg GAE/100g.

<sup>2</sup>mg C3G/100 g.

<sup>3</sup>μmol TE/g.

**Table 1.** Content of total polyphenols, monomeric anthocyanins and antioxidant activity in blue corn samples.

pelargonidin 3-(3'',6''-dimalonylglucoside), pelargonidin 3-(6''-malonyl glucoside)-5-(6'' acetyl glucoside), pelargonidin 3-glucoside-5-(6''-acetyl-glucoside), pelargonidin 3-(6''-malonylglucoside), and pelargonidin 3,5-diacetylglucoside. The data indicates that only acylated anthocyanins are present in blue corn from the Mixteco race. This could be due to genetic factors, farming practices, weather conditions, and soil type, which have an influence on the chemical composition of maize varieties.

### 3.3. Cytotoxic activity of the anthocyanin fraction from blue corn

In this study, the SRB assay was used to evaluate cytotoxic activity, and it was selected in order to avoid any interference of anthocyanins in the final reading. The effect of the blue corn extract at different concentrations on the prostate cancer cell line (PC3), neoplastic myelogenous leukemic cell line (K562), human colon cancer cell line (HCT-15), human breast cancer cell line (MCF-7), and lung cancer (SKLU-1) are shown in **Figure 4**. Generally speaking, it was observed that for blue maize extract, the percentage of growth inhibition of cancer cell lines improved with increased concentration; the analysis indicates that the blue corn extract causes growth inhibition in all cancer cell lines in a dose-dependent manner (**Figure 4**).




Figure 2. Polygon graph of total polyphenols, monomeric anthocyanins and antioxidant activity.




Figure 3. Anthocyanin profile of blue corn.

PC3 cells were selected due to their highly aggressive nature. Data showed 2.43% inhibition on prostate cancer cells at 500  $\mu\text{g}/\text{mL}$ . Previous studies have analyzed the anticancer properties of the anthocyanin fraction from potato extracts in prostate cancer cells (PC-3) showing cytotoxicity [22]. It has been reported that the anthocyanin profile has an effect on anticancer activity. For example, pomegranate extract has an abundance of delphinidin derivatives, a compound with anticancer activity on human prostate [23]. In the present




Peak	Retention time (min)	<i>m/z</i>	Tentative identification
1	1.2	621–287	Cyanidin-3-(3'', 6''-dimalonyl-glucoside)
2	1.9	358–271	Pelargonidin-3-glucoside dimalonate
3	2.8	639–271	Pelargonidin-3-(sinapoyl glucoside)-5-glucoside
4	3.4	605–271	Pelargonidin 3-(3'',6''-dimalonylglucoside)
5	3.9	740–519–433–271	Pelargonidin 3-(6''-malonyl glucoside)-5-(6'' acetyl glucoside)
6	5.0	654–595–434–271	Pelargonidin 3-glucoside-5-(6''-acetyl-glucoside)
7	6.6	519–271	Pelargonidin 3-(6''-malonylglucoside)
8	8.4	595–271	Pelargonidin 3,5-diacetylglucoside

**Table 2.** Chromatographic and mass spectral data of anthocyanins.

research, the anthocyanin profile of blue corn was composed only of cyanidin and pelargonidin derivatives; delphinidin was not detected.

In addition, the analysis indicates that the blue corn extract showed higher inhibition of cellular growth in MCF-7 cancer cells than SKLU-1 (**Figure 4**) at the same concentration



**Figure 4.** *In vitro* cytotoxicity of blue corn extract on several human cancer lines.

(500 µg/mL). Anthocyanin-rich extracts of cereals such as black rice and black sorghum have also showed the anticancer effects on MCF-7 cells. On the other hand, reports on the effects of anthocyanins on SKLU-1 cells are scarce; a study performed on kenaf seed extract showed cytotoxic activity toward SKLU-1 [24].

Interestingly, blue corn extract was able to inhibit 50.9% of lung cancer cells at 500 µg/mL, which suggests a potential for application in lung cancer treatment, one of the five cancer types most frequently diagnosed in male population worldwide; however, studies *in vivo* (animal experiments) and clinical trials are needed. In this regard, studies on blueberries report the presence of anthocyanins in lung tissue of mice fed with this anthocyanin-rich fruit (5% w/w) for 10 days, suggesting that fruits, vegetables, and cereals such as blue corn may be an important source of chemopreventive dietary components [25].

Likewise, blue corn extract also inhibited the growth of neoplastic myelogenous leukemic cells (K562) and colon cancer (HCT-15) cell lines on 46.7 and 62 % at 500 µg/mL, respectively. Given the above, the blue corn extract was more effective on the growth inhibitory activity on HCT-15 colon cancer cell lines as compared to other cancer cell lines. Current statistics indicate that in 2012 colorectal cancer was the third most common cancer in the world. For this reason, there is an increasing interest for chemoprevention as a cancer prevention strategy. Dietary agents such as anthocyanins have been explored for their chemopreventive effects against colon cancer [26]. *In vitro* data obtained in this research provides information for the future application of blue corn extract as a chemopreventive agent in colon cancer.

In summary, blue corn possesses antioxidant properties and its anthocyanin profile is constituted solely by acylated anthocyanins. These results are particularly important since corn is the basis of the Mexican diet; they suggest that corn anthocyanins may have anticancer activity. Further research is necessary to obtain deeper knowledge on specific molecular targets of blue corn and to ensure the safe use of these active compounds as therapeutic agents on lung and colon cancer.

## Acknowledgements

The authors would like to thank the SINAREFI (BEI-MAI-10-32) from México.

## Author details

Rosa Guzmán-Gerónimo<sup>1\*</sup>, Edna Alarcón Aparicio<sup>1</sup>, Oscar García Barradas<sup>2</sup>, Jose Chávez-Servia<sup>3</sup> and Tania Alarcón-Zavaleta<sup>1</sup>

\*Address all correspondence to: [roguzman@uv.mx](mailto:roguzman@uv.mx)

1 Basic Sciences Institute, University of Veracruz, Xalapa, Veracruz, México

2 Services in Analytical Resolution, SARA, University of Veracruz, Xalapa, Veracruz, México

3 The Interdisciplinary Research Center for Integrated Regional Development, Oaxaca Unit, National Polytechnic Institute, Oaxaca, México

## References

- [1] Kotesha R, Takami A, Espiniza JL. Dietary phytochemicals and cancer chemoprevention: a review. *Oncotarget*. 2016;**7**:52517-52529. DOI: 10.18632/oncotarget.9593
- [2] Mehta M, Shike M. Diet and physical activity in the prevention of colorectal cancer. *J Natl Compr Canc Netw*. 2014;**12**:1721-1726.
- [3] Howes MJ, Simmonds MS. The role of phytochemicals as micronutrients in health and disease. *Curr Opin Clin Nutr*. 2014;**17**:558-566. DOI: 10.1097/MCO.000000000000115.
- [4] Li H, Li HQ, Wang Y, Xu HX, Fan WT, Wang ML, Sun PH, Xie XY. An intervention study to prevent gastric cancer by micro-selenium and large dose of allitridum. *Chin Med J*. 2004;**117**:1155-1160.
- [5] Priyadarsini RV, Nagini S. Cancer chemoprevention by dietary phytochemicals: promises and pitfalls. *Curr Pharm Biotechnol*. 2012;**13**:125-136. DOI: 10.2174/138920112798868610
- [6] Wang LS1, Stoner GD. Anthocyanins and their role in cancer prevention. *Cancer Lett*. 2008;**269**:281-290. DOI: 10.1016/j.canlet.2008.05.020
- [7] Delgado-Vargas F, Jiménez AR, Paredes-López O. Natural pigments: carotenoids, anthocyanins, and betalains—characteristics, biosynthesis, processing, and stability. 1st ed. Boca Raton, Florida: CRC Press; 2003. 313 p. DOI: 10.1080/10408690091189257
- [8] Goufo, Trindade H. Rice antioxidants: phenolic acids, flavonoids, anthocyanins, proanthocyanidins, tocopherols, tocotrienols,  $\gamma$ -oryzanol, and phytic acid. *Food Sci Nutr*. 2014;**2**:75-104. DOI: 10.1002/fsn3.86
- [9] Awika JM1, Rooney LW.. Sorghum phytochemicals and their potential impact on human health. *Phytochemistry*. 2004;**65**:1199-1221. DOI: 10.1016/j.phytochem.2004.04.001
- [10] Guzmán-Gerónimo RI, Alarcón-Zavaleta TM, Oliart-Ros RM, Meza-Alvarado JE, Herrera-Meza S, Chávez-Servia JL. Blue maize extract improves blood pressure, lipid profiles, and adipose tissue in high-sucrose diet-Induced metabolic syndrome in rats. *J Med Food*. Forthcoming. DOI: 10.1089/jmf.2016.0087
- [11] Kato T, Mapes C, Mera, Serratos J, Bye R. Origin and diversification of maize: an analytical review. National Autonomous University of Mexico. 1st ed. Mexico, D.F.: CONABIO; 2009. 116 p.
- [12] Salinas-Moreno Y, Pérez-Alonso JJ, Vázquez-Carrill, G, Aragón-Cuevas F. Anthocyanins and antioxidant activity in maize grains (*Zea mays* L.) of chalqueño, elotes cónicos and bolita races. *Agroc*. 2012;**46**:693-706.
- [13] Salinas MY, Salas SG, Rubio HD, Ramos LN. Characterization of anthocyanin extracts from maize kernels. *J Chromatogr Sci*. 2005;**43**:483-487.
- [14] Ross NT, Wilson CJ. In vitro clinical trials: the future of cell-based profiling. *Front Pharmacol*. 2014;**5**:1-6. DOI: 10.3389/fphar.2014.00121

- [15] Singleton VL, Rossi JA. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. *Am J Enol Vitic.* 1965;**16**:144-150.
- [16] Giusti MM, Wrolstad RE. Characterization and measurement of anthocyanins by UV-visible spectroscopy. In: Wrolstad RE, Acree, TE; An H, Decker EA, Penner MH, Reid DS, Schwartz SJ, Shoemaker CF, Sporns P. (Eds.). *Current protocols in food analytical chemistry.* New York, N.Y.: John Wiley and Sons, Inc.; 2001. p. F1.2.1-F1.2.9. DOI: 0.1002/0471142913
- [17] Brand-Williams W, Cuvelier ME, Berset C. Use of a free radical method to evaluate antioxidant activity. *LWT—Food Sci Techn.* 1995;**28**:25-30. DOI: 10.1016/S0023-6438(95)80008-5
- [18] Welch CR, Wu Q, Simon JE. Recent advances in anthocyanin analysis and characterization. *Curr Anal Chem.* 2008;**4**:75-101. DOI: 10.2174/157341108784587795
- [19] Skehan P1, Storeng R, Scudiero D, Monks A, McMahon J, Vistica D, Warren JT, Bokesch H, Kenney S, Boyd MR. New colorimetric cytotoxicity assay for anticancer-drug screening. *J Natl Cancer Inst.* 1990;**82**:1107-1112.
- [20] Dao LT, Takeoka GR, Edwards RH, Berrios JDJ. Improved method for the stabilization of anthocyanidins. *J Agric Food Chem.* 1998;**46**:3564-3569.
- [21] Del Pozo-Insfran D, Brenes CH, Serna SO, Talcott ST. Polyphenolic and antioxidant content of white and blue corn (*Zea mays* L.) products. *Food Res Int.* 2006;**39**:696-703. DOI: 10.1016/j.foodres.2006.01.014
- [22] Reddivari L, Vanamala J, Chintharlapalli S, Safe SH, Miller JC. Anthocyanin fraction from potato extracts is cytotoxic to prostate cancer cells through activation of caspase-dependent and caspase-independent pathways. *Carcinog.* 2007;**28**:2227-2235. DOI: 10.1093/carcin/bgm117
- [23] Hafeez BB, Siddiqui IA, Asim M, Malik A, Afaq F, Adhami VM, Saleem M, Din M, Muhtar H. A dietary anthocyanidin delphinidin induces apoptosis of human prostate cancer PC3 cells in vitro and in vivo: involvement of nuclear factor-kappaB signaling. *Cancer Res.* 2008;**68**:8564-8572. DOI: 10.1158/0008-5472.CAN-08-2232
- [24] Wong YH, Tan WY, Tan CP, Long K., Nyam KL. Cytotoxic activity of kenaf (*Hibiscus cannabinus* L.) seed extract and oil against human cancer cell lines. *Asian Pac J Trop Biomed.* 2014;**4**:S510-S515. DOI: 10.12980/APJTB.4.2014C1090
- [25] Aqil F, Vadhanam MV, Jeyabalan J, Cai J, Singh IP, Gupta RC.. Detection of anthocyanins/anthocyanidins in animal tissues. *J Agric Food Chem.* 2014;**62**:3912-3918. DOI: 10.1021/jf500467b
- [26] Tramer F, Moze S, Ademosun AO, Passamonti S, Cvorovic J. Dietary anthocyanins: impact on colorectal cancer and mechanisms of action. In: Rajunor Ettarh, editor. *Colorectal cancer—from prevention to patient care.* 1st ed. Rijeka, Croatia: Intech; 2012. pp. 123-158. DOI: 10.5772/27678

