

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

**4,800**

Open access books available

**122,000**

International authors and editors

**135M**

Downloads

Our authors are among the

**154**

Countries delivered to

**TOP 1%**

most cited scientists

**12.2%**

Contributors from top 500 universities



**WEB OF SCIENCE™**

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Current Research of the Renin-Angiotensin System Effect on Stem Cell Therapy

---

Elham Ahmadian, Aziz Eftekhari and  
Ahmad Yari Khosroushahi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67726>

---

## Abstract

The renin-angiotensin system (RAS) is a chief regulator of the cardiovascular system and body fluid homeostasis. Stem/progenitor cell therapy has pointed towards a novel tool for medical and therapeutic intervention. In addition to the physiological and pathological role of the RAS and its pharmacological inhibitors, the proliferation, differentiation in stem cells is mediated through various cell-signalling pathways. This book chapter reviews the new role of RAS components, distinct from other common roles by considering its regulating impact on the several signalling pathways involved in different body tissues, as well as in stem cell therapy.

**Keywords:** stem cell, progenitor cell, renin-angiotensin system, pancreatic stem cells, cardiac stem cells

---

## 1. Introduction

The concept that the renin-angiotensin system (RAS) is involved in the regulation of stem (progenitor) cell function is novel. This is beyond the conventional notion of the RAS acting as a potent vasoconstrictor responsible for blood pressure regulation and body fluid homeostasis. The expression of RAS components during human embryonic development has been addressed in the literature. The existence of RAS components in different organs and tissues suggests the presence of local RAS in addition to the circulating common RAS, which has paracrine effects mediating stem (progenitor) cell function. Moreover, recent evidence has shown the expression of major RAS components such as angiotensinogen, renin, angiotensin-converting enzyme (ACE), angiotensin receptors type 1 and 2 and angiotensin-(1–7)

---

during growth, proliferation and differentiation of stem cells. Improvement of the stem cell functionality and making them ideal candidates in different kinds of disorders has been a new research field in the last decade. Meanwhile, the effect of RAS on stem cell growth, proliferation and function is an emerging attempt among researchers. Ang II receptor activation increases the proliferation of several progenitor cells, such as mouse bone marrow-derived stem cells and human cord blood cells. Accordingly, manipulation of the RAS may alter and/or have beneficial effects on the efficacy of stem cell therapy.

## 2. Stem cells

The term stem cell stands for the population of immature precursor cells, which are able to renew themselves and be the source of de novo replacement for many body tissues. Stem cells are classified into two main groups: embryonic stem cells (ESCs) and adult stem cells. ESCs can be obtained from the inner cell mass of the embryonal blastocyst. Although they are easily achieved, some disadvantages restrict their application. Adult stem cells such as mesenchymal and haematopoietic stem cells (HSCs) are obtained from mature tissues. Due to their plasticity, adult stem cells produce cell lineage different from their original organ. Thus, adult stem cells seem to be an appropriate candidate for organ regeneration in different kinds of diseases or lost/damaged organs. **Table 1** represents the main stem cell and their advantages and disadvantages.

Stem cell type	Origin	Advantages	Disadvantages
Embryonic stem cell	Blastocyst stage of an embryo	- High expansion - Pluripotent	- Ethical objection - Risk of rejection - Risk of teratocarcinoma
Adult stem cell	Mature tissue	- Easily obtained - No ethical objection - High compatibility	- Lack of specific identification markers

**Table 1.** Main stem cell and their advantages and disadvantages.

## 3. The RAS and ESCs

ESCs are pluripotent cells capable of differentiation into different cells such as cardiomyocytes, and endothelial cells have been considered as a source of regenerative medicine [1]. For instance, ESC-derived endothelial cells have therapeutic effects via the increment of angiogenesis and heart functionality [2]. PI3/Akt-signalling pathway has been shown to be linked with human ESC-derived cardiomyocyte proliferation in vitro [3]. RAS stimulation activates PI3/AKT pathway, while the inhibition of RAS increases Akt phosphorylation [4], which

might influence the proliferation of ESCs. High survival rate after transplantation is another main, noteworthy issue about ESCs [5].

RAS is a novel regulatory candidate, which controls the development of ESCs into different cell types. It has been reported that the expression AT1 receptors were detected in an early stage of human ESCs differentiation. Since the addition of Ang II results in the phosphorylation of extracellular signal-regulated kinase (ERK) 1/2 and Jun N-terminal (JNK) 1/2, this peptide is capable of acting as signalling molecules and thus it could regulate differentiation [6]. Moreover, Ang II has been shown to increase glucose uptake in ESCs [7] and induce mitogenic effect, possibly through protein kinase C and mitogen-activated protein kinase (MAPK)-signalling pathways. Interestingly, exposure to a high glucose niche in the presence of Ang II has been shown to produce a synergistic impact on ESC proliferation [8].

#### **4. The RAS and mesenchymal stem cells**

Mesenchymal stem cells (MSCs) as multipotent cells are mainly found in bone marrow and adipose tissue and can differentiate into various cell types [9]. Simple isolation, high immune prevalence and angiogenic-inducing properties have made MSCs suitable candidates for stem cell therapy of different kinds of diseases [10]. Besides, these MSCs exert paracrine effects causing the modulation of a large number of cellular responses, such as survival, proliferation, migration and gene expression [11]. Diminishing oxidative stress and suppression of the TGF- $\beta$ /Smad2-signalling pathway are some of these paracrine effects [12, 13]. In the rat pulmonary hypertension model, MSCs have shown superiority regarding the lowering of blood pressure and ventricular overload; hence, MSC transplantation in chronic lung disease with pulmonary hypertension has pointed towards a new therapeutic option [14]. Concomitant percutaneous trans-luminal renal angioplasty with MSC therapy has been reported to decrease inflammation, fibrinogenesis and vascular remodelling in atherosclerotic renal artery stenosis of swine [15]. Also, MSCs have the ability to recognize inflammation lesions, and Ang II effects the migration and homing of these cells to the site of injury [16]. Dysregulation of the RAS decreases the paracrine therapeutic potential of MSCs [17].

Expression of renin, AT1 and AT2 receptors has been implicated with the regulation of MSCs differentiation to adipocytes, while the differentiated cell produces significant amounts of endogenous RAS [18]. Indeed, endogenous blockade of AT1 receptor inhibited adipogenesis of MSCs [18]. These outcomes are in line with clinical observations that RAS blockade acts as a protective factor against the onset of obesity-induced diabetes mellitus type 2 [19]. Ang II also has been shown to stimulate the synthesis of vascular endothelial growth factor (VEGF) that is an angiogenic agent in MSCs [20]. Additionally, MSCs have been suggested as a promising regenerative medicine for treating ischaemic heart disease and diabetes [21]. Hence, it is probable that Ang II-induced production of VEGF might be a contributing underlying mechanism of the beneficial consequences obtained following MSC transplantation.

## 5. The RAS and endothelial progenitor cells

The identification of circulating endothelial progenitor cells (EPCs) has introduced the concept of postnatal vasculogenesis. EPCs could originate from haematopoietic stem cells (HSCs) or MSCs [22, 23]. Also, the EPCs existing in the adventitial layer of vessels have the ability to differentiate into adult endothelial cells [24]. Different factors such as ischaemia, vascular damage and even physical exercise result in the recruitment of circulating EPCs and thus neovascularization and restoration of endothelial functionality [25, 26]. In this context, the improvement of myocardial perfusion after EPC transplantation has been observed in clinical trials [27]. Several mechanisms have been suggested regarding EPCs mobilization. For instance, it was observed that ischaemic lesions release angiogenic factors like VEGF and activate MAPK or the RAS-signalling pathways [28], which increase EPCs migration.

Despite the important role of vascular endothelium in cardiovascular disease (CVD), their limited regeneration capacity remains a vital problem. EPCs improve angiogenesis and participate in endothelium recovery subsequent to vascular injuries [29]. Cardiovascular diseases (CVDs) are directly related to both the decline of EPC mobilization and the number of EPCs present in the damaged site. In this context, Ang II stimulates EPCs migration to ischaemic regions and commences vascularization through VEGF-associated endothelial nitric oxide synthase [30]. The activation of NADPH and subsequent ROS (reactive oxygen species) generation constitutes the stimulatory impact of Ang II on EPCs that is required for normal EPC function. However, the long-term activation of NADPH and oxidative stress is concomitant with cell senescence [31]. Moreover, acute high-dose exposure to Ang II has been shown to negatively modulate EPC function in the hind limb ischaemic rat model [32].

## 6. The vascular RAS and erythropoiesis

RAS has been shown to result in progenitor cell senescence and suppression of differentiation and adherence in bone marrow-derived EPCs in Ang II infusion models. This inhibitory impact could be attenuated by the administration of AT1 receptor antagonists [31]. Previous reports have proved the crucial role of Ang II during erythropoiesis [33]. In studies using transgenic mice expressing human renin and angiotensinogen, a drastic rise in levels of erythropoietin was observed, which is a glycoprotein hormone that controls erythropoiesis. Genetic ablation of AT1 receptor from these mice reduced erythropoietin levels and restored haematocrit levels [34]. Also, ACE blockade has been concomitant with haematocrit decrease *in vivo* [35]. The idea of ACE and/or Ang II being contributed to erythropoiesis was further confirmed by a recent research in which ACE marked haematopoietic stem cells from human embryonic, fetal and adult haematopoietic tissues [36]. However, the mechanism of Ang II-associated regulation of erythropoiesis is mainly unclear. Most of these effects are observed during early phases of erythropoiesis [37]. As mentioned above, some researchers imply that Ang II acts indirectly via its effect on erythropoietin levels [38], whereas others do not agree with this link [39]. The other possible mechanism is proposed to be the involvement of JAK (Janus kinase)/STAT (signal transducer and activator of transcription) pathway. JAK/STAT pathway is known to be activated by Ang II [40].

## 7. Current research on the RAS in pancreatic stem cells

The local RAS is not only involved in the physiology of pancreas, but it also influences the pancreatic stem cell (PSC) functionality. RAS has been shown to be associated with pancreatic islet cell function and proliferation and differentiation of PSCs/progenitor cells during development [41]. Different stem/progenitor cells have been reported to be differentiated into insulin-expressing cells, which make them appropriate candidates for islet cell transplantation. Regarding the potential role of RAS in stem cell differentiation, it is possible that RAS-modulated stem cell could be a new source of pancreatic  $\beta$ -cells. Both exocrine and endocrine pancreas are known to have local RAS components [42]. In exocrine part, AT1 receptor activation turns on signalling pathways such as ROS generation and activation of pro-inflammatory, vasoactive and growth factor receptors [43, 44]. Therefore, Ang II might result in fibrosis and inflammation of exocrine pancreas through the AT1 receptor. Hence, blockade of RAS has been considered a potential therapeutic opportunity for some pancreas disorders.

In the endocrine portion of pancreas, RAS has been shown to be a key regulator of insulin and islet physiology [43]. AT1 receptor stimulation leads to  $\beta$ -cells, decreased islet blood flow and insulin secretion, while AT2 receptor activation results in  $\beta$ -cell proliferation and islet blood flow and insulin secretion enhancement [19]. Moreover, the ACE2/Ang-(1-7)/Mas axis, which has been attracting more research attention recently, is present in several local tissues and mainly acts as a negative modulator of ACE/Ang II/AT1R signalling. Similar to AT2 receptor activation, ACE2 overexpression in the pancreas of type 2 diabetic animals restored glucose homeostasis, as evidenced by diminished blood glucose levels, elevated insulin secretion and  $\beta$ -cell proliferation [45].

PSCs exist in both developing and adult pancreas in three major pancreas sections, that is, ductal endothelium, islet and acinar tissues [46]. Embryo, foetus and adult pancreas as well as bone marrow-derived MSCs are probable sources for PSCs. Transplantation of mouse or human PSCs into diabetic mice has been revealed to reduce their diabetes [46].

A novel well-defined area of research is the developmental control of RAS on cell proliferation in tumours and in tissue regeneration. Both the ACE/AngII/AT1R signalling and the alternative RAS arm (ACE2/Ang-(1-7)/Mas) interact with different growth factors; hence, they might contribute to cell proliferation and angiogenesis in neoplasms, including pancreatic cancers [47-49]. It has been demonstrated that RAS inhibition seems to be a promising therapeutic approach for the mitigation of pathophysiological circumstances of the pancreas including diabetes [50], pancreatitis [43] and pancreatic cancer [51]. Transplantation of human fetal pancreatic progenitor cell has been shown to reverse hyperglycaemia and glucose intolerance in diabetic mice [52]. ROS production has a close relation with RAS activation, and ROS-signalling pathway is associated with stem/progenitor cell proliferation, differentiation and function [53]. So, it is an interesting probability that the elevation of RAS-induced differentiation of pancreatic progenitor cells towards an endocrine lineage might offer a basis for therapy in terms of islet replacement treatments for diabetes.

MSCs have been suggested as an appropriate substitute to islet transplantation for promoting regeneration of endogenous pancreatic progenitor cells to achieve permanent normal blood glucose level in patients with type 1 diabetes [54]. Local RAS in pancreatic islet could regulate PSC differentiation and thus lead to the beneficial outcomes following MSC transplantation. In a study, these kinds of pancreatic progenitor cells have shown to differentiate into insulin-secreting cells.

RAS components like angiotensinogen and renin are expressed after the beginning of pancreatic progenitor cells differentiation, but they are not present in undifferentiated cells. These results indicate that a functional RAS exists in pancreatic progenitor cells and in mature islets that could be modulating cellular differentiation. The mitogenic behaviour related to the Ang II bindings of AT1 receptors has been proposed to regulate reprogramming of pancreatic cells and the differentiation plasticity [55]. However, it is unclear whether AT2 receptor activation reveals counter-regulatory role in this context. Furthermore, it is hypothesized that the ACE2/Ang-(1-7)/Mas axis plays an essential role in pancreatic stem cell differentiation as previous studies have shown the involvement of ACE2 arm in the proliferation and differentiation of other stem cells [56].

## 8. Current research on the RAS in cardiac stem cells

Regarding the intracellular signalling pathway of Ang II, RAS effect on cardiovascular stem/progenitor cell transplantation has largely been investigated. Among the regenerative medicine-based therapies in the cardiovascular system, induced pluripotent stem cells (iPSCs), which are artificially derived from an adult non-differentiated somatic cell, are a field of research study. In spite of different origin, they resemble ESCs in their growth and gene expression profile [57]. Also, Ang II receptors are expressed in iPSCs, which induce the proliferation and differentiation of pluripotent stem cells to several kinds of stem cells. As mentioned before, Ang II stimulates cell-signalling cascade through ROS production which in turn instigates stem cell proliferation [58]. In a study, the administration of Tempol (ROS generation-blocking agent) in Ang II-treated pluripotent stem cells has attenuated the proliferation of stem cells and DNA synthesis suggesting the role of oxidative-signalling pathway in RAS-associated cell proliferation. The other signalling pathway linked to the differentiation of iPSCs and Ang II is JAK/STAT pathway [59].

Ang II is also able to induce ESCs differentiation. In this context, the effect of AT1 receptor activation on collagen IV protein has been investigated [18]. Collagen IV is an extracellular matrix protein having a role in cell adhesion, growth, migration and differentiation. Collagen IV has been shown to be involved in the differentiation of ESCs to smooth muscle cell.

Up-regulation of several transcription factors such as *egr-1*, *c-fos/c-jun*, *Stat91*, *NFκ-B*, which has a fundamental role in stem cell differentiation, is mediated through PI3/Akt pathway. Ang II is the upstream cascade of PI3/Akt [82-84]. *NFκ-B* is markedly up-regulated in Ang II-treated cells, proposing that there is *NFκ-B* involvement in ESC differentiation into the smooth muscle cells [60].

The TGF- $\beta$ /Smad pathway plays a key role in the cellular responses to Ang II. Ang II activates TGF- $\beta$  secretion in various tissues, such as fibroblasts and smooth muscle cells that induce interstitial fibrosis in the heart and kidney. Besides, TGF- $\beta$ /Smad pathway is highly engaged to vascular fibrosis and arteriosclerosis [61] and gives rise to the differentiation of MSCs to smooth muscle cell. Furthermore, TGF- $\beta$  secretion is connected with the MAPK/ERK cascade, and Ang II in this pathway interferes with TGF- $\beta$  production, thus leading to the differentiation of MSCs to smooth muscle cells [62].

Regarding owning various paracrine effects, MSC transplantation has gained great importance in cardiovascular disease [63]. The supportive effects of vascular VEGF have been recognized in the migration, invasion of extracellular matrix, proliferation, survival of MSCs, and they contribute to MSCs' paracrine effects [64, 65]. In this context, all pathways increasing VEGF would give rise to the function of MSCs. Ang II increases VEGF mRNA and protein expression in MSCs [20], which is associated with Akt-signalling pathway. Pre-treatment of MSCs with the Akt inhibitor (LY292002) has been shown to reduce Ang II-induced VEGF expression. So, local Ang II, as a cytokine, might boost VEGF generation in MSC grafts and upgrade the transplantation effectiveness.

The excess RAS expression is detected in CVDs such as myocardial infarction, hypertension, heart failure and atherosclerosis [66]. On the other hand, RAS inhibition via ACE inhibitors (ACEIs) and angiotensin receptor blockers (ARBs) has been widely examined in cardiovascular disease beyond their effects in lowering the blood pressure [67]. Despite the daily increasing use of regenerative medicine in treating different disorders, the functionality of stem cell transplantation is not sufficient in animal models and clinical trials [68].

Therefore, clarifying the mechanisms that enhance the graft efficacy seems important. Researchers have investigated the effect of RAS inhibition on stem cell therapy of cardiovascular system [69].

The insulin-like growth factor 1-1 secreted from stem cells has a close relationship with the RAS and down-regulates the local RAS through the attenuation of the p53 gene [70]. The IGF-1 has an anti-apoptotic effect on cardiomyocytes in ischaemic heart disease and also enhances differentiation and survival of stem cells after transplantation [71]. In acute MI in cardiomyocytes, ACEIs up-regulate the IGF-1 receptors; thus, the concurrent use of perindopril in bone marrow stem cell transplantation increases the paracrine effects of the IGF-1, which abolishes apoptosis through increased Bcl2 expression and improves cardiac function [72]. Also, pre-treatment of MSCs with ARBs before transplantation increases their trans-differentiation efficacy and also improves the systolic function of the heart [73].

## 9. Limitations and future directions

Comprehensive elucidation of the complexity of the regulatory network that drives stem cell therapy will require extensive effort and time. The accretion of daily increasing research and obtained ideas will undoubtedly assist the current research field of stem/progenitor cell



therapy. Regenerative progenitor cell therapy has emerged as a possible alternative for pharmacotherapy in different human diseases. A major problem in this field is insufficient efficacy during stem cell transplantation. In order to improve the efficiency of regenerative medicine, researchers examined the impact of the modulation of various cell-signalling pathways, including the RAS. Effects of Ang II in stem cell proliferation and differentiation have been documented in the literature. The presence of the RAS components in progenitor cells and many tissues may regulate growth and development and thus might contribute to the preparation of various progenitor cells for clinical transplantation.

## Acknowledgements

The moral support of the Faculty of Pharmacy and Drug Applied Research Center of Tabriz University of Medical Sciences in Tabriz, Iran, is gratefully acknowledged.

## Author details

Elham Ahmadian<sup>1,2,3</sup>, Aziz Eftekhari<sup>1,2</sup> and Ahmad Yari Khosroushahi<sup>2,4\*</sup>

\*Address all correspondence to: yarikhosroushahia@tbzmed.ac.ir

1 Department of Pharmacology and Toxicology, Tabriz University of Medical Science, Tabriz, Iran

2 Drug Applied Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

3 Student Research Committee, Tabriz University of Medical Science, Tabriz, Iran

4 Department of Pharmacognosy, Tabriz University of Medical Sciences, Tabriz, Iran

## References

- [1] Mummery, C.L., et al., Differentiation of human embryonic stem cells and induced pluripotent stem cells to cardiomyocytes a methods overview. *Circulation Research*, 2012. **111**(3): pp. 344-358.
- [2] Lundy, S.D., et al., Structural and functional maturation of cardiomyocytes derived from human pluripotent stem cells. *Stem Cells and Development*, 2013. **22**(14): pp. 1991-2002.
- [3] McDevitt, T.C., Laflamme, M.A., and Murry, C.E., Proliferation of cardiomyocytes derived from human embryonic stem cells is mediated via the IGF/PI 3-kinase/Akt signaling pathway. *Journal of Molecular and Cellular Cardiology*, 2005. **39**(6): pp. 865-873.
- [4] Cheng, W.H., et al., Renin activates PI3K-Akt-eNOS signalling through the angiotensin AT1 and Mas receptors to modulate central blood pressure control in the nucleus tractus solitarii. *British Journal of Pharmacology*, 2012. **166**(7): pp. 2024-2035.

- [5] van Laake, L.W., et al., Human embryonic stem cell-derived cardiomyocytes survive and mature in the mouse heart and transiently improve function after myocardial infarction. *Stem Cell Research*, 2007. **1**(1): pp. 9-24.
- [6] Huang, Z., et al., Angiotensin II type 1 and bradykinin B2 receptors expressed in early stage epithelial cells derived from human embryonic stem cells. *Journal of Cellular Physiology*, 2007. **211**(3): pp. 816-825.
- [7] Han, H.J., et al., ANG II-stimulated DNA synthesis is mediated by ANG II receptor-dependent Ca<sup>2+</sup>/PKC as well as EGF receptor-dependent PI3K/Akt/mTOR/p70S6K1 signal pathways in mouse embryonic stem cells. *Journal of Cellular Physiology*, 2007. **211**(3): pp. 618-629.
- [8] Kim, Y.H. and Han, H.J., Synergistic effect of high glucose and ANG II on proliferation of mouse embryonic stem cells: involvement of PKC and MAPKs as well as AT1 receptor. *Journal of Cellular Physiology*, 2008. **215**(2): pp. 374-382.
- [9] Trivedi, P., et al., Mesenchymal stem cell therapy for treatment of cardiovascular disease: helping people sooner or later. *Stem Cells and Development*, 2010. **19**(7): pp. 1109-1120.
- [10] Jiang, S., et al., Transcriptional profiling of young and old mesenchymal stem cells in response to oxygen deprivation and reparability of the infarcted myocardium. *Journal of Molecular and Cellular Cardiology*, 2008. **44**(3): pp. 582-596.
- [11] Choi, Y.-H., Kurtz, A., and Stamm, C., Mesenchymal stem cells for cardiac cell therapy. *Human Gene Therapy*, 2010. **22**(1): pp. 3-17.
- [12] Eirin, A., et al., Adipose tissue-derived mesenchymal stem cells improve revascularization outcomes to restore renal function in swine atherosclerotic renal artery stenosis. *Stem Cells*, 2012. **30**(5): pp. 1030-1041.
- [13] Ahmadian, E., Jafari, S., and Khosroushahi, A.Y., Role of angiotensin II in stem cell therapy of cardiac disease. *Journal of Renin-Angiotensin-Aldosterone System*, 2015. **16**(4): pp. 702-711.
- [14] Hansmann, G., et al., Mesenchymal stem cell-mediated reversal of bronchopulmonary dysplasia and associated pulmonary hypertension. *Pulmonary Circulation*, 2012. **2**(2): pp. 170-181.
- [15] Ebrahimi, B., et al., Mesenchymal stem cells improve medullary inflammation and fibrosis after revascularization of swine atherosclerotic renal artery stenosis. *PLoS One*, 2013. **8**(7): p. e67474.
- [16] de Resende, M.M., Stodola, T.J., and Greene, A.S., Role of the renin angiotensin system on bone marrow-derived stem cell function and its impact on skeletal muscle angiogenesis. *Physiological Genomics*, 2010. **42**(3): pp. 437-444.
- [17] Resende, M.M., Stodola, T.J., and Greene, A.S., Role of the renin angiotensin system on bone marrow-derived stem cell function and its impact on skeletal muscle angiogenesis after electrical stimulation. *The FASEB Journal*, 2009. **23**(1\_MeetingAbstracts): p. 1030.10.

- [18] Matsushita, K., et al., Local renin angiotensin expression regulates human mesenchymal stem cell differentiation to adipocytes. *Hypertension*, 2006. **48**(6): pp. 1095-1102.
- [19] Wang, L. and Leung, P.S., The role of renin-angiotensin system in cellular differentiation: implications in pancreatic islet cell development and islet transplantation. *Molecular and Cellular Endocrinology*, 2013. **381**(1): pp. 261-271.
- [20] Shi, R.-Z., et al., Angiotensin II induces vascular endothelial growth factor synthesis in mesenchymal stem cells. *Experimental Cell Research*, 2009. **315**(1): pp. 10-15.
- [21] Xu, Y.-X., et al., Mesenchymal stem cell therapy for diabetes through paracrine mechanisms. *Medical Hypotheses*, 2008. **71**(3): pp. 390-393.
- [22] Asahara, T., Kawamoto, A., and Masuda, H., Concise review: circulating endothelial progenitor cells for vascular medicine. *Stem Cells*, 2011. **29**(11): pp. 1650-1655.
- [23] Sata, M., et al., Hematopoietic stem cells differentiate into vascular cells that participate in the pathogenesis of atherosclerosis. *Nature Medicine*, 2002. **8**(4): pp. 403-409.
- [24] Richardson, M.R. and Yoder, M.C., Endothelial progenitor cells: quo vadis? *Journal of Molecular and Cellular Cardiology*, 2011. **50**(2): pp. 266-272.
- [25] Corselli, M., et al., The tunica adventitia of human arteries and veins as a source of mesenchymal stem cells. *Stem Cells and Development*, 2011. **21**(8): pp. 1299-1308.
- [26] Lam, C.-F., et al., Transplantation of endothelial progenitor cells improves pulmonary endothelial function and gas exchange in rabbits with endotoxin-induced acute lung injury. *Anesthesia & Analgesia*, 2011. **112**(3): pp. 620-627.
- [27] Traverse, J.H., et al., Effect of intracoronary delivery of autologous bone marrow mononuclear cells 2 to 3 weeks following acute myocardial infarction on left ventricular function: the LateTIME randomized trial. *JAMA*, 2011. **306**(19): pp. 2110-2119.
- [28] Jiang, H., et al., Palmitic acid promotes endothelial progenitor cells apoptosis via p38 and JNK mitogen-activated protein kinase pathways. *Atherosclerosis*, 2010. **210**(1): pp. 71-77.
- [29] Lee, P. and Poh, K.K., Endothelial progenitor cells in cardiovascular diseases. *World Journal of Stem Cells*, 2014. **6**(3): pp. 355-366.
- [30] Hiasa, K., et al., Gene transfer of stromal cell-derived factor-1a enhances ischemic vasculogenesis and angiogenesis via vascular endothelial growth factor/endothelial nitric oxide synthase-related pathway next-generation chemokine therapy for therapeutic neovascularization. *Circulation*, 2004. **109**(20): pp. 2454-2461.
- [31] Kobayashi, K., Imanishi, T., and Akasaka, T., Endothelial progenitor cell differentiation and senescence in an angiotensin II-infusion rat model. *Hypertension Research*, 2006. **29**(6): pp. 449-455.
- [32] You, D., et al., Combination of the angiotensin-converting enzyme inhibitor perindopril and the diuretic indapamide activate postnatal vasculogenesis in spontaneously hyper-

- tensive rats. *Journal of Pharmacology and Experimental Therapeutics*, 2008. **325**(3): pp. 766-773.
- [33] Hubert, C., et al., The hematopoietic system: a new niche for the renin–angiotensin system. *Nature Clinical Practice Cardiovascular Medicine*, 2006. **3**(2): pp. 80-85.
- [34] Kato, H., et al., Enhanced erythropoiesis mediated by activation of the renin-angiotensin system via angiotensin II type 1a receptor. *The FASEB Journal*, 2005. **19**(14): pp. 2023-2025.
- [35] Savary, K., et al., Role of the renin-angiotensin system in primitive erythropoiesis in the chick embryo. *Blood*, 2005. **105**(1): pp. 103-110.
- [36] Jokubaitis, V.J., et al., Angiotensin-converting enzyme (CD143) marks hematopoietic stem cells in human embryonic, fetal, and adult hematopoietic tissues. *Blood*, 2008. **111**(8): pp. 4055-4063.
- [37] Naito, M., et al., Effects of an angiotensin II receptor antagonist and angiotensin-converting enzyme inhibitors on burst forming units-erythroid in chronic hemodialysis patients. *American Journal of Nephrology*, 2003. **23**(5): pp. 287-293.
- [38] Freudenthaler, S., et al., Angiotensin II increases erythropoietin production in healthy human volunteers. *European Journal of Clinical Investigation*, 1999. **29**(10): pp. 816-823.
- [39] Chew, C., Weise, M., and Disney, A., The effect of angiotensin II receptor antagonist on the exogenous erythropoietin requirement of haemodialysis patients. *Nephrology Dialysis Transplantation*, 1999. **14**(8): pp. 2047-2049.
- [40] Durik, M., Pessôa, B.S., and Roks, A.J., The renin–angiotensin system, bone marrow and progenitor cells. *Clinical Science*, 2012. **123**(4): pp. 205-223.
- [41] Leung PS . Current research progress in the pancreas. In: *Advances in experimental medicine and biology book series. The renin-angiotensin system*. Dordrech: Springer; 2010. p. 1-207.
- [42] Leung, P.S. and Carlsson, P.-O., Pancreatic islet renin angiotensin system: its novel roles in islet function and in diabetes mellitus. *Pancreas*, 2005. **30**(4): pp. 293-298.
- [43] Chan, Y.C. and Leung, P.S., The renin–angiotensin system and reactive oxygen species: implications in pancreatitis. *Antioxidants & Redox Signaling*, 2011. **15**(10): pp. 2743-2755.
- [44] Iwai, M. and Horiuchi, M., Devil and angel in the renin–angiotensin system: ACE–angiotensin II–AT1 receptor axis vs. ACE2–angiotensin-(1-7)–Mas receptor axis. *Hypertension Research*, 2009. **32**(7): pp. 533-536.
- [45] Bindom, S.M., et al., Angiotensin I–converting enzyme type 2 (ACE2) gene therapy improves glycemic control in diabetic mice. *Diabetes*, 2010. **59**(10): pp. 2540-2548.
- [46] Jiang, F.-X. and Morahan, G., Pancreatic stem cells: from possible to probable. *Stem Cell Reviews and Reports*, 2012. **8**(3): pp. 647-657.

- [47] Ager, E.I., Neo, J., and Christophi, C., The renin–angiotensin system and malignancy. *Carcinogenesis*, 2008. **29**(9): pp. 1675-1684.
- [48] Ni, L., et al., Angiotensin-(1-7) inhibits the migration and invasion of A549 human lung adenocarcinoma cells through inactivation of the PI3K/Akt and MAPK signaling pathways. *Oncology Reports*, 2012. **27**(3): pp. 783-790.
- [49] Zhou, L., et al., Angiotensin-converting enzyme 2 acts as a potential molecular target for pancreatic cancer therapy. *Cancer Letters*, 2011. **307**(1): pp. 18-25.
- [50] Cheng, Q., et al., Combination of the dipeptidyl peptidase IV inhibitor LAF237 [(S)-1-[(3-hydroxy-1-adamantyl) ammo] acetyl-2-cyanopyrrolidine] with the angiotensin II type 1 receptor antagonist valsartan [N-(1-oxopentyl)-N-[[2'-(1H-tetrazol-5-yl)-[1, 1'-biphenyl]-4-yl] methyl]-L-valine] enhances pancreatic islet morphology and function in a mouse model of type 2 diabetes. *Journal of Pharmacology and Experimental Therapeutics*, 2008. **327**(3): pp. 683-691.
- [51] Lau, S.T., and Leung, P.S., Role of the RAS in pancreatic cancer. *Current Cancer Drug Targets*, 2011. **11**(4): pp. 412-420.
- [52] Suen, P., et al., PDZ-domain containing-2 (PDZD2) is a novel factor that affects the growth and differentiation of human fetal pancreatic progenitor cells. *The International Journal of Biochemistry & Cell Biology*, 2008. **40**(4): pp. 789-803.
- [53] Diehn, M., et al., Association of reactive oxygen species levels and radioresistance in cancer stem cells. *Nature*, 2009. **458**(7239): pp. 780-783.
- [54] Milanesi, A., et al.,  $\beta$ -Cell regeneration mediated by human bone marrow mesenchymal stem cells. *PloS One*, 2012. **7**(8): p. e42177.
- [55] Saleem, S., et al.,  $\beta$ 1 integrin/FAK/ERK signalling pathway is essential for human fetal islet cell differentiation and survival. *The Journal of Pathology*, 2009. **219**(2): pp. 182-192.
- [56] Burns, W.C., et al., Angiotensin II mediates epithelial-to-mesenchymal transformation in tubular cells by ANG 1-7/MAS-1-dependent pathways. *American Journal of Physiology-Renal Physiology*, 2010. **299**(3): pp. F585-F593.
- [57] Yu, J. and Thomson, J.A., Pluripotent stem cell lines. *Genes & Development*, 2008. **22**(15): pp. 1987-1997.
- [58] Chan, E.C., et al., Regulation of cell proliferation by NADPH oxidase-mediated signaling: potential roles in tissue repair, regenerative medicine and tissue engineering. *Pharmacology & Therapeutics*, 2009. **122**(2): pp. 97-108.
- [59] Kristensen, D.M., Kalisz, M., and Nielsen, J.H., Cytokine signalling in embryonic stem cells. *Apmis*, 2005. **113**(11-12): pp. 756-772.
- [60] Huang, D., et al., Angiotensin II promotes poly (ADP-ribosyl) ation of c-Jun/c-Fos in cardiac fibroblasts. *Journal of Molecular and Cellular Cardiology*, 2009. **46**(1): pp. 25-32.

- [61] Wang, W., et al., Essential role of Smad3 in angiotensin II-induced vascular fibrosis. *Circulation Research*, 2006. **98**(8): pp. 1032-1039.
- [62] Jeon, E.S., et al., Sphingosylphosphorylcholine induces differentiation of human mesenchymal stem cells into smooth-muscle-like cells through a TGF- $\beta$ -dependent mechanism. *Journal of Cell Science*, 2006. **119**(23): pp. 4994-5005.
- [63] Williams, A.R., et al., Intramyocardial stem cell injection in patients with ischemic cardiomyopathy functional recovery and reverse remodeling. *Circulation Research*, 2011. **108**(7): pp. 792-796.
- [64] Xu, M., et al., In vitro and in vivo effects of bone marrow stem cells on cardiac structure and function. *Journal of Molecular and Cellular Cardiology*, 2007. **42**(2): pp. 441-448.
- [65] Potapova, I.A., et al., Mesenchymal stem cells support migration, extracellular matrix invasion, proliferation, and survival of endothelial cells in vitro. *Stem Cells*, 2007. **25**(7): pp. 1761-1768.
- [66] Haznedaroglu, I.C. and Beyazit, Y., Review: Pathobiological aspects of the local bone marrow renin-angiotensin system: a review. *Journal of Renin-Angiotensin-Aldosterone System*, 2010. **11**(4): pp. 205-213.
- [67] Taylor, A.A., Siragy, H., and Nesbitt, S., Angiotensin receptor blockers: pharmacology, efficacy, and safety. *The Journal of Clinical Hypertension*, 2011. **13**(9): pp. 677-686.
- [68] Cleland, J.G., et al., Clinical trials update from the American Heart Association: REPAIR-AMI, ASTAMI, JELIS, MEGA, REVIVE-II, SURVIVE, and PROACTIVE. *European Journal of Heart Failure*, 2006. **8**(1): pp. 105-110.
- [69] Ebrahimian, T.G., et al., Dual effect of angiotensin-converting enzyme inhibition on angiogenesis in type 1 diabetic mice. *Arteriosclerosis, Thrombosis, and Vascular Biology*, 2005. **25**(1): pp. 65-70.
- [70] Huynh, K., et al., Cardiac-specific IGF-1 receptor transgenic expression protects against cardiac fibrosis and diastolic dysfunction in a mouse model of diabetic cardiomyopathy. *Diabetes*, 2010. **59**(6): pp. 1512-1520.
- [71] O'Sullivan, J.F., et al., Potent long-term cardioprotective effects of single low-dose insulin-like growth factor-1 treatment postmyocardial infarction. *Circulation: Cardiovascular Interventions*, 2011. **4**(4): pp. 327-335.
- [72] Huang, Y.-L., et al., Bone marrow stromal cell transplantation combined with angiotensin-converting enzyme inhibitor treatment in rat with acute myocardial infarction and the role of insulin-like growth factor-1. *Cytotherapy*, 2012. **14**(5): pp. 563-569.
- [73] Numasawa, Y., et al., Treatment of human mesenchymal stem cells with angiotensin receptor blocker improved efficiency of cardiomyogenic transdifferentiation and improved cardiac function via angiogenesis. *Stem Cells*, 2011. **29**(9): pp. 1405-1414.

