

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Comprehensive Physical Function Assessment in Elderly People

---

Tadashi Ito

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/67528>

---

## Abstract

In elderly people with mobility limitations, abnormalities in posture and gait contribute to the greater decline of physical motor function. The aim of the review article was to determine the comprehensive physical motor function assessment. Muscle function was assessed with the grip strength. Gait function was assessed with walking time tests conducted at a normal pace. Balance function was assessed with one-legged standing time. The 6-min walking distance test (6MD) was performed in a 10-m, straight corridor. Walking efficiency during the 6MD trials was measured using the Cosmed K4b2 (Rome, Italy), an indirect calorimetry system specifically designed to measure energy expenditure in nonlaboratory settings. The center of pressure was recorded using a balance board (Wii; Nintendo Co., Ltd., Kyoto, Japan). A vibratory stimulus was applied alternately to two muscles by fixing two vibrators from the vibration device onto the participant's gastrocnemius and lumbar multifidus muscle. These findings show that an assessment affecting postural control under proprioceptive stimulation might be a good indicator of elderly people. Also, the objective assessment of walking efficiency might be important for identifying the risk of external activity limitation or functional limitations among the late elderly.

**Keywords:** elderly people, physical function assessment, walking efficiency, proprioceptive system

---

## 1. Introduction

The ability to walk safely and postural control stabilization are vital for elderly people to decrease their risk of falling and to maintain their independence. A function of walking that accompanies aging is a higher energy cost of movement, often referred to as a lower economy

---

of walking [1, 2]. In addition, a defect or slowing of this mechanism has been suggested to explain the difficulties experienced by older persons when trying to control their posture [3, 4]. Moreover, both muscle strength and lower postural stability decline with aging [5, 6]. Therefore, aging is characterized by changes in the neuromuscular system that decreases muscle strength, balance, and proprioception.

Meanwhile, this progressive decline in physical capacities reduces the ability of elderly people to perform complex motor tasks and is associated with impaired mobility and a reduction in the ability [7]. Assessment of motor function contributes to the identification of factors that generate impairments to the performance of daily activities and an increase in the risk of falls for elderly people. Moreover, assessment of motor function by physical therapist provides important information about age-related progressive reduction in muscle strength, ability to walk, and postural control.

This chapter provides an overview of the assessment method on motor function in elderly people.

## **2. The relationship between walking efficiency and life-space assessment in elderly people**

### **2.1. Life-space assessment**

The increased metabolic cost of walking can detract from the activity and quality of life of elderly people as a decline in physical activity rapidly degrades physical and psychological functions [8, 9]. Life space has been reported to have good construct and criterion validity for measuring the severity of mobility limitations, and achieving efficient walking plays a crucial role in the extension of life space. Life-space assessment (LSA) is a tool that measures mobility and reflects participation in society based on the distance through which a person reports moving during the 4 weeks prior to the assessment [10, 11]. Life space, a relatively understudied concept in gerontology, can be defined as the size of the spatial area a person intentionally moves through in daily life, as well as the frequency of travel within a specific time frame [10]. Also, a reduced frequency of going outdoors could represent reduced life space, which has been hypothesized to be a risk factor in future physical disability. Murata et al. reported that life space was related not only to age or health status but also to environmental or psychosocial factors [12]. Shimada et al.'s reports confirm that LSA was more strongly associated with gait speed than the other gait variables [13] and also show that these declines in physical performances were apparent at age 80 years and over in women and at age 90 years and over in men [13].

LSA may reflect the physical activity status indirectly in elderly people because the LSA score is associated with physical performance, Activities of Daily Living (ADL), and sociodemographic factors [11].

Life-space mobility was measured using a Japanese translation of the LSA [14]. The LSA provided a score based on the distance a person reported moving during the 4 weeks before the assessment [10, 13].

The LSA scores range from 0 (“completely room-bound”) to 120 (“travel out of town every day without assistance”) [13].

## 2.2. Walking efficiency in elderly people

It may be that the decline in walking efficiency is a common result of the decrease of many bodily functions that capture the overall impact of life space and is an important indicator of life space. However, the mechanism by which older people’s life space and walking efficiency decline is not well investigated and remains poorly understood. Also, the specific mechanisms that explain the difference between the decline in walking efficiency and life space in early elderly and late elderly are not yet clear. Additionally, little is known about the relationship between the walking efficiency decline and confined life space to influence on aging.

In older adults with mobility limitations, abnormalities in posture and gait contribute to the greater energy cost of inefficient gait, with adjustments for age and gait speed [15]. Limitations in life space, as measured by the University of Alabama at Birmingham (UAB) life-space assessment, reflect lifestyle as well as walking efficiency and may be a valuable measure of functional decrease for community-dwelling elderly people, especially since life space specifically relates to mobility and a person’s participation in society.

### 2.2.1. Walking efficiency assessment

Walking efficiency during the 6MD trials was measured using the Cosmed K4b2 (Rome, Italy), an indirect calorimetry system specifically designed to measure energy expenditure in nonlaboratory settings (**Figure 1**).

In brief, it uses a breath-by-breath measurement of gas exchange through a rubberized facemask and a turbine for gas collection, secured by a head harness. A flexible facemask that the participants keep in place by a head harness covers the participants’ mouth and nose. Flexible facemask is attached to a digital turbine flowmeter to measure the volume of expired air and inspired. Sampling line from the turbine to analyzer unit system delivers expired air for the measurement ( $O_2$ ,  $CO_2$  content). Before test, the K4b2 was calibrated according to the manufacturer’s guidelines. After warming up the unit for 60 min, the  $CO_2$  and  $O_2$  analyzers were calibrated against room air as well as a reference gas of known composition (4.94%  $CO_2$ , 16.07%  $O_2$ ). Before walking efficiency assessment, each participant was required to sit quietly for 10 min as a rest period. The speed and distance of the 6-min walking distance test (6MD) was recorded using standardized procedures.




**Figure 1.** Expiration gas analyser.

Walking efficiency was calculated based on net efficiency as follows: Walking efficiency = (walking  $VO_2$  ml/kg - resting  $VO_2$  ml/kg)/6MD average walking speed (m/s) (**Figure 2**) [16]. The Cosmed K4b2 system uses the increased  $VO_2$ /kg ratio after noise processing to predict walking efficiency from  $O_2$  consumption.


Breath-by-breath values from the Excel spreadsheet were calculated from the increased ratio over 1-breath intervals. Outliers from the increased ratio were calculated from exploratory data analysis of participants. We defined outliers (noise) as values beyond an increased ratio of 100%. Data that were more than 100% of the increased ratio were removed, and a below-100% value was calculated from the remaining test data. This method contributes to noise processing of the  $VO_2$ /kg signal. For each subject, walking efficiency was calculated by recording the noise-processed  $VO_2$ /kg values.

### **2.3. Relationship between walking efficiency and life-space assessment**

Ito et al. reported the significant relationship between walking efficiency and LSA in late elderly [16]. Moreover, the data from this study suggest that as walking efficiency declines with age, life space increasingly declines (**Figure 3**). This suggests that the age-related decline of walking efficiency is caused by physiological changes of the late elderly. Other studies have suggested that as aerobic capacity declines with age, walking at a habitual speed becomes an increasingly more intense and therefore difficult activity, resulting in a slowing of walking speed in an effort to reduce fatigue [17]. Also, the previous study has shown that the walking energy cost for a comparable speed is generally higher for healthy



**Figure 2.** Walking efficiency = (walking  $VO_2$  ml/kg - resting  $VO_2$  ml/kg)/6MD average walking speed (m/s). Horizontal axis of blue shows the meaning of 10 min of the rest period and 6-min walking trial. Vertical axis of red shows the meaning of last 3 min of the 6-min walking trial. Following the transfer of data from the instrument, an Excel spreadsheet was used to calculate steady-state  $VO_2$ /kg and mean counts per minute during the last 3 min of the 6MD trial and 10 min of the rest period.



**Figure 3.** Schematic diagram of the incidence of walking efficiency decline in late elderly people.

elderly people, particularly those above 65 years, compared with younger people [18, 19]. Mechanisms related to the initiation and stepping patterns of gait, such as hip extension, step width, and cadence, have previously been reported to be related to the energy cost of walking in older adults with slow and variable gait [15]. Abe et al. reported that women of advanced age (75 years or older) have diminished pulmonary function, physical function, and mobility, and that diminished pulmonary function is associated with declining physical function [20]. Malatesta et al. reported that healthy octogenarians exhibited higher walking cost and greater stride time variability [21] and also reported that these declines in physical performances were apparent at age 80 years and over in women and at age 90 years and over in men [13]. Shimada reported that increased  $\text{VO}_2$  in older adults manifests as walking becomes inefficient and reduced endurance capacity occurs [22].

This suggests that going activity to extension of LSA may better impact walking efficiency than efforts to improve gait speed and muscle strength.

### 3. Physical function assessments in elderly people

#### 3.1. Performance assessment

The assessment measures are taken by well-trained staff who had nursing, physiotherapy, occupational therapy, or similar qualifications.

##### 3.1.1. Grip strength assessment

Muscle function is assessed by the grip strength (GS) (**Figure 4**). GS is measured in kilograms in the participant's dominant hand using a Smedley-type handheld dynamometer (GRIP-D; Takei Ltd, Niigata, Japan).

##### 3.1.2. Five-meter walk test

Five-meter walk test is assessed with walking conducted at a normal pace (usual walking speed; UWS). Two marks are used to indicate the path (start and end) of a 5-m walk path space. UWS is measured using a stopwatch system. Examiner tells the subjects to walk on a straight and flat surface at UWS. Moreover, examiner tells the subjects to maintain gait past the end of the walk path for a further 2.5 m.

##### 3.1.3. One-legged standing time test

Balance function is assessed with one-legged standing time (**Figure 5**). One-legged standing time is used to measure how long a subject remains standing on one leg (eyes open). Subjects are allowed to decide which leg to use as support leg. Next, examiner tells the subjects to lift the converse foot from flat floor. Moreover, examiner tells the subjects to confirm sure not to press the lifted leg against support leg. The test ended when lifted leg touched the support leg, the lifted leg touched the floor, or after 60 s of successful balance.



**Figure 4.** Grip strength assessment.



**Figure 5.** One-legged standing time test.

#### *3.1.4. The 6-min walking distance test*

6MD is performed in a 10-m, straight corridor. In the 6MD, participants are instructed to walk as quickly as possible to cover the longest distance possible within 6 min. This test measures the distance that a participant could quickly walk on a flat, hard surface in a period of 6 min.



### 3.1.5. Appendicular skeletal muscle mass assessment

In the elderly, muscle weakness is associated with the muscle atrophy aging (sarcopenia) progressive loss. Skeletal muscle mass loss may also have the potential to impact quality of life and ultimately the need for long-term care in elderly people [23]. Several studies developed equations for estimating skeletal muscle mass [24–26].

### 3.1.6. Prediction models

#### **Sanada K, et al., 2010: Prediction models of sarcopenia [24]**

Men: appendicular skeletal muscle mass =  $0.326 \times \text{body mass index (BMI)} - 0.047 \times \text{waist circumference} - 0.011 \times \text{Age} + 5.135$  ( $R^2 = 0.68$ ).

Women: appendicular skeletal muscle mass =  $0.156 \times \text{BMI} + 0.044 \times \text{hand grip strength} - 0.010 \times \text{waist circumference} + 2.747$  ( $R^2 = 0.57$ ).

#### **Yoshida D, et al., 2014: Bioelectrical impedance analysis [25]**

Men: appendicular skeletal muscle mass =  $0.197 \times (\text{impedance index}) + 0.179 \times (\text{weight}) - 0.019$  ( $R^2 = 0.87$ ).

Women: appendicular skeletal muscle mass =  $0.221 \times (\text{impedance index}) + 0.117 \times (\text{weight}) + 0.881$  ( $R^2 = 0.89$ ).

#### **Ito T, et al., 2016: Simple estimation of appendicular muscle mass [26]**

Appendicular skeletal muscle mass =  $5.051 \times (\text{gender: men} = 1, \text{ women} = 0) + 0.364 \times (\text{BMI}) + 0.168 \times (\text{maximum calf circumference}) - 0.815$  ( $R^2 = 0.80$ ).

## 3.2. Reduced physical activity

Past studies have provided the first evidence that slightly constricted life space may serve as an important marker and/or risk factor for the development of frailty, whereas severely constricted life space may indicate a high risk of mortality [27]. Webber et al. reported how mobility impairments can lead to limitations in accessing different life spaces and stressed the associations among determinants that influence mobility [28]. Previous study reported falls and reduced life space closely related to physical performance [29, 30].


## 4. Assessment of relative proprioceptive weighting ratio in elderly people with lumbar spondylosis

### 4.1. Sensory proprioceptive inputs in postural control

In daily life, an upright postural control task is essential for the activities of elderly people. After processing of the sensory inputs, individuals must integrate the respective contributions of the various sources of sensory information for regulating posture. Hay et al. reported

that older persons have difficulties in taking advantage of sensory redundancy in postural control [31]. In addition, a defect or slowing of this mechanism has been suggested to explain the difficulties experienced by older persons when trying to control their posture [32, 33]. Proprioceptive input from the muscles of the legs and trunk plays an important role in maintaining postural stability [34]. Previous studies have reported that proprioception and vibration sensation in the lower limbs decrease during normal aging [3, 4] and also reported that postural instability has been observed in elderly people [35]. Therefore, a vibratory stimulus that matches the response frequency of the receptors present in skeletal muscle may influence body sway (**Figure 6**).

Previous studies have reported that patients with recurrent low back pain (LBP) have impaired motor control [36] and altered lumbosacral proprioceptive acuity [37, 38]. LBP is a widespread pathological condition that is often related to impaired or degenerated trunk mobility, which becomes evident during common activities [39, 40]. Taimela et al. reported that lumbar muscle fatigue impaired lumbar positional sense in both patients with LBP and healthy subjects [41]. These impairments lead to pain and declines in postural strategy, muscle function, and proprioception [42, 43].



**Figure 6.** The representative receptor and response frequencies are 30 Hz in Meissner's corpuscles, 60 Hz in muscle spindles, and 240 Hz in Vater-Pacini corpuscles.

Muscle vibration, known to be a strong stimulus for muscle spindles and Vater-Pacini corpuscles, has been used to assess the role of proprioception [44, 45]. Both of these studies suggest that pain is a possible cause of decreased variability in postural strategy. These impairments lead to pain and declines in postural strategy, muscle function, and proprioception.

#### 4.2. Assessment of relative proprioceptive weighting ratio

The center of pressure (COP) was recorded using a balance board (Wii; Nintendo Co., Ltd., Kyoto, Japan) (**Figure 7**) [46–48]. A vibratory stimulus is applied alternately to each muscle by fixing vibrators from the vibration device onto the subjects' lumbar multifidus (LM) muscle and gastrocnemius (GS). The device consists of an amplifier, laptop computer, and four vibrators (**Figure 8**). This mechanical vibration test has been used to analyze the role of proprioception in the postural control strategy [49–53].

The subjects stood barefoot on the Wii Balance Board with their feet together and their eyes closed. They were instructed to remain still and relax in the standing posture with their arms hanging loosely at their sides (**Figure 9**).


To provide information regarding relative proprioceptive sensation dominance, the relative proprioceptive weighting (RPW) ratio was calculated using following computation expression.

$$RPW = (Abs\ GS)/(Abs\ LM + Abs\ GS) \times 100[\%]$$


RPW of near to 0 conform to 100 [%] dependence on trunk strategy, whereas RPW of near to 100 conform to 100 [%] dependence on lower limb strategy [54–56].



**Figure 7.** Recently, the Wii Balance Board has been much used in the field of medical research, and it has been reported that results from the Wii Balance Board correlate closely with those of commercially available force plates.



**Figure 8.** Photograph and block diagram of variable-frequency vibratory stimulation device.



**Figure 9.** Experimental setup: paraspinous muscles and gastrocnemius muscles in vibration trial.

#### **4.3. Proprioceptive input decline from the muscles of the legs or trunk**

Recent studies in which a vibratory stimulation of 60 Hz was used have suggested that people with LBP adopt a lower leg-derived postural control strategy [57]. A possible explanation

is that these participants were exploiting this strategy to its maximum effect during vibratory stimulation of 60 Hz. Also, according to other study, the impairment of back muscle strength leads to the motor function and sensory deficit that affects balance performance [58]. Moreover, according to another study, the lower leg's response to balance control under 30 Hz proprioceptive stimulation might be a good indicator of declining gait function [59].

These differences may result from differences between the measurement conditions and physical status of the participants.

## 5. Conclusions

Comprehensive physical function assessment has been proved useful for understanding the ability of elderly people. Walking efficiency is closely correlated with life space, and it can be used for comparing physical function. Performance assessment, such as the grip strength, skeletal muscle mass, and postural control stabilization, may help slow the events that ultimately can lead to motor function disorder and disability. This chapter suggests that comprehensive physical function assessment is a useful method to evaluate the physical function and therefore aids rehabilitation programs.

## Acknowledgements

We also thank Yoshihito Sakai, Yoshifumi Morita, Noritaka Sato, and Kazunori Yamazaki for their help in this chapter.

## Author details

Tadashi Ito

Address all correspondence to: sanjigen@mikawa-aoitori.jp

Three-dimensional motion analysis room, Aichi Prefecture Mikawa Aoitori Medical and Rehabilitation Center for Developmental Disabilities, Okazaki-shi, Aichi, Japan

## References

- [1] Wert DM, Brach JS, Perera S, et al. The association between energy cost of walking and physical function in older adults. *Arch Gerontol Geriatr.* 2013;**57**(2):198–203.
- [2] Schrack JA, Simonsick EM, Ferrucci L. The relationship of the energetic cost of slow walking and peak energy expenditure to gait speed in mid-to-late life. *Am J Phys Med Rehabil.* 2013;**92**(1):28–35.

- [3] Skinner HB, Barrack RL, Cook SD. Age-related decline in proprioception. *Clin Orthop Relat Res.* 1984;**184**:208–11.
- [4] Steinberg FU, Graber AL. The effect of age and peripheral circulation on the perception of vibration. *Arch Phys Med Rehabil.* 1963;**44**:645–50.
- [5] Thelen DG, Schultz AB, Alexander NB, et al. Effects of age on rapid ankle torque development. *J Gerontol A Biol Sci Med Sci.* 1996;**51**(5):M226–32.
- [6] Woodhull-McNeal AP. Changes in posture and balance with age. *Aging (Milano).* 1992;**4**(3):219–25.
- [7] Meuleman JR, Brechue WF, Kubilis PS, et al. Exercise training in the debilitated aged: strength and functional outcomes. *Arch Phys Med Rehabil.* 2000;**81**(3):312–18.
- [8] Bäckmand H, Kaprio J, Kujala UM, et al. Physical and psychological functioning of daily living in relation to physical activity. A longitudinal study among former elite male athletes and controls. *Aging Clin Exp Res.* 2006;**18**(1):40–9.
- [9] Young DR, Masaki KH, Curb JD. Associations of physical activity with performance-based and self-reported physical functioning in older men: the Honolulu Heart Program. *J Am Geriatr Soc.* 1995;**43**(8):845–54.
- [10] Baker PS, Bodner EV, Allman RM. Measuring life-space mobility in community-dwelling older adults. *J Am Geriatr Soc.* 2003;**51**(11):1610–4.
- [11] Peel C, Sawyer Baker P, Roth DL, et al. Assessing mobility in older adults: The UAB Study of Aging Life-Space Assessment. *Phys Ther.* 2005;**85**(10):1008–19.
- [12] Murata C, Kondo T, Tamakoshi K, et al. Factors associated with life space among community-living rural elders in Japan. *Public Health Nurs.* 2006;**23**(4):324–31.
- [13] Shimada H, Kim H, Yoshida H, et al. Relationship between age-associated changes of gait and falls and life-space in elderly people. *J Phys Ther Sci.* 2010;**22**(4):419–24.
- [14] Shimada H, Sawyer P, Harada K, et al. Predictive validity of the classification schema for functional mobility tests in instrumental activities of daily living decline among older adults. *Arch Phys Med Rehabil.* 2010;**91**(2):241–6.
- [15] Wert DM, Brach J, Perera S, et al. Gait biomechanics, spatial and temporal characteristics, and the energy cost of walking in older adults with impaired mobility. *Phys Ther.* 2010;**90**(7):988–85.
- [16] Ito T, Shimada H, Yoshida D, et al. The relationship between walking efficiency and Life-space assessment in elderly people. In: *The 49nd Congress of Japanese Society of Physical Therapy*; May 31; Kanagawa Japan. Rigakuryohogaku: Japanese Physical Therapy Association; 2014. p. 1246.
- [17] Fiser WM, Hays NP, Rogers SC, et al. Energetics of walking in elderly people: factors related to gait speed. *J Gerontol A Biol Sci Med Sci.* 2010;**65**(12):1332–7.

- [18] Waters RL, Hislop HJ, Perry J, et al. Comparative cost of walking in young and old adults. *J Orthop Res*. 1983;1(1):73–6.
- [19] McCann DJ, Adams WC. A dimensional paradigm for identifying the size-independent cost of walking. *Med Sci Sports Exerc*. 2002;34(6):1009–17.
- [20] Abe T, Suzuki T, Yoshida H, et al. The relationship between pulmonary function and physical function and mobility in community-dwelling elderly women aged 75 years or older. *J Phys Ther Sci*. 2011;23(3):443–9.
- [21] Malatesta D, Simar D, Dauvilliers Y, et al. Energy cost of walking and gait instability in healthy 65- and 80-yr-olds. *J Appl Physiol*. 2003;95(6):2248–56.
- [22] Shimada H, Sturnieks D, Endo Y, et al. Relationship between whole body oxygen consumption and skeletal muscle glucose metabolism during walking in older adults: FDG PET study. *Aging Clin Exp Res*. 2011;23(3):175–82.
- [23] Janssen I, Heymsfield SB, Ross R. Low relative skeletal muscle mass (sarcopenia) in older persons is associated with functional impairment and physical disability. *J Am Geriatr Soc*. 2002;50(5):889–96.
- [24] Sanada K, Miyachi M, Yamamoto K, et al. Prediction models of sarcopenia in Japanese adult men and women. *Tairyoku Kagaku*. 2010;59(3):291–301.
- [25] Yoshida D, Shimada H, Park H, et al. Development of an equation for estimating appendicular skeletal muscle mass in Japanese older adults using bioelectrical impedance analysis. *Geriatr Gerontol Int*. 2014;14(4):851–7.
- [26] Ito T, Sakai Y, Morita Y, et al. Simple estimation of appendicular muscle mass using the maximum calf circumference of elderly inpatients. *Rigakuryoho Kagaku*. 2016;31(4):511–5.
- [27] Xue QL, Fried LP, Glass TA, et al. Life-space constriction, development of frailty, and the competing risk of mortality: the Women’s Health And Aging Study I. *Am J Epidemiol*. 2008;167(2):240–8.
- [28] Webber SC, Porter MM, Menec VH. Mobility in older adults: a comprehensive framework. *Gerontologist*. 2010;50(4):443–50.
- [29] Imms FJ, Edholm OG. Studies of gait and mobility in the elderly. *Age Ageing*. 1981;10(3):147–56.
- [30] Bootsma-van der Wiel A, Gussekloo J, De Craen AJ, et al. Common chronic diseases and general impairments as determinants of walking disability in the oldest-old population. *J Am Geriatr Soc*. 2002;50(8):1405–10.
- [31] Hay L, Bard C, Fleury M, et al. Availability of visual and proprioceptive afferent messages and postural control in elderly adults. *Exp Brain Res*. 1996;108(1):129–39.
- [32] Shumway-Cook A, Woollacott M. Attentional demands and postural control: the effect of sensory context. *J Gerontol A Biol Sci Med Sci*. 2000;55(1):M10–6.

- [33] Teasdale N, Stelmach GE, Breunig A, et al. Age differences in visual sensory integration. *Exp Brain Res.* 1991;**85**(3):691–6.
- [34] Bloem BR, Allum JH, Carpenter MG, et al. Is lower leg proprioception essential for triggering human automatic postural responses?. *Exp Brain Res.* 2000;**130**(3):375–91.
- [35] Pyykkö I, Jäntti P, Aalto H. Postural control in elderly subjects. *Age Ageing.* 1990;**19**(3):215–21.
- [36] Hodges PW, Richardson CA. Inefficient muscular stabilization of the lumbar spine associated with low back pain. A motor control evaluation of transversus abdominis. *Spine (Phila Pa 1976).* 1996;**21**(22):2640–50.
- [37] Brumagne S, Cordo P, Lysens R, et al. The role of paraspinal muscle spindles in lumbosacral position sense in individuals with and without low back pain. *Spine (Phila Pa 1976).* 2000;**25**(8):989–94.
- [38] Newcomer KL, Laskowski ER, Yu B, et al. Differences in repositioning error among patients with low back pain compared with control subjects. *Spine (Phila Pa 1976).* 2000;**25**(19):2488–93.
- [39] Luoto S, Aalto H, Taimela S, et al. One-footed and externally disturbed two-footed postural control in patients with chronic low back pain and healthy control subjects. A controlled study with follow-up. *Spine (Phila Pa 1976).* 1998;**23**(19):2081–9; discussion 2089–90.
- [40] Radebold A, Cholewicki J, Polzhofer GK, et al. Impaired postural control of the lumbar spine is associated with delayed muscle response times in patients with chronic idiopathic low back pain. *Spine (Phila Pa 1976).* 2001;**26**(7):724–30.
- [41] Taimela S, Kankaanpää M, Luoto S. The effect of lumbar fatigue on the ability to sense a change in lumbar position. A controlled study. *Spine (Phila Pa 1976).* 1999;**24**(13):1322–7.
- [42] Moseley GL, Hodges PW. Are the changes in postural control associated with low back pain caused by pain interference?. *Clin J Pain.* 2005;**21**(4):323–9.
- [43] Brumagne S, Lysens R, Swinnen S, et al. Effect of paraspinal muscle vibration on position sense of the lumbosacral spine. *Spine (Phila Pa 1976).* 1999;**24**(13):1328–31.
- [44] Roll JP, Vedel JP. Kinaesthetic role of muscle afferents in man, studied by tendon vibration and microneurography. *Exp Brain Res.* 1982;**47**(2):177–90.
- [45] Frost LR, Bijman M, Strzalkowski ND, et al. Deficits in foot skin sensation are related to alterations in balance control in chronic low back patients experiencing clinical signs of lumbar nerve root impingement. *Gait Posture.* 2015;**41**(4):923–8.
- [46] Clark RA, Bryant AL, Pua Y, et al. Validity and reliability of the Nintendo Wii Balance Board for assessment of standing balance. *Gait Posture.* 2010;**31**(3):307–10.
- [47] Young W, Ferguson S, Brault S, et al. Assessing and training standing balance in older adults: a novel approach using the 'Nintendo Wii' Balance Board. *Gait Posture.* 2011;**33**(2):303–5.



- [48] Holmes JD, Jenkins ME, Johnson AM, et al. Validity of the Nintendo Wii® balance board for the assessment of standing balance in Parkinson's disease. *Clin Rehabil.* 2013;**27**(4):361–6.
- [49] Capicíková N, Rocchi L, Hlavacka F, et al. Human postural response to lower leg muscle vibration of different duration. *Physiol Res.* 2006;**55**(Suppl 1):S129–34.
- [50] Barbieri G, Gissot AS, Fouque F, et al. Does proprioception contribute to the sense of verticality?. *Exp Brain Res.* 2008;**185**(4):545–52.
- [51] Thompson C, Bélanger M, Fung J. Effects of bilateral Achilles tendon vibration on postural orientation and balance during standing. *Clin Neurophysiol.* 2007;**118**(11):2456–67.
- [52] Ivanenko YP, Solopova IA, Levik YS. The direction of postural instability affects postural reactions to ankle muscle vibration in humans. *Neurosci Lett.* 2000;**292**(2):103–6.
- [53] Ceyte H, Cian C, Zory R, et al. Effect of Achilles tendon vibration on postural orientation. *Neurosci Lett.* 2007;**416**(1):71–5.
- [54] Brumagne S, Janssens L, Janssens E, et al. Altered postural control in anticipation of postural instability in persons with recurrent low back pain. *Gait Posture.* 2008;**28**(4):657–62.
- [55] Janssens L, Brumagne S, Polspoel K, et al. The effect of inspiratory muscles fatigue on postural control in people with and without recurrent low back pain. *Spine (Phila Pa 1976).* 2010;**35**(10):1088–94.
- [56] Janssens L, Brumagne S, McConnell AK, et al. Proprioceptive changes impair balance control in individuals with chronic obstructive pulmonary disease. *PLoS One.* 2013;**8**(3):e57949.
- [57] Brumagne S, Janssens L, Knapen S, et al. Persons with recurrent low back pain exhibit a rigid postural control strategy. *Eur Spine J.* 2008;**17**(9):1177–84.
- [58] Ito T, Sakai Y, Nakamura E, et al. Relationship between paraspinal muscle cross-sectional area and relative proprioceptive weighting ratio of older persons with lumbar spondylosis. *J Phys Ther Sci.* 2015;**27**(7):2247–51.
- [59] Ito T, Sakai Y, Kubo A, et al. The relationship between physical function and postural sway during local vibratory stimulation of middle-aged people in the standing position. *J Phys Ther Sci.* 2014;**26**(10):1627–30.