

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Molecular Phylogeny and Taxonomy of *Lepidoptera* with Special Reference to Influence of *Wolbachia* Infection in the Genus *Polytremis*

Weibin Jiang

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.69958>

Abstract

This chapter provides a case of genus *Polytremis* Mabille, 1904 (*Lepidoptera*: HesperIIDae), to explain the molecular phylogeny and taxonomy of *Lepidoptera* and the influence of *Wolbachia* infection. Earlier studies of *Lepidoptera* were focused mainly on the morphological classification, population distribution, and identification of new species. As the supplementary to morphological research, analysis of DNA has been widely used in the phylogenetic studies of *Lepidoptera*. The study provides a conservative estimate that the *Wolbachia* infection rate in *Polytremis nascens* Leech (1893) is 31%, and no significant difference in the prevalence is found between the sexes. The *Wolbachia* infection mainly prevails in populations of *P. nascens* in southern China, which influence the diversity of mtDNA in *P. nascens* by a *Wolbachia*-induced sweep. The *Wolbachia* infection rate in *Polytremis fukia* Evans (1940) is 47% and shows a weak association existed between mitochondrial DNA haplotypes and wFuk1 infection status.

Keywords: *Lepidoptera*, microsatellite, mitochondrial genome, molecular phylogeny, taxonomy, *Wolbachia*

1. Introduction to molecular phylogeny and taxonomy of *Lepidoptera*

Butterflies and moths (*Lepidoptera*) have long served as a model system for ecological and evolutionary studies on the basis of the high degree of diversity and complexity, which constitute one of the most diverse insect orders with more than 157,000 described species. Earlier studies of *Lepidoptera* were focused mainly on the morphological classification, population distribution, and identification of new species. As the supplementary to morphological research, analysis of

DNA has been widely used in the phylogenetic and taxonomic studies of *Lepidoptera*. The case of genus *Polytremis* will be discussed as follows.

2. The phylogeny of the butterfly genus *Polytremis*


The family HesperIIDae includes more than 4000 species, commonly known as “skippers,” of which HesperIIDae is the largest subfamily. *Polytremis* Mabille (1904) is a genus of subfamily HesperIIDae, which has 18 members and is restricted geographically to the continental part of the southeastern Palaearctic and northern oriental regions. They have a thick body and relatively small wings. These wings are commonly dark brown or yellowish brown [1]. The main external features are characterized by the unspined mesotibia, the absence of a cell spot on the underside of each hind wing, and the serial, linear, and semi-hyaline spots. Male genitalia are distinguished by the elongated harpe, swollen tegumen, and bifid uncus [2].

2.1. The method of constructing and analyzing phylogenetic tree

The specimens from 15 of the estimated 18 species in the genus *Polytremis* were collected, from different localities. The DNA was isolated from leg tissue. The mitochondrial cytochrome c oxidase I (COI) gene, recommended as the universal and standard barcoding marker for species identification [3], was amplified approximately 490 bp. For nuclear DNA, three expansion segments of 18S rDNA and 28S rDNA were chosen, the slowly evolving genes used normally in higher classification studies [4]. The haplotype sequence matrix was used for all subsequent phylogenetic analyses. MEGA v4.0 was used to calculate the intra- and interspecific genetic divergences based on the K2P model [5]. Phylogenetic trees were constructed by the maximum-likelihood (ML) methods using PAUP 4.0b10 [6]. Relationships among the mitochondrial COI and concatenated sequence (mitochondrial COI + nuclear rDNA) haplotypes were represented as a haplotype network obtained by the software DnaSP4.90 [7] and Network4.5 using the median-joining method [8].

2.2. Genetic divergence, phylogenetics, and network of genus *Polytremis*

Figure 1A reveals five main clades and shows the ML tree based on the data set of COI. Clade I contained eight species: *Polytremis suprema* and *Polytremis gigantean*, *Polytremis caerulescens*, *Polytremis kiraizana* Sonan, 1938, and *Polytremis matsuii* Sugiyama, 1999 are first clustered with a strong support value, followed by clustering of *Polytremis gotama* and *Polytremis nascens* Leach, 1893. Then *Polytremis jigongi* Zhu, 2012 is revealed [2]. Two haplotypes of *P. jigongi* are clearly separated from other species form and strongly supported lineages. Clade II contains only *Polytremis theca*. It was reported to include three subspecies. They show a greater intraspecific genetic distance than some interspecific genetic distances in the genus *Polytremis*. Furthermore, the sister group relationship between *Polytremis zina* Evans, 1932 and *Polytremis pellucida* Murray, 1875 in Clade III is confirmed. Clade IV contains only one species, *Polytremis mencia* Moore, 1877. Clade V contains three species: *Polytremis discreta* Elwes & Edwards, 1897, *Polytremis lubricans* Herrich-Schäffer, 1869, and *Polytremis eltola* Hewitson, 1869, which are distributed sympatrically in the oriental region throughout Malaya and India.



_10

Figure 1. (A) Maximum-likelihood phylogeny on the basis of the mitochondrial COI sequences and (B) network on the basis of the mitochondrial COI sequences.

The level of DNA sequence divergence reflected the taxonomic hierarchy of the original species. The lowest intraspecific COI genetic distance was observed between *P. suprema* and *P. gigantea* (K2P distance 1.7%). Except for *P. theca*, the intraspecific distances were shorter. The COI data confirmed the sister group relationship between *P. suprema* and *P. gigantea*, which form a monophyletic group together with *P. matsuii*, *P. caerulescens*, and *P. kiraizana*. All of their interspecific distances were smaller than 3% (K2P distance). We can infer from morphological traits. These five species shared many morphological traits including ear-like uncus with a pair of processes, the absence of a cornuti, and thin coecum penis. *P. theca* was the only species for which the intraspecific genetic distance was greater than some interspecific

genetic distances based on COI in the genus *Polytremis*. However, the distance was much less than the average interspecific genetic distances of the genus *Polytremis* (K2P distance 7.9%) [9]. *P. theca* was widely distributed in the south of the Qinling Mountains in China, except in the Hainan Province and the southern tropical regions of Yunnan Province [10]. Three subspecies of *P. theca* were reported on the basis of morphological features of the wings. Our specimens included two of them, namely, *Polytremis theca theca* and *Polytremis theca fukia*. The COI tree revealed these two distinct haplotype lineages without intermediates (K2P distance 4.2%). Additionally, the subspecies were separated by nuclear rDNA sequence, and the K2P distance was 0.3%, suggesting the possible existence of a sibling species paired with allopatric distribution.

The average interspecific rDNA genetic distance (K2P distance 1.0%) was far less than that of COI (K2P distance 7.9%). Except for *P. caerulescens* and *P. gotama*, other species in *Polytremis* could be distinguished with rDNA. These two species could be separated in the COI, and K2P distance was 1.9% but showed no variation in the rDNA. Based on mitochondrial and nuclear markers, the differences observed between results may contribute to recent separation, introgressive hybridization, or incomplete lineage sorting [11]. Because *Polytremis* is a fairly old genus and the splits of COI of the two lineages are also quite old, it seems that incomplete lineage sorting may not be an appealing explanation for the discordance. Additionally, *P. gotama* has been described as an independent species by COI data and morphological features [12]. A specimen of *P. gotama* and three specimens of *P. caerulescens* (from two populations) revealed two distinct haplotype lines without intermediates (K2P distances 4.9%). The observation indicates that they were two species based on the molecular and morphological level. Nevertheless, more specimens of the two species from different population need to be collected and analyzed in the future to see if this pattern is recovered consistently and further confirm the relationship of them. Instead, some arguments favor the assumption of recent separation [9]. As far as their morphological characteristics were concerned, *P. caerulescens* was considered to be closely related to *P. gotama* on the basis of the structural similarity of the cell spot on the upper side of the hind wing and the male genitalia, which were not found in the other *Polytremis* species [13]. Additionally, only these two species were observed and captured at altitudes higher than 2000 m. *P. gotama* was a little smaller than *P. caerulescens*. They both varied in other morphological traits which clearly support the existence of two closely related but distinct species, including male stigma on the upper side of the forewing and the ground color of the wings. rDNA showed no sequence variation, whereas K2P distances of the COI fragments reached 4.9%, suggesting a possible recent separation of *P. gotama* and *P. caerulescens*.

In *Lepidoptera*, thresholds have been proposed as 3% for COI [3]. The intraspecific and interspecific genetic divergences did not fall into separate intervals, and an obvious "barcode gap" did not occur in COI in our study of *Polytremis* [9]. It was entailed by two factors. Firstly, all intraspecific distances were less than 3% except for *P. theca*. However, we inferred *P. t. theca* and *P. t. fukia* could be a sibling species pair according the morphological and molecular data in the study. Secondly, the interspecific genetic distances among five sister species (*P. gigantean*, *P. suprema*, *P. caerulescens*, *P. kiraizana*, and *P. matsuii*) were less than 3%, which caused the interspecific and intraspecific genetic divergences to overlap from 1.7 to 3%. Regardless, for COI, the overlaps between intraspecific and interspecific variations would not affect

identification in a thoroughly sampled evaluation [14]. The *Polytremis* species could be clustered with a well support and distinguished by tree-based methods, suggesting that the COI sequence could be used to correctly identify almost all species in the genus as a DNA barcode (**Figure 1A**). Additionally, the markers of the nuclear rDNA sequences used in our studies, three expansion segments of 18S and 28S rDNA, have been proposed as a reasonable alternative to mitochondrial COI. These markers could avoid problems of mitochondrial markers such as introgression and pseudogenes and identify or delimit species or species-like units as they are not inherited maternally [9].

2.3. Conclusion: combined morphological and molecular analysis

A total of 20 morphological characters yield a two-cluster solution with hierarchical cluster analysis. The first cluster includes 12 species of *Polytremis*, and the second includes the remaining 3 species and outgroups. All supported clades from the combined data matrix are also appearing in the molecular data matrix. ML tree based on COI constructed in this study showed that individuals belonging to the same species formed a monophyletic cluster. Meanwhile, there was considerable congruence in topology of the interspecies level for both mtDNA COI and concatenated sequences of ML trees indicating certain clades were well differentiated phylogenetically (**Figure 1**) [9]. The strong support for the monophyly of *Polytremis* was found in the analyses of the concatenated alignments and COI [9]. In genus *Polytremis*, the results obtained by hierarchical cluster analysis showed traditional classification was basically consistent with molecular phylogeny. However, because the morphological characters and character states were commonly homologous in *Polytremis*, the morphological analysis resulted in only limited resolution based on just 20 morphological characters. On contrary, molecular classification provided a lower artificial and more precise taxonomic rank [15]. Thus, the combination of the morphological and molecular matrix was better resolved for understanding of the phylogeny in the genus.

3. Taxonomic status of two sibling species of *Polytremis* (*Lepidoptera*: Hesperiiidae)

The skipper *P. theca* is widely distributed in south China, except Taiwan, Hainan, and the southern tropical regions of Yunnan Province. Three subspecies have been recorded: *P. t. theca* [9] (west Sichuan and south Shaanxi Province), *P. t. fukia* [16] (Zhejiang to west Sichuan Province), and *Polytremis theca macrotheca* Huang, 2002 (Northwest Yunnan Province). In a preliminary study of molecular phylogeny of the genus *Polytremis* Mabilbe, 1904 using mitochondrial cytochrome c oxidase I (COI), we found the inter-subspecific distance between *P. t. theca* and *P. t. fukia* ranged up to 4.2%, which is higher than some interspecific genetic distances in *Polytremis*. Additionally, *P. theca* is also the only species whose intraspecific distance is more than 3%; thresholds of species identification have been proposed in *Lepidoptera* for COI, in genus *Polytremis* [9]. Thus, we suspected the possible existence of a sibling species paired or the cryptic diversity in the species.

3.1. Genetic divergences and haplotype networks

All 46 samples yielded high quality of DNA. A total of 19 haplotypes were identified in all 46 samples, and the haplotype network was constructed and presented in **Figure 2**. There was no shared haplotype among the four taxa. Haplotypes of the same taxon differed from each other by no more than five mutation distance. The five mutation distances existed between the haplotype Ptt I and Ptt III of *P. t. theca*. The potential ancestral haplotype of *P. t. fukia*, defined by its central position in the network, was designated as Ptf I, which was found in three samples from Tianmushan, one from Jinggangshan, and one from Anjiangping. Ptf II was the most common haplotype in *P. t. fukia* and shared with 10 samples. Haplotype Ptf III was found in two samples from Wuyishan. Haplotype Ptf XIII was identified in two samples from Maoershan and one sample from Anjiangping. The remaining haplotypes of *P. t. fukia* occurred in only one individual (**Figure 2A**).

The data set of nuclear *wingless* contains 390 nucleotide positions without gaps or stop codons, of which 18 positions are variable and 9 are parsimony informative. In total, 10 haplotypes were found in all samples, in which 2 haplotypes were found in *P. t. theca*, 4 in *P. t. fukia*, 3 in *P. nascens*, and 1 in *P. mencia*. Haplotypes of the same taxon differed from each other by no more than two mutation distances, in particular, the 2 haplotypes in 13 samples of *P. t. theca* differed by only one-mutation distance. Five nucleotide substitutions were observed between the potential ancestral haplotypes of *P. t. theca* (Ptt I) and that of *P. t. fukia* (Ptf I) (**Figure 2B**).

Overall, *P. t. theca* had a lower diversity than *P. t. fukia* according to the result of analysis of both mitochondrial COI and nuclear *wingless*. The haplotype diversity (Hd) and nucleotide diversity (π) for *P. t. theca* and *P. t. fukia* are given in **Table 1**. Additionally, they differed from each other by $5.07 \pm 0.49\%$ (4.3–5.9% divergence) for the COI sequences and by $1.70 \pm 0.27\%$ (1.3–2.1% divergence) for the *wingless* sequences.

3.2. Population structure and phylogenetic analysis

The analysis of molecular variance (AMOVA) for the COI sequences of *P. t. theca* and *P. t. fukia* revealed that 88.53% of the genetic variation was among populations and 11.47% was within populations (**Table 2**). The average Φ_{ST} value is 0.896 ($p < 0.01$), suggesting significant genetic variation among the populations. Pair-wise estimates of F_{ST} (0.885) and gene flow ($Nm = 0.065$) between *P. t. theca* and *P. t. fukia* suggest that the subspecies in this species are highly differentiated [16].

Mitochondrial haplotypes sampled from *P. theca* form well-supported clades that closely correspond with subspecific boundaries delimited primarily on the basis of wing color and pattern (**Figure 3A**). The haplotype clades associated with both subspecies are deeply genetically divergent, differing from each other by $5.07 \pm 0.49\%$ (4.3–5.9% divergence). This degree of divergence suggests that evolutionary separation of both subspecies occurred about 0.81 highest probability density (HPD = 0.53–1.28) Mya, likely sometime during the Pleistocene based on a molecular clock calibration of 3.54% pair-wise divergence per million years for




Figure 2. Network profile of (A) COI and (B) *wingless* gene haplotypes based on the nucleotide sequences of *P. t. theca*, *P. t. fukia*, *P. nascens*, and *P. mencia*.

	Ns	Nh	Hd	Nv	π	SD (π)	D	F
All <i>P. theca</i> samples (COI)	33	12	0.875	38	0.0207	0.0019	0.281	2.368
<i>P. t. theca</i> samples (COI)	8	3	0.750	6	0.0064	0.0048	1.598	2.631
<i>P. t. fukia</i> samples (COI)	25	9	0.803	13	0.0050	0.0070	-1.014	-1.886*
All <i>P. theca</i> samples (<i>wingless</i>)	33	5	0.773	9	0.0078	0.0057	1.122	1.509
<i>P. t. theca</i> samples (<i>wingless</i>)	8	2	0.571	1	0.0015	0.0010	1.444	1.100
<i>P. t. fukia</i> samples (<i>wingless</i>)	25	3	0.640	3	0.0030	0.0020	1.080	1.159

Ns, number of samples; Nh, number of haplotypes; Hd, haplotype diversity; Nv, number of variable sites; π , nucleotide diversity; SD, standard deviation; D, Tajima's D statistic; F, Fu's F statistic.

*Significant difference.

Table 1. Genetic diversity and neutrality tests calculated for *P. t. theca* and *P. t. fukia*.

a homologous mtDNA fragment in other insect species [17]. It is noteworthy for the nuclear *wingless* sequences that *P. t. theca* and *P. mencia* are considered distinct species with a genetic divergence of $0.65 \pm 0.15\%$, while the *P. t. fukia* is currently considered a subspecies of the *P. t. theca* despite $1.70 \pm 0.27\%$ sequence divergence. The phylogenetic of *wingless* gene indicates that the *P. theca* is paraphyly with three species sisters to the clade of *P. t. theca* (**Figure 3B**). While 100% of *P. t. fukia* constitutes one separate clade, the clade consisting of *P. t. theca* also includes *P. pellucid*, *P. zina*, and *P. mencia*. The clade consisting of *P. t. theca* is not monophyletic, but complex. This suggests that *P. t. theca* and *P. t. fukia* differ from each other, as evident from the COI tree where they form two separate clades (**Figure 3A**). Concordance between strongly differentiated mtDNA, nuclear haplotype clades, and phenotypic variation supports the hypothesis that both subspecies of *P. theca* deserve recognition at the species level under the general lineage concept of species.

3.3. Demographic inference and estimation of divergence times

Demographic history changes were analyzed for *P. t. theca* and *P. t. fukia* populations through neutrality tests and mismatch distribution. The neutrality tests reveal that the mitochondrial COI appear to be not evolving neutrally as Fu's F values in *P. t. fukia* group are negative significantly (**Table 1**). The Tajima's D and Fu's F values were nonsignificantly positive in

Source of variation	df	Sum of squares	Variance components	Percentage variation	Φ statistic
Among populations	1	169.650	9.98266 Va	88.53	–
Within populations	11	45.269	1.29341 Vb	11.47	0.896 ($p < 0.01$)
Total	12	214.919	11.27607		
Fixation index	0.8853				

Table 2. Analysis of molecular variance (AMOVA) for the COI sequences of *P. t. theca* and *P. t. fukia*.




Figure 3. The maximum-likelihood tree for (A) mitochondrial COI and (B) nuclear *wingless* haplotypes of *Polytreemis*.

P. t. theca group and all *P. theca* sample group. The mismatch analysis yielded a unimodal distribution of pair-wise differences for *P. t. fukia* compared to the multimodal distribution of *P. t. theca* samples and the pooled samples. According to Rogers and Harpending [18], the observed curves with unimodal represent population expansion and the observed curves with many peaks or resemblance to expected curves mean equilibrium population, which further elucidates the demographic history of *P. theca*. The results suggest population expansion in *P. t. fukia* and population equilibrium in *P. t. theca*. We still confirm the result of the population size change in haplotype network (Figure 2). Statistical parsimony network reflects genealogical relationships of the mtDNA haplotypes, that is, single mutation steps separate adjacent haplotypes in the network and older haplotypes are placed at internal branching points, whereas younger ones occur toward the tip positions [19]. The haplotype network of *P. t. fukia* displays a star-like pattern (Figure 2). Haplotype I, the second most common and geographically widespread in central-west of China, is in the star's center, and derivatives are connected to it by short branches [16]. Based on coalescence theory, the star-like topologies for this cluster strongly suggest the effect of a population expansion [20]. Divergence time analysis with an uncorrelated lognormal relaxed clock run in Bayesian MCMC analysis of molecular sequences (BEAST) produced a tree with a topology similar to ML tree (Figure 4).




Figure 4. Bayesian inference (BI) tree of mtDNA datasets for *Polytremsis* using uncorrelated lognormal relaxed clock.

P. t. theca diverged from *P. t. fukia* around 0.81 (HPD = 0.53–1.28) million years ago (Mya) during the Pleistocene (node a in **Figure 4**). *P. theca* diverged from other congeners included in the analysis about 1.36 (HPD = 1.02–1.53) Mya during the Pleistocene eras (node b in **Figure 4**). In our study, a higher F_{ST} value indicated a lower level of gene flow (N_m) and higher genetic differentiation among populations. The results of two-level AMOVA show that significant genetic variation exists among the examined populations. These results provide a second line of support to a conclusion that the *P. t. fukia* is a different species [16].

3.4. Conclusion

There is a small region of overlap in west Sichuan province in the distribution of *P. t. theca* and *P. t. fukia*, but otherwise they are not sympatric. *P. t. theca* inhabits the higher elevations of west Sichuan and south Shaanxi Province [21]. *P. t. fukia* occurs in the whole Southern China except Taiwan, Hainan, and south Yunnan Province, from Zhejiang to west Sichuan Province [22]. According to the description on morphological variation between *P. t. theca* and *P. t. fukia* [23], we found a different morphological feature existing in female genitalia except for the different colors and spot numbers in some part of wings (**Table 3**). The lateral edge of lamella antevaginalis of female *P. t. fukia* is more rounded than that of *P. t. theca*. We suspected the taxonomic status of the subspecies from their geographic separation and the morphological variation. Our investigations and analyses revealed significant molecular and biogeographical differences between *P. t. theca* and *P. t. fukia*. We propose that *P. t. fukia* should be treated as a distinct species called *Polytremsis fukia* [16] under the phylogenetic species concept. In fact, it has been

	Wing			Genital		
	Color of cilia of wings	Color of underside ground	Number of spots in space Cu2 of the forewing	Color of scales scattered in costa and subapical area of forewing	Color of scales scattered in discal area and dorsum of hind wing	Ductus bursae
<i>P. t. theca</i>	Brown	Yellowish brown	0	Greenish ochreous	Greenish ochreous	Thin
<i>P. t. fukia</i>	Grayish white	Greenish ochreous	1 or 2	Grayish white	Grayish white	Thick

Table 3. Different morphological features of genitals and wings between *P. t. theca* and *P. t. fukia*.

recently found that other species previously considered subspecies based on morphology are in fact sibling species that passed unnoticed until the advent of molecular techniques [24]. Results from our study strengthen information about the *Polytremis* species complex and help in developing appropriate integrated pest management strategies for these insect pests.

4. The application of mitochondrial genome and microsatellite DNA in *Polytremis*

With the development of the research, the single molecular fragment cannot meet the research requirements. New molecular markers need to be explored. Recently, the mitochondrial genome has become one of the important molecular markers to explore different categories of *Lepidoptera*. Additionally, the ideal molecular marker, microsatellite DNA, has become very prevalent in molecular ecological studies including inferring population genetic structure, exploring taxonomic status, and studying reproductive ecology.

4.1. Complete mitochondrial DNA genome of *P. nascens* and *Polytremis jigongji*

Lepidoptera is the second largest insect order in the world and contains more than 160,000 species. However, the information currently existing on lepidopteran mitogenomes is limited. To date, only 236 complete or nearly complete mitogenome sequences have been determined to belong to six superfamilies. The phylogenetic inference based on the variations at such short gene sequences is not always robust and may lack sufficient resolution compared to the phylogenies based on longer mitogenome sequences [25]. Additionally, mitogenomes are also applied to the studies on comparative and evolutionary genomics [26], molecular evolution [27], phylogeography [28], etc. Thus, further insight into *Lepidopteran* phylogeny and evolution awaits more related species sequences to be determined [29].

The complete mtDNA genome of *P. nascens* was a circular molecule of 15,392 bp in length, including the standard 13 protein-coding genes (PCGs), 2 ribosomal RNA (rRNA) genes, 22

transfer RNA (tRNA) genes, and 1 noncoding region. The overall-based composition was 39.7% A, 40.7% T, 7.7% G, and 11.9% C, with a slight A+T bias of 80.4%. Thirteen PCGs and two rRNA were first identified using an open reading frame (ORF) finder, specifying the invertebrate mitochondrial genetic code. Then, they were calibrated by sequence similarity using published lepidopteran mitogenome sequences. All PCGs use standard ATN (ATT, ATG, or ATA) as the start codon except COX1 that uses CGA. Eight PCGs (ND2, ATP8, ATP6, COIII, ND3, ND4, ND6, and ND1) employ the typical stop codon TAA, while the remaining five PCGs terminate with a single T 33 [22].

The complete mtDNA genome of *P. jigongi* was a circular molecule of 15,353 bp in length, including the standard 13 protein-coding genes (PCGs), 2 ribosomal RNA (rRNA) genes, 22 transfer RNA (tRNA) genes, and 1 noncoding region. Its organization and arrangement are identical to other skippers [30, 31]. The overall-based composition was 39.8% A, 41.1% T, 7.6% G, and 11.5% C, with a slight A+T bias of 80.9%. Thirteen PCGs and two rRNA were identified using an open reading frame (ORF) finder and calibrated by sequence similarity using published lepidopteran mitogenome sequences. All PCGs use standard ATN (ATT, ATG, or ATA) as the start codon except COX1 that uses CGA. Eight PCGs (ND2, ATP8, ATP6, COIII, ND3, ND4, ND6, and ND1) employ the typical stop codon TAA, while the remaining five PCGs terminate with TA or T [22]. Alignment of amino acid sequences of each of individual 13 PCGs was performed through Clustal X [32], and the phylogenetic analysis was carried out using neighbor-joining (NJ) method with MEGA version 5.0 program [33] (Figure 5).




Figure 5. Phylogenetic tree of the lepidopterans based on 13 PCG nucleotide sequences of the mitogenome using NJ analysis.

4.2. Isolation and characterization of microsatellite loci in *P. nascens* and *P. fukia*

Microsatellites are highly polymorphic and codominant molecular markers based on simple repeated and frequent sequences common in the all-living organisms, which have proven to be a powerful tool available in population genetic and evolutionary studies [34]. The ideal molecular marker has become very prevalent in studies of insects over the last 10 years [35]. Variability of the 12 polymorphic microsatellites was surveyed in 53 individuals of *P. nascens*. We found 53 different multilocus genotypes, and the number of alleles per locus ranged from 3 to 12. Observed (HO) and expected heterozygosity (HE) values ranged from 0.33 to 0.71 and from 0.61 to 0.90. Only one locus (Polynyas 13) ($p < 0.01$) that deviated from HWE showed significant heterozygote deficit in the populations of Lianglu, Shengtangshan, and Maoershan [33]. Analysis performed with Micro-Checker showed that the deviation was attributed to the homozygote excess with null allele frequency of 0.2041, 0.2914, and 0.3533 in the three populations, respectively [33]. No linkage disequilibrium was detected for any pair of loci ($p > 0.01$) in any populations following Bonferroni correction [33]. Comparisons of pair-wise F_{ST} among six populations show the genetic difference based on 12 microsatellite loci (Table 4). The population of Baishanzu showed the largest pair-wise F_{ST} values, corresponding to relatively geographic isolation from the other five populations [33].

The variability of the 11 polymorphic microsatellites was surveyed in 21 individuals of *P. fukia*. We found 21 different multilocus genotypes, and the number of alleles per locus ranged from 5 to 10. Observed heterozygosity (H_o) values ranged from 0.48 to 0.65 and expected heterozygosity (H_e) values ranged from 0.69 to 0.86. Polymorphic information content (PIC) ranged from 0.59 to 0.88 per locus, and all markers were highly informative ($PIC > 0.5$) [33]. All loci were in Hardy-Weinberg equilibrium, consistent with inbreeding and/or Wahlund effects. No linkage disequilibrium was detected in any pair of loci [33]. Micro-Checker software found no evidence of scoring error due to stuttering or large allele dropout. We also tested the selected primers for amplification on eight other *Polytremis* species: *P. caerulescens* Mabille, *P. jigongi* Zhu, *P. theca* Evans, *P. zina* Evans, *P. mencia* Moore, *P. lubricans* Herrich-Schäffer, *P. eltola* Hewitson, and *P. discrete* Elwes & Edwards. Of the 11 markers, we found that cross-amplification was successful for 10 loci in at least one conge-

	Baishanzu	Hailuogou	Lianglu	Shengtangshan	Maoershan	Kuankuoshui
Baishanzu						
Hailuogou	0.044					
Lianglu	0.050	0.003				
Shengtangshan	0.029	0.025	0.022			
Maoershan	0.023	0.021	0.020	0.013		
Kuankuoshui	0.026	0.012	0.014	0.022	0.017	

Table 4. Comparisons of multilocus pair-wise F_{ST} values ($p < 0.05$) among 6 regional populations of *P. nascens* based on 12 microsatellite loci.

neric species [33]. Two primer loci amplified satisfactorily among all congeneric species, indicating that these loci may be useful for population genetics, including phylogeography, species cohesion and delimitation, and barriers to gene flow for other congeneric and related species [33].

5. *Wolbachia* infection and influence in *Polytremis* species

Wolbachia may be the most widespread endosymbiont in terrestrial ecosystems. The maternally inherited endosymbiotic bacteria infect perhaps two-thirds of present-day insect species [36, 37]. The transmission of *Wolbachia* is primarily vertical and secondarily horizontal [38]. The bacteria manipulate the reproduction of their host to ensure their vertical transmission by cytoplasmic incompatibility, feminization, male killing, and parthenogenesis [39]. *Wolbachia* can potentially influence mitochondrial variation of their hosts. The linkage disequilibrium is expected to occur between them since they are co-transmitted maternally. The rapidly spread of *Wolbachia* in the host populations can result in the hitch-hiking effect of mitochondrial DNA [31]. One particular mitochondrial haplotype can reduce mtDNA polymorphism in the infected population and sweep through a population [19]. *Wolbachia* may also drive introgression of mtDNA following hybridization events between sibling species: the introduction and spread of *Wolbachia* in a novel species result in spread of the mtDNA from the neighboring species [40].

Wolbachia infections have been reported in various *Lepidoptera* families such as Papilionidae, Lycaenidae, Pieridae, Nymphalidae, Hesperidae, Pyralidae, Noctuidae, and Lasiocampidae. A few butterfly species harboring the bacteria have been thoroughly studied. We have reported the molecular phylogeny of the genus in a prior study [9]. Meanwhile, we have preliminarily screened for the presence of *Wolbachia* in *Polytremis* and found at least three species (*P. nascens* Leech, 1893, *P. theca* Evans, 1937 and *P. pellucid* Murray, 1875) are infected with *Wolbachia*.

5.1. *Wolbachia* infection status and genetic structure in natural populations of *P. nascens*

We surveyed *Wolbachia* infection rates in 14 regional populations of *P. nascens*. Twenty-one specimens (31%) infected with *Wolbachia* in seven regional populations were detected. However, all specimens are free from infection with *Wolbachia* in the other seven regional populations. The infection rates between the female and male butterflies show no significant difference ($\chi^2 = 0.65$, $p > 0.05$). The further rearing experiment would be made to reveal whether there was a sex-ratio distortion in *P. nascens* induced by the strain of *Wolbachia*. A positive relationship of *Wolbachia* infections and latitudinal distribution has also been found. The lower or no *Wolbachia* infection rates in central-west of China may indicate that incidence is apparently lower in regions experiencing longer dry seasons and higher average daily temperatures. The higher *Wolbachia* prevalence occurs in more southerly moist and temperate populations (**Figure 6**). This has been observed in the beetle *Chelymorpha alternans* and in ants of the genus *Solenopsis* [41, 42].




Figure 6. Distribution of mtDNA haplotypes among populations of *P. nascens* collected in China.

Many explanations have been proposed that deviation from neutral evolution in a species and the absence of diversity in mtDNA can be associated with either a genome-wide bottleneck effect or a selective sweep on mtDNA [43]. In our study, the uninfected butterflies show higher mtDNA polymorphism than butterflies infected with *Wolbachia* (Table 5). The perfect concordance of *Wolbachia* infection status and mtDNA polymorphism suggests that the mitochondrial genetic structure of the host insects may be strongly affected by the *Wolbachia* infection. Decreased mtDNA polymorphism as a consequence of *Wolbachia* infection has also been reported in several other insects [38, 40]. We also found that the mtDNA genes of the butterflies infected with *Wolbachia* deviated significantly from neutral evolution according to both D (-2.3303 , $p < 0.05$) and F values (-3.7068 , $p < 0.05$), while this was not so for uninfected ones [31]. These results suggest that the populations of *P. nascens* have

	N	Number of haplotypes	Haplotype diversity	Number of variable sites (S)	π	SD (π)	D	F
All sequences	67	12	0.797	119	0.0197	0.0012	1.4795	2.0907*
Infection	21	2	0.095	12	0.0006	0.0005	-2.3303*	-3.7068*
Free from infection	46	10	0.750	77	0.0122	0.0015	0.8835	1.9320

N, number of samples; π , nucleotide diversity; SD, standard deviation; D, Tajima's D statistic; F, Fu's F statistic.
 *Significant difference.

Table 5. Mt haplotype and nucleotide diversity estimates from infected and uninfected samples.

recently been subjected to a *Wolbachia*-induced sweep, making the mtDNA undergo purifying selection. Additionally, we analyzed the population size change of *P. nascens* infected with *Wolbachia* and all *P. nascens*, respectively, by the software DnaSP4.90 [7] and got no evidence for population expansion in *Wolbachia*-infected group [18]. However, we got unimodal curves representing population expansion in *P. nascens*. A selective sweep would erase variability in the population even under population expansion, potentially eliminating any evidence of past demographic processes. Moreover, recent studies also revealed that there are some other endosymbionts known to manipulate host reproduction like *Wolbachia*, for example, *Arsenophonus*, *Cardinium*, and *Rickettsia* [44]. A wide range of insect species can host more than one endosymbiont [42]. The infection status of the secondary endosymbionts in *P. nascens* needs to be further studied.

Although there seems to be a strong association existed between mtDNA haplotypes and *Wolbachia* infection status, the association between mtDNA haplotypes, *Wolbachia* infection, and geographical distribution is weak [31]. In our preliminary experiment, we found that two sympatrically distributed sister species of *P. nascens* (*P. theca* and *P. pellucid*) are infected with *Wolbachia*. We could not eliminate the possibility of the multiple introgression events and hybridizations between the species pairs [40]. Thus, we excluded the infected group (Clade I) in the analysis of biogeographical implications of *P. nascens*. It is notable that if we only take the uninfected group into account, the conclusion can be drawn that *P. nascens* probably has two genetically diverse and geographically localized clades in China based on the mtDNA haplotype phylogeny and networks (**Figure 6**) [31]. They are the central-west of China clade (H1–H3) and the Eastern and Southern China clade (H4–H10). These two clades are isolated mainly by the Yangtze River, with the exception of a specimen in HH [31]. Our results were similar to those obtained from the striped stem borer, *Chilo suppressalis* [45], and the melon fly, *Bactrocera cucurbitae* [46], which suggest that the Yangtze River Range has acted as a substantial barrier to gene flow. This contrasts with studies of the beet armyworm, *Spodoptera exigua*, and the migratory locust, *Locusta migratoria*, and the cotton bollworm *Helicoverpa armigera*, which showed little or no evidence that the Yangtze River Range limits gene flow [47–49].

The network analysis reflects genealogical relationships of the mtDNA haplotypes [31]. The single mutation steps separate adjacent haplotypes in the network and older haplotypes are placed at internal branching points, whereas younger ones occur toward the tip positions [50]. The haplotype network of *P. nascens* displays a star-like pattern (**Figure 6**). Haplotype 1 (H1) is in the star's center, and derivatives are connected to it by short branches. The haplotype is the most common and geographically widespread in central-west of China. The star-like topologies for this cluster strongly suggest the effect of a population expansion based on coalescence theory [20]. We still confirm the result of the population size change of *P. nascens* with the software DnaSP4.90 and got unimodal curves representing population expansion [31]. The most common H1 had strong support as the ancestral haplotype due to its representation in a significant proportion of individuals in all populations and its basal location in the network [31]. A reticulation connecting multiple haplotypes from the Eastern and Southern China clade was formed in the network [31].

5.2. A prevalence survey of *Wolbachia* in *P. fukia*

Of the butterflies examined by diagnostic PCR for 16S rRNA, 47% (15/32) were *Wolbachia* positive. The infection rates in female and male are 69% (11/16) and 25% (4/16). For characterization of *Wolbachia* strains, we amplified a segment of the *Wolbachia* cell cycle gene *ftsZ*. BLAST searches of the *ftsZ* sequences yielded a significant sequence similarity between the *Wolbachia* strains infecting *P. fukia* and the *Wolbachia* supergroup A typically found in insects. Phylogenetic analyses suggested that the bacteria are subdivided into two strains with a genetic divergence of approximately 2.8%. These strains will be referred to as *wFuk1* and *wFuk2* in the following.

Figure 7A shows the ML tree based on the data set of the concatenated sequences and supports the monophyly of *P. fukia*. On the phylogeny, the sequences are split into three clades supported by high bootstrap values. Clade I exclusively consists of the *P. fukia* from two geographical populations in Southwest side of their distribution area (Maoershan and Anjiangpin). These butterflies are either free from *Wolbachia* infection or infected with *wFuk2*. Clade II exclusively consists of the butterflies from four geographical populations (West Tianmushan, Qingliangfeng, Wuyishan, and Jinggangshan), which are either free from *Wolbachia* infection or infected with *wFuk2*. These two clades are consistent with the distribution of geographical population. Eight females constitute Clade III are invariably infected with *wFuk1*. In our data set comprising 32 individuals (16♀ and 16♂), we did not detect any male *P. fukia* infected




Figure 7. (A) Maximum-likelihood phylogeny on the basis of the concatenated mitochondrial sequences and (B) network on the basis of the concatenated mitochondrial sequences constructed with software Network4.5.

	<i>N</i>	Number of haplotypes	Hd	SD (Hd)	Number of variable sites (S)	π	SD (π)	D	F
All sequences	32	23	0.980	0.012	50	0.008	0.001	0.568	-3.913
Infection with <i>wFul1</i>	8	4	0.821	0.101	5	0.001	0.000	0.840	0.428
Infection with <i>wFul2</i>	7	7	1.000	0.076	32	0.008	0.001	0.077	-1.085
Free from infection	17	14	0.978	0.027	36	0.007	0.002	0.522	-2.510

N, number of samples; Hd, haplotype diversity; π , nucleotide diversity; SD, standard deviation; D, Tajima's D statistic; F, Fu's F statistic.

Table 6. Mt haplotype and nucleotide diversity estimates from infected and uninfected samples.

with *wFuk1*. All eight individuals infected with *wFuk1* are female. Males belong to Clade I or Clade II and, if infected, the strains *wFuk2*. All mtDNA of *P. fukia* were used for network construction with the software Network4.5 using the median-joining method (**Figure 7B**). Twenty-three haplotypes are clustered into three groups in accordance with three clades in ML tree (see **Figure 7A**). Two haplotypes were shared by the individuals infected with *Wolbachia* and free from *Wolbachia*. The *Wolbachia*-induced sweep has been shown in *P. nascens* [51]. However, such *Wolbachia* effects on mtDNA variation in *P. fukia* examined in this study were not as conspicuous as in the above studies. Still, mtDNA variation in the *P. fukia* is likely to be weakly associated with the presence of *wFuk1*, although the sample size was not large enough to draw a firm conclusion [37]. The nucleotide diversity in *wFuk1*-infected butterflies is smaller than that of uninfected butterflies (**Table 6**) and their mt concatenated sequences form a clade solely (**Figure 7**), presumably reflecting a *Wolbachia* sweep. This tendency was consistently seen in all the mitochondrial regions examined. A larger sample will reveal more in detail the association of *Wolbachia* infection status with variation of mtDNA in *P. fukia*.

Acknowledgements

This study was financially supported by grants from National Natural Science Foundation of China (no. 31401997) and Young Teacher Training Scheme for Colleges and Universities in Shanghai (no. ZZssd15068).

Author details

Weibin Jiang

Address all correspondence to: jiangwb@shnu.edu.cn

College of Life and Environmental Sciences, Shanghai Normal University, Shanghai, People's Republic of China

References

- [1] Huang H. Some new butterflies from China-2. *Atalanta*. 2002;**33**:109-122
- [2] Zhu JQ, Chen ZB, Li LZ. *Polytremis jigongi*: A new skipper from China (*Lepidoptera*: *Hesperiidae*). *Zootaxa*. 2012;**3274**:63-68
- [3] Hebert PDN, Penton E, Burns J. Ten species in one: DNA barcoding reveals cryptic species in the neotropical skipper butterfly *Astrartes fulgerator*. *Proceedings of National Academy Sciences*. 2004;**101**:14812-14817
- [4] Hwang UW, Ree HI, Won K. Evolution of hypervariable regions, V4 and V7, of insect 18S rRNA and their phylogenetic implications. *Zoological Science*. 2000;**17**:111-121
- [5] Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular evolutionary genetics analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*. 2007;**24**:1596-1599
- [6] Swofford DL. PAUP* Phylogenetic Analysis Using Parsimony (*and other methods). Version 4.0b10 ed. Massachusetts: Sunderland Sinauer Associates; 2002
- [7] Rozas J, Sánchez-Delbarrio JC, Messeguer X, Rozas R. DnaSP, DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics*. 2003;**19**:2496-2497
- [8] Bandelt HJ, Foster P, Röhl A. Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution*. 1999;**16**:37-48
- [9] Jiang WB, Zhu JQ, Song C, Li XY, Yang Y, Yu WD. Molecular phylogeny of the butterfly genus *Polytremis* (*Hesperiidae*, *Hesperiinae*, Baorini) in China. *PLoS ONE*. 2013;**8**:e84098
- [10] Huang H. A list of butterflies collected from Nujiang and Dulongjiang, China with descriptions of new species, new subspecies and revisional notes. *Neue Entomologische Nachrichten*. 2003;**55**:3-114
- [11] Raupach MJ, Astrin JJ, Hannig K, Peters MK, Stoeckle MY, Wägele JW. Molecular species identification of Central European ground beetles (*Coleoptera*: *Carabidae*) using nuclear rDNA expansion segments and DNA barcodes. *Frontiers in Zoology*. 2010;**7**:26
- [12] Sugiyama H. New butterflies from western China. *Pallarge*. 1999;**6**:1-14
- [13] Zhu JQ. A Taxonomic Study on the Tribe Baorini Doherty, 1886 (*Lepidoptera*: *Hesperiidae*: *Hesperiinae*) of China. Shanghai: Shanghai Normal University. Dissertation; 2012
- [14] Meyer CP, Paulay G. DNA barcoding: Error rates based on comprehensive sampling. *PLoS Biology*. 2005;**3**:2229-2238
- [15] Tautz D, Arctander P, Minelli A, Thomas RH, Vogler AP. A plea for DNA taxonomy. *Trends in Ecology and Evolution*. 2003;**18**:70-74

- [16] Jiang WB, He HY, Li YD, Ren MY, Ma YZ, Zheng LL, Zhu JQ, Yu WD*. Taxonomic status and molecular phylogeography of two sibling species of *Polytremis* (*Lepidoptera: HesperIIDae*). *Scientific Reports*. 2016;**6**:20820
- [17] Papadopoulou A, Anastasiou I, Vogler AP. Revisiting the insect mitochondrial molecular clock: The mid-Aegean trench calibration. *Molecular Biology and Evolution*. 2010;**27**:1659-1672
- [18] Rogers AR, Harpending H. Population growth makes waves in the distribution of pairwise genetic differences. *Molecular Biology and Evolution*. 1992;**9**:552-569
- [19] Narita S, Nomura M, Kato Y, Fukatsu T. Genetic structure of sibling butterfly species affected by *Wolbachia* infection sweep: Evolutionary and biogeographical implications. *Molecular Ecology*. 2006;**15**:1095-1108
- [20] Slatkin M, Hudson RR. Pairwise comparisons of mitochondrial DNA sequences in stable and exponentially growing populations. *Genetics*. 1991;**129**:555-562
- [21] Chou I. In: *Monographia Rhopalocerorum Sinensium*. Zhengzhou: Henan Scientific and Technological Press; 1994
- [22] Jiang WB, Zhu JQ*, Yang QC, Zhao HD, Chen MH, He HY, Yu WD*. Complete mitochondrial DNA genome of *Polytremis nascens* (*Lepidoptera: HesperIIDae*). *Mitochondrial DNA Part A*. 2016;**27**(5):3131-3132
- [23] Eliot JN, Corbet AS, Pendlebury HM. In: *The butterflies of the Malay Peninsula*. 4th ed. Kuala Lumpur: United Selangor Press; 1992
- [24] Evans III EL, Martinez-Gomez JE, Sehgal RNM. Phylogenetic relationships and taxonomic status of the endemic Socorro Warbler (*Setophaga pitiayumi graysoni*). *Journal of Ornithology*. 2015;**156**:363-370
- [25] Powell AFLA, Barker FK, Lanyon SM. Empirical evaluation of partitioning schemes for phylogenetic analyses of mitogenomic data: an avian case study. *Molecular Phylogenetics and Evolution*. 2013;**66**:69-79
- [26] Ballard JWO. Comparative genomics of mitochondrial DNA in *Drosophila simulans*. *Journal of Molecular Evolution*. 2000;**51**:64-75
- [27] Zakharov EV, Caterino MS, Sperling FAH. Molecular phylogeny, historical biogeography, and divergence time estimates for swallowtail butterflies of the genus *Papilio* (*Lepidoptera: Papilionidae*). *Systematic Biology*. 2004;**53**:193-215
- [28] Malyarchuk B, Derenko M, Denisova G. A mitogenomic phylogeny and genetic history of sable (*Martes zibellina*). *Gene*. 2014;**550**:56-67
- [29] Wang JP, Cao TW, Xuan SB, Wang H, Zhang M, Ma EB. The complete mitochondrial genome of *Sasakia funebris* (Leech) (*Lepidoptera: Nymphalidae*) and comparison with other Apaturinae insects. *Gene*. 2013;**526**:336-343

- [30] Hao J, Sun Q, Zhao H, Sun X, Gai Y, Yang Q. The complete mitochondrial genome of *Ctenoptilum vasava* (*Lepidoptera: Hesperidae: Pyrginae*) and its phylogenetic implication. *Comparative and Functional Genomics*. 2012;**2012**:328049. DOI: 10.1155/2012/328049
- [31] Jiang WB, Zhu JQ, Chen MH, Yang QC, Du X, Chen SY, Zhang LN, Yu YM, Yu WD. *Wolbachia* infection status and genetic structure in natural populations of *Polytremis nascens* (*Lepidoptera: Hesperidae*). *Infection, Genetics and Evolution*. 2014;**27**:202-211
- [32] Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG. The CLUSTAL_X windows interface: Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Research*. 1997;**25**:4876-4882
- [33] Jiang WB, Chen MH, He HY, Wu YJ, Zhu JQ*, Yu WD*. Eleven microsatellite loci developed from the Chinese Hesperid butterflies, *Polytremis fukia* (*Lepidoptera: Hesperidae*), and cross-amplification in other species of *Polytremis*. *Applied Entomology and Zoology*. 2016;**51**(4):669-672
- [34] Goldstein D, Schlötterer C. *Microsatellites: Evolution and Applications*. Oxford: Oxford University Press; 1999
- [35] Zakharov EV, Lobo NF, Nowak C, Hellmann JJ. Introgression as a likely cause of mtDNA paraphyly in two allopatric skippers (*Lepidoptera: Hesperidae*). *Heredity*. 2009;**102**:590-599
- [36] Hilgenboecker K, Hammerstein P, Schlattmann P, Telschow A, Werren JH. How many species are infected with *Wolbachia*?—A statistical analysis of current data. *FEMS Microbiology Letters*. 2008;**281**:215-220
- [37] Jiang WB, Wu YJ, He HY, Zhu JQ, Yu WD*. A prevalence survey of *Wolbachia* in *Polytremis fukia* (*Lepidoptera: Hesperidae*). *Journal of Asia-Pacific Entomology*. 2016;**19**(2):399-404
- [38] Raychoudhury R, Grillenberger BK, Gadau J. Phylogeography of *Nasonia vitripennis* (Hymenoptera) indicates a mitochondrial-*Wolbachia* sweep in North America. *Heredity*. 2010;**104**:318-326
- [39] Stouthamer R, Breeuwer JA, Hurst GD. *Wolbachia pipientis*: Microbial manipulator of arthropod reproduction. *Annual Review of Microbiology*. 1999;**53**:71-102
- [40] Jackel R, Mora D, Dobler S. Evidence for selective sweeps by *Wolbachia* infections: Phylogeny of *Altica* leaf beetles and their reproductive parasites. *Molecular Ecology*. 2013;**22**:4241-4255
- [41] Keller GP, Windsor DM, Saucedo JM, Werren JH. Reproductive effects and geographical distributions of two *Wolbachia* strains infecting the neotropical beetle, *Chelymorpha alternans* Boh. (*Chrysomelidae, Cassidinae*). *Molecular Ecology*. 2004;**13**:2405-2420
- [42] Martins C, Souza RF, Bueno OC. Presence and distribution of the endosymbiont *Wolbachia* among *Solenopsis* spp. (*Hymenoptera: Formicidae*) from Brazil and its evolutionary history. *Journal of Invertebrate Pathology*. 2012;**109**:287-296

- [43] Shoemaker DD, Dyer KA, Ahrens M, McAbee K, Jaenike J. Decreased diversity but increased substitution rate in host mtDNA as a consequence of *Wolbachia* endosymbiont infection. *Genetics*. 2004;**168**:2049-2058
- [44] Duron O, Bouchon D, Boutin S, Bellamy L, Zhou L, Engelstädter J, Hurst GD. The diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC Biology*. 2008;**6**:27
- [45] Meng XF, Shi M, Chen XX. Population genetic structure of *Chilo suppressalis* (Walker) (*Lepidoptera*: Crambidae): Strong subdivision in China inferred from microsatellite markers and mtDNA gene sequences. *Molecular Ecology*. 2008;**17**:2880-2897
- [46] Zhu ZH, Ye H, Zhang Z. Genetic relationships among four *Bactrocera cucurbitae* geographic populations in Yunnan Province. *Chinese Journal of Applied Ecology*. 2005;**16**:1889-1892
- [47] Niu CW, Zhang QW, Ye ZH, Luo LZ. Analysis of genetic diversity in different geographic populations of the beet armyworm *Spodoptera exigua* (*Lepidoptera*: Noctuidae) with AFLP technique. *Acta Entomologica Sinica*. 2006;**49**:867-873
- [48] Zhang MZ, Kang L. The genetic differentiation among geographic populations of *Locusta migratoria* in China. *Scichina*. 2005;**35**:220-230
- [49] Xu G, Wang GR, Wu KM, Guo YY. Gene flow analysis among different geographical populations of *Helicoverpa armigera* (Hübner) by restriction fragment length polymorphisms. *Cotton Science*. 2002;**14**:352-355
- [50] Yang QC, Zhao HD, Chang Y, Liu L, Zhu JQ, Yu WD, Jiang WB*. Complete mitochondrial genome of the Chinese skipper, *Polytremis jigongi* (*Lepidoptera*: Hesperiiidae). Mitochondrial DNA Part A. 2016;**27**(4):2375-2376. (SCI)
- [51] Jiang WB, Zhu JQ, Zhan L, Chen MH, Song C, Yu WD. Isolation and characterization of microsatellite loci in *Polytremis nascens* (*Lepidoptera*: Hesperiiidae), and their cross-amplification in related species. *Applied Entomology and Zoology*. 2014;**49**:177-181