

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



The Purview of Phytotherapy in the Management of Gastric Ulcer

Sabiu Saheed, Ajani Emmanuel Oladipo,
Taofik Olatunde Sunmonu,
Fatai Oladunni Balogun and
Anofi Omotayo Tom Ashafa

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.70007>

Abstract

Stomach/gastric ulcer is a debilitating disease affecting more than 10% of the global population. Sufferers often have chronic pains with life-threatening gastrointestinal haemorrhage or perforation. Since the first diagnosis of stomach ulcer (SU) in the 19th century, excessive gastric juice that eroded the mucosa of the stomach was opined as its major cause. Efforts were channelled toward effective control of the resulting acid build-up through the use of antiulcer medications and reduction in stress-induced activities, which may aggravate gastric hyperacidity. An intense treatment option involved vagotomy (surgically severing the nerves surrounding an ulcer) to prevent hyperacidity and further perforation of the stomach epithelium. Despite these interventions, SU disease remained an impediment to clinical practice. Literatures revealed that many botanicals have been used to treat SU and this is hinged on their being endowed with antiulcerogenic phytonutrients of therapeutic significance. In this review, attempts have been made to highlight the main mechanisms of action and limitations of the conventional antiulcerogenic drugs, various antiulcerogenic experimental models, as well as compile selected medicinal plants and their implicated phytonutrients that will ultimately and eventually present effective and globally competitive exciting opportunities for the development of new lead therapeutics for the management of SU disorders.

Keywords: antiulcerogenic, gastric ulcer, gastropathy, hemorrhage, *Helicobacter pylori*, pepsin, perforation, phytonutrients, vagotomy

1. Introduction

Ulcer is an open sore of the biological membrane characterized by sloughing of inflamed dead tissue [1]. More specifically, it could either occur as a lesion on the surface of the skin

or a mucous membrane with significant superficial loss of tissue. Although, ulcers may be encountered at almost any part of the body, they are mostly found on the skin of the lower extremities and in the gastrointestinal tract [2]. There are many types of ulcer including mouth, esophageal, peptic and genital ulcer. Of these, peptic ulcer (PU) is the most prevalent [2]. The PUs are erosion of lining of either the stomach or duodenum [3] and this has availed the two most common types of PU as the gastric/stomach ulcer (SU) and duodenal ulcer. A person may have both gastric and duodenal ulcers at the same time. SUs are located in the stomach and mainly characterized by hemorrhage and pain. Other symptoms may include nausea, vomiting, and weight loss. Although patients with SU have normal or diminished acid production, yet ulcers may occur even in complete absence of acid [3]. Generally, pain occurs when the stomach is empty and relieves after eating. In some cases, SU can be life threatening with symptoms like bloody stool, severe abdominal pain, and cramps coupled with blood vomiting [4].

Under normal conditions, a physiologic balance exists between gastric acid secretion and mucosal defense. The epithelial cells of the stomach secrete mucus in response to irritation of the epithelial lining and as a result of cholinergic stimulation [5]. Ordinarily, the superficial portion of the gastric mucosa is jelly-like and impermeable to acid and pepsin. Other gastric cells secrete bicarbonate, which buffers acid that lies penultimate to the mucosa. Also, the prostaglandins of the E (PGE) type of the epithelia offered significant protection by increasing the bicarbonate content and consequently strengthening the mucous layer. However, when the acid and pepsin enters the epithelial cells, further fortifying mechanisms are triggered to ameliorate injury [5]. This may be best observed within the epithelial cells, where ion pumps in the basolateral cell membrane aids intracellular pH regulation through removal of excess H^+ and subsequent migration of healthy cells to the site of injury. By so doing, the acid that diffused through the injured mucosa is effectively removed by the flow of blood in the mucosa. This also provides bicarbonate to the superficial epithelial cells. However, when there is disequilibrium between offensive (acid, pepsin, and *Helicobacter pylori*) and defensive factors (mucin, prostaglandin, bicarbonate, nitric oxide, and growth factors), the pathophysiological features of SU become evident [6]. Although, SU was once believed to be caused by spicy food and stress, they have however been established to be mere aggravating factors while the real causes include reaction to various medications, particularly nonsteroidal anti-inflammatory drugs (NSAIDs) and *H. pylori* infection [7]. *H. pylori*, NSAIDs, emotional stress, alcohol abuse, and smoking are the principal etiological factors associated with SU [8]. Usually, SU occur as breaks across the entire length of the stomach epithelia and in some cases, the deeper layers of the muscle wall are considerably affected. This disruption of the mucosal integrity may potentiate perforation, bleeding, obstruction, pain and death, if proficient treatments are not timely administered. The exclusive production of urease by *H. pylori* renders its microenvironment alkaline and allows its long-lasting survival in the hostile acidic environment of the stomach, where it either worsens the severity of SU disease or merely causes mucosal inflammation. More often in SU cases, inflammation is secondary to the colonizing action of *H. pylori* on the gastric mucosa [5]. In patients infected with *H. pylori*, high levels of gastrin and pepsinogen

and reduced levels of somatostatin have been measured [5]. Most patients with SU have impaired gastric bicarbonate secretion, which has also proven to be caused by *H. pylori* because its eradication reverses the defect [9]. Similarly, pepsin (a proteolytic enzyme) and HCl are both essential for food digestion but at the same time have the tendency to erode the cell linings of the digestive system if secreted in excess. Although, the stomach defends itself from these aggravating factors by creating a mucus coating and producing bicarbonates, *H. pylori* infection and NSAIDs can impair the protective functions and make the linings of the gastrointestinal tract susceptible to HCl and pepsin action and consequently results in the formation of ulceration [10].

Globally, SU is the most prevalent gastrointestinal disorder ever known, affecting more than 10% of people and accounting for an estimated 15,000 mortality yearly [11]. The annual incidence of SU perforation and hemorrhage were 3.8–14.0 and 19.4–57.0 per 100,000 persons, respectively and this is anticipated to further worsen if no practical and viable alternatives are sought [12]. Specifically in the United States, PU disease affects approximately 4.5 million people annually and about 10% of the US population has evidence of a SU at some point in their life time [5]. Initially, PU diseases were more prevalent in the male populations but the current statistics suggest quite similar figures in both males and females with the overall lifetime prevalence tending toward 8–11% and 11–14% in women and men, respectively [5]. The age trends for the occurrence of PU show appreciably declining rates in younger males, while it is increasing steadily in older women. The statistics for SU disease in other countries is variable and is hinged primarily on the major causes of the disease: *H. pylori* and NSAIDs [13].

Although, orthodox medicine has provided succor in the management of SU disease over the years, a significant percentage of the global population still use traditional systems of medicine to manage and treat SU due to better cultural acceptability, improved compatibility, affordability, and lesser side effects [14]. The present study was conducted to review medicinal plants considered as gastroprotective and healing agents on SUs with particular focus on selected antiulcerogenic botanicals and their implicated phytonutrients. To achieve this, information were retrieved from online databases (Google, Pubmed, MEDLINE, Science Direct, Scopus and SID) in form of published articles, books, conference proceedings and other high profile intellectual resources. The retrieved studies either showed effectiveness of these plants or indirectly their efficacy on the involved mechanisms in the treatment of SU.

2. The conventional therapy for stomach ulcer management

SU therapy has witnessed many strides over the last decades and a number of drugs are now available for its treatment. Since the occurrence of SU disease is attributable to either pepsin action and hyperacidity or inadequate mucosal resistance, hence, its effective management lies exclusively in stemming the aggressive factors or fortify the defensive mechanisms. A

corner stone in this approach was the advent of histamine H_2 -receptors and the respective antagonist [15], which signified a landmark achievement in the management of disorders characterized by gastric hypersecretion. The H_2 -antagonists were instantaneously identified as potent and safe agents which may replace the previously used drugs. For instance, the appearance of cimetidine (an H_2 -antagonist agent), led to the virtual disappearance of drugs of unknown mechanism like gefarnate, sulphoglycopeptide, amilopectine, zolimidine, xylamide, etc. Despite the inherent improvement witnessed with the use of the H_2 -antagonist over the previously used drugs, their untoward reactions have undermined their appreciable application. Generally, the antiulcer drugs may be classified according to the site and/or mechanism of action as: (a) antacids; (b) gastric muscle stimulants; (c) agents which protect the mucosa, increase mucosal resistance or coat ulcer craters; (d) antisecretory drugs, which may be anticholinergic agents; (e) corticohypothalamic drugs; and (f) proton pump inhibitors.

2.1. Antacids

It is noteworthy that either antacid mixtures or combinations of antacids with other compounds are more commonly used than single-entity antacids [15]. These kinds of formulation have been made to elicit a better neutralizing effect and to extenuate side effects associated with single constituent entities. The calcium and aluminium combinations reduce diarrhea while magnesium caters well for constipation. Similarly, a combination of slow- and fast-acting agents could increase the total buffering time. More improved benefits are claimed for mixtures having alginic acid (a foam-forming agent), which floats above the gastric juice with eventual aiding of contact between the antacid and the mucosa. In the event of gastroesophageal reflux, the alginic acid appears to prevent reflux by being the first to come in contact with the esophagus. By so doing, further erosion by the gastric acid is effectively prevented. Also, when simethicone (dimethylpolysiloxanes with characteristics antifoaming and water-repellant properties) is used, defoaming the gastric juice to reduce flatulence and gastroesophageal reflux is achieved. However, the resulting contributory adverse effects of the respective constituent in the antacid especially gastric irritation are a major challenge consistent with the use of antacids.

2.2. Gastric muscle stimulants

In individuals with stomach atony where there is consequential prolonged contact time between acid and mucosa due to the delay in gastric emptying, stimulants of gastric motility produce satisfactory outcomes. Domperidone and metoclopramide are the two most widely used compounds in this category. While both accelerates gastric emptying in experimental animals and humans, the metoclopramide still acts via other four important mechanisms that have made it more potent: (i) a cholinergic effect on muscarinic receptors; (ii) a direct effect on smooth muscle; (iii) an effect on specific centers regulating gastrointestinal motility; and (iv) a release of motilin which could be strongly responsible for the effects on the proximal bowel [15].

2.3. Mucosa protecting agents

Compounds like sucralfate, carbenoxolone and chelated bismuthate belong to this class of drugs. The sucralfate (a complex of sucrose and aluminium hydroxide) potentiates its action by inhibiting gastric hydrolysis with significant affinity for ulcerated mucosa. It forms a complex with susceptible proteins (fibrinogen, albumin etc) that adhere to the ulcerated area, thereby shielding against acid, pepsin action and penetration of bile acid. The merits of sucralfate over other antiulcer drugs relate to a lack of systemic effect due to the poor absorption of the compound [16]. The disadvantages include a certain delay in gastric emptying and constipation which affects almost 2% of the sufferers. Furthermore, compliance is another issue of concern as sucralfate must be taken four times daily (q.i.d.) and at least one hour before meals to be optimally effective. While chelated bismuthate and carbenoxolone works quite similar to sucralfate, appreciable incidence of aldosterone-like adverse effects (sodium and fluid retention, hypertension etc) have hampered their use.

2.4. Drugs with antisecretory effect

2.4.1. Anticholinergics

Despite that the drugs in this class have been used for many years, information on their ulcer healing capacity is still elusive. This may be attributable to the radiological method that was initially used but later found not to be very effective in wound healing assessment. Hyoscyamine, atropine, phentonium, propantheline and methantheline with characteristic ganglion-blocking and antimuscarinic effects are the classical examples of anticholinergics. Subsequent to the discovery and acceptance of the H₂-antagonists, the anticholinergics virtually disappeared due to inherent adverse effects of dry mouth, blurring of vision, delay in gastric emptying, tachycardia, possible constipation, and urinary retention.

2.4.2. H₂-antagonists

Undoubtedly, the H₂-antagonists are the most significant agents for the management and treatment of SU and for pathological conditions characterized by hyperacidity. Burimamide (the prototype of H₂-antagonists) was faced-out because of low oral activity and toxicological concerns. Similarly, metiamide (the 2nd agent in the series), was orally active but potentiated significant bone marrow toxicity in clinical trials. Cimetidine (still the most widely used antiulcer drug), is the 3rd drug in this class and well over 70 million sufferers have so far benefitted from its pharmacological efficacy till date. This was closely followed by ranitidine, which was the first agent with an alkyl furan ring, which substituted the imidazole ring of the preceding H₂-antagonists. More recent reports however suggest that optimum H₂-antagonism may also be achieved also with other different agents. For this class of drugs, not only could potency and efficacy be optimized, the pharmacokinetics could also be modified with the known duration of action of 4–6 h, exclusive to the 'short-acting' H₂-blockers [17], optimized to and beyond 24–48 h to have 'long-acting' H₂-blockers. Generally, over the last

three decades, the most frequently used H₂-antagonists in clinical practice are cimetidine, ranitidine and famotidine [2]. However, many adverse reactions such as effects on the endocrine, cardiovascular and central nervous systems (CNS) have been associated with these drugs.

2.5. Corticohypothalamic drugs

A role has been established for the CNS in the regulation of gastric secretion and in the pathogenesis of peptic ulcer, although clarification is required in many areas. It is not surprising that drugs which act specifically on the CNS may exert a beneficial effect on SU, which sometimes is significantly better than other drugs. Trimipramine is a tricyclic antidepressant which causes a slight decrease in gastric secretion that is apparently not connected with its anticholinergic action. Like other traditional tricyclic antidepressants, it may have some H₂-antagonistic effects and may also act on α -adrenoceptors enhancing catecholamine availability at central synapses [18], or may depress central vagal function. However the use of trimipramine in non-depressed ulcer patients is questionable.

2.6. Proton pump inhibitors (PPIs)

At present, PPIs are the most commonly prescribed class of antiulcer drugs. Their mode of action involves blockage of the site of gastric acid secretion in the parietal cell of the stomach [19]. However, because the parietal cells are constantly reproducing in millions, effective inhibition of gastric acid secretion is almost unachievable and this partly explains their relative safety compared to other groups of antiulcer drugs. In general, the incidence of short-term adverse effects subsequent to PPI usage is relatively low and this may be the reason for their being well tolerated. Their long-term use has not been frequently studied and the dearth of information in this regards has made it difficult to make definitive statements [20]. For all the PPIs (omeprazole, lansoprazole, dexlanprazole, esomeprazole, pantoprazole, rabeprazole and ilaprazole), the occurrence of adverse effects are similar, though they have been reported more frequently with omeprazole. This may be due to its longer availability. The common adverse effects with PPIs include headache, nausea, diarrhea, abdominal pain, fatigue and dizziness. Infrequent adverse effects may also include rash, itch, flatulence, constipation, anxiety, depression, myopathies and rhabdomyolysis [21].

3. Ulcer inducing agents/models

The pathological mechanisms of SU disease that compromises its functional capability and the structural integrity has been established to arise mainly through either production of too much acid and pepsin, or weakening of the gastric epithelia that consequently results in too little mucosal resistance [2]. **Table 1** shows some of the known ulcerogenic agents/models and the pathological mechanism involved in their ulcer pathogenesis. A good understanding of the pathogenic mechanism of action of these models is crucial to spotting and either managing or preventing ulceration and the associated disorders.

Agent/model	Underlying mechanism	Reference(s)
NSAIDs (aspirin, indomethacin and ibuprofen)	Gastric acid secretion and inhibition of prostaglandin synthetase activity	[22]
Water-immersion/cold-restraint stress	Release of histamine resulting in increased acid secretion, decreased mucus production and poor flow of gastric blood	[23, 24]
Ethanol	Solubilizes mucous membrane and renders it vulnerable to the proteolytic and hydrolytic actions of HCl and pepsin	[25]
Acetic acid	Induces round, deep ulcers in the stomach through over production of acid secretion	[26]
Histamine	Acid stimulating and vasodilating effect that results to increased vascular permeability of the gastric mucosa	[27]
Reserpine	Degranulation of gastric mast cells consequent to histamine liberation that is facilitated by cholinergic system	[28]
Serotonin	Causes vasoconstriction thereby reducing gastric mucosal blood flow resulting to acute mucosal injury	[29]
Pylorous-ligation	Accumulation of gastric acid that consequently produces ulcer subsequent to the breakdown of gastric mucosal barriers	[30]
Ischemia-reperfusion	Causes erosion of the gastric epithelia due to free radicals formation	[31]
Acetic acid- <i>H. pylori</i>	Increased acid secretion and decreased mucus production	[32]
Iron-ascorbic acid	Linked with lipid peroxidation mediated by oxygen radicals	[33]

Table 1. Commonly used experimental models for ulcer induction.

4. Some scientifically validated antiulcerogenic medicinal plants

Despite the rapidly changing concept of SU disease management from conventional vagotomy, H₂-receptor antagonists and antacids to proton pump inhibitors, gastrointestinal toxicity and other inherent adverse effects remain significant impediments to their application in clinical practice. Investigation on the phytotherapeutic applications of medicinal plants that are highly valued and widely used in the traditional systems of medicine have been and still providing efficient formulation for better management of SU [2, 22].

The under-listed medicinal plants have been pharmacological reported to possess antiulcer activity as previously compiled [34–39]. They are:

Acacia arabica (Family: Mimosaceae); *Abutilon indicum* L. (Family: Malvaceae); *Adansonia digitate*; *Aegle marmelos* (Family: Rutaceae); *Allium sativum* (Family: Liliaceae); *Allophylus serratus* Kurz (Family: Sapindaceae); *Aloe vera* (Family: Liliaceae); *Alstonia scholaris*; *Annona squamosa* (Family: Annonaceae); *Asparagus racemosus*; *Azadirachta indica* (Family: Meliaceae); *Bacopa monnieri*; *Benincasa hispida*; *Bauhinia purpurea* (Family: Leguminosae); *Bauhinia variegata* (Family: Caesalpiniaceae); *Berberis aristata*; *Beta vulgaris*; *Buchanania lanzan* Spreng. (Family:

Anacardiaceae); *Butea frondosa* Roxb. (Family: Fabaceae); *Boswellia serrata* (Family: Burseraceae); *Careya arborea* (Family: Myrtaceae); *Carica papaya* (Family: Caricaceae); *Capsicum annuum* L. (Family: Solanaceae); *Centella asiatica*; *Cissus quadrangularis* L. (Family: Vitaceae); *Curcuma longa* L. (Family: Zingiberaceae); *Desmostachya bipinnata* (L.) Stapf (Family: Gramineae); *Desmodium gangeticum*; *Emblica officinalis* (Family: Euphorbiaceae); *Excoecaria agallocha* (Family: Euphorbiaceae); *Garcinia cambogia*; *Glycyrrhiza glabra* (Family: Leguminosae); *Ficus arnottiana*; *Ficus religiosa* (Family: Urticaceae); *Hemidesmus indicus*; *Hibiscus rosa sinensis* (Family: Malvaceae); *Ipomoea batatas* L. (Family: Convolvulaceae); *Ixora pavetta* (Family: Rubiaceae); *Kielmeyera coriacea* Mart (Family: Guttiferae); *Lagenaria siceraria* (Family: Cucurbitaceae); *Leucas lavandulifolia* Sm. (Family: Labiatae); *Mangifera indica* (Family: Anacardiaceae); *Mimosa pudica* (Family: Fabaceae); *Mentha arvensis* L. (Family: Lamiaceae); *Momordica charantia* (Family: Cucurbitaceae); *Momordica cymbalaria* Hook. (Family: Cucurbitaceae); *Morinda citrifolia*; *Moringa oleifera* (Family: Moringaceae); *Musa sapientum*; *Myrtus communis* (Family: Myrtaceae); *Ocimum sanctum* (Family: Lamiaceae); *Oryza sativa* (Family: Gramineae); *Phyllanthus niruri* (Family: Euphorbiaceae); *Plectranthus amboinicus*; *Polyalthia longifolia* (Family: Annonaceae); *Psidium guajava* (Family: Myrtaceae); *Rhus coriaria* (Family: Anacardiaceae); *Rhizophora mangle* L. (Family: Rhizophoraceae); *Sapindus trifoliatus* L. (Family: Sapindaceae); *Sesbania grandiflora* (Fabaceae); *Shorea robusta* (Family: Dipterocarpaceae); *Solanum nigrum* (Family: Solanaceae); *Tamarindus indica* (Family: Caesalpiniaceae); *Tecomaria capensis* (Family: Bignoniaceae); *Terminalia chebula* (Family: Combretaceae); *Terminalia pallida*; *Utleria salicifolia* Bedd. Ex.Hook. F. (Family: Periplocaceae); *Vinca minor* L. (Family: Apocynaceae). A comprehensive list of some selected plants being embraced as antiulcerogenic agents is presented in **Table 2**.

Plant	Family	Plant used	Phytonutrients	Reference(s)
<i>Acacia arabica</i>	Mimosaceae	Gum, leaves	Arabic acid, malate, sugar, mineral elements, tannins	[40, 41]
<i>Achyranthus aspera</i>	Amaranthaceae	Root, seeds	Saponin, glycosides	[42]
<i>Adansonia digitata</i>	Malvaceae	Leaves	Mucilage, glucose, albuminoids, adansonin, tannin	[43]
<i>Aegle marmelos</i>	Rutaceae	Leaves	Flavonoid, tannins, saponin	[44]
<i>Aleo vera</i>	Liliaceae	Whole plant	Aloin, isobarbaloin, emodin, saponin	[45]
<i>Alhagi maurorum</i>	Fabaceae	Root	Terpenes, saponin, tannins	[46]
<i>Allium sativum</i>	Liliaceae	Bulb	Mucilage, starch, albumen, vitamins, sugar, allicin, alliin	[47]
<i>Annona squamosa</i>	Annonaceae	Leaves	Alkaloids, flavonoids, saponin, tannins	[48]
<i>Azadirachta indica</i>	Meliaceae	Leaves	Saponin, flavonoids, phenolics, tannin	[49]
<i>Bauhinia variegata</i>	Caesalpiniaceae	Stem-bark, root	Rutin, quercetin, apigenin, tannin	[50]
<i>Berberis aristata</i>	Berberidaceae	Root	Alkaloids	[51]
<i>Bata vulgaris</i>	Chenopodiaceae	Root	Betin	[52]
<i>Carica papaya</i>	Caricaceae	Fruit, seeds	Papain, pectin, carpaine, carposide	[53]

Plant	Family	Plant used	Phytonutrients	Reference(s)
<i>Centella asiatica</i>	Apiaceae	Whole plant	Flavonoids, narigin, alkaloids, saponin, asiatic acid	[54]
<i>Cordial myxa</i>		Fruit	Tannins, carbohydrate, saponin	[55]
<i>Ficus exasperata</i>	Moraceae	Leaves, stem-bark	Flavonoids, saponin, alkaloids, glycosides, tannins	[2]
<i>Ficus religiosa</i>	Urticaceae	Stem-bark	Tannins, wax, cochtone	[56]
<i>Gossypium barbadense</i>	Malvaceae	Leaves	Gossypol, saponin, steroids, cardiac glycosides	[22]
<i>Gossypium herbaceous</i>	Malvaceae	Flowers	Flavonoids, phenolics, saponin	[57]
<i>Hibiscus rosa sinensis</i>	Malvaceae	Root	Phenolics, cyanidin, hydrocitrates	[58]
<i>Langeneria breviflora</i>	Cucurbitaceae	Fruit, leaves	Saponin, phenolics, cucurbitacin	[59]
<i>Langeneria siceraria</i>	Cucurbitaceae	Fruit	Flavonoids, steroids, phenols, saponin	[42]
<i>Mangifera indica</i>	Anacardiaceae	Leaves	Alkaloids, sterols, saponin, tannins, flavonoids, mangiferin	[60]
<i>Momordica tuberosa</i>	Cucurbitaceae	Tubers	Alkaloids, tannins, saponin	[61]
<i>Moringa oleifera</i>	Moringaceae	Leaves	Quercetin, β -sitosterol, β -carotene, alkaloids, tannins, saponin	[62]
<i>Musa paradisiacal</i>	Musaceae	Root, leaves, trunk	Tannins, starch, vitamin C, albuminoids	[42]
<i>Myrtus communis</i>	Myrtaceae	Leaves	Resin, tannins, citrate, malate, sugar	[63]
<i>Nerium indicum</i>	Apocynaceae	Flowers	Alkaloids, glycosides, flavonoids, phenolics, tannins	[64]
<i>Ocimum sanctum</i>	Lamiaceae	Leaves	Alkaloids, tannins, saponin, flavonoids, sterols	[65]
<i>Oryza sativa</i>	Gramineae	Grain, bran	Starch, mineral matter, protein	[66]
<i>Phyllanthus niruri</i>	Euphorbiaceae	Whole plant	Alkaloids, tannins, flavonoids, carbohydrate, glycosides	[67]
<i>Prunus amygdalus</i>	Rosaceae	Seeds, fruit	Ursolic acid, quercetin, flavonoids	[42]
<i>Psidium guyava</i>	Myrtaceae	Leaves, stem-bark	Resin, tannins, cellulose, flavonoids, quercetin, quajaverin	[68]
<i>Rhus coriaria</i>	Anacardiaceae	Whole plant	Tannins, flavonoids	[69]
<i>Sesbania grandiflora</i>	Fabaceae	Leaves	Saponin, tannins, triterpenes	[70]
<i>Smilax china</i>	Smilacaceae	Root	Tannins, resin, saponin, flavonoids	[71]
<i>Solanum nigrum</i>	Solanaceae	Leaves	Flavonoids, saponin, alkaloids, phytosterols	[72]
<i>Spondias mombin</i>	Anacardiaceae	Leaves	Tannins, saponin, flavonoids, phenolics, glycosides	[73]
<i>Tamarindus indica</i>	Caesalpinaceae	Leaves, seeds	Albuminoids, fiber, pectin, tannins	[74]
<i>Terminalia chebula</i>	Combretaceae	Leaves	Gallic acid, sorbitol, tannins, mucilage	[75]

Table 2. Some selected medicinal plants with antiulcerogenic properties.

Following the experimental demonstration that many medicinal plants are endowed with good antiulcerogenic activity with relatively lesser adverse effect compared with the conventional drugs, further steps have been taken in presenting a good number of them for clinical trials. Despite this giant stride, not many of the medicinal plants have passed market entry stage. To the best of our knowledge, of the many presented for developmental evaluations in 2004, only *Azadirachta indica* (Family: Meliaceae) received remarkable attention at its advance stage of clinical trial. It exhibited significant therapeutic potency by reducing gastric hypersecretion, gastroesophageal and gastroduodenal ulcers [76].

5. Phytonutrients associated with antiulcerogenic activity

Several phytonutrients have proven health benefits and have been reported to elicit significant antiulcerogenic potential in both humans and experimental animal models [73]. While steroid glycosides, tannins, terpenoids and flavonoids have been shown to preserve gastric mucosal against oxidative insults of reactive metabolites and oxidative stress [22, 59], the tendency of phenolic compounds and alkaloids to regulate gastric acid secretion and protect the gastric mucosal epithelia against erosion and other aggressive factors in different ulcer models have been demonstrated [77]. While **Table 1** also presents some of these phytonutrients as being responsible for the elicited antiulcerogenic properties of the plants, several others have also been identified and isolated from diverse plants. Some of these include; saponins, phobaphenes, glucose, luvangetin, tartarate, potash, nimbodin, quercetin, apigenin, papain, chymopapain, pectin, carposide, carotenoids, antheraxanthin, carpaine, resin, euphorbon, caoutchouc, rutin, anthocyanins, cyanindin, kaempferol, sterols, mucilage, terpenoids, kaepferom, ash, starch, fats, proteins, glycosides, ellagic acid, beta sitosterol, gallic acid, limonene, pinene, albuminous matter, cellulose, chlorophyll, mineral salts, myricitin, triterpenes, and sorbitol [37]. These compounds have either been elucidated to decrease acid/pepsin secretion or confer cytoprotection via effective modulation on mucosal defensive factors.

6. Some selected medicinal plants with antiulcerogenic properties in Nigeria

In Nigeria, SU remains a significant public health challenge affecting people of all ages. While its management through orthodox medicine has recorded substantial successes over the years, a considerable proportion of the populace still rely exclusively on complementary and alternative medicine (CAM) in seeking aid to treat and manage SU. This may be due to the ease of accessibility, affordability and minimal side effect associated with the use of medicinal plants [22]. A compilation of selected antiulcerogenic medicinal plants in Nigeria revealed that the most widely used plants in the management of SU are *Occimum basillicum*, *M. paradisiaca*, *Aloe vera*, *Azadiracter indica*, *Brassica oleracae* and *Carica papaya* [78]. Others include but not limited to the following: *Diodia sarmentosa*, *Cassia nigricans*, *Ficus exasperate*, *Synclisia*

scabrida, *Artocarpus heterophyllus* Lam., *Blighia sapida* Konig., *Dialium guineense* Willd., *Embllica officinalis* Gaertn., *Gongronema latifolium*, *Ageratum conyzoides*, *Aloe vera*, *Artocarpus altilis*, *Aspilia africana*, *Bryophyllum pinnatum*, *Fluerya aestuans*, *Musa paradisiaca*, *Musa sapientum*, *Persea Americana*, *Talinum triangulare*, *Fluerya aetuans*, *Brassica oleraceae*, *Acacia nilotica* L., *Alchornea cordifolia* Schum & thonn, *Anacardium occidentale* L., *Balanites aegyptiaca* L., *Bridelia ferruginea* Benth, *Carica papaya* Linn, *Ficus thonningii* Blume, *Guiera senegalensis* J. F. Gmel, *Hibiscus sabbdariffa* L., *Mangifera indica* L., *Momordica charantia* L., *Ocimum gratissimum* L., *Piliostigma reticulatum* (DC) Hochst, *Pisidium guajava* L., *Scoparia dulcis*, *Vernonia kotschyana* Sch. Bip., *Zingiber officinale* Rosc [79].

7. The role of medicinal plants in oxidative gastropathy

Reactive oxygen species (ROS) are a by-product of normal metabolism and have roles in cell signaling and homeostasis [80]. Mechanisms exist that regulate cellular levels of ROS, as their reactive nature may otherwise cause damage to key cellular components including DNA, protein, and lipids [81]. A good number of NSAIDs have been implicated in cellular toxicity leading to oxidative gastropathy [82]. Despite the use of NSAIDs as antipyretic and anti-inflammation agents, and in the treatment of rheumatic, musculoskeletal, and cardiovascular diseases [83], gastrointestinal toxicity through ROS formation has limited their application [84, 85]. It has been proposed that NSAID-mediated gastrointestinal lesions involve the uncoupling of oxidative phosphorylation and inhibition of electron transport chain causing incomplete reduction of oxygen [82]. This they do by tenaciously binding to a site near complex I and ubiquinone, thus facilitating events leading to ROS generation [86, 87]. Subsequently, when the gastric antioxidant capacity is overwhelmed, the epithelia mitochondrial aconitase is inhibited, resulting in the release of iron that reacts with H_2O_2 , producing hydroxyl radical. These cascades of event amplify gastric oxidative stress whose consequential effect is manifested as gastropathy [88]. Oxidative stress-induced functional loss is well correlated with numerous disease states including cardiovascular, neurological, cancer, aging processes and gastropathy [83] and is also implicated in a variety of drug-induced toxicities such as SU [2]. Antioxidative and free radical scavenging mechanisms play an important role in the protection against ROS mediated toxicities [89]. Over the past decades, interests in medicinal plants, especially the antioxidative ones, have increased appreciably and they have been elucidated to significantly either protect against or ameliorate ROS-mediated oxidative gastropathy [22, 90]. Such annihilation of ROS in SU diseases have been achieved through induction of enzymic antioxidants (superoxide dismutase, catalase, glutathione reductase and peroxidase) and optimization of reduced glutathione (GSH) contents [2, 22, 59]. Harmonizing the foregoing, a probable mechanism of antioxidative and gastroprotective activities of medicinal plants may be idealized as illustrated in **Figure 1**. This ultimately involves induction and optimization of preventive (catalase, glutathione peroxidase) and chain-breaking (superoxide dismutase, glutathione reductase) antioxidants that subsequently improve gastric GSH level, annihilate liberated reactive metabolite and effectively scavenge ROS (O_2^- , OH^-) (**Figure 1**). This may also be opined to regulate mucosal fluidity and strengthens defensive mechanisms against oxidative gastric damage.




Figure 1. Probable mechanism of antioxidative and gastroprotective capabilities of medicinal plants. The dotted arrows represent sites of induction and optimization by the plants that consequently promote scavenging of O₂⁻ and OH⁻. This will normalize and increase gastric reduced glutathione (GSH) content and promotes its mobilization toward detoxification of the liberated reactive metabolites. NSAIDs, nonsteroidal anti-inflammatory drugs; MP, medicinal plants; GPx, glutathione peroxidase; GRx, glutathione reductase; GSSG, oxidized glutathione.

8. Conclusion

Globally, SU is a devastating disease posing serious threat to the quality of life of humans. It affects significant proportion of the populace in both developed and developing countries. Although, conventional drugs have been used to manage and treat SU sufferers, affordability and inherent side effects have limited their application. Consequently, alternatives are being sought in medicinal plants, which provide a potential source of antiulcerogenic drugs and are widely used in traditional systems of medicine. Several medicinal plants have been investigated for their proven health benefits in SU management with their phytonutrients playing significant roles. Of the phytonutrients, tannins seem to top the list and has suggested probable focus on their characterization for antiulcer therapy. In spite of the impressive experimental evaluation of medicinal plants for the treatment of SU, very few have reached clinical trials and not very many have been marketed. This indicates that the intended benefits of CAM research are not yet having far-reaching effect. Nevertheless, the continuous search for

antiulcerogenic agents of plant origin (available as gifts of nature) is imperative. This will ultimately and eventually present effective and globally competitive exciting opportunities for the development of new lead therapeutics for SU and other related disorders.

Author details

Sabiu Saheed^{1,2,3*}, Ajani Emmanuel Oladipo¹, Taofik Olatunde Sunmonu⁴, Fatai Oladunni Balogun² and Anofi Omotayo Tom Ashafa²

*Address all correspondence to: saeed.sabiu@gmail.com

1 Phytomedicine, Food Factors and Toxicology Research Laboratory, Biochemistry Department, School of Basic Medical Sciences, Kwara State University, Ilorin, Nigeria

2 Phytomedicine and Phytopharmacology Research Group, Plant Sciences Department, University of the Free State, Phuthaditjhaba, South Africa

3 Department of Microbial, Biochemical, and Food Biotechnology, University of the Free State, Bloemfontein, South Africa

4 Phytomedicine and Plant Biochemistry Research Laboratory, Biochemistry Unit, Department of Biological Sciences, Al-Hikmah University, Ilorin, Nigeria

References

- [1] Chan FKL, Graham DY. Review article: prevention of non-steroidal anti-inflammatory drug gastrointestinal complications—Review and recommendations based on risk assessment. *Alimentary Pharmacology and Therapeutics*. 2004;**19**(10):1051-1061
- [2] Sabiu S, Sulyman AO, Garuba T, Sunmonu TO, Abdulrahman AA. Combined administration of *Spondias mombin* and *ficus exasperata* leaf extracts stall indomethacin-mediated gastric mucosal onslaught in rats. *African Journal of Traditional Complementary Alternative Medicine*. 2015;**12**:45-51
- [3] Debjit B, Chiranjib C, Tripathi KK, Pankaj J, Sampath KP. Recent trends of treatment and medication peptic ulcerative disorder. *International Journal of PharmTech Research*. 2010;**2**(1):970-980
- [4] Vyawahare NS, Deshmukh VV, Godkari MR, Kagathara VG. Plants with anti-ulcer activity. *Pharmacognosy Review*. 2009;**3**:108-115
- [5] Anand BS. Peptic ulcer diseases. *Gastroenterology*. 2017;**1**:23-35
- [6] Hoogerwerf WA, Pasricha PJ. Agents Used for Control of Gastric Acidity and Treatment of Peptic Ulcers and Gastro Esophageal Reflux Disease, 10th ed. New York, NY, USA: McGraw-Hill; 2001. pp. 1005-1019

- [7] Marshall BJ, Warren JR. Unidentified curved bacilli in the stomach of patients with gastritis and peptic ulceration. *The Lancet*. 1984;1(8390):1311-1315
- [8] Malfertheiner P, Chan FK, McColl KE. Peptic ulcer disease. *The Lancet*. 2009;374(9699):1449-1461
- [9] Sung JJ, Tsoi KK, Ma YK, Yung MY, Lau JY, Chiu PW. Causes of mortality in patients with peptic ulcer diseases: A prospective cohort study of 10428 cases. *American Journal of Gastroenterology*. 2010;105(1):84-89
- [10] Bansal, et al. Herbal approach to peptic ulcer disease: Review. *Journal of Bioscience Technology*. 2009;1:52-58
- [11] Shristi B, Neha J, Indu BP, Rajesh G. A review on some Indian medicinal plants for anti-ulcer activity. *Journal of Science Research and Pharmacy*. 2012;1:6-9
- [12] Lau JY, Sung J, Hill C, Henderson C, Howden CW, Metz DC. Systematic review of the epidemiology of complicated peptic ulcer disease: Incidence, recurrence, risk factors and mortality. *Digestion*. 2011;84(2):102-113
- [13] Cai S, Garcia RLA, Masso-Gonzalez EL, Hernandez-Diaz S. Uncomplicated peptic ulcer disease in the United Kingdom: Trends from 1997 to 2005. *Alimentary Pharmacology Therapeutics*. 2009;30(10):1039-1048
- [14] Kumar R. A review on medicinal plants for peptic ulcer. *Der Pharmacia Lettre*. 2011;3(3):414-420
- [15] Bertaccini G, Coruzzi G. Pharmacology of the treatment of peptic ulcer disease. *Digestive Diseases and Sciences*. 1985;30(11):43s-51s
- [16] Martin F, Farley A, Gagnon M, Bensemana D. Comparison of healing capacities of sucralfate and cimetidine in short treatment of duodenal ulcer. Double blind randomized trial. *Gastroenterology*. 1982;82:401-405
- [17] Buyniski JP, Cavanagh RL, Pircio AW, Agieri AA, Crenshaw RR. In: Melchiorre C, Gianella M, editors. *Highlights in Receptor Chemistry*. Amsterdam: Elsevier; 1984. pp. 195-213
- [18] Pendleton RG, McCafferty JP, Roesler JM, Hieble JP. Mechanism for gastric antisecretory effects of desmethylinipramine in rats. *European Journal of Pharmacology*. 1981;75:171-178
- [19] Sachar H, Vaidya K, Laine L. Intermittent vs continuous proton pump inhibitor therapy for high-risk bleeding ulcers: a systematic review and meta-analysis. *JAMA Internal Medicine*. 2014;174(11):1755-1762
- [20] Corleto VD. Proton pump inhibitor therapy and potential long-term harm. *Current Opinion in Endocrinology, Diabetes and Obesity*. 2016;21(1):3-8
- [21] Rossi S, editor. *Australian Medicines Handbook 2006*. Adelaide: Australian Medicines Handbook 2006

- [22] Sabiu S, Ajani EO, Ajao AA, Sunmonu TO, Ibraheem AS, Ibrahim R, Mustapha H, Adekeye AO. Biomembrane stabilization and antiulcerogenic properties of aqueous leaf extract of *Gossypium barbadense* L. (Malvaceae). Beni-Suef University Journal of Basic and Applied Sciences. 2017. <http://dx.doi.org/10.1016/j.bjbas.2017.05.007>
- [23] Senay EC, Levine RJ. Synergism between cold and restraint for rapid production of stress ulcers in rats. Proceedings of the Society for Experimental Biology and Medicine. 1967;**124**(4):1221-1223
- [24] Kitagawa H, Fujiwara M, Osumi Y. Effects of water immersion stress on gastric secretion and mucosal blood flow in rats. Gastroenterology. 1979;**77**(2):298-302
- [25] Oates PJ, Hakkinen JP. Studies on the mechanism of ethanol-induced gastric damage in rats. Gastroenterology. 1988;**94**(1):10-21
- [26] Okabe S, Amagase K. An overview of acetic acid ulcer models: The history and state of the art of peptic ulcer research. Biological and Pharmaceutical Bulletin. 2005;**28**(8):1321-1341
- [27] Cho CH, Pfeiffer CJ. Gastrointestinal ulceration in the guinea pig in response to dimaprit, histamine, and H₁- and H₂-blocking agents. Digestive Diseases and Sciences. 1981;**26**(4):306-311
- [28] Singh S. Evaluation of gastric anti-ulcer activity of fixed oil of *Ocimum basilicum* Linn. and its possible mechanism of action. Indian Journal of Experimental Biology. 1999;**37**(3):253-257
- [29] LePard KJ, Stephens RL. Serotonin inhibits gastric acid secretion through a 5-hydroxytryptamine₁-like receptor in the rat. Journal of Pharmacology and Experimental Therapeutics. 1994;**270**(3):1139-1147
- [30] Shay H, Komarov SA, Fels SS, Meranze D, Gruenstein M, Sipler H. A simple method for the uniform production of gastric ulceration in the rat, Gastroenterology. 1945;**5**:43-61
- [31] Wada K, Kamisaki Y, Kitano M, Kishimoto Y, Nakamoto K, Itoh T. A new gastric ulcer model induced by ischemia reperfusion in the rat: Role of leukocytes on ulceration in rat stomach. Life Sciences. 1996;**59**(19):295-301
- [32] Konturek PC, Brzozowski T, Kania J. Pioglitazone, a specific ligand of peroxisome proliferator-activated receptor γ , accelerates gastric ulcer healing in rat. European Journal of Pharmacology. 2003;**472**(3):213-220
- [33] Naito Y, Yoshikawa T, Yoneta T. A new gastric ulcer model in rats produced by ferrous iron and ascorbic acid injection. Digestion. 1995;**56**(6):472-478
- [34] Kaur D, Rana AC, Sharma N, Kumar S. Herbal drugs with antiulcer activity. Journal of Applied Pharmaceutical Science. 2012;**2**(3):160-165
- [35] Srinivas TL, Lakshmi SM, Shama SN, Reddy GK, Prasanna KR. Medicinal plants as anti-ulcer agents. Journal of Pharmacognosy and Phytochemistry. 2013;**2**(4):91-97
- [36] Schmeda-Hirschmann, G, Yesilada E. Traditional medicine and gastroprotective crude drugs. Journal of Ethnopharmacology. 2005;**100**(1-2):61-66

- [37] Vimala G, Shoba GF. A Review on antiulcer activity of few Indian medicinal plants. *International Journal of Microbiology*. 2014;Article ID 519590, 14 pages
- [38] Priyanka A. Some of the medicinal plants with anti-ulcer activity: A review. *Journal of Pharmaceutical Science and Research*. 2015;7(9):772-775
- [39] Mehta D. Ulcer-Review on types, anti-ulcer drugs, anti-ulcer medicinal plants, anti-ulcer drug market, diagnostics and current global clinical trials status. *Invention and Rapid Pharmacy Practice*. 2016;2:1-8
- [40] Nadkarni's KM. *Indian Materia Medica*. Popular Prakashan, Mumbai, India. 1976;1:9-10
- [41] Abd El-Mawla AMA, Osman HEH. Effects of *Gum acacia* aqueous extract on the histology of the intestine and enzymes of both the intestine and the pancreas of albino rats treated with meloxicam. *Pharmacognosy Research*. 2011;3(2):114-121
- [42] Maury PK, Jain SK, Lal N, Alok S. A review on antiulcer activity. *International Journal of Pharmaceutical Science Research*. 2012;3(8):2487-2493
- [43] Nadkarni's KM. *Indian Materia Medica*. Popular Prakashan, Mumbai, India. 1976;1:38-39
- [44] Ilavarasan JR, Monideen S, Vijayalakshmi M. Antiulcer activity of *Aegle marmelos*. *Ancient Science of Life*. 2002;21(4):23-26
- [45] Borra SK, Lagisetty RK, Mallela GR. Anti-ulcer effect of *Aloe vera* in non-steroidal anti-inflammatory drug induced peptic ulcers in rats. *African Journal of Pharmacy and Pharmacology*. 2011;5(16):1867-1871
- [46] Shaker E, Mahmoud H, Mnaa S. Anti-inflammatory and anti-ulcer activity of the extract from *Alhagi maurorum* (camelthorn). *Food and Chemical Toxicology*. 2010;48:2785-2790
- [47] Azamthulla M, Asad M, Prasad VS. Antiulcer activity of *Allium sativum* bulb juice in rats. *Saudi Pharmaceutical Journal*. 2009;17(1):70-77
- [48] Mohamed TSS, Pradeep RK, Priyanka N, Madhuri M, Sravanti V, Sarala K. Anti-ulcerogenic effect of aqueous extract of *Annona squamosa* (Linn). *International Journal of Research in Phytochemistry and Pharmacology*. 2012;2(3):157-159
- [49] Divakar MC, Rao SB, Nair GRN, Hisham A. The role of fatty acids on the ulcer healing property of the nimbidin fraction of the neem oil. *Journal of Medicinal and Aromatic Plants Science*. 2001;23(3):404-408
- [50] Kumar YR, Rajani GP. Analgesic and anti-ulcer activities of ethanol and aqueous extracts of root of *Bauhinia variegata* Linn. *International Journal of Pharmacology*. 2011;7(5):616-622
- [51] Nadkarni's KM. *Indian Materia Medica*. Popular Prakashan, Mumbai, India. 1976;1:187-188
- [52] Nadkarni's KM. *Indian Materia Medica*. Popular Prakashan, Mumbai, India. 1976;1:197

- [53] Indran M, Mahmood AA, Kuppusamy VR. Protective effect of *Carica papaya* L leaf extract against alcohol induced acute gastric damage and blood oxidative stress in rats. *West Indian Medical Journal*. 2008;**57**(4):323-326
- [54] Sairam K, Rao CHV, Goel RK. Effect of *Centella asiatica* Linn on physical and chemical factors induced gastric ulceration and secretion. *Indian Journal of Experimental Biology*. 2001;**39**:137-142
- [55] Inas ZA, Abdallah-Hala AH, Khattab H, Gehan HH. Gastroprotective effect of *Cordia Myxa* L. fruit extract against indomethacin-induced gastric ulceration in rats. *Life Sciences Journal*. 2011;**8**:433-445
- [56] Gregory M, Divya B, Mary RA, Hipolith MMV, Kalaichelvan VK, Palanivel V. Anti-ulcer activity of *Ficus religiosa* leaf ethanolic extract. *Asian Pacific Journal of Tropical Biomedicine*. 2013;**3**(7):554-556
- [57] Khalid MS, Hassan SK, Suresh DK, Hasan R, Saleem MA, Farooqui Z. Antiulcer activity of ethanolic extract of *Gossypium herbaceus* flowers. *RGUH. Journal of Pharmaceutical Sciences*. 2011;**1**:79S-84S
- [58] Srivastava JJ, Gautam H, Sharma S, Rao CV. Anti-ulcer activity of methanol extract of *Hibiscus rosa sinensis* leaves. *International Journal of Pharmacy and Pharmaceutical Sciences*. 2013;**5**(3):829-830
- [59] Ajani EO, Sabiu S, Bamisaye FA, Salami MR, Nurain IO. Ethanolic leaf extract of *Lagenaria brevifolia* (Bitter gourd) inhibits gastric onslaught in indomethacin ulcerated rats. *Journal of Applied Biosciences*. 2015;**85**:7871-7880
- [60] Neelima MS, Patil MB, Lakshmi BVS. Anti-ulcer activity and HPTLC analysis of *Mangifera indica* leaves. *International Journal of Pharmaceutical and Phytopharmacological Research*. 2012;**1**(4):146-155
- [61] Pramod K, Devala RG, Lakshmayya A, Ramachandra SS. GC-MS analysis and antiulcer activity of ethanol extract of tubers of *Momordica tuberosa* Cogn. (cucurbitaceae) in rats. *Journal of Applied Pharmacy*. 2011;**4**:359-369
- [62] Verma VK, Singh N, Saxena P, Singh R. Anti-ulcer and anti-oxidant activity of *Moringa oleifera* (Lam) leaves against aspirin and ethanol induced gastric ulcer in rats. *International Research Journal of Pharmacy*. 2002;**2**:46-57
- [63] Sumbul S, Ahmad MA, Asif M, Saud I, Akhtar M. Evaluation of *Myrtus communis* Linn. berries in experimental ulcer models in rats. *Human and Experimental Toxicology*. 2010;**29**(11):935-944
- [64] Kamble RD, Maske KS, Shaikh HM, Balghare KC, Shikare ON. A review of antiulcer medicinal plants. *International Journal of Pharmacy*. 2013;**4**(1):78-81
- [65] Singh S, Majumdar DK. Evaluation of the gastric antiulcer activity of fixed oil of *Ocimum sanctum* (Holy Basil). *Journal of Ethnopharmacology*. 1999;**65**(1):13-19

- [66] Panchal S, Goswami SS, Shan MB, Santani DD. Effect of *Oryza sativa* bran oil in various experimental ulcer models. *Indian Journal of Pharmacology*. 2008;**40**(2):146-148
- [67] Okoli CO, Ezike AC, Akah PA. Studies on wound healing and antiulcer activities of extract of aerial parts of *Phyllanthus niruri* L. (Euphorbiaceae). *American Journal of Pharmacology and Toxicology*. 2009;**4**(4):118-126
- [68] Umana EU, Timbuak JA, Musa SA, Ikyembe DT, Abdurrashid S, Hamman WO. Ulceroprotective effect of methanol extract of *Psidium guajava* leaves on ethanol induced gastric ulcer in adult wistar rats. *Asian Journal of Medical Sciences*. 2012;**4**(2):75-78
- [69] Ahmad H, Wadud A, Jahan N, Khazir M, Sofi G. Evaluation of anti-ulcer activity of hydro alcoholic extract of Post Sumaq (*Rhus coriaria* Linn.) in ethanol induced gastric ulcer in experimental rats. *International Research Journal of Medical Sciences*. 2013;**1**(10):7-12
- [70] Bhalke RD, Giri MA, Anarthe SJ, Pal SC. Antiulcer activity of the ethanol extract of leaves of *Sesbania grandiflora* (Linn.). *International Journal of Pharmacy and Pharmaceutical Sciences*. 2010;**2**(4):206-208
- [71] Khan I, Nisar M, Ebad F, Nadeem S, Saeed M, Khan H. Anti-inflammatory activities of sieboldogenin from *Smilax china* Linn.: Experimental and computational studies. *Journal of Ethnopharmacology*. 2009;**121**:175-177
- [72] Kavitha GGS, Parvathi S, Ramkumar PSS, Shanmuga SP. Pharmacological and phytochemical evaluation of anti-ulcerogenic potential of *Solanum nigrum*. *Indian Journal of Pharmaceutical Science and Research*. 2012;**3**(8):2837-2840
- [73] Saheed S, Taofeeq G, Taofik OS, Emmanuel OA, Abdulhakeem S, Ismaila NI, Abdulazeez B. Indomethacin-induced gastric ulceration in rats: Protective roles of *Spondias mombin* and *Ficus exasperate*. *Toxicology Report*. 2015;**2**:261-267
- [74] Kumar S, Suman S, Sharma S, Kalra P. Antiulcer effect of the methanolic extract of *Tamarindus indica* seeds in different experimental models. *Journal of Pharmacy and Bioallied Sciences*. 2011;**2**(3):236-241
- [75] Raju D, Ilango K, Chitra V, Ashish K. Evaluation of antiulcer activity of methanolic extract of *Terminalia chebula* fruits in experimental rats. *Journal of Pharmaceutical Sciences and Research*. 2009;**1**(3):101-107
- [76] Bandyopadhyay U, Biswas K, Sengupta A, Moitra P, Dutta P, Sarkar D, Debnath P, Ganguly CK, Banerjee RK. Clinical studies on the effect of Neem (*Azadirachta indica*) bark extract on gastric secretion and gastroduodenal ulcer. *Life Sciences*. 2004;**75**(24):2867-2878
- [77] Michael BA, Charles A, Isaac G, Alexander N. *In vivo* models used for evaluation of potential antigastroduodenal ulcer agents. *Ulcers*, 2013. Article ID 796405, 12 pages
- [78] Kadiri M, Ojewumi AW, Kokumo YO. Ethnobotanical study of plants used in managing ulcer in Abeokuta metropolis, Ogun State, Nigeria. *Journal of Natural Science, Engineering and Technology*. 2013;**12**:76-88

- [79] Alebiosu CO, Ugwah OM, Ugwah-Oguejiofor CJ, Njan AA. Ethno botanical studies of medicinal plants used in the management of Peptic ulcer disease in Sokoto State, North Western Nigeria. *International Research Journal of Pharmacy and Pharmacology*. 2012;**2**(9):225-230
- [80] Sabiu S, O'Neill FH, Ashafa AOT. *Zea mays*, *Stigma maydis* prevents and extenuates acetaminophen-perturbed oxidative onslaughts in rat hepatocytes. *Pharmaceutical Biology*, 2016;**54**(11): 2664-2673. DOI: <http://dx.doi.org/10.1080/13880209.20161178307>
- [81] Ajani EO, Sabiu S, Bamisaye FA, Adenigba BV, Awomoyi DD, Adeyanju MM. Hepatoprotective and antioxidative effect of ethanolic leaf extract of *Langenaria breviflora* (bitter gourd) on indomethacin-ulcerated rats. *IOSR Journal of Pharmacy and Biological Science*. 2014;**9**(5 V I):61-68
- [82] Hong Y, Xingchang P, Zhixiu S, Shaokang W, Ligang Y, Guiju S. Protective effect of wheat peptides against indomethacin-induced oxidative stress in IEC-6 cells. *Nutrients*. 2014;**6**:564-574
- [83] Dae YC, Jin IK, Soo-Heon P, Jae KK. Proanthocyanidin from grape seed extracts protects indomethacin-induced small intestinal mucosal injury. *Gastroenterology Research Practice*. 2014;**14**:618-626
- [84] Bjornsson E. Review article: Drug-induced liver injury in clinical practice. *Alimentary Pharmacology Therapeutics*. 2013;**32**:3-13
- [85] Jarupongprapa S, Ussavasodhi P, Katchamart W. Comparison of gastrointestinal adverse effects between cyclooxygenase-2 inhibitors and non-selective, non-steroidal anti-inflammatory drugs plus proton pump inhibitors: A systematic review and meta-analysis. *Journal of Gastroenterology*. 2013;**48**:830-838
- [86] Kang HS, Ock J, Lee HJ, Lee YJ, Kwon BM, Hong SH. Early growth response protein-1-upregulation and nuclear translocation by 2'-benzoyloxycinnamaldehyde induces prostate cancer cell death. *Cancer Letter*. 2013;**329**:217-227
- [87] Vaish V, Piplani H, Rana C, Vaiphei K, Sanyal SN. NSAIDs may regulate EGR-1-mediated induction of reactive oxygen species and non-steroidal anti-inflammatory drug-induced gene (NAG)-1 to initiate intrinsic pathway of apoptosis for the chemoprevention of colorectal cancer. *Molecular Cell Biochemistry*. 2013;**378**:47-64
- [88] Valko M, Leibfritz D, Moncol J, Cronin MTD, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *International Journal of Biochemistry and Cell Biology*. 2007;**39**:44-84
- [89] Sabiu S, Wudil AM. Comparative effects of *Telfaira occidentalis* and *Vernonia amygdalina* on garlic-induced hepatotoxicity in rats. *Biological and Environmental Sciences Journal for the Tropics*. 2011;**8**(4):193-197
- [90] Shu YZ. Recent natural products based drug development: A pharmaceutical industry perspective. *Journal of Natural Product*. 1998;**61**:1053-1107

