

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Current Therapeutic Options for HCV-HIV Coinfection

Ljiljana Perić and Dario Sabadi

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.70730>

Abstract

Due to shared risk factors for transmission, coinfection with human immunodeficiency virus (HIV) and hepatitis C virus (HCV) is a very common event. The prevalence of HCV infection among HIV-positive patients averages about 35%. In HIV/HCV co-infected patients, liver-related morbidity and mortality is a prominent non-AIDS-defining complication: up to 90% of liver-related deaths in HIV-infected patients are attributable to HCV. The progression of liver fibrosis is accelerated in HIV/HCV-coinfected patients, particularly in individuals with low CD4 counts (≤ 350 cells/mm³). Antiretroviral therapy may slow liver disease progression in HIV/HCV-coinfected patients and should, therefore, be considered for all coinfecting patients regardless of CD4 cell count. Most patients with HIV/HCV coinfection are taking multi-drug antiretroviral therapy, which may pose a problem with drug–drug interactions when initiating therapy with HCV medications. Rapid advances in HCV drug development led to the discovery of new classes of direct-acting antiviral (DAA) agents that target the HCV replication cycle. Several studies demonstrated comparable rates of sustained virological response (SVR) in coinfecting and mono-infected patients with new DAA-based therapy.

Keywords: HCV/HIV-coinfection, liver cirrhosis, CD4 T lymphocytes, antiretroviral therapy (ART), direct-acting antiviral (DAA) agents, drug–drug interaction

1. Introduction

By the Global AIDS Update: 2016, around 36.7 million people are living with human immunodeficiency virus (HIV) in the world today [1]. Five million of them are also infected with hepatitis C virus (HCV) [1]. HIV accelerates the progression of hepatitis C, inducing increased morbidity and mortality [2]. HIV-infected people are on average six times more likely than HIV-uninfected people to have HCV infection [3].

HIV and HCV share modes of transmission: often occurring by exposure to blood, sexual intercourse or by mother-to-child transmission.

2. Epidemiology

The prevalence of HCV antibodies varies widely among HIV transmission groups, ranging from 7–8% in men who have sex with men to 60–70% in hemophiliacs and 80–90% in intravenous drug users (IDUs), the most important group (**Figure 1**) [4].




Figure 1. Prevalence of HCV antibodies in different transmission groups. IDU, intravenous drug users; HCV, hepatitis C virus. Inspired by Management of Hepatitis C and HIV coinfection, Clinical Protocol for the WHO European Region. Available at: http://www.euro.who.int/data/assets/pdf_file/0008/78146/E90840Chapter6.pdf, Version September 1th, 2015.

For HIV-infected patients with HCV coinfection, liver-related morbidity and mortality is a prominent non-AIDS-defining complication [5]. Up to 90% of liver-related deaths in HIV-infected patients are attributable to HCV [5].

Among patients with chronic HCV infection, approximately one-third progress to cirrhosis, at a median time of 20 years [6–8]. The risk of progression is even greater in HCV/HIV-coinfected patients with low CD4 T lymphocyte (CD4) cell counts (≤ 350 cells/mm³) [6, 9, 10]. Cirrhosis has been observed to occur 12–16 years earlier in HIV/HCV-coinfected patients compared with those who have HCV mono-infection [11].

3. Antiretroviral therapy (ART) in HIV/HCV-coinfected patients

Antiretroviral therapy (ART) may slow liver disease progression in HIV/HCV-coinfected patients and should, therefore, be considered for all coinfecting patients regardless of CD4

NRTIs	NNRTIs	Protease inhibitors	Entry inhibitors	Integrase inhibitors
Abacavir	Efavirenz	Atazanavir, atazanavir/ritonavir	Enfuvirtide	Dolutegravir
Didanosine	Etravirine	Darunavir/ritonavir Darunavir/cobicistat	Maraviroc	Raltegravir
Emtricitabine	Nevirapine	Fosamprenavir		
Lamivudine	Rilpivirine	Lopinavir		
Stavudine		Saquinavir		
Tenofovir				
Zidovudine				

Table 1. Standard recommended treatments for naive patients with HIV-1 infection.

cell count [12]. This recommendation is supported by observational studies that suggest that antiretroviral therapy may reduce the risk of liver-related morbidity. The key issues in the clinical management of HIV/HCV-coinfected patients are which treatment for each condition and when to initiate it [12].

Classes of antiretroviral agents are nucleoside reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), protease inhibitors (PIs), integrase inhibitors (INSTIs), entry/fusion inhibitors (FIs) and chemokine receptor antagonists (CCR5 antagonists).

Standard recommended treatments for naive patients with HIV-1 infection (**Table 1**) generally consist of two nucleoside reverse transcriptase inhibitors (NRTIs) in combination with a third active antiretroviral drug from one of three drug classes: an integrase strand transfer inhibitor (INSTI), a non-nucleoside reverse transcriptase inhibitor (NNRTI) or a protease inhibitor (PI) with a pharmacokinetic (PK) enhancer (booster) (cobicistat or ritonavir) [12].

Antiretroviral therapy (ART) associated with liver injury is more common in HIV/HCV-coinfected patients than in those with HIV mono-infection [6, 13]. Some older ART have been associated with higher rates of liver injury in patients with chronic HCV infection, but newer ART drugs currently in use appear to be less hepatotoxic [6, 13]. Patients with significant alanine aminotransferase (ALT) and/or aspartate aminotransferase (AST) elevation should be carefully evaluated for signs and symptoms of liver insufficiency and for alternative causes of liver injury (e.g., acute HAV or HBV infection, hepatobiliary disease, or alcoholic hepatitis) [6, 14]. Short-term interruption of the ART regimen or of the specific drug suspected of causing the liver injury may be required [6, 14].

4. Concurrent treatment of HIV and HCV by the Office of AIDS Research advisory council (OARAC 2016)

If the decision is made to treat HCV, the antiretroviral regimen may need to be modified before HCV treatment is initiated to reduce the potential drug–drug interactions and/or toxicities that may develop during the period of concurrent HIV and HCV treatment [6].

In patients with suppressed plasma HIV RNA and modified antiretroviral therapy, HIV RNA should be measured within 4–8 weeks after changing antiretroviral therapy to confirm the effectiveness of the new regimen [6]. After completion of HCV treatment, the modified ART regimen should be continued for at least 2 weeks before reinitiating the original regimen [6]. This is necessary because of the prolonged half-life of some HCV drugs and the potential risk of drug–drug interactions if a prior HIV regimen is resumed soon after HCV treatment is completed [6].

5. HCV therapy in HIV/HCV-coinfected patients by EASL recommendations on treatment of hepatitis C, 2016

With direct-acting antivirals (DAAs), HCV cure rates of both HCV mono and HIV/HCV-coinfected persons are greater than 95% [15]. Current treatment guidelines no longer separate these two groups. Indications for HCV treatment and choice of direct-acting antiviral (DAA) agents combination are now the same for all HCV patients. In HIV/HCV co-infection, drug interactions between HIV and HCV agents need be checked prior to starting HCV therapy [15]. The higher risk of hepatic decompensation in HIV/HCV-coinfected patients, including those receiving successful antiretroviral therapy, continues to make these patients a high priority group for receiving access to direct-acting antiviral (DAA) agents as combination therapy [15].

6. Key studies for treatment of HCV with HIV coinfection

Using DAA therapy, several studies demonstrated comparable rates of sustained virological response (SVR) in coinfecting and mono-infected patients.

These trials, however, have primarily included individuals with CD4 counts >200 cells/mm³, and most patients in these trials did not have cirrhosis.

6.1. Sofosbuvir for genotype 1–4 in HIV coinfection by Rodriguez-Torres et al.

In an open-label trial, 23 HCV/HIV-coinfected treatment-naive patients with genotype 1–4 received the 12-week triple therapy of peginterferon alfa-2a, ribavirin (weight-based) and sofosbuvir [16]. Mean CD4 count was 562 cells/mm³, and all were on antiretroviral therapy (tenofovir-emtricitabine plus one of the following: efavirenz, atazanavir plus ritonavir, darunavir plus ritonavir, rilpivirine or raltegravir) [16]. The overall SVR12 rate was 91%; of the 19 patients with genotype 1, 89% achieved an SVR12 (**Figure 2**) [16].

6.2. TURQUOISE-I by Wyles et al.

This open-label study randomized treatment-naive and experienced patients with chronic HCV genotype 1 and HIV coinfection to receive a 12- or 24-week course of ombitasvir-paritaprevir-ritonavir and dasabuvir plus ribavirin [17]. Patients were required to have




Figure 2. Sofosbuvir for genotype 1-4 in HIV coinfection. PEG, peginterferon alfa-2a; RBV, ribavirin; SVR12, sustained viral response 12 weeks after the end of treatment; GT, genotype. Inspired by http://slides.hcvonline.org/uploads/151/sofosbuvir_for_genotype_14_in_hiv_coinfection.pdf

a CD4 > 200 cells/mm³ and an HIV RNA level < 40 copies while receiving an atazanavir- or raltegravir-based regimen [17]. The sustained virological response (SVR) 12 rates were 93.5% (29 of 31) in the 12-week group and 90.6% (29 of 32) in the 24-week group (**Figure 3**) [17].




Figure 3. TURQUOISE-I. 3 D, Ombitasvir-Paritaprevir-Ritonavir and Dasabuvir; RBV, ribavirin; GT, genotype; SVR, sustained viral response. Inspired by <https://depts.washington.edu/hepstudy/presentations/uploads/137/turquoise13d.pdf>

6.3. ALLY-2 study (daclatasvir + sofosbuvir in HCV GT 1–4 and HIV coinfection) by Wyles et al.

Among HIV/HCV-coinfected patients who received 12 weeks of daclatasvir plus sofosbuvir, sustained virologic response across all genotypes was 97.0% (including black patients and those with cirrhosis) and 76.0% after 8 weeks [18].

6.4. ION-4 study (ledipasvir and sofosbuvir for HCV genotype 1 or 4 in patients coinfecting with HIV-1) by Naggie et al.

In this multicenter, open-label, single-group study, 12 weeks of treatment with the once-daily, single-tablet regimen of ledipasvir-sofosbuvir resulted in a sustained virologic response in 96% of patients [19]. In exploratory subgroup analyses, rates of sustained virologic response 12 weeks after the end of therapy (the primary efficacy end point) were similar across all subgroups except that black patients, who made up 34% of the study population, had lower rates of sustained virologic response (**Figure 4**) [19].




Figure 4. Ledipasvir and sofosbuvir for HCV genotype 1 or 4 in patients coinfecting with HIV-1. GT 1 or 4, genotype 1 or 4; SVR 12, sustained viral response 12 weeks after the end of treatment. Inspired by http://slides.hcvonline.org/uploads/149/ion4_ls.pdf

7. Conclusions

Due to shared risk factors for transmission, HIV/HCV coinfection is a very common event, the prevalence averages about 35% in the United States and Europe [20, 21].

The progression of liver fibrosis is accelerated in HIV/HCV-coinfected patients. HCV guidance recommends using the same HCV treatment approach for patients coinfecting with HIV as those with HCV mono-infection.

DAA and interferon-free combination therapy has changed the landscape of therapy for HIV/HCV-coinfected patients.

Multiple studies demonstrating comparable rates of SVR in coinfecting and mono-infected patients.

Author details

Ljiljana Perić^{1,2*} and Dario Sabadi^{1,2}

*Address all correspondence to: ljiljana.peric@mefos.hr

1 Department of Infectious Diseases and Dermatology, Faculty of Medicine Osijek, University of Osijek, Osijek, Croatia

2 Clinic for Infectious Diseases, University Hospital Center Osijek, Osijek, Croatia

References

- [1] Global AIDS Response Progress Reporting (GARPR) 2016. Geneva, Switzerland: UNAIDS 2016 Estimates; 2016
- [2] Operskalski EA, Kovacs A. HIV/HCV co-infection: Pathogenesis, clinical complications, treatment, and new therapeutic technologies. *Current HIV/AIDS Reports*. 2011;8(1):12-22
- [3] Platt L, Easterbrook P, Gower E, McDonald B, Sabin K, McGowan C, Yanny I, Razavi H, Vickerman P. Prevalence and burden of HCV co-infection in people living with HIV: A global systematic review and meta-analysis. 2016.
- [4] Management of hepatitis C and HIV coinfection, Clinical Protocol for the WHO European Region, Chapter6.pdf, Version 1 September 2015. Available from: http://www.euro.who.int/data/assets/pdf_file/0008/78146/E90840
- [5] Spach DH, Nina Kim H. Treatment of Chronic Hepatitis C Infection Overview. Available from: <http://www.hepatitisc.uw.edu/go/treatment-infection>
- [6] Guidelines for the Use of Antiretroviral Agents in HIV-1-Infected Adults and Adolescents, Developed by the DHHS Panel on Antiretroviral Guidelines for Adults and Adolescents – A Working Group of the Office of AIDS Research Advisory Council (OARAC), 8/25/2017. Available from: <https://aidsinfo.nih.gov/contentfiles/lvguidelines/AdultandAdolescentGL.pdf>
- [7] Alter MJ, Margolis HS, Krawczynski K, et al. The natural history of community-acquired hepatitis C in the United States. The Sentinel Counties Chronic non-A, non-B Hepatitis

- Study Team. The New England Journal of Medicine 1992;**327** (27):1899-1905. Available at: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=1280771
- [8] Thomas DL, Astemborski J, Rai RM, et al. The natural history of hepatitis C virus infection: Host, viral, and environmental factors. *Journal of the American Medical Association*. 2000;**284**(4):450-456 Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=10904508
- [9] Considerations for Antiretroviral Use in Patients with Coinfections, Hepatitis C (HCV)/ HIV Coinfection, Last Updated: July 14, 2016; Last Reviewed: July 14, 2016. Developed by the DHHS Panel on Antiretroviral Guidelines for Adults and Adolescents – A Working Group of the Office of AIDS Research Advisory Council (OARAC) , Available at: <https://aidsinfo.nih.gov/guidelines/html/1/adult-and-adolescent-arv-guidelines/26/hcv-hiv>
- [10] Graham CS, Baden LR, Yu E, et al. Influence of human immunodeficiency virus infection on the course of hepatitis C virus infection: A meta-analysis. *Clinical Infectious Diseases*. 2001;**33**(4):562-569 Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=11462196
- [11] Benhamou Y, Bochet M, Di Martino V, Charlotte F, Azria F, Coutellier A, Vidaud M, Bricaire F, Opolon P, Katlama C, Poynard T. Liver fibrosis progression in human immunodeficiency virus and hepatitis C virus coinfecting patients. *Hepatology*. 1999;**30**:1054-1058
- [12] Considerations for Antiretroviral Use in Patients with Coinfections, Hepatitis C (HCV)/ HIV Coinfection, Last Updated: 14 July 2016; Last Reviewed: 14 July 2016. Available from: <https://aidsinfo.nih.gov/guidelines/html/1/adult-and-adolescent-arv-guidelines/26/hcv-hiv>
- [13] Aranzabal L, Casado JL, Moya J, et al. Influence of liver fibrosis on highly active antiretroviral therapy-associated hepatotoxicity in patients with HIV and hepatitis C virus coinfection. *Clinical Infectious Diseases*. 2005;**40**(4):588-593 Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=15712082
- [14] Sulkowski MS, Thomas DL. Hepatitis C in the HIV-infected patient. *Clinics in Liver Disease*. 2003;**7**(1):179-194 Available from: http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citation&list_uids=12691466
- [15] Rockstroh JK, Hardy WD. Current treatment options for hepatitis C patients co-infected with HIV, *Expert Rev Gastroenterol Hepatol*. 2016;**10**:689-695. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26799571>
- [16] Rodriguez-Torres M et al. Sofosbuvir for chronic hepatitis C virus infection genotype 1-4 in patients coinfecting with HIV. *Journal of Acquired Immune Deficiency Syndrome*. 2015;**68**:543-549 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/25622055>, http://slides.hcvonline.org/uploads/151/sofosbuvir_for_genotype_14_in_hiv_coinfection.pdf

- [17] Wyles D et al. TURQUOISE-I Part 1b: Ombitasvir/Paritaprevir/Ritonavir and Dasabuvir with Ribavirin for Hepatitis C Virus Infection in HIV-1 Coinfected Patients on Darunavir. *The Journal of Infectious Disease*. 2017;**215**(4):599-605. doi: 10.1093/infdis/jiw597. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/28329334> and <https://depts.washington.edu/hepstudy/presentations/uploads/137/turquoise13d.pdf>
- [18] Wyles DL et al. Daclatasvir in combination with sofosbuvir for HIV/HCV coinfection: ALLY-2 Study. *The New England Journal of Medicine*. 2015;**373**:714-725 Available from: http://slides.hcvonline.org/uploads/163/ally2_daclatasvir.pdf and <http://www.nejm.org/doi/full/10.1056/NEJMoa1503153#t=article>
- [19] Naggie S et al. Ledipasvir and sofosbuvir for HCV in patients coinfecting with HIV-1-ION-4 Study. *The New England Journal of Medicine*. 2015;**378**:705-713 Available at <http://www.nejm.org/doi/full/10.1056/NEJMoa1501315?rss=searchAndBrowse#t=article> and http://slides.hcvonline.org/uploads/149/ion4_ls.pdf
- [20] Nina Kim H, Spach DH. Treatment of Hepatitis C in Patients with HIV Coinfection. Available from: <http://www.hepatitisc.uw.edu/go/special-populations-situations/treatment-hiv-coinfection/core-concept/all#drug-drug-interactions-hiv-hcv-coinfection-treatment>
- [21] Verucchi G et al. Human immunodeficiency virus and hepatitis C virus coinfection: Epidemiology, natural history, therapeutic options and clinical management. *Infection*. 2004;**32**(1):33-46 Available from: <https://www.ncbi.nlm.nih.gov/pubmed/15007741>

