

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Alternative Therapeutic Approach for Cartilage Repair

Marina Cristina Akuri, Mariana Ricci Barion,
Sandra Maria Barbalho and Élen Landgraf Guiguer

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.72478>

Abstract

The cartilage is a flexible tissue, which supports the adjacent soft tissues. The damages that cause degenerative articular diseases are marked by the increase of cytokines such as tumor necrosis factor- α (TNF- α), IL-1 β , IL-6, IL-18, and IL-17, which cause intense inflammatory process and release of metalloproteinases and disintegrin enzymes that lead to cartilage degradation. The *Curcuma longa* possesses bioactive compounds designated as curcuminoids that display therapeutic potential in several pathologies. Curcumin is one of these compounds that may exhibit anti-inflammatory, antioxidant, antiviral, antibacterial, and antitumor effects. It may promote decrease of IL-1 β , IL-6, IL-8, TNF- α , COX-2, and reactive oxygen species. Furthermore, curcumin inhibits the activity of several kinases related to the degradation of the cartilage, including tyrosine kinase, p21-activated kinase, mitogen-activated protein kinase, protein kinase C, the activator protein 1 pathway, and NF- κ B leading to the suppression of the production of metalloproteinases and inflammatory cytokines. Curcumin has also been related to the stimulation of the production of type II collagen and glycosaminoglycan by chondrocytes. Studies have shown that this compound may alleviate joint pain and crepitation, reduce the use of other drugs for pain relief, stimulate the production of type II collagen and glycosaminoglycan resulting in a protective and anti-inflammatory action of cartilage and bones, and improve the quality of life of the patients.

Keywords: cartilage, inflammation, *Curcuma longa*, curcumin

1. Introduction

The articular cartilage is a flexible tissue, which supports the adjacent soft tissues and possesses the extracellular matrix (ECM), collagen, chondrocyte, proteoglycans, and water [1]. This tissue is alymphatic, avascular, and aneural, and for these reasons, when a severe damage occurs, the self-repair is a highly difficult process [2–4].

The damages that cause degenerative articular diseases are marked by the increase of cytokines that cause intense inflammatory process and enzymes that cause cartilage degradation [5, 6].

The osteoarthritis (OA) is an example of a progressive degenerative disease characterized by a chronic inflammatory process, joint pain, and loss of function and injury of adjacent tissue. The great destruction of the articular cartilage is the main characteristic of this disease [7–9], and therefore, it is used, in this chapter, as a prototype of cartilage destruction and regeneration.

Drugs such as nonsteroidal anti-inflammatory drugs (NSAIDs) and acetaminophen are the therapeutic approaches for the pharmacological treatment of degenerative diseases. However, this kind of medications is associated with gastrointestinal, cardiovascular, and renal adverse effects and do not effectively inhibit the disease progression and destruction of cartilage [7, 8, 10, 11]. Furthermore, corticosteroids, another therapeutic option due to their potent anti-inflammatory action and ability to reduce symptoms, should also not be used for an extended period because they can lead to a more rapid progression of OA [12].

The development of therapeutic alternatives that do not cause adverse effects and inhibit the progression of the disease is urgent and, therefore, has been widely studied. *Curcuma longa*, herbal medicine, has been shown to be one of these possible alternatives because it presents significant benefits in degenerative diseases such as OA, and this plant may play a crucial role in the reduction of the inflammatory pathways [13].

2. Physiopathology and cartilage destruction

In healthy cartilage, chondrocytes can form ECM components and enzymes that degrade cartilage in equilibrium. Although the pathophysiology of OA and its triggers have not yet been fully elucidated, it is known that inflammation, joint destruction, synovitis, and osteoclastogenesis are involved [9].

In OA, there is an increase in the enzymes involved in the cartilage degradation such as disintegrin and metalloproteinase (MMP) with a thrombospondin motif (ADAMTS). This enzymatic increase occurs due to stimulation by interleukins (IL) and inflammatory mediators such as tumor necrosis factor- α (TNF- α), IL-1 β , IL-6, IL-18, and IL-17 [6, 14].

The most widely studied and related to the destruction of cartilage are ADAMTS4 and ADAMTS5 that are released after stimulation of inflammatory cytokines such as IL-1 β . They are aggrecanases and aggrecan aggregation of proteoglycans and one of the components of ECM. After degradation of aggrecans by these enzymes, MMP-3 acts in synergism in the degradation of proteoglycans [8, 15–17].

MMPs are enzymes implicated primarily in the destruction of type II collagen and therefore play a fundamental role in the destruction of cartilage. MMP-1, MMP-3, MMP-9, and MMP-13 are the most involved enzymes in this process, and the last one is not found in adult cartilage without OA. Fragments from cleavage of collagen type 2 by MMPs amplify the destruction of ECM and amplify the release of more MMPs [5–9].

Activated synoviocytes also produce inflammatory cytokines such as IL-1, IL-6, and TNF- α , which act by amplifying the inflammatory process and cartilaginous destruction. There is increased release of reactive oxygen species (ROS), mainly nitric oxide (NO), peroxynitrite (ONOO⁻), and superoxide anion radicals (O₂⁻). Other inflammatory mediators such as cyclooxygenase-2 (COX-2), produced by synovial monocytes, and prostaglandins 2 (PGE₂) are also involved in the pathophysiology of the disease [6, 15].

The nuclear factor-kappa B (NF- κ B) pathway is responsible for the production of various cytokines and induction of inflammation. When stimulated by interleukins IL-1 β and TNF- α , there is activation of I kappa beta kinase (IKK), which promotes the phosphorylation of IKB- α . Thus, IKB- α is degraded by ubiquitination, and the dimers compounded by p50 and p65 reach the nucleus and can stimulate the expression of more than 400 genes, of which some are pro-inflammatory and pro-apoptotic genes [8, 18]. Therefore, there is production of various interleukins, including IL-1, IL-6, IL-8, and IL-10 [8, 15, 16].

Besides the destruction of cartilage in OA, an intense process of bone resorption occurs. This process is a result of osteoclast activation known as osteoclastogenesis [19]. The receptor activator NF-kappa ligand (RANKL) is produced by some cells as the osteoblast and has an affinity for RANK, which is present in the membrane of osteoclast precursor cells [20]. When RANKL binds to RANK, a phosphorylation process occurs, culminating in the activation of NF-kB [5, 21, 22]. The osteoprotegerin also has an affinity for RANK, thus competing with RANKL, inducing apoptosis of mature osteoclasts [5]. In the OA, the increase of RANKL and the decrease of OPG are observed [23, 24]. **Figure 1** summarizes the inflammatory process in the cartilage.




Figure 1. The activation of NF- κ B is related to the release of TGF β , IL-1, IL-6, and IL-8 and further activation of TH17 that leads to the stimulation of several cells and expression of other inflammatory cytokines and metalloproteinases (MMP), and further development of features characteristic of inflammation and degradation of cartilage and bone. NF- κ B: nuclear factor κ B; IL: interleukin; TH17: T-helper 17; MMP: matrix metalloproteinase; RANKL: receptor activator of nuclear factor κ B ligand; TH17: type 17 T-helper; TNF- α : tumor necrosis factor- α ; TGF- β , transforming growth factor- β ; ROS: reactive oxygen species; and NO: nitric oxide.

3. *Curcuma longa*

C. longa, or turmeric or saffron, is native to Asia and India and belongs to the Zingiberaceae family, and its rhizome has been used as a seasoning and in the traditional medicine since ancient times [18, 25].

The bioactive compounds derived from turmeric are called curcuminoids and have shown therapeutic potential in various pathologies. The three most important compounds originated from this rhizome are curcumin (diferuloylmethane), bisdemethoxycurcumin, and demethoxycurcumin, which are present, respectively, in concentrations of 77, 17, and 3%. Curcumin gives the typical yellowish coloration of the rhizome, and this part of the plant is the most widely studied [18, 26, 27].

Several studies have been conducted in order to show their actions *in vitro* and *in vivo*. Curcumin acts with different mechanisms and in different cell types and pathways. It shows anti-inflammatory, antioxidant, antiviral, antibacterial, and antitumor effects. Its therapeutic potential covers diseases such as cancer, Alzheimer's disease, osteoporosis, inflammatory bowel disease, depression, arthritis, diabetes, vitiligo, endometriosis, and several others. **Figure 2** shows some effects of curcumin [5, 28–30].

Studies on the action of curcumin and its analogs show that it can act directly or indirectly in the decrease of the formation of inflammatory molecules and pro-inflammatory transcription factors. Under its action, there is a reduction of IL-1 β , IL-6, IL-8, TNF- α , NF-kB, COX-2, and reactive oxygen species (ROS). Apart from that, curcumin has been shown to inhibit the activity of several kinases related to the degradation of the cartilage, including a tyrosine




Figure 2. Some benefits of curcumin on human health.

kinase, p21-activated kinase 1 (PAK1), mitogen-activated protein kinase (MAPK), and protein kinase C (PKC). **Figure 3** shows the process of cartilage inflammation and the effects of curcumin in the healing process [29, 31, 32].

Many studies have shown that curcumin has potent effects on the induction of apoptosis and decreased tumor cell proliferation and may promote the inhibition of important angiogenesis regulators, signal transducers and activators of transcription 3 (STAT3), and vascular endothelial growth factor (VEGF). Besides, it downregulates the expression of differentiated embryochondrocyte expressed gene 1 (DEC1) and hypoxia-inducible factor-1- α (HIF-1 α) [33–36].

Furthermore, several authors have shown that the supplementation with curcumin may bring a plethora of benefits in the treatment and prevention of the osteopenia [37]. This compound has been demonstrated to be able to avert the suppression of osteoblasts proliferation and to enhance the index of osteoprotegerin and RANKL, which indicates osteoblastogenesis [38].

As mentioned earlier, the actions of curcumin vary from potent anti-inflammatory and anti-apoptotic to antioxidant [39]. The wide variety of sites of actions and consequently decrease in the inflammation markers make this compound and its analogs extremely promising in chronic inflammatory diseases such as OA [5, 28]. Also, this herbal medicine inhibits the phosphorylation of IKB- α and thereby reduces cartilage degradation, as shown in **Figure 4**.

Conventional OA therapies are restricted to the reduction of symptoms in patients, but they do not decrease the degradation of cartilage and, consequently, do not alter the progression of the disease. For these reasons, the need for new therapies is striking, and curcumin and its analogs have become extremely promising in this context [13, 28].




Figure 3. The inflammation of the cartilage may occur due to several processes such as an increase in the expression of enzymes, increase in the formation of ROS, and release of cytokines. The consequence is the loss of type II collagen and glycosaminoglycan resulting in the degradation of the cartilage. Curcumin interferes in this scenario and may help in the healing process. ROS: reactive oxygen species; IL: interleukin; TNF- α : tumor necrosis factor- α ; PGE2: prostaglandin E2.




Figure 4. Effects of curcumin on the inhibition of the process involving the degradation of cartilage. IKK: I kappa B kinase; IKB α : inhibitor of kappa B; NF- κ B: nuclear factor κ B.

4. The potential effects of *Curcuma longa*

In articular cartilage, the ECM is composed of different compounds such as collagen, proteoglycans (glycosaminoglycan and proteins) mainly aggrecan and non-collagenous proteins [28, 40].

In the intra-articular space, there is the synovial fluid, which is enveloped by the synovial membrane and is also responsible for the nutrition of articular cartilage cells. The main cells in the synovial membrane are synoviocytes that have phagocytic functions and are responsible for the production of synovial fluid [41] and contribute to the inflammatory process when it releases several cytokines and proteases which contribute to joint destruction [42, 43].

Some degenerative diseases are involved with synovial inflammation and destruction of ECM of articular cartilage [44, 45]. According to the World Health Organization (WHO), musculo-skeletal or rheumatic conditions consist over 150 syndromes and diseases. These ailments are liable for chronic pain, disability, and dysfunction. Among these diseases, rheumatoid arthritis, osteoarthritis, spinal disorders, and severe limb trauma deserve special mention because of the greatest impact on society such as healthcare expenditures. In developed countries, OA is one of the most disabling diseases [45].

OA is a degenerative disorder involving synovial inflammation and destruction of ECM leading to several symptoms such as pain, disability, and significant morbidity, requiring

many medications that, in most cases, do not show effective actions resulting in the damage of the synovial tissue. For these reasons, new pharmacotherapies and therapies for this illness are essential [28].

As pointed earlier, curcumin may act in many different locals of inflammation resulting, directly or indirectly, in the reduction of the production of inflammatory mediators and interleukins, resulting in less destruction of cartilage. Besides that, patients treated with curcumin have decreased C-reactive protein, a marker of inflammation [46, 47].

Moreover, curcumin has been shown to inhibit the activator protein 1 (AP-1) pathway and NF- κ B leading to the suppression of the production of MMP-3, MMP-9, and MMP-13 [15, 19]. Zhang et al. [9] demonstrated in a mouse model that the production of MMP-1, MMP-3, MMP-13, IL-1 β , TNF- α , and ADAMTS5 was decreased when the animals were treated with curcumin. They also showed an increase in the expression of the chondroprotective gene CITED 2 (Cbp/P300 interacting transactivator with Glu/Asp rich carboxy terminal domain 2), which seems to be involved in the suppression of NF- κ B activity [9, 19]. Curcumin has also been related to the stimulation of the production of type II collagen and glycosaminoglycan by chondrocytes [5].

Curcumin inhibits the activation of I kappa B kinase (IKK) in chondrocytes, osteoblasts, and synovial cells [15, 48]. By inhibiting the phosphorylation of this kinase, curcumin prevents the activation of NF- κ B. Consequently, it inhibits the expression of pro-apoptotic genes in chondrocytes (caspase-3) and the formation of inflammatory mediators [18]. Thus, it is responsible for the downregulation of lipoxygenases, COX-2, phospholipase A2, prostaglandin E2 (PGE2), IL-1 β , IL-6, and IL-8 [19, 30]. Wherefore, curcumin blocks the signaling by NF- κ B, leading to the inhibition of this factor resulting in the decrease of the degradation of collagen. This pathway is induced by the activation of the chondrocytes stimulated by IL-1 [15, 16].

Curcumin inhibits TNF- α , which is associated with increased cartilage reabsorption. This cytokine associated with IL-6 and IL-1 inhibits the proteoglycan synthesis [5, 49, 50].

Studies have shown that compounds from *Curcuma* sp. can alleviate joint pain and crepitation, which lead to improved scores on WOMAC (Western Ontario and McMaster Universities Osteoarthritis Index), improve function, reduce the use of other drugs for pain relief, and is as effective as the use of ibuprofen [51–59].

Therefore, curcumin acts on the NF- κ B system, in addition to the stimulation of the production of type II collagen and glycosaminoglycan resulting in a protective and anti-inflammatory action of cartilage and bones, reducing pain and improving the quality of life of patients with degenerative diseases [18].

5. Disadvantages of *Curcuma longa*

The major problem of curcumin is that it is extremely hydrophobic and thus has low oral bioavailability, thus decreasing their beneficial effects. Another problem is the rapid metabolism of curcuminoids considering the extensive biotransformation and consequent reduction in the plasmatic levels [25, 60].

Some techniques, such as nanoparticles, phospholipid complexes, and liposomes, have been used as drug delivery systems to improve the bioavailability of these substances [61, 62]. Some compounds, such as folic acid, piperine, phosphatidylcholine, galactose, and the complex arginine-glycine-aspartic acid, are also used to improve this bioavailability and effects. Green tea and collagen associated with curcumin extracts may also enhance its effects [8, 30, 63, 64].

6. Conclusions

The curcumin has been used as an alternative therapy in the control of cartilage healing once it may interfere with the inflammatory pathways reducing the release of pro-inflammatory cytokines. Nevertheless, the use of curcumin and its analogs need to be more extensively studied and tested to determine the bioavailability, the therapeutic properties, adequate delivery formulations, doses, and possible risks of use.

Author details

Marina Cristina Akuri¹, Mariana Ricci Barion¹, Sandra Maria Barbalho^{1,2*} and Élen Landgraf Guiguer^{1,2}

*Address all correspondence to: sbarbalho@gmail.com

1 Medical School of Marília, UNIMAR, Marília, São Paulo, Brazil

2 Food Technology School, Marília, São Paulo, Brazil

References

- [1] Fernández-Torres J, Martínez-Nava GA, Gutiérrez-Ruiz MC, Gómez-Quiroz LE, Gutiérrez M. Role of HIF-1 α signaling pathway in osteoarthritis: A systematic review. *Revista Brasileira de Reumatologia English Edition*. 2017;**57**:162-173. DOI: 10.1016/j.rbre.2016.07.008
- [2] Zhou F, Zhang X, Cai D, Li J, Mu Q, Zhang W, Zhu S, Jiang Y, Shen W, Zhang S, Ouyang HW. Silk fibroin-chondroitin sulfate scaffold with immuno-inhibition property for articular cartilage repair. *Acta Biomaterialia*. 2017;**63**:64-75. DOI: 10.1016/j.actbio.2017.09.005 pii: S1742-7061(17)30569-X
- [3] Ramezanifard R, Kabiri M, Hanaee Ahvaz H. Effects of platelet rich plasma and chondrocyte co-culture on MSC chondrogenesis, hypertrophy and pathological responses. *EXCLI Journal*. 2017;**16**:1031-1045. DOI: 10.17179/excli2017-453
- [4] Xu J, Zhang C. In vitro isolation and cultivation of human chondrocytes for osteoarthritis renovation. *In Vitro Cellular & Developmental Biology. Animal*. 2014;**50**:623-629. DOI: 10.1007/s11626-014-9742-5

- [5] Akuri MC, Barbalho SM, Val RM, Guiguer EL. Reflections about osteoarthritis and *Curcuma longa*. *Pharmacognosy Reviews*. 2017;**11**:8-12. DOI: 10.4103/phrev.phrev_54_16
- [6] Schadow S, Simons VS, Lochnit G, Kordelle J, Gazova Z, Siebert HC, Steinmeyer J. Metabolic response of human osteoarthritic cartilage to biochemically characterized collagen hydrolysates. *International Journal of Molecular Sciences*. 2017;**18**(1):207. DOI: 10.3390/ijms18010207 pii: E207
- [7] Xie XW, Wan RZ, Liu ZP. Recent research advances in selective matrix metalloproteinase-13 inhibitors as anti-osteoarthritis agents. *ChemMedChem*. 2017;**12**:1157-1168. DOI: 10.1002/cmdc.201700349
- [8] Comblain F, Sanchez C, Lespoune I, Balligand M, Serisier S, Henrotin Y. Curcuminoids extract, hydrolyzed collagen and green tea extract synergically inhibit inflammatory and catabolic mediator's synthesis by normal bovine and osteoarthritic human chondrocytes in monolayer. *PLoS One*. 2015;**10**:e0121654. DOI: 10.1371/journal.pone.0121654
- [9] Zhang Z, Leong DJ, Xu L, He Z, Wang A, Navati M, Kim SJ, Hirsh DM, Hardin JA, Cobelli NJ, Friedman JM, Sun HB. Curcumin slows osteoarthritis progression and relieves osteoarthritis-associated pain symptoms in a post-traumatic osteoarthritis mouse model. *Arthritis Research & Therapy*. 2016;**18**:128. DOI: 10.1186/s13075-016-1025-y
- [10] Rannou F, Pelletier JP, Martel-Pelletier J. Efficacy and safety of topical NSAIDs in the management of osteoarthritis: Evidence from real-life setting trials and surveys. *Seminars in Arthritis and Rheumatism*. 2016;**45**:S18-S21. DOI: 10.1016/j.semarthrit
- [11] Li P, Zheng Y, Chen X. Drugs for autoimmune inflammatory diseases: From small molecule compounds to anti-TNF biologics. *Frontiers in Pharmacology*. 2017;**8**:460. DOI: 10.3389/fphar.2017.00460
- [12] Euppayo T, Punyapornwithaya V, Chomdej S, Ongchai S, Nganvongpanit K. Effects of hyaluronic acid combined with anti-inflammatory drugs compared with hyaluronic acid alone, in clinical trials and experiments in osteoarthritis: A systematic review and meta-analysis. *BMC Musculoskeletal Disorders*. 2017;**18**:387. DOI: 10.1186/s12891-017-1743-6
- [13] Del Grossi Moura M, Lopes LC, Biavatti MW, Kennedy SA, de Oliveira E, Silva MC, Silva MT, de Cássia Bergamaschi C. Oral herbal medicines marketed in Brazil for the treatment of osteoarthritis: A systematic review and meta-analysis. *Phytotherapy Research*. 2017;**31**(11):1676-1685. DOI: 10.1002/ptr.5910
- [14] Johnson JL. Metalloproteinases in atherosclerosis. *European Journal of Pharmacology*. 2017;**816**:93-106. DOI: 10.1016/j.ejphar.2017.09.007 pii: S0014-2999(17)30591-5
- [15] Henrotin Y, Clutterbuck AL, Allaway D, Lodwig EM, Harris P, Mathy-Hartert M, Shakibaei M, Mobasher A. Biological actions of curcumin on articular chondrocytes. *Osteoarthritis and Cartilage*. 2010;**18**:141-149. DOI: 10.1016/j.joca.2009.10.002
- [16] Wojdasiewicz P, Poniatowski LA, Szukiewicz D. The role of inflammatory and anti-inflammatory cytokines in the pathogenesis of osteoarthritis. *Mediators of Inflammation*. 2014;**2014**:561459. DOI: 10.1155/2014/561459

- [17] Yang CY, Chanalaris A, Troeberg L. ADAMTS and ADAM metalloproteinases in osteoarthritis—Looking beyond the 'usual suspects. *Osteoarthritis and Cartilage*. 2017;**25**:1000-1009. DOI: 10.1016/j.joca.2017.02.791
- [18] Chin KY. The spice for joint inflammation: Anti-inflammatory role of curcumin in treating osteoarthritis. *Drug Design, Development and Therapy*. 2016;**10**:3029-3042. DOI: 10.2147/DDDT.S117432
- [19] Yeh CC, YH S, Lin YJ, Chen PJ, Shi CS, Chen CN, et al. Evaluation of the protective effects of curcuminoid (curcumin and bisdemethoxycurcumin)-loaded liposomes against bone turnover in a cell-based model of osteoarthritis. *Drug Design, Development and Therapy*. 2015;**9**:2285-2300. DOI: 10.2147/DDDT.S78277
- [20] Tamma R, Zallone A. Osteoblast and osteoclast crosstalks: From OAF to Ephrin. *Inflammation & Allergy Drug Targets*. 2012;**11**:196-200
- [21] Bharti AC, Takada Y, Aggarwal BB. Curcumin (diferuloylmethane) inhibits receptor activator of NF-kappa B ligand-induced NF-kappa B activation in osteoclast precursors and suppresses osteoclastogenesis. *Journal of Immunology*. 2004;**172**:5940-5947. DOI: <https://doi.org/10.4049/jimmunol.172.10.5940>
- [22] Wei S, Teitelbaum SL, Wang MW, Ross FP. Receptor activator of nuclear factor-kappa b ligand activates nuclear factor-kappa b in osteoclast precursors. *Endocrinology*. 2001;**142**:1290-1295. DOI: 10.1210/endo.142.3.8031
- [23] Liu YD, Yang HX, Liao LF, Jiao K, Zhang HY, Lu L, et al. Systemic administration of strontium or NBD peptide ameliorates early stage cartilage degradation of mouse mandibular condyles. *Osteoarthritis and Cartilage*. 2016;**24**:178-187. DOI: 10.1016/j.joca.2015.07.022
- [24] Xu L, Guo H, Li C, Xu J, Fang W, Long XA. Time-dependent degeneration manner of condyle in rat CFA-induced inflamed TMJ. *American Journal of Translational Research*. 2016;**8**:556-567
- [25] Mahmood K, Zia KM, Zuber M, Salman M, Anjum MN. Recent developments in curcumin and curcumin based polymeric materials for biomedical applications: A review. *International Journal of Biological Macromolecules*. 2015;**81**:877-890. DOI: 10.1016/j.ijbiomac.2015.09.026
- [26] Yadav SK, Sah AK, Jha RK, Sah P, Shah DK. Turmeric (curcumin) remedies gastroprotective action. *Pharmacognosy Reviews*. 2013;**7**:42-46. DOI: 10.4103/0973-7847.112843
- [27] Aggarwal BB, Harikumar KB. Potential therapeutic effects of curcumin, the anti-inflammatory agent, against neurodegenerative, cardiovascular, pulmonary, metabolic, autoimmune and neoplastic diseases. *The International Journal of Biochemistry & Cell Biology*. 2009;**41**:40-59. DOI: 10.1016/j.biocel.2008.06.010
- [28] A M, Henrotin Y, Biesalski HK, Shakibaei M. Scientific evidence and rationale for the development of curcumin and resveratrol as nutraceuticals for joint health. *International Journal of Molecular Sciences*. 2012;**13**:4202-4232. DOI: 10.3390/ijms13044202

- [29] Prasad S, Tyagi AK. Curcumin and its analogues: A potential natural compound against HIV infection and AIDS. *Food & Function*. 2015;**6**:3412-3419. DOI: 10.1039/c5fo00485c
- [30] J Wang, J Ma, JH Gu, FY Wang, XS Shang, HR Tao, X Wang. Regulation of type II collagen, matrix metalloproteinase-13 and cell proliferation by interleukin-1 β is mediated by curcumin via inhibition of NF- κ B signaling in rat chondrocytes. *Molecular Medicine Reports*. 2017;**16**:1837-1845. DOI: 10.3892/mmr.2017.6771
- [31] Oliviero F, Scanu A, Zamudio-Cuevas Y, Punzi L, Spinella P. Anti-inflammatory effects of polyphenols in arthritis. *Journal of the Science of Food and Agriculture*. 2017. DOI: 10.1002/jsfa.8664
- [32] Agrawal DK, Mishra PK. Curcumin and its analogues: Potential anticancer agents. *Medicinal Research Reviews*. 2010;**30**:818-860. DOI: 10.1002/med.20188
- [33] Wang XP, Wang QX, Lin HP, Chang N. Anti-tumor bioactivities of curcumin on mice loaded with gastric carcinoma. *Food & Function*. 2017;**8**(9):3319-3326. DOI: 10.1039/c7fo00555e
- [34] Feng T, Wei Y, Lee RJ, Zhao L. Liposomal curcumin and its application in cancer. *International Journal of Nanomedicine*. 2017;**21**(12):6027-6044. DOI: 10.2147/IJN.S132434
- [35] Xu X, Zhu Y. Curcumin inhibits human non-small cell lung cancer xenografts by targeting STAT3 pathway. *American Journal of Translational Research*. 2017;**9**(8):3633-3641
- [36] Saberi-Karimian M, Katsiki N, Caraglia M, Boccellino M, Majeed M, Sahebkar A. Vascular endothelial growth factor: An important molecular target of curcumin. *Critical Reviews in Food Science and Nutrition*. 2017;**30**:1-14. DOI: 10.1080/10408398.2017.1366892
- [37] Riva A, Togni S, Giacomelli L, Franceschi F, Eggenhoffner R, Feragalli B, Belcaro G, Cacchio M, Shu H, Dugall M. Effects of a curcumin-based supplementation in asymptomatic subjects with low bone density: A preliminary 24-week supplement study. *European Review for Medical and Pharmacological Sciences*. 2017;**21**:1684-1689
- [38] Chen Z, Xue J, Shen T, Mu S, Fu Q. Curcumin alleviates glucocorticoid-induced osteoporosis through the regulation of the Wnt signaling pathway. *International Journal of Molecular Medicine*. 2016;**37**:329-338. DOI: 10.3892/ijmm.2015.2432
- [39] Flores G. *Curcuma longa* L. extract improves the cortical neural connectivity during the aging process. *Neural Regeneration Research*. 2017 Jun;**12**:875-880. DOI: 10.4103/1673-5374.208542
- [40] Mittelstaedt D, Xia Y. Depth-dependent glycosaminoglycan concentration in articular cartilage by quantitative contrast-enhanced micro-computed tomography. *Cartilage*. 2015;**6**:216-225. DOI: 10.1177/1947603515596418
- [41] Lovell S, Burchell RK, Roady PJ, Fredrickson RL. Gal. Canine intrathoracic sarcoma with ultrastructural characteristics of human synovial sarcoma - case report. *BMC Veterinary Research*. 2017;**13**:247. DOI: 10.1186/s12917-017-1181-6

- [42] Hao L, Wan Y, Xiao J, Tang Q, Deng H, Chen L. A study of Sirt1 regulation and the effect of resveratrol on synoviocyte invasion and associated joint destruction in rheumatoid arthritis. *Molecular Medicine Reports*. 2017;**16**:5099-5106. DOI: 10.3892/mmr.2017.7299
- [43] Benedetti G, Bonaventura P, Lavocat F, Miossec P. IL-17A and TNF- α increase the expression of the antiapoptotic adhesion molecule amigo-2 in arthritis synoviocytes. *Frontiers in Immunology*. 2016;**7**:254. DOI: 10.3389/fimmu.2016.00254
- [44] Buckwalter JA, Mankin HJ. Articular cartilage: Degeneration and osteoarthritis, repair, regeneration, and transplantation. *Instructional Course Lectures*. 1998;**47**:487-504
- [45] Mobasheri A, Airley R, Foster CS, Schulze-Tanzil G, Shakibaei M. Post-genomic applications of tissue microarrays: Basic research, prognostic oncology, clinical genomics and drug discovery. *Histology and Histopathology*. 2004;**19**:325-335. DOI: 10.14670/HH-19.325.
- [46] Mollazadeh H, Cicero AFG, Blesso CN, Pirro M, Majeed M, Sahebkar A. Immune modulation by curcumin: The role of interleukin-10. *Critical Reviews in Food Science and Nutrition*. 2017;**11**:1-13. DOI: 10.1080/10408398.2017.1358139
- [47] Jagetia GC, Aggarwal BB. "spicing up" of the immune system by curcumin. *Journal of Clinical Immunology*. 2007;**27**:19-35. DOI: 10.1007/s10875-006-9066-7
- [48] Buhrmann C, Mobasheri A, Matis U, Shakibaei M. Curcumin mediated suppression of nuclear factor-kB promotes chondrogenic differentiation of mesenchymal stem cells in a high-density co-culture microenvironment. *Arthritis Research & Therapy*. 2010;**12**:R127. DOI: 10.1186/ar3065
- [49] Saklatvala J. Tumour necrosis factor alpha stimulates resorption and inhibits synthesis of proteoglycan in cartilage. *Nature*. 1986;**322**:547-549. DOI: 10.1038/322547a0
- [50] Séguin CA, Bernier SM. TNF α suppresses link protein and type II collagen expression in chondrocytes: Role of MEK1/2 and NF-kappaB signaling pathways. *Journal of Cellular Physiology*. 2003;**197**:356-369. DOI: 10.1002/jcp.10371
- [51] Perkins K, Sahy W, Beckett RD. Efficacy of *Curcuma* for treatment of osteoarthritis. *Journal of Evidence-Based Complementary & Alternative Medicine*. 2017;**22**:156-165. DOI: 10.1177/2156587216636747
- [52] Panahi Y, Alishiri GH, Parvin S, Sahebkar A. Mitigation of systemic oxidative stress by curcuminoids in osteoarthritis: Results of a randomized controlled trial. *Journal of Dietary Supplements*. 2016;**13**:209-220. DOI: 10.3109/19390211.2015.1008611
- [53] Kuptniratsaikul V, Dajpratham P, Taechaarpornkul W, Buntragulpoontawe M, Lukkanapichonchut P, Chootip C, et al. Efficacy and safety of *Curcuma domestica* extracts compared with ibuprofen in patients with knee osteoarthritis: A multicenter study. *Clinical Interventions in Aging*. 2014;**9**:451-458. DOI: 10.2147/CIA.S58535
- [54] Appelboom T, Maes N, Albert A. A new *Curcuma* extract (flexofytol[®]) in osteoarthritis: Results from a Belgian real-life experience. *Open Rheumatology Journal*. 2014;**8**:77-81. DOI: 10.2174/1874312901408010077

- [55] Kizhakkedath R. Clinical evaluation of a formulation containing *Curcuma longa* and *Boswellia serrata* extracts in the management of knee osteoarthritis. *Molecular Medicine Reports*. 2013;**8**:1542-1548. DOI: 10.3892/mmr.2013.1661
- [56] Madhu K, Chanda K, Saji MJ. Safety and efficacy of *Curcuma longa* extract in the treatment of painful knee osteoarthritis: A randomized placebo-controlled trial. *Inflammopharmacology*. 2013;**21**:129-136. DOI: 10.1007/s10787-012-0163-3
- [57] Kertia N, Asdie AH, Rochmah W, Marsetyawan. Ability of curcuminoid compared to diclofenac sodium in reducing the secretion of cyclooxygenase-2 enzyme by synovial fluid's monocytes of patients with osteoarthritis. *Acta Medica Indonesiana*. 2012;**44**:105-113
- [58] Chopra A, Lavin P, Patwardhan B, Chitre D. A 32-week randomized, placebo-controlled clinical evaluation of RA-11, an Ayurvedic drug, on osteoarthritis of the knees. *Journal of Clinical Rheumatology*. 2004;**10**:236-245. DOI: 10.1097/01.rhu.0000138087.47382.6d
- [59] Chandrasekaran CV, Sundarajan K, Edwin JR, Gururaja GM, Mundkinajeddu D, Agarwal A. Immune-stimulatory and anti-inflammatory activities of *Curcuma longa* extract and its polysaccharide fraction. *Pharmacognosy Research*. 2013;**5**:71-79. DOI: 10.4103/0974-8490.110527
- [60] Szymusiak M, Hu X, Leon Plata PA, Ciupinski P, Wang ZJ, Liu Y. Bioavailability of curcumin and curcumin glucuronide in the central nervous system of mice after oral delivery of nano-curcumin. *International Journal of Pharmaceutics*. 2016;**511**:415-423. DOI: 10.1016/j.ijpharm.2016.07.027
- [61] Zamarioli CM, Martins RM, Carvalho EC, Freitas Luis AP. Nanoparticles containing curcuminoids (*Curcuma longa*): Development of topical delivery formulation. *Revista Brasileira de Farmacognosia*. 2015;**25**:53-60. DOI: 10.1016/j.bjp.2014.11.010
- [62] Mirzaei H, Naseri G, Rezaee R, Mohammadi M, Banikazemi Z, Mirzaei HR, et al. Curcumin: A new candidate for melanoma therapy. *International Journal of Cancer*. 2016;**139**:1683-1695. DOI: 10.1002/ijc.30224
- [63] Gupta NK, Dixit VK. Bioavailability enhancement of curcumin by complexation with phosphatidyl choline. *Journal of Pharmaceutical Sciences*. 2011;**100**:1987-1995. DOI: 10.1002/jps.22393
- [64] Bhardwaj RK, Glaeser H, Becquemont L, Klotz U, Gupta SK, Fromm MF. Piperine, a major constituent of black pepper, inhibits human P-glycoprotein and CYP3A4. *The Journal of Pharmacology and Experimental Therapeutics*. 2002;**302**:645-650. DOI: 10.1124/jpet.102.034728

