

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Toward Future Engineering of the *N*-Glycosylation Pathways in Microalgae for Optimizing the Production of Biopharmaceuticals

Rodolphe Dumontier, Alain Mareck,
Narimane Mati-Baouche, Patrice Lerouge and
Muriel Bardor

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.73401>

Abstract

Microalgae are eukaryotic and photosynthetic organisms which are commonly used in biotechnology to produce high added value molecules. Recently, biopharmaceuticals such as monoclonal antibodies have been successfully produced in microalgae such as *Chlamydomonas reinhardtii* and *Phaeodactylum tricornutum*. Most of these recombinant proteins are indeed glycosylated proteins, and it is well established that their glycan structures are essential for the bioactivity of the biopharmaceuticals. Therefore, prior to any commercial usage of such algae-made biopharmaceuticals, it is necessary to characterize their glycan structures and erase glycosylation differences that may occur in comparison with their human counterpart. In this context, the chapter summarizes successful attempts to produce biopharmaceuticals in microalgae and underlines current information regarding glycosylation pathways in microalgae. Finally, genome editing strategies that would be essential in the future to optimize the microalgae glycosylation pathways are highlighted.

Keywords: antibodies, biopharmaceuticals, genome editing tools, glycosylation, microalgae

1. Introduction

Microalgae are currently used for a broad spectrum of industrial applications including food and livestock feed industries, bioenergy, cosmetics, healthcare and environment [1–4]. Recently, due to their numerous advantages (high growth rate, easy cultivation, low production cost, etc.), microalgae have emerged as a solar-fueled green alternative cell factories for the production

of recombinant proteins [4–7]. Among different attempts to produce vaccines and biopharmaceuticals in microalgae, the production of monoclonal antibodies (mAbs) represents the most extensive work [7, 8]. Indeed, the first significant effort to produce recombinant mAb fragments was made in the green microalga *Chlamydomonas reinhardtii* with the synthesis and accumulation in its chloroplast of a human single chain antibody directed against the herpes simplex virus glycoprotein D (HSV8-lsc) [8]. Later, a full-length human IgG1 directed against anthrax was produced successfully in the chloroplast of *C. reinhardtii* [9]. The *Chlamydomonas*-made mAb was able to bind the anthrax protective antigen 83 (PA83) [9]. In another study, a series of complex chimeric proteins was expressed in the chloroplast of *C. reinhardtii*. Such chimeric proteins were composed of a single chain antibody fragment (*scFv*) targeting the B-cell surface antigen CD22, genetically fused either to the eukaryotic ribosome inactivating protein, gelonin, from *Gelonium multiflora* [10] or to *Pseudomonas aeruginosa* exotoxin A domains 2 and 3 [11]. These molecules, termed immunotoxins, were encoded by a single gene that produces an antibody-toxin chimeric protein. Such algae-made immunotoxins are able to bind target B cells and efficiently kill them *in vitro* [11]. Full-length mAbs have also been expressed in the diatom *Phaeodactylum tricornutum* through nuclear transformation [12–14]. Those mAbs correspond respectively to a recombinant mAb directed against the nucleoprotein of Marburg virus, a close relative of Ebola virus [14] and to a human IgG1 directed against the Hepatitis B virus Antigen (HBsAg) [12, 13]. The latter has been biochemically characterized in order to check the quality of the diatom-made mAb as well as its *N*-glycosylation profile [15]. Moreover, it has been demonstrated that this glycosylated antibody is able to bind human Fcγ receptors [16], thus suggesting that it could be efficient in human therapy.

When the production of biopharmaceuticals is considered, their *N*-glycosylation has to be investigated. Indeed, among the biopharmaceuticals that were approved in 2016 and 2017, 96% were glycosylated [17]. The glycosylation of the approved biopharmaceutical represents a critical quality attribute (CQA) that may affect its safety and biological activities [18–20]. In addition, introduction by the expression system of nonhuman epitopes on the recombinant protein may induce immune response after injection to patients [21]. Thus, the *N*-glycosylation of biopharmaceuticals is a real challenge for the commercial production of biopharmaceuticals. The glycosylation state of therapeutic proteins has to be accurately identified and characterized as per the World Health Organization and International Conference on Harmonization Q6B guidelines [17]. Therefore, in the context of developing the microalgae as alternative platforms for the production of biopharmaceuticals, the capability of these unicellular eukaryotic cells to introduce *N*-glycans on their endogenous proteins and on recombinant proteins, as well as their regulation, have to be considered and understood.

2. *N*-glycosylation in microalgae

2.1. General aspects

N-glycosylation is a major post-translational modification of proteins in eukaryotes. Protein *N*-glycosylation first starts by the synthesis of a lipid-linked oligosaccharide formed by transfer of monosaccharides on a dolichol pyrophosphate (PP-Dol) anchored in the membrane of

the endoplasmic reticulum (ER) *via* the action of a set of enzymes named asparagine-linked glycosylation (ALG) [22, 23]. The final Glc₃Man₉GlcNAc₂ precursor is transferred *en bloc* by the oligosaccharyltransferase (OST) complex onto the asparagine residues of the consensus Asn-X-Ser/Thr sequences of a protein [22] (**Figure 1**). Alternative consensus sequences, such as Asn-X-Cys and Asn-X-Val, have also been found to be glycosylated in some proteins [24–26].




Figure 1. Comparison of protein *N*-glycosylation pathways in eukaryotes. Biosynthesis steps occurring in the ER are gathered in the box. Mature *N*-glycan structures observed in mammals, plants, insects, yeasts and filamentous fungi are drawn according to [33]. ■, N-acetylglucosamine; ☆, xylose; ●, mannose; ▲, fucose; ●, galactose; ◆, sialic acid; Asn, asparagine; PP-Dol, pyroPhosphate dolichol; FuT, fucosyltransferase; GalT, galactosyltransferase; SiaT, sialyltransferase; XylT, xylosyltransferase; ALG, asparagine-linked glycosylation; OST, oligosaccharyltransferase.

In the ER, neo-synthesized glycoproteins are then submitted to a quality control process through the deglycosylation by glucosidases and reglycosylation by an UDP-glucose: glycoprotein glucosyltransferase (UGGT) of the *N*-glycans. This allows the synthesis of monoglycosylated glycan intermediates that interact with ER-resident chaperones, thus ensuring proper folding of the glycoproteins [27]. When the glycoprotein is correctly folded, α -glucosidase II would finally remove the last glucose residue, and ER-mannosidase will eventually remove one mannose residue that leads to the formation of an oligomannoside $\text{Man}_{9/8}\text{GlcNAc}_2$. The quality control events are conserved in eukaryotes because they are crucial for the secretion of well-folded proteins [28]. As a consequence, whatever the expression system used, a recombinant therapeutic protein leaving the ER compartment exhibits a *N*-glycosylation similar to one of the reference proteins with unique oligomannoside $\text{Man}_{9/8}\text{GlcNAc}_2$ attached to the Asn residue of the *N*-glycosylation consensus site.

After transfer to the Golgi apparatus, oligomannosides resulting from the ER processing are modified by the action of specific mannosidases and glycosyltransferases [29]. These Golgi cell-specific repertoires give rise to various organism-specific oligosaccharides. In most eukaryotes, a *N*-acetylglucosaminyltransferase I (GnT I)-dependent *N*-glycan processing occurs (**Figure 1**). In this pathway, the α -mannosidase I converts $\text{Man}_{9/8}\text{GlcNAc}_2$ into the branched isomer of $\text{Man}_5\text{GlcNAc}_2$. Then, actions of GnT I, α -mannosidase II and GnT II, respectively, give rise to the core $\text{GlcNAc}_2\text{Man}_3\text{GlcNAc}_2$ that is common to most eukaryotes [27–31] (**Figure 1**). This core is then decorated by the action of specific glycosyltransferases that differ from one organism to another. This allows the protein to be decorated by organism-specific *N*-glycans that confer to the mature protein *in vivo* bioactivities [32]. It is worth noting that GnT I-independent *N*-glycan processing also occurs in some eukaryotes such as filamentous fungi and yeasts in which *N*-glycosylation in the Golgi apparatus results in the synthesis of high mannose and hypermannose *N*-glycans, respectively (**Figure 1**). As a consequence, in the context of the production of biopharmaceuticals by genetic engineering, such a diversity of mature *N*-linked glycans is a limitation because the expression system used may introduce inappropriate epitopes and heterogeneous glycosylation on the therapeutics and may also fail in introducing glycan sequences that are required for *in vivo* bioactivity of the biopharmaceuticals.

2.2. Protein *N*-glycosylation in microalgae

Overall, protein *N*-glycosylation in microalgae received little attention. Few studies, published in the 1990s have demonstrated that proteins secreted by green microalgae carry mainly oligomannosides or xylose-containing *N*-glycans based on affinodetection or enzymatic sequencing [34–36]. More recently, analysis by mass spectrometry of glycans *N*-linked to microalgae endogenous proteins has been reported. First, the 66 kDa cell wall glycoprotein from the red microalga *Porphyridium* sp. has been found to carry $\text{Man}_8\text{GlcNAc}_2$ and $\text{Man}_9\text{GlcNAc}_2$ oligomannosides containing 6-*O*-methyl mannose residues and substituted by one or two xylose residues [37, 38] (**Figure 2**). Investigation of *C. reinhardtii* has demonstrated that proteins in this green microalga carry oligomannosides ranging from $\text{Man}_2\text{GlcNAc}_2$ to $\text{Man}_5\text{GlcNAc}_2$ as well as $\text{Man}_4\text{GlcNAc}_2$ and $\text{Man}_5\text{GlcNAc}_2$ *N*-glycans containing 6-*O*-methyl mannoses and substituted by one or two xylose residues (**Figure 2**) [39]. Initially reported as branched




Figure 2. Major mature *N*-linked glycans from the green microalga *Chlamydomonas reinhardtii* and *Botryococcus braunii*, the red microalga *Porphyridium* sp. and the diatom *Phaeodactylum tricorutum*. *N*-glycan structures are drawn according to [33]. ■, *N*-acetylglucosamine; ☆, xylose; ●, mannose; ▲, fucose; ●, galactose; Asn, asparagine; Me, methyl.

oligomannosides, the structure of $\text{Man}_5\text{GlcNAc}_2$ was re-evaluated in 2017 as being linear sequences based on ESI- MS^n analyses [40]. Although mature *N*-glycans from *Porphyridium* sp. and *C. reinhardtii* share common structural features, the location of the xylose residues on the *N*-glycan differs between these two microalgae (**Figure 2**). As mature *N*-glycans do not exhibit any terminal GlcNAc residues, they were proposed to result from Golgi xylosylation and *O*-methylation of oligomannosides deriving from the precursor synthesized in the ER in a GnT I-independent processing, even if this needs to be completely elucidated and that methylation occurring in the ER cannot be ruled out yet [38].

N-glycan profile from *P. tricorutum* has been described to contain $\text{Man}_3\text{GlcNAc}_2$ to $\text{Man}_5\text{GlcNAc}_2$ oligomannosides and also minute amount of paucimannosidic fucosylated *N*-glycans (**Figure 2**) [41]. In contrast to *Porphyridium* sp. and *C. reinhardtii*, these *N*-glycans result from a GnT I-dependent pathway (**Figure 2**) [41]. As evidence, GnT I gene predicted in the *P. tricorutum* genome encodes an enzyme able to restore the maturation of complex-type *N*-glycans in the CHO Lec1 mutant that lacks endogenous GnT I activity [41]. *N*-glycans arising from a GnT I-dependent pathway have also been recently reported in the green microalga *Botryococcus braunii* through a glycoproteomic approach [42]. In contrast to *P. tricorutum*, these *N*-glycans harbor a GlcNAc residue at the nonreducing end as well as mono- and di-*O*-methylations of the core mannose residue. Moreover, this *N*-glycan bearing a terminal GlcNAc resulting from the GnT I activity could be further elongated with an additional hexose or methyl-hexose residue. In addition, proteins from this green microalga also exhibit methylated *N*-linked oligomannosides carrying core fucose and core xylose residues (**Figure 2**) [42].

In support to these biochemical data, protein *N*-glycosylation in microalgae can be drawn on the basis of public genomic databases. Microalgae genomes from different phyla are available

to date (<https://genome.jgi.doe.gov/pages/tree-of-life.jsf>) [4, 43]. Since protein *N*-glycosylation occurs in the ER and the Golgi apparatus, bioinformatics analyses of microalgae genomes must be investigated independently for the two compartments: search for gene encoding proteins involved in the precursor biosynthesis and the ER protein quality control on the one hand, and search for Golgi glycosidases and glycosyltransferases involved in the synthesis of mature *N*-glycans on the other hand.

Genes encoding subunits of OST, glucosidases, as well as ER-resident UGGT and chaperones are predicted in microalgae genomes suggesting that the process of ER quality control in these unicellular organisms is similar to the one described in other eukaryotic cells [41, 44, 45]. Among these putative ER candidates, only the activity of the $\alpha(1,3)$ -glucosidase, also called glucosidase II, from the red microalga *Porphyridium* sp. has been biochemically confirmed [44]. Most ALG genes are also predicted in microalgae genomes [39, 41, 44] suggesting that the synthesis of the oligosaccharide precursor is overall conserved. However, some of these ALG, that is ALG3, ALG9 and ALG12, are not predicted in *C. reinhardtii* [39, 45]. These ER enzymes are involved in the completion of the biosynthesis of the precursor Man₉GlcNAc₂-PP-Dol, prior to its glucosylation, by addition of mannose residues on the $\alpha(1,6)$ -mannose arm of the core (**Figure 1**). Reinvestigation in *C. reinhardtii* of the structure of oligomannosides and analysis of the ER *N*-glycan precursor [40] confirmed the absence of ALG3, ALG9 and ALG12 activities and the synthesis in this green microalga of linear oligomannoside sequences instead of branched isomer initially proposed in [39]. It is worth noting that in this truncated ER pathway, the presence of the triglucosyl extension is likely sufficient to ensure interaction of the *N*-glycan precursor with chaperones of the ER quality control process. In addition to the lack of the ALG3, ALG9 and ALG12 in *C. reinhardtii*, other microalgae genomes lack genes encoding ALG10 and GCS1, an $\alpha(1,2)$ -glucosidase [44]. Because ALG10 is the $\alpha(1, 2)$ -glucosyltransferase responsible for the addition of the $\alpha(1, 2)$ -glucose residue on the precursor *N*-glycan and GCS1 is responsible for trimming this residue, we hypothesize that the ER quality control in these microalgae involved only diglucosylated *N*-glycan intermediates.

With regard to Golgi *N*-glycosylation events, the presence of GnT I is predicted in some microalgae including haptophytes and cryptophytes, but not in *C. reinhardtii*, *Volvox* and *Ostreococcus* [41, 42]. As mentioned previously, *P. tricornutum* GnT I activity was confirmed by the complementation of CHO Lec 1 mutant cell line [41]. A recent study of *B. braunii* [42] confirmed the involvement of this transferase in this green microalga *N*-glycosylation pathway. Concerning other Golgi enzymes, α -mannosidases (CAZy GH 47) and $\alpha(1,3)$ -fucosyltransferases (CAZy GT10) are also predicted in microalgae genomes studied so far [41, 44, 45]. These enzymes are respectively involved in the trimming of mannose residue of oligomannosides and the transfer of fucose on the proximal GlcNAc. These sequences exhibit peptide motifs that were demonstrated to be required for activities of such Golgi enzymes, but, in contrast to GnT I, no biochemical data of their activity and specificity are available yet.

As depicted, protein *N*-glycosylation occurring in microalgae is specific and largely differs from the one described in mammals (**Figures 1 and 2**). Therefore, production in microalgae of biopharmaceuticals exhibiting *N*-glycans compatible with their use in human therapy

would be challenging and requiring metabolic engineering of the *N*-glycosylation pathway in microalgae. This will include the inactivation of enzymes that introduce nonhuman glyco-epitopes onto *N*-linked glycans and complementation of microalgae with appropriate glycosyltransferases to introduce missing glycan sequences. These strategies have already been successfully carried out for the engineering of the *N*-glycan pathways in plants and yeasts [46, 47]. In addition, the success of the complementation with human glycosyltransferases requires the availability in the Golgi apparatus of appropriate nucleotide-activated sugars [48]. For instance, sialic acids that terminate bi-antennary *N*-glycans in mammals have not been reported in microalgae such as *P. tricornutum* and *Porphyridium* sp. [38, 41]. As well, there is no evidence for the import of GlcNAc in the Golgi apparatus in microalgae exhibiting a GnT I-independent *N*-glycan pathway, even if putative candidates for UDP-GlcNAc transporter have been identified in microalgae such as *C. reinhardtii* [49]. Indeed, the two GlcNAc of the chitobiose unit of *N*-linked glycans are transferred onto the PP-Dol lipid in the cytosolic face of the ER membrane. Currently, metabolic engineering strategies are now feasible due to the recent development of transgene expression and gene inactivation in microalgae as summarized in Section 3.

3. Genetic engineering tools now available to envision future *N*-glycosylation engineering in microalgae

3.1. Different tools to generate genome-modified organisms

Classical strategies of genetic engineering involve the modulation of gene expression including overexpression and inactivation by RNA interference [50–52]. The most used engineering methods are based on random insertional mutagenesis obtained by various processes such as conjugation, agitation with glass beads, electroporation, biolistic microparticle bombardment, agrobacterium-mediated transformation or multipulse electroporation. The transformation step is followed by phenotypic selection using antibiotics to generate genome-modified organisms [53]. Those processes present the advantage to be simple and reach a high level of transformed cells. For example, *P. tricornutum* transformation reached 1 per 10⁶ cells with biolistic bombardment system [54]. However, cell-wall-less strains are required for almost all the classical methods quoted above [50, 55]. Furthermore, genetic stability of the mutagenesis obtained after transformation by random insertion depends on microalgae species [53]. For example, a high mutagenesis stability has been shown in *C. reinhardtii* [55]. Unlike, mutagenesis was unstable in *Thalassiosira weissflogii* [56]. More recently, new tools have been developed in order to knock in, knock out, modify, replace, or insert genes. These new genetic engineering tools consist of the action of nucleases effecting their molecular scissor activities in specific loci [52]. A break in the DNA causes activation of DNA repair mechanisms, which can be either the homologous-recombination (HR) or the non-homologous end-joining (NHEJ) [52]. The HR results in sequence modification in the target locus [57]. In the NHEJ process, the two ends of the broken chromosome are stuck together causing small deletions or small insertions [57]. These events confer several modifications of the target gene such as gene inactivations or

insertions. Very little is known about these mechanisms in microalgae due to their complexity as reported by Daboussi in 2017 [53].

Several researches have recently contributed to demonstrate that particular nucleases could be used for targeting stable modifications by acting like molecular scissors. Among these nucleases, we can quote meganucleases (MNs), zinc finger nucleases (ZFNs), transcription activator-like effector nucleases (TALENs) and finally, the famous clustered regularly interspaced short palindromic repeats (CRISPR)/nuclease Cas9 system. These four cited nucleases are described in the following paragraphs.

Meganuclease is an engineered endonuclease able to recognize and cleave a long specific DNA sequence from 18 to 30 base pairs. The meganuclease strategy requires to design a homing endonuclease from the LAGLIDADG family especially the I-CreI enzyme from *C. reinhardtii* implied in the targeting of interesting gene sequences that need to be modified [58]. This was tested for the first time in 2014 using *P. tricornutum* as a model [59]. In this study, two engineered meganucleases targeting genes involved in the lipid metabolism are allowed to obtain 29% of targeted mutagenesis [59]. Even successful, this strategy is time-consuming as compared to the other alternatives [52].

Zinc finger nucleases (ZFNs) are hybrid proteins composed of a restriction enzyme *FokI* with a designed zinc-finger DNA-binding domains [60]. These *FokI* enzymes are inactive in a homodimer conformation [61]. Therefore, cleavage of a typical DNA-target sequence requires to design two different ZFNs for binding to adjacent half-sites of a specific locus. Each designed ZFN is able to recognize a sequence of 9–12 nucleotides in the genome [52]. A set of zinc finger nucleases has been recently used to modify by insertion of template DNA, the *Cop3* gene locus encoding a light-activated channel in *C. reinhardtii* [62]. Moreover, in 2017, the genome editing was reliably performed using the ZFN strategy in order to inhibit and modify nuclear photoreceptor genes in this same microalga [63]. Despite these promising results, the ZFN system is barely used because of its low specificity. Indeed, cleavage of DNA requires both ZFN monomers to recognize a homologous target in the genome in the proper spatial orientation to assemble a functional ZFN [64]. Also, ZFN system is time-consuming implementation [64]. Nowadays, other designed nucleases like TALENs or CRISPR/cas9 are emerging in the scientific community to perform genome editing in microalgae.

Transcriptional activator-like effector nuclease (TALEN) system is similar to ZFN because it uses nucleases composed of a restriction enzyme domain fused to a DNA-binding domain (here the TAL effector domain) and a nonspecific DNA cleavage domain *FokI* [65]. TALEN proteins are characterized by a repeated 34-amino acid sequence that recognizes specific DNA sequences [66]. *P. tricornutum* lipid metabolism was recently modified using TALEN [59]. In this study, seven genes involved in this metabolism were modified. Each genome modification had a high frequency reaching up to 56% of colonies with targeted mutagenesis [59]. This genetic engineering allowed creation of a high lipid-producing strain by inactivating a key gene for carbohydrate energy storage [59]. Another team has inactivated successfully the urease gene in *P. tricornutum* with 24% of transformed colonies [67]. In addition, TALEN system has also been used in order to inactivate red/far-red light-sensing *phytochrome* gene of this diatom [68].

The clustered regularly interspaced short palindromic repeats (CRISPR)/cas9 system is the most famous engineered nuclease system of this decade because it is a powerful and precise tool applied in numerous eukaryotic organisms [69]. This system is based on the RNA-guided DNA cleavage defense system from archaea and many bacteria. Indeed, these organisms are able to store bacteriophage DNA fragments along a previous bacteriophage infection in the CRISPR locus, which is formed of DNA repeat sequences spaced by a unique DNA sequence. This system establishes the basis of a bacterial defense as a response to bacteriophage attacks [70]. This defense mechanism has been highlighted for the first time by Pr Emmanuelle Charpentier and her team in 2011 [70, 71]. The CRISPR/Cas9 system has been developed into a simple toolkit based on a custom single guide RNA (sgRNA) that contains a targeting sequence (crRNA sequence) and a cas9 nuclease-recruiting sequence (tracrRNA) [52]. In microalgae, CRISPR/cas9 has been used in *C. reinhardtii* [72]. However, the Cas9 nuclease production seemed to be toxic for the microalga limiting efficiency to obtain genome-modified strains [72]. Two years later, a new assay has been performed in this same microalga using another strategy avoiding toxicity [73]. Indeed, the authors succeeded to generate CRISPR/cas9-induced NHEJ-mediated knock-in mutant strains in three loci [73]. In the same year, CRISPR/cas9 gene knockout technology has been used in *P. tricornutum* to induce mutant for the *CpSRP43* gene, a member of the chloroplast signal recognition particle pathway. Using

	MN system	ZFN system	TALEN system	CRISPR/cas9
Actor(s) of gene targeting	Chimeric endonuclease	Chimeric endonuclease	Chimeric endonuclease	RNA guide and cas9 nuclease
Engineered protein origin	<i>Chlamydomonas reinhardtii</i> <i>Saccharomyces cerevisiae</i> [59, 77]	Xanthomonas [78]	Xanthomonas [78]	Bacteria and Archaea [70]
Nuclease specificity	Low	Moderate	Moderate	High
Mutagenesis frequency in microalgae	Up to 29% [59]	Not reported	Up to 56% [59]	Up to 63% [74]
Toxicity in cells	Low	Moderate ¹	Moderate ¹	Moderate [72]
Time investment	Very high [52]	Very high [64]	Moderate	Low
Possibility of multiple gene targeting	No	No	Yes	Yes
System cost ²	Not reported	Expensive (4000–7000\$)	Expensive (3000–5000\$)	Cheap (500\$)

¹Source: <https://www.news-medical.net/life-sciences/How-Does-CRISPR-Compare-to-Other-Genome-Editing-Techniques.aspx> visited [Accessed: 2017-12-06].

²Source: <http://www.biocompare.com/Editorial-Articles/144186-Genome-Editing-with-CRISPRs-TALENs-and-ZFNs/> [Accessed: 2017-12-06].

Table 1. Comparison of four specific genomic tools based on nuclease systems in order to generate genomic-modified species in microalgae.

this strategy, the authors obtained 31% of mutation efficiency [74]. This team targeted two other genes of the diatom using this technology and obtained from 25 to 63% of mutation level [74]. Adaptability of the CRISPR/Cas9 system has been demonstrated in other diatoms like *Thalassiosira pseudonana* [75] as well as in the heterokont, *Nannochloropsis oceanica* in order to knock out the nitrate reductase activity [76]. In conclusion, CRISPR/cas9 system is a promising technology to generate genome-modified organisms in microalgae. **Table 1** compares this system with the other nuclease systems cited above in terms of their technical characteristics and highlights their advantages and disadvantages.

3.2. Mutant libraries

The study of mutants impaired in a glycosidase or a glycosyltransferase implied in the *N*-glycan pathway is of great interest. Indeed, the synthesis of oligosaccharides is a sequential process. Inactivation of an enzyme usually results in the accumulation of its *N*-glycan substrate which enables the step-by-step dissection of the entire pathway. Moreover, mutant phenotyping of the glycosylation pathway allows to investigate to which extent the protein *N*-glycan processing is required for normal growth and development. An indexed and mapped mutant library has been created in *C. reinhardtii* by single random insertional mutagenesis of gene cassettes in 2016 [79]. This library already envisioned to study the function of genes encoding putative glycosyltransferases, glycosidases or even putative translocators in microalgae and to confirm their physiological role from reverse genetic studies.

4. Conclusion

The production of biopharmaceuticals in microalgae currently requires a better understanding of the *N*-glycosylation pathway mechanism and regulation. Such information can be gained by the use of mutant libraries like the one recently developed for *C. reinhardtii*. Indeed, characterization of each individual mutants will allow an understanding of a specific step of the *N*-glycan processing, and mutant cells could represent interesting cell lines for the production of biopharmaceuticals bearing a chosen *N*-glycan profiling.

Once these pathways would be completely deciphered in the microalgae model intended to be used for the production of biopharmaceuticals, the humanization of the *N*-glycosylation pathway could be initiated using designed engineered nucleases strategies recently developed in microalgae. We can now consider that transformed microalgae by these innovative new genomic tools will constitute in a near future one of the most suitable green cell factories for the production of humanized biopharmaceuticals.

Acknowledgements

The authors are indebted to all coworkers and students at the Glyco-MEV laboratory who are currently contributing to the microalgae research project or did so in the past. They are also

thankful to the University of Rouen Normandy, the region Normandie and the I.U.F. for their financial support.

Conflict of interest

The authors have declared no conflict of interest.

Author details

Rodolphe Dumontier¹, Alain Mareck¹, Narimane Mati-Baouche¹, Patrice Lerouge¹ and Muriel Bardor^{1,2*}

*Address all correspondence to: muriel.bardor@univ-rouen.fr

1 Normandie Univ, UniRouen, Laboratoire de Glycobiologie et Matrice Extracellulaire Végétale, Rouen, France

2 I.U.F. (Institut Universitaire de France), Paris Cedex 05, France

References

- [1] Sasso S, Pohnert G, Lohr M, Mittag M, Hertweck C. Microalgae in the postgenomic era: A blooming reservoir for new natural products. *FEMS Microbiology Reviews*. Jul 1, 2012;**36**(4):761-785
- [2] Spolaore P, Joannis-Cassan C, Duran E, Isambert A. Commercial applications of microalgae. *Journal of Bioscience and Bioengineering*. Feb 1, 2006;**101**(2):87-96
- [3] Mata TM, Martins AA, Caetano NS. Microalgae for biodiesel production and other applications: A review. *Renewable and Sustainable Energy Reviews*. Jan 1, 2010;**14**(1):217-232
- [4] Cadoret J-P, Garnier M, Saint-Jean B. Microalgae, functional genomics and biotechnology. *Advances in Botanical Research*. Jan 1, 2012;**64**:285-341
- [5] León-Bañares R, González-Ballester D, Galván A, Fernández E. Transgenic microalgae as green cell-factories. *Trends in Biotechnology*. Jan 1, 2004;**22**(1):45-52
- [6] Barrera DJ, Mayfield SP. High-value recombinant protein production in microalgae. In: Emeritus ARPD, Hu Q, editors. *Handbook of Microalgal Culture* [Internet]. Oxford, UK: John Wiley & Sons, Ltd; 2013. pp. 532-544. Available from: <http://onlinelibrary.wiley.com/doi/10.1002/9781118567166.ch27/summary>
- [7] Hempel F, Maier UG. Microalgae as solar-powered protein factories. *Advances in Experimental Medicine and Biology*. 2016;**896**:241-262

- [8] Rosales-Mendoza S. Algae-Made Antibodies and Immunotoxins. In: Algae-Based Biopharmaceuticals [Internet]. Springer, Cham; 2016 [cited 2017 Nov 17]. pp. 77-93. Available from: https://link.springer.com/chapter/10.1007/978-3-319-32232-2_5
- [9] Tran M, Zhou B, Pettersson PL, Gonzalez MJ, Mayfield SP. Synthesis and assembly of a full-length human monoclonal antibody in algal chloroplasts. *Biotechnology and Bioengineering*. Nov 1, 2009;**104**(4):663-673
- [10] Tran M, Henry RE, Siefker D, Van C, Newkirk G, Kim J, et al. Production of anti-cancer immunotoxins in algae: Ribosome inactivating proteins as fusion partners. *Biotechnology and Bioengineering*. Nov 1, 2013;**110**(11):2826-2835
- [11] Tran M, Van C, Barrera DJ, Pettersson PL, Peinado CD, Bui J, et al. Production of unique immunotoxin cancer therapeutics in algal chloroplasts. *Proceedings of the National Academy of Sciences of the United States of America*. Jan 2, 2013;**110**(1):E15-E22
- [12] Hempel F, Lau J, Klingl A, Maier UG. Algae as protein factories: Expression of a human antibody and the respective antigen in the diatom *Phaeodactylum tricornutum*. *PLoS One*. Dec 2, 2011;**6**(12):e28424
- [13] Hempel F, Maier UG. An engineered diatom acting like a plasma cell secreting human IgG antibodies with high efficiency. *Microbial Cell Factories*. Sep 13, 2012;**11**:126
- [14] Hempel F, Maurer M, Brockmann B, Mayer C, Biedenkopf N, Kelterbaum A, et al. From hybridomas to a robust microalgal-based production platform: Molecular design of a diatom secreting monoclonal antibodies directed against the Marburg virus nucleoprotein. *Microbial Cell Factories*. Jul 27, 2017;**16**:131
- [15] Vanier G, Hempel F, Chan P, Rodamer M, Vaudry D, Maier UG, et al. Biochemical characterization of human anti-hepatitis B monoclonal antibody produced in the microalgae *Phaeodactylum tricornutum*. *PLoS One*. 2015;**10**(10):e0139282
- [16] Vanier G, Stelter S, Vanier J, Hempel F, Maier UG, Lerouge P, et al. Alga-made anti-hepatitis B antibody binds to human Fc γ receptors. *Biotechnology Journal*. 1700496. DOI: 10.1002/biot.201700496
- [17] O'Flaherty R, Trbojević-Akmačić I, Greville G, Rudd PM, Lauc G. The sweet spot for biologics: Recent advances in characterization of biotherapeutic glycoproteins. *Expert Review of Proteomics*. 2018;**15**(1):13-29
- [18] Lingg N, Zhang P, Song Z, Bardor M. The sweet tooth of biopharmaceuticals: Importance of recombinant protein glycosylation analysis. *Biotechnology Journal*. Dec 1, 2012;**7**(12):1462-1472
- [19] Brorson K, Jia AY. Therapeutic monoclonal antibodies and consistent ends: Terminal heterogeneity, detection, and impact on quality. *Current Opinion in Biotechnology*. Dec, 2014;**30**:140-146
- [20] Beyer B, Schuster M, Jungbauer A, Lingg N. Microheterogeneity of recombinant antibodies: Analytics and functional impact. *Biotechnol J*. 2018 Jan;**13**(1). DOI: 10.1002/biot.201700476. PMID: 28862393

- [21] van Beers MMC, Bardor M. Minimizing immunogenicity of biopharmaceuticals by controlling critical quality attributes of proteins. *Biotechnology Journal*. 2012 Dec;7(12):1473-1484
- [22] Burda P, Aebi M. The dolichol pathway of N-linked glycosylation. *Biochimica et Biophysica Acta (BBA)—General Subjects*. Jan 6, 1999;1426(2):239-257
- [23] Weerapana E, Imperiali B. Asparagine-linked protein glycosylation: From eukaryotic to prokaryotic systems. *Glycobiology*. Jun 1, 2006;16(6):91R-101R
- [24] Gil G-C, Velandar WH, Van Cott KE. N-glycosylation microheterogeneity and site occupancy of an Asn-X-Cys sequon in plasma-derived and recombinant protein C. *Proteomics*. May, 2009;9(9):2555-2567
- [25] Zielinska DF, Gnad F, Wiśniewski JR, Mann M. Precision mapping of an in vivo N-Glycoproteome reveals rigid topological and sequence constraints. *Cell*. May 28, 2010;141(5):897-907
- [26] Matsui T, Takita E, Sato T, Kinjo S, Aizawa M, Sugiura Y, et al. N-glycosylation at non-canonical Asn-X-Cys sequences in plant cells. *Glycobiology*. Aug 1, 2011;21(8):994-999
- [27] Williams DB. Beyond lectins: The calnexin/calreticulin chaperone system of the endoplasmic reticulum. *Journal of Cell Science*. Feb 15, 2006;119(4):615-623
- [28] Helenius A, Aebi M. Intracellular functions of N-linked glycans. *Science*. Mar 23, 2001;291(5512):2364-2369
- [29] Varki A. Loss of N-glycolylneuraminic acid in humans: Mechanisms, consequences, and implications for hominid evolution. *American Journal of Physical Anthropology*. Jan 1, 2001;116(S33):54-69
- [30] Lerouge P, Cabanes-Macheteau M, Rayon C, Fischette-Lainé A-C, Gomord V, Faye L. N-glycoprotein biosynthesis in plants: Recent developments and future trends. *Plant Molecular Biology*. Sep 1, 1998;38(1-2):31-48
- [31] Wilson IBH, Zeleny R, Kolarich D, Staudacher E, Stroop CJM, Kamerling JP, et al. Analysis of Asn-linked glycans from vegetable foodstuffs: Widespread occurrence of Lewis a, core α 1,3-linked fucose and xylose substitutions. *Glycobiology*. Apr 1, 2001;11(4):261-274
- [32] Gagneux P, Varki A. Evolutionary considerations in relating oligosaccharide diversity to biological function. *Glycobiology*. Aug 1, 1999;9(8):747-755
- [33] Varki A, Cummings RD, Esko JD, Freeze HH, Stanley P, Bertozzi CR, et al., editors. *Essentials of Glycobiology* [Internet]. 2nd ed. Cold Spring Harbor (NY): Cold Spring Harbor Laboratory Press; 2009. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK1908/>
- [34] Balshüsemann D, Jaenicke L. The oligosaccharides of the glycoprotein pheromone of *Volvox carteri* f. *nagariensis* Iyengar (Chlorophyceae). *European Journal of Biochemistry*. Aug 1, 1990;192(1):231-237

- [35] Grunow A, Becker B, Melkonian M. Isolation and characterization of the Golgi apparatus of a flagellate scaly green alga. *European Journal of Cell Biology*. Jun, 1993;**61**(1):10-20
- [36] Gödel S, Becker B, Melkonian M. Flagellar membrane proteins of *Tetraselmis striata* butcher (Chlorophyta). *Protist*. Aug, 2000;**151**(2):147-159
- [37] Levy-Ontman O, Arad SM, Harvey DJ, Parsons TB, Fairbanks A, Tekoah Y. Unique N-glycan moieties of the 66-kDa cell wall glycoprotein from the red microalga *Porphyridium* sp. *The Journal of Biological Chemistry*. Jun 17, 2011;**286**(24):21340-21352
- [38] Levy-Ontman O. N-Glycosylation of the 66-kDa Cell-Wall Glycoprotein of a Red Microalga. 2012 [cited 2017 Nov 17]; Available from: <http://www.intechopen.com/books/glycosylation/n-glycosylation-of-the-66-kda-cell-wall-glycoprotein-of-a-red-microalga>
- [39] Mathieu-Rivet E, Scholz M, Arias C, Dardelle F, Schulze S, Le Mauff F, et al. Exploring the N-glycosylation pathway in *Chlamydomonas reinhardtii* unravels novel complex structures. *Molecular & Cellular Proteomics*. Nov, 2013;**12**(11):3160-3183
- [40] Vanier G, Lucas P-L, Loutelier-Bourhis C, Vanier J, Plasson C, Walet-Balieu M-L, et al. Heterologous expression of the N-acetylglucosaminyltransferase I dictates a reinvestigation of the N-glycosylation pathway in *Chlamydomonas reinhardtii*. *Scientific Reports*. Aug 31, 2017;**7**(1):10156
- [41] Baïet B, Burel C, Saint-Jean B, Louvet R, Menu-Bouaouiche L, Kiefer-Meyer M-C, et al. N-Glycans of *Phaeodactylum tricornutum* diatom and functional characterization of its N-acetylglucosaminyltransferase I enzyme. *The Journal of Biological Chemistry*. Feb 25, 2011;**286**(8):6152-6164
- [42] Schulze S, Urzica E, Reijnders MJMF, van de Geest H, Warris S, Bakker LV, et al. Identification of methylated GnTI-dependent N-glycans in *Botryococcus brauni*. *The New Phytologist*. Sep, 2017;**215**(4):1361-1369
- [43] Casabianca S, Cornetti L, Capellacci S, Vernesi C, Penna A. Genome complexity of harmful microalgae. *Harmful Algae*. Mar 1, 2017;**63**(Supplement C):7-12
- [44] Levy-Ontman O, Fisher M, Shotland Y, Weinstein Y, Tekoah Y, Arad SM. Genes involved in the endoplasmic reticulum N-glycosylation pathway of the red microalga *Porphyridium* sp.: A bioinformatic study. *International Journal of Molecular Sciences*. Feb 7, 2014;**15**(2):2305-2326
- [45] Mathieu-Rivet E, Kiefer-Meyer M-C, Vanier G, Ovide C, Burel C, Lerouge P, et al. Protein N-glycosylation in eukaryotic microalgae and its impact on the production of nuclear expressed biopharmaceuticals. *Frontiers in Plant Science* [Internet]. Jul 28, 2014 [cited 2016 Nov 16]:5. Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4135232/>
- [46] Kallolimath S, Castilho A, Strasser R, Grünwald-Gruber C, Altmann F, Strubl S, et al. Engineering of complex protein sialylation in plants. *Proceedings of the National Academy of Sciences of the United States of America*. Aug 23, 2016;**113**(34):9498-9503

- [47] Hamilton SR, Davidson RC, Sethuraman N, Nett JH, Jiang Y, Rios S, et al. Humanization of yeast to produce complex terminally Sialylated glycoproteins. *Science*. Sep 8, 2006;**313**(5792):1441-1443
- [48] Gügi B, Le Costaouec T, Burel C, Lerouge P, Helbert W, Bardor M. Diatom-specific oligosaccharide and polysaccharide structures help to unravel biosynthetic capabilities in diatoms. *Marine Drugs*. Sep 18, 2015;**13**(9):5993-6018
- [49] Mathieu-Rivet E, Lerouge P, Bardor M. *Chlamydomonas reinhardtii*: Protein glycosylation and production of biopharmaceuticals. In: Hippler M, editor. *Chlamydomonas: Biotechnology and Biomedicine*. Microbiology Monographs. Vol. 31. Cham: Springer; 2017. DOI: https://doi.org/10.1007/978-3-319-66360-9_3
- [50] Galván A, González-Ballester D, Fernández E. Insertional mutagenesis as a tool to study genes/functions in *Chlamydomonas*. *Advances in Experimental Medicine and Biology*. 2007;**616**:77-89
- [51] Doron L, Segal N, Shapira M. Transgene expression in microalgae-from tools to applications. *Frontiers in Plant Science*. 2016;**7**:505
- [52] Daboussi F. Advances in editing microalgae genomes. *Perspectives in Phycology*. 2017;**4**(1):17-23
- [53] Ng I-S, Tan S-I, Kao P-H, Chang Y-K, Chang J-S. Recent developments on genetic engineering of microalgae for biofuels and bio-based chemicals. *Biotechnology Journal*. Oct 2017;**12**(10):1600644. DOI: 10.1002/biot.201600644
- [54] Falciatore A, Casotti R, Leblanc C, Abrescia C, Bowler C. Transformation of nonselectable reporter genes in marine diatoms. *Marine Biotechnology (New York, NY)*. May, 1999;**1**(3):239-251
- [55] Kindle KL. High-frequency nuclear transformation of *Chlamydomonas reinhardtii*. *Proceedings of the National Academy of Sciences*. Jan 2, 1990;**87**(3):1228-1232
- [56] Armbrust EV. Identification of a new gene family expressed during the onset of sexual reproduction in the centric Diatom *Thalassiosira weissflogii*. *Applied and Environmental Microbiology*. Jan 7, 1999;**65**(7):3121-3128
- [57] Takata M, Sasaki MS, Sonoda E, Morrison C, Hashimoto M, Utsumi H, et al. Homologous recombination and non-homologous end-joining pathways of DNA double-strand break repair have overlapping roles in the maintenance of chromosomal integrity in vertebrate cells. *The EMBO Journal*. Sep 15, 1998;**17**(18):5497-5508
- [58] Maeder ML, Gersbach CA. Genome-editing technologies for gene and cell therapy. *Molecular Therapy*. Mar 1, 2016;**24**(3):430-446
- [59] Daboussi F, Leduc S, Maréchal A, Dubois G, Guyot V, Perez-Michaut C, et al. Genome engineering empowers the diatom *Phaeodactylum tricornutum* for biotechnology. *Nature Communications*. May 29, 2014;**5**:3831

- [60] Miller JC, Holmes MC, Wang J, Guschin DY, Lee Y-L, Rupniewski I, et al. An improved zinc-finger nuclease architecture for highly specific genome editing. *Nature Biotechnology*. Jul 1, 2007;**25**(7):nbt1319
- [61] Smith J, Bibikova M, Whitby FG, Reddy AR, Chandrasegaran S, Carroll D. Requirements for double-strand cleavage by chimeric restriction enzymes with zinc finger DNA-recognition domains. *Nucleic Acids Research*. Sep 1, 2000;**28**(17):3361-3369
- [62] Sizova I, Greiner A, Awasthi M, Kateriya S, Hegemann P. Nuclear gene targeting in *Chlamydomonas* using engineered zinc-finger nucleases. *The Plant Journal*. 2013 Mar 1; **73**(5):873-882
- [63] Greiner A, Kelterborn S, Evers H, Kreimer G, Sizova I, Hegemann P. Targeting of Photoreceptor Genes in *Chlamydomonas reinhardtii* via Zinc-finger Nucleases and CRISPR/Cas9. *Plant Cell*. Jan 1, 2017;tpc.00659.2017
- [64] Gabriel R, Lombardo A, Arens A, Miller JC, Genovese P, Kaeppl C, et al. An unbiased genome-wide analysis of zinc-finger nuclease specificity. *Nature Biotechnology*. Aug 7, 2011;**29**(9):nbt.1948
- [65] Christian M, Cermak T, Doyle EL, Schmidt C, Zhang F, Hummel A, et al. Targeting DNA double-strand breaks with TAL effector nucleases. *Genetics*. Oct 1, 2010;**186**(2):757-761
- [66] Kay S, Hahn S, Marois E, Hause G, Bonas U. A bacterial effector acts as a plant transcription factor and induces a cell size regulator. *Science*. Oct 26, 2007;**318**(5850):648-651
- [67] Weyman PD, Beeri K, Lefebvre SC, Rivera J, McCarthy JK, Heuberger AL, et al. Inactivation of *Phaeodactylum tricornutum* urease gene using transcription activator-like effector nuclease-based targeted mutagenesis. *Plant Biotechnology Journal*. May, 2015;**13**(4):460-470
- [68] Fortunato AE, Jaubert M, Enomoto G, Bouly J-P, Raniello R, Thaler M, et al. Diatom phytochromes reveal the existence of far-red light based sensing in the ocean. *Plant Cell*. Jan 1, 2016;tpc.00928.2015
- [69] Liao H-K, Hatanaka F, Araoka T, Reddy P, Wu M-Z, Sui Y, et al. In vivo target gene activation via CRISPR/Cas9-mediated trans-epigenetic modulation. *Cell* [Internet]. Dec 7, 2017 [cited 2017 Dec 9]. Available from: [http://www.cell.com/cell/abstract/S0092-8674\(17\)31247-3](http://www.cell.com/cell/abstract/S0092-8674(17)31247-3)
- [70] Makarova KS, Haft DH, Barrangou R, Brouns SJJ, Charpentier E, Horvath P, et al. Evolution and classification of the CRISPR–Cas systems. *Nature Reviews. Microbiology*. Jun, 2011;**9**(6):467-477
- [71] Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E. A programmable dual-RNA–guided DNA endonuclease in adaptive bacterial immunity. *Science*. Aug 17, 2012;**337**(6096):816-821
- [72] Jiang W, Brueggeman AJ, Horken KM, Plucinak TM, Weeks DP. Successful transient expression of Cas9 and single guide RNA genes in *Chlamydomonas reinhardtii*. *Eukaryotic Cell*. Jan 11, 2014;**13**(11):1465-1469

- [73] Shin S-E, Lim J-M, Koh HG, Kim EK, Kang NK, Jeon S, et al. CRISPR/Cas9-induced knockout and knock-in mutations in *Chlamydomonas reinhardtii*. Scientific Reports. Jun 13, 2016;**6**:27810
- [74] Nymark M, Sharma AK, Sparstad T, Bones AM, Winge P. A CRISPR/Cas9 system adapted for gene editing in marine algae. Scientific Reports. Apr 25, 2016;**6**:24951
- [75] Hopes A, Nekrasov V, Kamoun S, Mock T. Editing of the urease gene by CRISPR-Cas in the diatom *Thalassiosira pseudonana*. Plant Methods. Nov 24, 2016;**12**:49
- [76] Wang Q, Lu Y, Xin Y, Wei L, Huang S, Xu J. Genome editing of model oleaginous microalgae *Nannochloropsis* spp. by CRISPR/Cas9. The Plant Journal. Dec 1, 2016;**88**(6):1071-1081
- [77] Chevalier BS, Stoddard BL. Homing endonucleases: Structural and functional insight into the catalysts of intron/intein mobility. Nucleic Acids Research. Sep 15, 2001;**29**(18): 3757-3774
- [78] Joung JK, Sander JD. TALENs: A widely applicable technology for targeted genome editing. Nature Reviews. Molecular Cell Biology. 2013 Jan;**14**(1):49-55
- [79] Li X, Zhang R, Patena W, Gang SS, Blum SR, Ivanova N, et al. An indexed, mapped mutant library enables reverse genetics studies of biological processes in *Chlamydomonas reinhardtii*. The Plant Cell. 2016 Feb;**28**(2):367-387

