

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

5,300

Open access books available

130,000

International authors and editors

155M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Biomarkers of Common Childhood Renal Diseases

Samuel N. Uwaezuoke

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.74016>

Abstract

Novel biomarkers are now used in the diagnostic and prognostic evaluation of common kidney diseases in children. The increased scientific interest in these biomarkers is partly due to the remarkable progress in their discovery techniques, and their validation in clinical subjects. However, the wide variation in their sensitivity and specificity is still a major concern. In the identification of biomarkers of kidney injury, an ideal biomarker should be produced after organ injury in concentrations which directly correlates with the degree of injury; should be easily measured in body fluids; and should serve as a potential tool to monitor therapeutic response which is predicated upon a post-injury decrease in its concentration. This book chapter aims to highlight and discuss the novel biomarkers used in the diagnostic and prognostic evaluation of common acute diseases of the kidney in children, such as urinary tract infection (UTI) and acute kidney injury (AKI), as well as chronic kidney disease (CKD) secondary to idiopathic nephrotic syndrome (INS) and diabetic nephropathy (DN).

Keywords: acute diseases, chronic diseases, kidney, children, biomarkers

1. Introduction

Novel biomarkers are now utilized in the diagnostic and prognostic evaluation of common kidney diseases in children [1–3]. Unlike in developing countries where they are scarcely used in medical diagnostics, they are currently used as site-of-care and laboratory tools for disease evaluation in developed settings although some of them have not been fully validated for routine clinical use. The increased scientific interest in novel biomarkers is nevertheless due to the remarkable progress in their discovery techniques, their validation in clinical populations and their utility as diagnostic tests. Despite the reliability of the conventional diagnostic methods for some of these diseases, their limitations as diagnostic tools have underscored the

need for a paradigm shift to biomarkers. The attraction of these novel biomarkers lies in the non-invasive nature of their use in disease evaluation. However, the wide variation in their sensitivity and specificity is still a major challenge.

According to the definition by the *National Institute of Health Biomarkers Definitions Working Group*, a biomarker refers to 'a characteristic that is objectively measured and evaluated as an indicator of normal biological processes, pathogenic processes, or pharmacologic responses to a therapeutic intervention' [4, 5]. In other words, it is a measurable indicator of a biological state or condition. Although there are diverse perspectives to the definition of a biomarker, for the purpose of the current topic, a biomarker can be defined as a substance whose detection indicates a particular disease-state, or a change in expression/state of a protein which correlates with the risk/progression of a disease, or with disease susceptibility to a given therapy. In fact, the most actively studied biomarkers presently are the biological macromolecules. For instance, proteins are most easily measured by immunoassays or by mass spectrometry combined with high-resolution separation techniques, while nucleic acids can be amplified by the polymerase chain reaction (PCR) and related techniques.

In the identification of biomarkers of kidney injury, an ideal biomarker should be produced after organ injury in concentrations which directly correlate with the degree of injury; should be easily measured in body fluids; and should serve as a potential tool to monitor therapeutic response, predicated upon a post-injury decrease in its concentration [6].

In this book chapter, novel biomarkers which essentially meet these requirements with respect to evaluating common kidney diseases in children are highlighted and discussed.

2. Acute diseases of the kidney and biomarkers

Biomarkers have been identified for the following acute diseases of the kidney: acute pyelonephritis or upper urinary tract infection (UTI) and acute cystitis/urethritis or lower UTI, as well as acute kidney injury (AKI). UTI ranks third in the hierarchy of common childhood infections in developing countries [7], and remains one of the commonest causes of febrile illness in the pediatric age group [8, 9].

2.1. Biomarkers of UTI

Serum procalcitonin (PCT), urine/serum interleukin 6 (IL-6) and 8 (IL-8), and urine neutrophil gelatinase-associated lipocalin (NGAL) are currently the major biomarkers identified in the evaluation of the disease. Other novel biomarkers include urine interleukin 1-beta (IL-1 β), urine 8-hydroxy-2-deoxyguanosine (8-oxodG), and urine total anti-oxidant capacity (TAC) (**Figure 1**).

PCT is one of the inflammatory markers recognized as a biomarker of severe bacterial infection [10]. It is essentially a peptide precursor of the hormone- calcitonin - produced by the thyroid parafollicular cells, as well as the neuroendocrine cells of the lungs and intestines. Its serum level in healthy subjects is below the limit of detection (0.01 $\mu\text{g/L}$) of laboratory assays [11]. Notably, serum PCT level is not elevated significantly with viral or non-infectious inflammatory process, but may rise up to 100 $\mu\text{g/L}$ in systemic inflammatory response from severe bacterial infection. In fact, estimation of PCT can be used as a marker of severe bacterial




Figure 1. Schematic representation of the list of biomarkers used in the three-fold evaluation of urinary tract infection (UTI).

sepsis, as its levels correlate with the severity of sepsis [12]. In comparison with other acute phase reactants such as IL-2, IL-6, IL-8, C-reactive protein (CRP) and tumor necrosis factor- α (TNF- α), PCT has a high sensitivity (85%) and specificity (91%) for discriminating between patients with systemic inflammatory response syndrome (SIRS) and those with sepsis [13]. It is now well established that serum PCT has a good diagnostic accuracy for the diagnosis of acute pyelonephritis than for lower UTI, with reported high sensitivity and specificity values [14–16]. For instance, sensitivity and specificity values of 94.1 and 89.7% respectively, and sensitivity and specificity values of 83.3 and 93.7% respectively, have been documented [15, 16]. PCT also has a good predictive ability for vesico-ureteric reflux (VUR), as high levels of this biomarker are significantly related to high-grade VUR [17–19]. Again, children with VUR reportedly presented with significantly higher median PCT levels than those without VUR; sensitivity and negative predictive values of 94.3 and 95.4% respectively for predicting high-grade VUR (using the biomarker alone), as well as 97.1 and 97.8% respectively (using a combination of the biomarker and ultrasound studies) have been recorded [18]. Finally, PCT can also serve as an early predictor of renal parenchymal damage in children with UTI [20]. The implication is that the prognostic utility of PCT may in future preclude the need for dimer-captosuccinic acid (DMSA) scan, or better still, may serve as an adjunct to this imaging study.

Similarly, other biomarkers such as urine 8-hydroxy-2'-deoxyguanosine (8-oxodG) and total antioxidant capacity (TAC) are useful in the prediction of renal parenchymal injury in children with UTI [21]. Interestingly, those patients with positive DMSA scan had higher levels of

urine 8-oxodG and higher urine TAC than the patients with normal DMSA scan: showing that urine levels of 8-oxodG may directly correlate with renal parenchymal injury.

Urine NGAL is another biomarker for the diagnostic evaluation of UTI. NGAL (also called lipocalin-2) is a protein involved in innate immunity as it is involved in the sequestration of iron which ultimately blocks bacterial growth [22]. It is expressed mainly in neutrophils and in low concentrations in the kidney, prostate, as well as respiratory and gastrointestinal epithelia [23]. NGAL is currently employed as a biomarker of renal injury [24], given that it is secreted in high concentrations into the blood and urine within 2 hours of AKI [25]. In addition, the biomarker is easily excreted and detected in the urine because it is protease-resistant. Thus, urine NGAL appears to be a specific and sensitive biomarker for the diagnosis of UTI and for the prediction of renal parenchymal injury in the disease [26, 27].

Finally, some interleukins also play a role in disease diagnosis and in differentiating upper from lower UTI. Generally, the infection stimulates both local and systemic cytokine responses [28, 29]. For instance, *Escherichia coli* is known to activate a cytokine response in the uroepithelial cell lines to produce IL-6 and IL-8, and in peripheral blood monocytes to produce interleukin 1-alpha (IL-1 α), IL-1 β , IL-8, IL-6, and TNF- α [28]. Adherence of this bacterium on the mucosal sites of the urinary tract triggers inflammation, including a mucosal cytokine-response which results in the secretion of IL-6 and IL-8 by activated uroepithelial cells [29, 30]. Thus, since IL-6 and IL-8 are expressed early after a UTI episode and are secreted by uroepithelial cells, these cytokines have clearly fulfilled some of the five characteristics of an ideal biomarker outlined in the introductory section. Reports indeed suggest that urine IL-6 and IL-8 are proven biomarkers for UTI in children [31]. The serum and urine levels of IL-6 were more sensitive and specific for upper UTI than levels of IL-8 [32]. Sensitivity and specificity values for serum IL-6 were noted to be 88 and 83% respectively, while for urine IL-6, sensitivity and specificity values were 86 and 81%, respectively [32]. Furthermore, serum IL-6 was able to differentiate upper from lower UTI with a sensitivity of 88% and specificity of 74%, while serum IL-1 β was known to have a sensitivity of 97% and a specificity of 59% for detecting upper UTI [14].

2.2. Biomarkers of AKI

Serum creatinine represents a poor traditional biomarker for AKI due to some limitations. First, its levels are only altered when renal function diminishes by 50% [33]. Second, there are well-known confounders to serum creatinine levels such as muscle size, chronologic age, gender, drugs and state of hydration [34]. Third, a sudden reduction in renal function may not be shown by a rise in serum creatinine until after 24–48 hours. Finally, it provides little information about the underlying cause and nature of kidney injury, and is less accurate for patients with small muscle mass and unusual diets [1]. These draw-backs have resulted in a paradigm shift to novel biomarkers.

Generally, there are two main types of AKI biomarkers: *biomarkers of kidney function* and *biomarkers of kidney injury*. Irrespective of the type of kidney injury and the clinical scenario, an inflammatory response appears to play a significant role in the pathogenesis of AKI [34]. The major triggers of AKI (ischemia, nephrotoxins, and bacterial endotoxins) stimulate the release of inflammatory mediators from renal endothelial and tubular cells. As AKI progresses, a number of causative factors result in the accumulation of biomarkers in plasma and urine, and possibly indicate different pathophysiologic events during the process of kidney

injury and repair [1]. For example, biomarkers like NGAL, interleukin-18 (IL-18), N-acetyl- β -D-glucosaminidase (NAG), kidney injury molecule-1 (KIM-1), accumulate in urine due to an induced tubular epithelial synthesis in different parts of the nephron, or as a consequence of reduced reabsorption of the filtered load in the proximal tubule (NGAL, cystatin C) [34]. Furthermore, production of biomarkers from transmigrated, activated immune cells into the tubular lumen may be contributory (NGAL, IL-18), while increased production of some biomarkers in other tissues (NGAL, IL-18) also occurs, thus raising concerns about their diagnostic value in the disease [35]. Biomarkers of AKI have been classified as follows: *functional markers* (serum cystatin C, urine albumin and NGAL), *up-regulated proteins* (KIM-1, liver-type fatty-acid binding protein [L-FABP], IL-18, β -trace protein and asymmetric dimethyl arginine), *low-molecular weight proteins* (urine cystatin C, NAG, glutathione S-transferase, γ -glutamyl-trans peptidase [γ GT]) and *enzymes* (alanine amino-peptidase and lactate dehydrogenase) [36]. Other markers which have shown diagnostic ability include hepatocyte growth factor (HGF), vasoactive endothelial growth factor (VEGF), interferon gamma-induced protein 10 (IP-10) and total protein [37]. Some of the novel biomarkers are highlighted as below.

First, KIM-1 is a transmembrane tubular glycoprotein which is upregulated approximately 50- to 100-fold in the kidney, and is secreted into the urine after proximal tubular injury. KIM-1 is highly expressed in renal tubules, and is typically seen in areas of fibrosis and inflammation. The results of a meta-analysis indicate that urine KIM-1 represents a promising biomarker for early detection of AKI, with a good predictive value, especially in cardiac surgery patients [38]. Furthermore, in comparison to other biomarkers used as indicators of drug toxicity, KIM-1 significantly performed better than serum creatinine and blood urea nitrogen at detecting renal tubular injury in murine models; this makes it a useful marker for determining drug toxicity [39].

Second, NAG (another biomarker of kidney injury) is a large (>130 kDa) lysosomal enzyme which is located in several human cells including the renal tubule [34]. Its size hinders glomerular filtration, and elevated urine levels are thus presumed to emanate from the tubules, indicating tubular injury. In fact, during active renal disease, urine NAG activity is increased [40]. This feature makes it a potential and sensitive biomarker of AKI.

Third, cystatin C is a low-molecular-weight protein produced by all nucleated cells in the body at a constant rate, freely filtered by the glomeruli but completely reabsorbed and catabolized by the renal tubule [41]. Thus, its elevated urine levels are also seen in a tubulopathy because of the reduced re-absorptive capacity of the proximal tubules: making it a non-specific biomarker of AKI [42]. However, urine cystatin C could correctly predict the need for dialysis in intensive care unit (ICU) patients with established AKI [43]; could be an early predictor of AKI in children and in pediatric RIFLE classification, as well as a predictor of reduced estimated GFR (eGFR) after cardiac surgery [44]; while serum cystatin C could be used alone or in combination with serum creatinine and eGFR for early and accurate diagnosis of AKI in patients at emergency settings [45].

Finally, NGAL (which plays a role in the evaluation of UTI as previously mentioned) is a universal iron-transporter protein expressed in the tubular epithelium of the distal nephron and released into the blood and urine following tubular injury. This biomarker was first identified as a 25 kDa protein in the secondary granules of human neutrophils which is released into the bloodstream in response to bacterial infection. Interestingly, its elevated level in the urine may be diagnostic of AKI using the Acute Kidney Injury Network (AKIN) criteria: although

with a moderate predictive value [46, 47]. In addition, its combination with another urine biomarker such as L-FABP resulted in early detection of AKI after cardiac surgery in a sample of adult patients before a rise in serum creatinine was noted [48]. Similarly, in murine models of ischemic and toxic AKI, NGAL was identified as one of the most speedily-induced proteins; its level was elevated by several folds in both serum and urine within hours of the insult [49].

3. Chronic kidney disease (CKD) and biomarkers

Several novel biomarkers have also been identified in the evaluation of childhood CKD: especially CKD secondary to idiopathic nephrotic syndrome and diabetic nephropathy (DN) [2, 50]. Remarkably, an overlap exists between some of the biomarkers of CKD and those of AKI and acute pyelonephritis. For instance, urine NAG, NGAL, cystatin C and L-FABP all play a role in the evaluation of both CKD and these acute kidney diseases. Similar to the grouping of biomarkers of AKI, biomarkers of CKD can be broadly classified into *biomarkers of kidney function* and *biomarkers of kidney damage*.

GFR is the most important marker of kidney function in CKD, although it is poorly estimated in most clinical and research settings. Thus, equations for its estimation are predicated upon filtration biomarkers such as serum creatinine and serum cystatin C [51]. However, equations for eGFR based on cystatin C appear more reliable because this biomarker is not affected by muscle mass and gives a better representation of GFR, in addition to having a more stable rate of production compared to creatinine [52].

Although albuminuria represents the traditional biomarker of kidney damage, it may only be present after significant damage has occurred, or may be absent in other types of kidney damage such as tubulointerstitial disease and hypertensive kidney disease. Therefore, there is now a paradigm shift to novel biomarkers which could identify patients with CKD early enough so that prompt interventions could slow down the progression of the disease.

3.1. Biomarkers of diabetic nephropathy (DN) and other causes of CKD

The major pathogenic components of DN consist of renal fibrosis, mesangial expansion, glomerular hypertrophy, oxidative stress and tubular inflammation [53]. Myriad novel biomarkers of DN have now been identified. An attempt to categorize them has also been made [50]. *Glomerular biomarkers* include transferrin, immunoglobulin G (IgG), ceruloplasmin, type IV collagen, laminin, glycosaminoglycans (GAGs), lipocalin-type prostaglandin D synthase (L-PGDS), fibronectin, podocytes-podocalyxin, and VEGF. *Tubular biomarkers* include NGAL, α -1-microglobulin, KIM-1, NAG, cystatin C, and L-FABP. *Biomarkers of inflammation* comprise TNF- α , IL-1 β , IL-18, IP-10, monocyte chemoattractant protein 1 (MCP-1), granulocyte colony-stimulating factor (G-CSF), eotaxins, RANTES (regulated on activation, normal T cell expressed and secreted) or CCL-5, and orosomucoid. A typical example of *biomarkers of oxidative stress* is 8oHdG, while *miscellaneous biomarkers* include some tubular markers such as urine heart fatty acid-binding protein and urine retinol-binding protein 4 (RBP4), and podocyte biomarkers such as podocalyxin, nephrin, and VEGF, and urine advanced glycation end products (AGEs). Notably, these podocyte biomarkers are also regarded as glomerular markers. Some of the biomarkers of DN are shown in **Table 1**, and will be briefly discussed below.

Biomarkers	Clinical significance	Comments
• Kidney injury molecule-1	• Diabetic nephropathy [‡]	• Predicts CKD progression to ESKD
• Neutrophil gelatinase-associated lipocalin	• Ig A nephropathy • Polycystic kidney disease	• Predicts CKD progression from early stages to ESKD
• N-acetyl-β-D-glucosaminidase	• Diabetic nephropathy ^{‡‡} • Idiopathic glomerulonephritis (MN, FSGS, and MCN)	• Predicts early and late stages of CKD in type 1 diabetics, and early CKD in type 2 diabetics • Predicts proteinuria-induced tubular damage in early stage of disease
• Liver fatty acid-binding protein	• Diabetic nephropathy ^{‡‡}	• Predicts early CKD and progression to ESKD in type 1 diabetics and early CKD in type 2 diabetics
• Cystatin C	• Diabetic nephropathy [‡]	• Predicts early CKD in type 2 diabetics
• α-1-microglobulin	• Diabetic nephropathy [‡]	• Predicts early CKD in type 2 diabetics
• Monocyte chemoattractant protein-1 (MCP-1)	• Diabetic nephropathy [‡]	• Predicts CKD and cardiovascular disease risk in type 2 diabetics
• Interleukin-18	• Lupus nephritis • Diabetic nephropathy [‡]	• Elevated urine levels • Same as in MCP-1
• Retinol-binding protein 4	• Tubulointerstitial diseases	• Elevated urine levels

MN, membranous nephropathy; FSGS, focal segmental glomerulosclerosis; MCN, minimal change nephropathy.

[‡]Among type 1 diabetics.

^{‡‡}Among type 2 diabetics.

Table 1. Novel biomarkers of chronic kidney diseases.

NAG has the ability to predict the onset and progression of CKD in diabetes mellitus. For instance, baseline urine levels of this tubular biomarker can independently predict microalbuminuria and macroalbuminuria in type 1 diabetes mellitus [54]. Furthermore, it is a sensitive biomarker for the detection of early renal damage in type 2 diabetes mellitus [55], while elevated urine levels can precede microalbuminuria in type 1 diabetes mellitus [56].

Furthermore, high levels of KIM-1 have been observed in DN: a disease characterized by renal fibrosis and tubular inflammation, among other components that have been previously

mentioned. For instance, majority of type 1 diabetic subjects with DN (stage 1 to 3 CKD) who had higher plasma KIM-1 levels reportedly ended up with end-stage kidney disease (ESKD); only a few of their counterparts who had lower plasma KIM-1 levels developed ESKD [57]. Baseline plasma KIM-1 levels also correlated with the rate of eGFR decline after adjusting for baseline urine albumin-to-creatinine ratio, eGFR, and glycated hemoglobin (Hb1Ac).

Another biomarker of DN is L-FABP, whose baseline urine levels in newly-diagnosed type 1 diabetics not only predicted the development of microalbuminuria but also the progression of microalbuminuria to macroalbuminuria [58]. In type 2 diabetes mellitus, elevated urine level of this biomarker also plays a role in predicting early CKD [59], and also serves as an independent predictor of CKD progression in type 1 diabetics [60]. Thus, L-FABP has the advantage of predicting kidney injury before albuminuria, especially in type 1 diabetics. Serum and urine cystatin C levels are useful biomarkers for early prediction of nephropathy in type 2 diabetes mellitus [61], and for progression of diabetic kidney disease [62]. Furthermore, α -1-microglobulin has been identified as an inexpensive biomarker for the early prediction of DN [63]. Elevated urine levels occur in patients with normoalbuminuric type 2 diabetes mellitus: preceding the onset of microalbuminuria and confirming this tubular biomarker as a more sensitive urine biomarker of CKD [64].

Expression of MCP-1 is upregulated in kidney diseases which present with a continued inflammatory response, such as in DN and lupus nephritis. Some reports indeed indicate that elevated levels of urine MCP-1 were observed in DN [65], as well as in active lupus nephritis [66]. Serum and urine levels of IL-18 were positively correlated with albumin excretion rate, whereas its serum levels were positively correlated with the development of carotid intima-media thickness in type 2 diabetics, and may therefore be a predictor of DN progression and cardiovascular diseases [67].

Finally, urine RBP4 is elevated in patients with tubulointerstitial disease, and may constitute a risk factor for long-term allograft loss, independent of the histology of renal biopsy, as well as for albuminuria [68].

3.2. Biomarkers of idiopathic nephrotic syndrome

The use of biomarkers in childhood idiopathic nephrotic syndrome has been well documented [2]. It represents a non-invasive approach in diagnostic nephrology, as these markers can be used in the prediction and prognostic evaluation of the disease, as well as in differentiating steroid-resistant nephrotic syndrome (SRNS) from steroid-sensitive nephrotic syndrome (SSNS). **Table 2** summarizes the list of identified biomarkers reported for childhood idiopathic nephrotic syndrome. Adiponectin (ADPN) – one of the adipokines, neopterin, β 2-microglobulin, and NAG were reported to be diagnostic markers [69–71]. In addition to neopterin and NAG, urine vitamin D-binding protein (u VDBP) and α 1 β -glycoprotein were able to differentiate SRNS from SSNS [72, 73] while NAG and β 2-microglobulin could also predict steroid responsiveness and renal outcome in SRNS [74]. Some of these biomarkers are further discussed as follows. First, elevated serum ADPN levels were documented in SRNS patients in relapse compared to those in remission [69]. Specifically, strong positive correlations were observed between serum ADPN levels and lipid parameters/proteinuria, whereas negative correlations were noted between ADPN levels and serum protein/albumin levels. Second, serum neopterin levels were found to be significantly elevated among SSNS and

Novel biomarkers (body fluids)	Reported role of biomarkers (references)
• Adiponectin (serum)	• Diagnostic* (Bakkalöglu et al. [69])
• Neopterin (serum)	• Diagnostic** (Bakr et al. [70]) • Discriminatory† (Bakr et al. [70])
• Vitamin D binding protein (urine)	• Discriminatory† (Bennett et al. [72])
• N-acetyl-beta-D glucosaminidase (urine)	• Discriminatory† (Calişkan et al. [71]) • Prognostic‡ (Fede et al. [74])
• α 1- β glycoprotein (13.8 kDa fragment) (urine)	• Discriminatory† (Piyaphanee et al. [73])
• beta2-microglobulin (urine)	• Prognostic*** (Fede et al. [74])
• N-acetyl-beta-D glucosaminidase (urine)	
• beta2-microglobulin (urine)	• Diagnostic [§] (Calişkan et al. [71])
• N-acetyl-beta-D glucosaminidase (urine)	

*Raised serum levels in steroid-resistant nephrotic syndrome (SRNS) relapse.

**Raised serum levels in primary active nephrotic syndrome.

†Differentiates SRNS from steroid-sensitive nephrotic syndrome (SSNS).

‡Predicts steroid-responsiveness.

***Predicts tubular injury and dysfunction in SRNS.

§Elevated levels in SRNS and SSNS.

Table 2. Role of some novel biomarkers in childhood idiopathic nephrotic syndrome.

SRNS patients with the active disease when compared to those in remission and the controls [70]. The diagnostic utility of neopterin for active idiopathic nephrotic syndrome was thus highlighted, but its poor discriminatory ability for SSNS and SRNS were also noted in the report. Third, uVDBP was reported to have a high discriminatory ability in distinguishing SRNS from SSNS [72]. For instance, levels of uVDBP were significantly higher in patients with SRNS than in patients with SSNS and in the controls. Despite the direct correlation between microalbuminuria and uVDBP, the latter exhibited a higher discriminatory ability for differentiating SRNS from SSNS than the former.

4. Conclusion

Many biomarkers have now been identified for the diagnostic and prognostic evaluation of acute and chronic diseases of the kidney in children. However, more evidence-based studies are still required to validate some of the novel biomarkers. Remarkably, a biomarker-panel comprising several of the markers potentially improves their sensitivity and specificity in disease evaluation. Inequities in the availability and accessibility of the laboratory tools between the developed and developing world still remain a challenge. Biotechnology firms should urgently prioritize the mass production of tools for identifying these biomarkers in order to bridge this gap.

Acknowledgements

The author acknowledges the invaluable information obtained from his previous published works in Refs. [1–3], during the preparation of this book chapter.

Conflict of interest

The author declares no conflict of interest in this work.

Author details

Samuel N. Uwaezuoke

Address all correspondence to: samuel.uwaezuoke@unn.edu.ng

College of Medicine, University of Nigeria/University of Nigeria Teaching Hospital, Enugu, Nigeria

References

- [1] Uwaezuoke SN. Acute kidney injury in children: Enhancing diagnosis with novel biomarkers. *Journal of Acute Disease*. 2016;**5**:267-270
- [2] Uwaezuoke SN. The role of novel biomarkers in childhood idiopathic nephrotic syndrome: A narrative review of published evidence. *International Journal of Nephrology and Renovascular Disease*. 2017;**10**:123-128
- [3] Uwaezuoke SN. Urinary tract infection in children: Diagnostic and prognostic utility of biomarkers. *Journal of Comprehensive Pediatrics*. 2017;**8**:e59248. DOI: 10.5812/compreped.59248
- [4] Strimbu K, Jorge T. What are biomarkers? *Current Opinion in HIV and AIDS*. 2010;**5**:463-466. DOI: 10.1097/COH.0b013e32833ed177
- [5] Atkinson AJ Jr, Colburn WA, DeGruttola VG, et al. Biomarkers and surrogate endpoints: Preferred definitions and conceptual framework. *Clinical Pharmacology and Therapeutics*. 2001;**69**:89-95. DOI: 10.1067/mcp.2001.113989
- [6] Cruz DN, Goh CY, Haase-Fielitz A, Ronco C, Haase M. Early biomarkers of renal injury. *Congestive Heart Failure*. 2010;**16**:S25-S31. DOI: 10.1111/j.1751-7133.2010.00163.x
- [7] Srivastava RN, Bagga A. Urinary tract infection. In: Srivastava RN, Bagga A, editors. *Pediatric Nephrology*. 4th ed. New Delhi: Jaypee Brothers Medical Publishers Ltd; 2005. pp. 235-264. DOI: 10.5005/jp/books/11308

- [8] Shaikh N, Morone NE, Bost JE, Farrell MH. Prevalence of urinary tract infection in childhood: A meta-analysis. *The Pediatric Infectious Disease Journal*. 2008;**27**(4):302-308. DOI: 10.1097/INF.0b013e31815e4122
- [9] Chang SL, Shortliffe LD. Pediatric urinary tract infections. *Pediatric Clinics of North America*. 2006;**53**:379-400. DOI: 10.1016/j.pcl.2006.02.011
- [10] Assicot M, Gendrel D, Carsin H, Raymond J, Guilbaud J, Bohuon C. High serum procalcitonin concentrations in patients with sepsis and infection. *Lancet*. 1993;**341**:515-518. DOI: 10.1016/0140-6736(93)90277-N
- [11] Dandona P, Nix D, Wilson MF, et al. Procalcitonin increase after endotoxin injection in normal subjects. *Journal of Clinical Endocrinology and Metabolism*. 1994;**79**:1065-1608. DOI: 10.1210/jcem.79.6.7989463
- [12] Meisner M, Tsachikowsky K, Palmaers T, Schmidt J. Comparison of procalcitonin (PCT) and C-reactive protein (CRP) plasma concentrations at different SOFA scores during the course of sepsis and MODS. *Critical Care*. 1999;**3**:45-50. DOI: 10.1186/cc306
- [13] Balci C, Sungurtekin H, Gürses E, Sungurtekin U, Kaptanoğlu B. Usefulness of procalcitonin for diagnosis of sepsis in the intensive care unit. *Critical Care*. 2003;**7**:85-90. DOI: 10.1186/cc1843
- [14] Gürgöze MK, Akarsu S, Yilmaz E, Gödekmerdan A, Akça Z, Çiftçi I, et al. Proinflammatory cytokines and procalcitonin in children with acute pyelonephritis. *Pediatric Nephrology*. 2005;**20**:1445-1448. DOI: 10.1007/s00467-005-1941-6
- [15] Smolkin V, Koren A, Raz R, Colodner R, Sakran W, Halevy R. Procalcitonin as a marker of acute pyelonephritis in infants and children. *Pediatric Nephrology*. 2002;**17**:409-412. PMID: 12107804
- [16] Pecile P, Miorin E, Romanello C, Falletti E, Valent F, Giacomuzzi F, et al. Procalcitonin: A marker of severity of acute pyelonephritis among children. *Pediatrics*. 2004;**114**:e249-e254. DOI: 10.1542/peds.114.2.e249
- [17] Leroy S, Adamsbaum C, Marc E, Moulin F, Raymond J, Gendrel D, et al. Procalcitonin as a predictor of vesicoureteral reflux in children with a first febrile urinary tract infection. *Pediatrics*. 2005;**115**:e706-e709. DOI: 10.1542/peds.2004-1631
- [18] Sun HL, Wu KH, Chen SM, Chao YH, Ku MS, Hung TW, et al. Role of procalcitonin in predicting dilating vesicoureteral reflux in young children hospitalized with a first febrile urinary tract infection. *The Pediatric Infectious Disease Journal*. 2013;**32**:e348-e354. DOI: 10.1097/INF.0b013e3182905d83
- [19] Rahimzadeh N, Otukesh H, Hoseini R, Shadani S, Hooman N. Serum procalcitonin level for prediction of high-grade vesicoureteral reflux in urinary tract infection. *Iranian Journal of Kidney Diseases*. 2014;**8**:105-108
- [20] Barati L, Safaeian B, Mehrjerdian M, Vakili MA. Early prediction of renal parenchymal injury with serum procalcitonin. *Journal of Renal Injury Prevention*. 2016;**5**:108-111. DOI: 10.15171/jrip.2016.23

- [21] Chien JW, Wang LY, Cheng YS, Tsai YG, Liu CS. Urinary 8-hydroxy-2'-deoxyguanosine (8-oxodG) level can predict acute renal damage in young children with urinary tract infection. *Biomarkers*. 2014;**19**:326-331. DOI: 10.3109/1354750X.2014.910552
- [22] Yang J, Goetz D, Li JY, Wang W, Mori K, Setlik D, et al. An iron delivery pathway mediated by a lipocalin. *Molecular Cell*. 2002;**10**:1045-1056. DOI: 10.1016/s1097-2765(02)00710-4
- [23] Cowland JB, Borregaard N. Molecular characterization and pattern of tissue expression of the gene for neutrophil gelatinase-associated lipocalin from humans. *Genomics*. 1997;**45**:17-23. DOI: 10.1006/geno.1997.4896
- [24] Devarajan P. Neutrophil gelatinase-associated lipocalin: A troponin-like biomarker for human acute kidney injury. *Nephrology (Carlton)*. 2010;**15**:419-428. DOI: 10.1111/j.1440-1797.2010.01317.x
- [25] Bennett M, Dent CL, Ma Q, Dastrala S, Grenier F, Workman R, et al. Urine NGAL predicts severity of acute kidney injury after cardiac surgery: A prospective study. *Clinical Journal of the American Society of Nephrology*. 2008;**3**:665-673. DOI: 10.2215/CJN.04010907
- [26] Ghasemi K, Esteghamati M, Borzoo S, Parvaneh E. Predictive accuracy of urinary neutrophil gelatinase associated lipocalin (NGAL) for renal parenchymal involvement in children with acute pyelonephritis. *Electronic Physician*. 2016;**8**:1911-1917. DOI: 10.19082/1911
- [27] Yilmaz A, Sevketoglu E, Gedikbasi A, Karyagar S, Kiyak A, Mulazimoglu M, et al. Early prediction of urinary tract infection with urinary neutrophil gelatinase associated lipocalin. *Pediatric Nephrology*. 2009;**24**:2387-2392. DOI: 10.1007/s00467-009-1279-6
- [28] Agace W, Hedges S, Andersson U, Andersson J, Ceska M, Svanborg C. Selective cytokine production by epithelial cells following exposure to *Escherichia coli*. *Infection and Immunity*. 1993;**61**:602-609. PMID: PMC302770
- [29] Agace W, Hedges S, Ceska M, Svanborg C. Interleukin-8 and the neutrophil response to mucosal gram-negative infection. *The Journal of Clinical Investigation*. 1993;**92**:780-785. DOI: 10.1172/JCI116650
- [30] Svanborg C, Agace W, Hedges S, Lindstedt R, Svensson ML. Bacterial adherence and mucosal cytokine production. *Annals of the New York Academy of Sciences*. 1994;**730**:162-181. DOI: 10.1111/j.1749-6632.1994.tb44247.x
- [31] Jantusch BA, O'Donnell R, Wiedermann BL. Urinary interleukin-6 and interleukin-8 in children with urinary tract infection. *Pediatric Nephrology*. 2000;**15**:236-240. DOI: 10.1007/s004670000456
- [32] Sheu JN, Chen MC, Lue KH, Cheng SL, Lee IC, Chen SM, et al. Serum and urine levels of interleukin-6 and interleukin-8 in children with acute pyelonephritis. *Cytokine*. 2006;**36**:276-282. DOI: 10.1016/j.cyto.2007.02.006
- [33] Coca SG, Yalavarthy R, Concato J, Parikh CR. Biomarkers for the diagnosis and risk stratification of acute kidney injury: A systematic review. *Kidney International*. 2008;**73**:1008-1016. DOI: 10.1038/sj.ki.5002729

- [34] Mårtensson J, Martling CR, Bell M. Novel biomarkers of acute kidney injury and failure: Clinical applicability. *British Journal of Anaesthesia*. 2012;**109**:843-850. DOI: 10.1093/bja/aes357
- [35] Mårtensson J, Bellomo R. The rise and fall of NGAL in acute kidney injury. *Blood Purification*. 2014;**37**:304-310. DOI: 10.1159/000364937
- [36] de Geus HR, Betjes MG, Bakker J. Biomarkers for the prediction of acute kidney injury: A narrative review on current status and future challenges. *Clinical Kidney Journal*. 2012;**5**:102-108. DOI: 10.1093/ckj/sfs008
- [37] Vanmassenhove J, Vanholder R, Nagler E, Van Biesen W. Urinary and serum biomarkers for the diagnosis of acute kidney injury: In-depth review of the literature. *Nephrology, Dialysis, Transplantation*. 2013;**28**:254-273. DOI: 10.1093/ndt/gfs380
- [38] Shao X, Tian L, Xu W, Zhang Z, Wang C, Qi C, et al. Diagnostic value of urinary kidney injury molecule 1 for acute kidney injury: A meta-analysis. *PLoS One*. 2014;**9**:e84131. DOI: 10.1371/journal.pone.0084131
- [39] Bonventre JV, Vaidya VS, Schmouder R, Feig P, Dieterle F. Next generation biomarkers for detecting kidney toxicity. *Nature Biotechnology*. 2010;**28**:436-440. DOI: 10.1038/nbt0510-436
- [40] Endre ZH, Pickering JW, Walker RJ, Devarajan P, Edelstein CL, Bonventre JV, et al. Improved performance of urinary biomarkers of acute kidney injury in the critically ill by stratification for injury duration and baseline renal function. *Kidney International*. 2011;**79**:1119-1130. DOI: 10.1038/ki.2010.555
- [41] Srivastava RN, Bagga A. Evaluation of renal function. In: Srivastava RN, Bagga A, editors. *Pediatric Nephrology*. 4th ed. New Delhi: Jaypee Brothers Medical Publishers Ltd; 2005. pp. 20-29. DOI: 10.5005/jp/books/11308
- [42] Ghys L, Paepe D, Smets P, Lefebvre H, Delanghe J, Daminet S. Cystatin C: A new renal marker and its potential use in small animal medicine. *Journal of Veterinary Internal Medicine*. 2014;**28**(4):1152-1164. DOI: 10.1111/jvim.12366
- [43] Nejat M, Pickering JW, Walker RJ, Westhuyzen J, Shaw GM, Frampton CM, et al. Urinary cystatin C is diagnostic of acute kidney injury and sepsis, and predicts mortality in the intensive care unit. *Critical Care*. 2010;**14**:R85. DOI: 10.1186/cc9014
- [44] Hassinger AB, Backer CL, Lane JC, Haymond S, Wang D, Wald EL, et al. Predictive power of serum cystatin C to detect acute kidney injury and pediatric-modified RIFLE class in children undergoing cardiac surgery. *Pediatric Critical Care Medicine*. 2012;**13**:435-440. DOI: 10.1097/PCC.0b013e318238b43c
- [45] Bongiovanni C, Magrini L, Salerno G, Gori CS, Cardelli P, Hur M, et al. Serum cystatin C for the diagnosis of acute kidney injury in patients admitted in the emergency department. *Disease Markers*. 2015;**2015**:416059. DOI: 10.1155/2015/416059
- [46] Schinstock CA, Semret MH, Wagner SJ, Borland TM, Bryant SC, Kashani KB, et al. Urinalysis is more specific and urinary neutrophil gelatinase-associated lipocalin

- is more sensitive for early detection of acute kidney injury. *Nephrology, Dialysis, Transplantation*. 2013;**28**(5):1175-1185. DOI: 10.1093/ndt/gfs127
- [47] Xu SY, Carlson M, Engström A, Garcia R, Peterson CG, Venge P. Purification and characterization of a human neutrophil lipocalin (HNL) from the secondary granules of human neutrophils. *Scandinavian Journal of Clinical and Laboratory Investigation*. 1994;**54**:365-376. PMID: 7997842
- [48] Liu S, Che M, Xue S, Xie B, Zhu M, Lu R, et al. Urinary L-FABP and its combination with urinary NGAL in early diagnosis of acute kidney injury after cardiac surgery in adult patients. *Biomarkers*. 2013;**18**:95-101. DOI: 10.3109/1354750X.2012.740687
- [49] Obermüller N, Geiger H, Weipert C, Urbschat A. Current developments in early diagnosis of acute kidney injury. *International Urology and Nephrology*. 2014;**46**:1-7. DOI: 10.1007/s11255-013-0448-5
- [50] Uwaezuoke SN. The role of novel biomarkers in predicting diabetic nephropathy: A review. *International Journal of Nephrology and Renovascular Disease*. 2017;**10**:221-231. DOI: 10.2147/IJNRD.S143186
- [51] Lopez-Giacoman S, Madero M. Biomarkers in chronic kidney disease, from kidney function to kidney damage. *World Journal of Nephrology*. 2015;**4**:57-73. DOI: 10.5527/wjn.v4.i1.57
- [52] Shlipak MG, Matsushita K, Ärnlöv J, et al. Cystatin C versus creatinine in determining risk based on kidney function. *The New England Journal of Medicine*. 2013;**369**:932-943. DOI: 10.1056/NEJMoa1214234
- [53] Arora MK, Singh UK. Molecular mechanisms in the pathogenesis of diabetic nephropathy: An update. *Vascular Pharmacology*. 2013;**58**:259-271. DOI: 10.1016/j.vph.2013.01.001
- [54] Kern EF, Erhard P, Sun W, Genuth S, Weiss MF. Early urinary markers of diabetic kidney disease: A nested case-control study from the Diabetes Control and Complications Trial (DCCT). *American Journal of Kidney Diseases*. 2010;**55**:824-834. DOI: 10.1053/j.ajkd.2009.11.009
- [55] Patel DN, Kalia K. Efficacy of urinary N-acetyl- β -D glucosaminidase to evaluate early renal tubular damage as a consequence of type 2 diabetes mellitus: A cross-sectional study. *International Journal of Diabetes in Developing Countries*. 2015;**35**:449-457. DOI: 10.1007/s13410-015-0404-2
- [56] Jones AP, Lock S, Griffiths KD. Urinary N-acetyl- β -glucosaminidase activity in type 1 diabetes mellitus. *Annals of Clinical Biochemistry*. 2006;**21**:142-148. DOI: 10.1177/000456329503200104
- [57] Krolewski AS, Niewczas MA, Skupien J, et al. Early progressive renal decline precedes the onset of microalbuminuria and its progression to macroalbuminuria. *Diabetes Care*. 2014;**37**:226-234. DOI: 10.2337/dc13-0985
- [58] Nielsen SE, Sugaya T, Hovind P, Baba T, Parving HH, Rossing P. Urinary liver-type fatty acid-binding protein predicts progression to nephropathy in type 1 diabetic patients. *Diabetes Care*. 2010;**33**:1320-1324. DOI: 10.2337/dc09-2242

- [59] Kamijo-Ikemori A, Sugaya T, Yasuda T, et al. Clinical significance of urinary liver-type fatty acid-binding protein in diabetic nephropathy of type 2 diabetic patients. *Diabetes Care*. 2011;**34**:691-696. DOI: 10.2337/dc10-1392
- [60] Panduru NM, Forsblom C, Saraheimo M, et al.; Finn Diane Study Group. Urinary liver-type fatty acid-binding protein and progression of diabetic nephropathy in type 1 diabetics. *Diabetes Care*. 2013;**36**:2077-2083. DOI: 10.2337/dc12-1868
- [61] Jeon YK, Kim MR, Huh JE, et al. Cystatin C as an early biomarker of nephropathy in patients with type 2 diabetes. *Journal of Korean Medical Science*. 2011;**26**:258-263. DOI: 10.3346/jkms.2011.26.2.258
- [62] Kim SS, Song SH, Kim IJ, et al. Urinary cystatin C and tubular proteinuria predict progression of diabetic nephropathy. *Diabetes Care*. 2013;**36**:656-661. DOI: 10.2337/dc12-0849
- [63] Shore N, Khurshid R, Saleem M. Alpha-1 microglobulin: A marker for early detection of tubular disorders in diabetic nephropathy. *Journal of Ayub Medical College, Abbottabad*. 2010;**22**:53-55. <http://www.ayubmed.edu.pk/JAMC/PAST/22-4/Najla.pdf>
- [64] Hong CY, Hughes K, Chia KS, Ng V, Ling SL. Urinary alpha-1-microglobulin as a marker of nephropathy in type 2 diabetic Asian subjects in Singapore. *Diabetes Care*. 2003;**26**:338-342. DOI: 10.2337/diacare.26.2.338
- [65] Takebayashi K, Matsumoto S, Aso Y, Inukai T. Aldosterone blockade attenuates urinary monocyte chemoattractant protein-1 and oxidative stress in patients with type 2 diabetes complicated by diabetic nephropathy. *The Journal of Clinical Endocrinology and Metabolism*. 2006;**91**:2214-2217. DOI: 10.1097/MNH.0b013e32833ce6d5
- [66] Noris M, Bernasconi S, Casiraghi F, Sozzani S, Gotti E, Remuzzi G, Mantovani A. Monocyte chemoattractant protein-1 is excreted in excessive amounts in the urine of patients with lupus nephritis. *Laboratory Investigation*. 1995;**73**:804-809. PMID: 8558841
- [67] Nakamura A, Shikata K, Hiramatsu M, et al. Serum interleukin-18 levels are associated with nephropathy and atherosclerosis in Japanese patients with type 2 diabetes. *Diabetes Care*. 2005;**28**:2890-2895. DOI: 10.2337/diacare.28.12.2890
- [68] Amer H, Lieske JC, Rule AD, et al. Urine high and low molecular weight proteins one-year post-kidney transplant: Relationship to histology and graft survival. *American Journal of Transplantation*. 2013;**13**:676-684. DOI: 10.1111/ajt.12044
- [69] Bakkalöglu SA, Soylemezoglu O, Buyan N, et al. High serum adiponectin levels during steroid-responsive nephrotic syndrome relapse. *Pediatric Nephrology*. 2005;**20**:474-477. DOI: 10.1007/s00467-004-1770-z
- [70] Bakr A, Rageh I, El-Azouny M, Deyab S, Lotfy H. Serum neopterin levels in children with primary nephrotic syndrome. *Acta Paediatrica*. 2006;**95**:854-856
- [71] Calışkan S, Hacibekiroğlu M, Sever L, Ozbay G, Arisoy N. Urinary N-acetyl-beta-D-glucosaminidase and beta 2-microglobulin excretion in primary nephrotic children. *Nephron*. 1996;**74**:401-404. DOI: 10.1159/000189442

- [72] Bennett MR, Pordal A, Haffner C, Pleasant L, Ma Q, Devarajan P. Urinary vitamin D-binding protein as a biomarker of steroid-resistant nephrotic syndrome. *Biomarker Insights*. 2016;**11**:1-6. DOI: 10.4137/BMI.S31633
- [73] Piyaphanee N, Ma Q, Kremen O, et al. Discovery and initial validation of α 1-B glycoprotein fragmentation as a differential urinary biomarker in pediatric steroid-resistant nephrotic syndrome. *Proteomics. Clinical Applications*. 2011;**5**:334-342. DOI: 10.1002/prca.201000110
- [74] Fede C, Conti G, Chimenz R, Ricca M. N-acetyl-beta-D-glucosaminidase and beta2-microglobulin: Prognostic markers in idiopathic nephrotic syndrome. *Journal of Nephrology*. 1999;**12**:51-55. PMID: 10203005