

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.  
For more information visit [www.intechopen.com](http://www.intechopen.com)



---

## Geriatric Trauma

---

Banu Arslan

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.77151>

---

### Abstract

Worldwide, the proportion of elderly people is constantly increasing. The aging of the baby boomers (people born between 1946 and 1964) and longer life spans (the maximum number of years that a human can live) result in a substantial increase in the number and proportion of older adults (whose age is  $\geq 65$ ). The older population is projected to more than double from 40.3 million in 2010 to 83.7 million in 2050 and, by 2050, it is estimated that older adults will represent 20.9% of the US population. In the early twentieth century, the average life expectancy at birth was 47.3 whereas it was 76.9 in 2000. With the increase in life expectancy due to improvement in quality of medical care, additionally, the oldest old age (age  $\geq 85$ ) forms a rapidly growing group within the older population. The rapid growth of these populations has many significant impacts on public health, emergency room visits, and economy.

**Keywords:** geriatric trauma, ATLS, trauma

---

### 1. Introduction and epidemiology

Worldwide, the number and proportion of elderly people is constantly increasing. The aging of the baby boomers (people born between 1946 and 1964) and longer life spans (the maximum number of years that a human can live) result in a substantial increase in the number and proportion of older adults (who is age  $\geq 65$ ). The elderly population is projected to reach 83, 7 million in the year of 2050 and, by 2050, it is estimated that older adults will represent 20.9% of the U.S. population [1]. With the surge of the elderly population, there will be an increasing number of geriatric trauma patients admit to the emergency departments. Additionally, the rapid growth of these populations will have many significant impacts on public health and economy.

Geriatric trauma patients are less likely to be injured than younger people; however, they are more likely to have fatal outcomes. Death rates for Americans have decreased in the last century. Although there is a dramatic decline in deaths from cardiovascular diseases, heart diseases remain the leading cause of deaths in the elderly. Also, trauma became the more common cause of death. According to the National Center for Health Statistics 2015 report, unintentional injuries became the seventh common cause of death in the elderly [2].

## 2. Pathophysiology

### 2.1. What is aging?

Aging is characterized by a progressive loss of physiological integrity, leading to impaired function and increased vulnerability to death [3]. This multifactorial and extremely complex process results in significant anatomic and functional changes in all major organ systems. Most important systems which are affected are seen in **Table 1**.

### 2.2. Age-related alterations and clinical consequences

**Airway:** The anatomy and physiology of the airway are affected with the aging process. Tooth decay which is common in elderly may cause loose, dislodged and subsequently aspiration of the teeth during emergency procedures such as endotracheal intubation (ET). Esthetic operations and loss of teeth interfere with achieving a good face-mask seal. Pharynx becomes more dry and fragile and care must be taken to prevent profuse bleeding while using laryngoscope. Oral cavity tumors and macroglossia may limit visualization of the vocal cords. Usage of Miller blade can be considered [4]. Also, cervical osteoarthritis increases the risk for spinal cord injury. Excessive movement of the neck should be avoided.

**Age-related alterations:** See **Table 1**.

### 2.3. Common mechanisms of injury

The common causes of geriatric trauma include falls, motor vehicle collisions, pedestrian injuries and thermal injuries and elder abuse (**Figure 1**).

**Falls** remain the leading cause of geriatric trauma and affect approximately 30% of persons aged  $\geq 65$  years each year [17]. Approximately 50% of people living in long-term care institutions fall each year, and 40% of them experienced recurrent falls [18]. Women experience significantly more fall-related injuries than men (35.7 vs.24.6%, respectively) [19]. Falls account for 40% of all injury-associated deaths [20]. Predisposing risk factors include age-related changes in muscle strength, gait and balance, poor vision and home hazards. In addition, drugs and alcohol may contribute to falls. Anticoagulants usage are frequent in elderly and it may cause potentially lethal injuries even with minor traumas. Osteoporosis and the tendency to fall increase the risk of hip fractures. Also, falls are the most common cause of traumatic brain injury in the elderly. Even when those injuries are minor, they seriously affect older

Organ system	Age-related alterations	Clinical consequences
Respiratory system	<p>Elastin component of the lung matrix ↑</p> <p>type III collagen ↓ [5].</p> <p>Pulmonary compliance ↓</p> <p>Osteoporosis</p> <p>Stiffness of the thoracic cage</p> <p>Outward recoil ↑</p> <p>Kyphosis [6].</p> <p>Chest wall compliance ↓</p> <p>Thickening of the alveolar basement membrane [7].</p> <p>Diffusion capacity ↓</p> <p>Gas change ↓</p> <p>Muscle atrophy</p> <p>Respiratory muscle weakness [8].</p>	<p>Risk for pneumonia ↑ [5]</p> <p>Poor tolerance to rib fractures</p> <p>Work of breathing ↑</p> <p>Risk for respiratory failure ↑ [8].</p> <p>Forced expiratory volume in one second (FEV1) ↓</p> <p>Forced vital capacity (FVC) ↓</p> <p>FEV1/FVC ↓ [9].</p> <p>Functional residual capacity (FRC) ↓</p> <p>Residual volume (RV) ↑</p> <p>Vital capacity (VC) ↓</p>
Circulatory system	<p>Vascular stiffness</p> <p>Left ventricle (LV) wall thickness ↑ [10].</p> <p>Retarded early diastolic cardiac filling and LV diastolic function ↓</p> <p>Afterload ↑</p> <p>Left atrial size ↑</p> <p>myocyte mass with ↓</p> <p>deposition of amyloid and collagen</p> <p>Deterioration of the cardiac conduction</p> <p>Decreased sensitivity to catecholamines</p> <p>Maximal heart rate ↓</p> <p>Maximum tachycardia response ↓ [11].</p>	<p>Elevated baseline blood pressure</p> <p>Atherosclerosis of coronary vessel</p> <p>Risk for cardiac ischemia ↑</p> <p>Increased risk of dysrhythmias</p> <p>Impaired cardiac reserve</p> <p>Cardiac index ↓</p> <p>Lack of classic response to hypovolemia</p>

Organ system	Age-related alterations	Clinical consequences
Musculoskeletal system	Stiffening of structural instruments (tendons, ligaments, cartilage) Spontaneous rupture Joint stability ↓ ineffective repair of cartilage tissue Osteoarthritis (cervical, temporomandibular) Bone volume-mass ↓ Muscle size-number ↓ [12]. Osteoporosis Sarcopenia	Risk of injury ↑ Risk of fracture ↑ Difficulty for oral intubation Risk of falls ↑ Mobility ↓
Nutrition and metabolism	Taste acuity, smell and appetite decrease Poor dentition Inability to eat independently	Food intake ↓ [13].
Central nervous system	Brain volume decreases Replaced by cerebrospinal fluid [14]. Protection against contusions Blood can be collected Parasagittal bridging veins stretch More prone to tear injury Demyelination ↑ Peripheral conduction velocity slows Dura adheres to the skull more tightly Cerebral blood flow Cerebral oxygen consumption [15]. Degeneration of vertebrae, intervertebral disks and facet joints.	Less contusions Clinical signs may manifest late More subdural hematoma Vision and auditory functions ↓ [16]. Reaction time ↑ Attention span ↓ Less epidural hematoma Risk for spine and spinal cord injury

**Table 1.** Age-related alterations and their clinical consequences.




Figure 1. Common causes of falls in the elderly [64].

adults' quality of life by inducing a fear of falling, which can lead to self-imposed activity restrictions, anxiety, social withdraw and depression [21].

**Motor vehicle collision** involving elderly continue to increase. Age-related changes that include vision and hearing impairment, decreased night vision and glare resistance are the prominent factors on the incidence of injury and death. Additionally, medical conditions and medications may distort the reaction time, attention and judgment which increase the risk for the collision.

**Pedestrian injuries:** according to the 2015 pedestrian data, 19% of all pedestrian fatalities and an estimated 13% of all pedestrians injured were people aged 65 and older in the United States, and pedestrian-motor vehicle collisions are one of the most lethal mechanisms of injury in this age group with a 53% case fatality rate [22].

**Thermal injuries:** There is a direct relationship between age and burn mortality, as evidenced by the traditionally taught BauxScore. The empiric formula is clearly the simplest, whereby the sum of the patient's age and burn size predict mortality. Based on the data from the American Burn Association (ABA) National Burn Repository (NBR) from 2000 to 2009, overall

mortality was 4% in all age groups and 17% in seniors [23]. Moreover, for seniors there is a greater increase in mortality risk for every 1% increase in burn size and 1-year increase in age than among adults [23].

### 3. Clinical features and the management of injured elderly patients

The management of injured elderly requires the rapid assessment and rapid intervention of life-threatening situations. The assessment sequence should be same as in adults and pediatric population and includes the following elements:

#### 3.1. Prehospital management and triage

The triage decision can be made through “field triage decision scheme” which was published by the American College of Surgeons Committee on Trauma (ACS-COT) to provide a guidance for the field triage process (**Figure 2**). Under triage, inaccurate triage which results in an assignment of lower triage level is more common among the elderly patients [24]. In order to avoid high under-triage rates in elderly, two important statements were added to Step Four of the scheme:

- SBP <110 might represent shock after age 65.
- Low impact mechanisms (e.g., ground-level falls) might result in severe injury.

Furthermore, we recommend that the injured elderly who met Step Four criteria should be transported to the trauma center [25]. Moreover, elderly seem to benefit more from triage to trauma center with improved outcomes [26]. Also, it is important that the transferring and receiving facilities develop transfer agreements in advance.

#### 3.2. Primary survey

Primary survey of geriatric trauma patients includes rapid and efficient assessment of vital functions, assessment of the ABCDs, and identification and therapeutic intervention of life-threatening conditions as those for adults. Establishing and maintaining a patent airway to provide adequate oxygenation within-line cervical stabilization is the first objective. Avoiding excessive movement of the neck is crucial to prevent spinal cord injury. Because geriatric patients have limited respiratory reserve, early administration of supplemental oxygen is crucial. Early intubation should be considered if geriatric trauma patients present shock or chest wall injury/altered level of consciousness. For geriatric trauma patients, it is more challenging to recognize the early symptoms of shock. The aging process diminishes the physiologic reserve and chronic diseases can impair their ability to respond to injury; a tachycardic response may be absent or blunted. Also, medications such as  $\beta$ -blockers may mask tachycardia. Blood pressures are also misleading in the elderly patients. Due to increased incidence of underlying hypertension, the clinician must use a higher cutoff for hypotension than in younger patients [27]. In addition, frequently repeated measurement and interpreting the results according to baseline and previous ones may help the clinicians. Early and close monitoring must be instituted. Resuscitation of the elderly warrants special attention. Fluid




Figure 2. Guidelines for field triage of injured patients-United States, 2011 [65].

resuscitation is often challenging in geriatric trauma patients because of underlying cardiac dysfunction and concerns about precipitating heart failure. Primary survey also includes urinary and gastric catheters, arterial blood gas levels and X-rays (e.g., chest and pelvis). In the



elderly, due to alterations with aging, anticoagulant usage may increase the chance of profuse bleeding during the catheter procedure. Special care should be taken during this procedure.

### 3.3. Secondary survey

Secondary survey includes head-to-toe evaluation, reassessment of all vital signs, diagnostic tests and expanded history of the geriatric trauma patients. A detailed description of the secondary survey is provided separately; special circumstances in geriatric trauma patients are discussed here. Clinicians should focus on identifying and treating injuries which were not discovered during the primary survey. Geriatric trauma patients often present with significant occult injury mostly caused by minor mechanism such as ground-level falls. It is demonstrated that the elderly with blunt head trauma are more likely to present in occult fashion than youngsters, even if they have significant intracranial injury. Moreover, persistent vomiting and headache were less likely to occur in elderly with any intracranial injury [28]. Also, initially stable geriatric trauma patients may deteriorate rapidly and without warning. During the secondary survey it is essential to assess the alterations in mental status, especially compared to presentation.

### 3.4. High-risk injuries

The risk of complications increases with the severity of the *trauma*; however, even minor traumas such as ground-level falls or slipping while walking off a curb may seem relatively harmless in elderly patients, they can lead to severe injury and death [29].

### 3.5. Head injury

Traumatic brain injury (TBI) is a significant problem among the elderly. For the age of 65 years and older, falls are the primary mechanisms of TBI-related ED visits (81.8%) and TBI-related deaths (54.4%) [30]. In the review of the literature, it is recognized that older adults with moderate–severe TBI have poor outcomes with high rates of significant disability and mortality. Two major factors put geriatric trauma patients at a greater risk for increased incidence of TBI: age-related structural changes and preinjury anticoagulant-antiplatelet usage. First, with aging, parasagittal bridging veins stretch and make the elderly more susceptible to traumatic tears. Thus, the elderly have a higher incidence of subdural hematoma. Also, cerebral atrophy leads to a significant amount of blood accumulating in subdural area before clinical signs manifest. Rapid neurologic decline should be considered in these patients. Second, an increased incidence of the anticoagulant and antiplatelet therapy in the elderly may have detrimental consequences. It is suggested that taking anticoagulant therapy at the time of the injury increases the risk of intracranial hemorrhage [31] and is related with worse outcomes [32, 33]. One of the most frequently prescribed anticoagulant medications is warfarin. Also, Franko et al. concluded that warfarin use at the time of injury also makes mortality significantly higher after the age 70 [32]. Thus, immediate noncontrast head computed tomography (CT) is recommended for the elderly patients who take anticoagulant or antiplatelet therapy, even if their trauma seems minor. Additionally, rapid screening for anticoagulant use, INR value and subsequent correction with blood component therapy may improve outcomes.

### 3.6. Spine injury

**Cervical spine** injuries are more common in the elderly and the incidence appears to be increasing [34, 35]. The most commonly seen injury site is upper cervical spine (UCS) especially the odontoid process [36] and caused by falls. The UCS injuries are associated with a high rate of mortality and morbidity. Elderly patients tend to sustain more C-spine fracture following simple falls such as ground-level falls [37]. It is attributed to increased frequency of preexisting cervical spine pathology such as osteoporosis and osteoarthritis [36]. It may also result in occult presentation, delayed diagnosis, increased risk for spinal cord injuries and difficulty in interpreting plain radiographs. Moreover, mild extension injuries followed by fall or rear-end motor vehicle crashes may cause central cord syndrome in the presence of preexisting spinal canal stenosis [37].

**Thoracolumbar spine** fractures in the elderly are usually associated with osteoporosis. Osteoporosis affects almost 50% of these individuals and contributes to the occurrence of spontaneous vertebral compression fractures. The majority of the osteoporotic vertebral fractures are situated in thoracolumbar spine, and the anterior wedge compression fractures are the most common site.

Treatment of diagnosed vertebral fractures in these individuals is still controversial. Two options are available: conservative therapy and surgery. Unstable fractures, flexion distraction injuries and severe burst fractures causing neurologic deficit mostly indicate surgical intervention. However, in the patient who is neurologically intact, conservative treatment including bed-rest and bracing seems a more viable option depending on the type of fracture [38]. Consequently, we recommended that apparently low-energy level injuries should be considered as a high-risk for spine injury and investigated elaborately. CT scan is the preferred initial modality for assessing the geriatric cervical spine because The Canadian Cervical-Spine Rule, but not the National Emergency X-Radiography Utilization Study criteria, excludes patients aged  $\geq 65$  years from being considered low risk for cervical spine injury.

### 3.7. Chest trauma

Chest traumas account for ~796,000 emergency department (ED) visits annually in the USA [39]. For blunt chest trauma, the most prominent factors in etiology are falls and motor vehicle collisions. The elderly are more prone to incur chest injuries following blunt chest trauma, and this is associated with a high risk of mortality and morbidity [40]. Rib fractures and pulmonary contusions are more common in this population due to preexisting osteoporosis, loss of muscle mass and comorbidities [41]. The mortality and risk for pneumonia following blunt chest trauma significantly increase after 65 years [40, 41] and it is correlated with the increased number of rib fractures [40, 42]. In the presence of pulmonary contusion, clinicians should consider early ventilatory support because these patients are highly vulnerable to respiratory compromise. Given these risks, detailed physical examination, close observation and early administration of supplemental oxygen with adequate pain medication are highly recommended for elderly patients with even one rib fracture. Also, advanced imaging is warranted in older patients with multiple rib fractures. CT may be necessary to assess the extent of

injuries that might not be seen on plain radiographs. Simple pneumothorax and hemothorax are poorly tolerated by elderly patients. Thus, geriatric patients with life-threatening chest trauma should be considered for intensive care unit (ICU) observation.

### 3.8. Abdominal trauma

Abdominal examination can be less reliable and more difficult because of decreased pain sensation and increased laxity of abdominal wall musculature. Also, guarding and rigidity may be lacking in the elderly. Tachycardia response to hemorrhagic shock may not be seen even in the setting of significant blood loss. High index of suspicion and close observation must be continued to be avoided under-diagnosis. The Focused Assessment with Sonography for Trauma (FAST) can be used to detect intraperitoneal fluid in patients who sustain blunt abdominal trauma. CT remains the gold standard to diagnose intra-abdominal injuries. Retroperitoneum is an occult source of bleeding. Also, the risk of occult retroperitoneal bleeding is higher with chronic anticoagulant usage. Therefore, CT with contrast should be considered to evaluate hemorrhage, especially for the elderly patients who have pelvis or hip fracture.

### 3.9. Musculoskeletal trauma

Fractures are frequent in the elderly and can cause severe pain, disability and loss of independence. The increased risk of fracture with age may attribute to increased risk of fall, osteoporosis, sarcopenia and frailty.

**Pelvis fractures:** In the elderly, low-energy traumas such as ground level falls may result in pelvic fractures [43]. Although patients with pelvic fractures due to minor trauma generally do not present complications, mortality and morbidity increase with accompanied hemorrhage and other associated injuries. The portable AP pelvic X-ray should be obtained as a part of the primary survey. However, posterior ring fractures can be missed. Patients who have pelvic tenderness following pelvic trauma must be assessed regarding pelvic fracture. CT of the pelvis can be obtained in stable patients. If an active bleeding is suspected, pelvic contrast CT is recommended considering the risk of contrast-induced nephropathy. If an active bleeding is identified, arteriography and embolization can be performed for the patients in danger of life. Consequently, expeditious hemorrhage control with simultaneous emergency skeletal stabilization and resuscitation is crucial for the management of pelvic fractures in the elderly.

**Proximal femur fractures:** In elderly patients, hip fractures should be considered as a serious injury. They may lead to immobility, permanent dependence and death. According to several epidemiological studies, the incidence of proximal femoral fractures increases with age, starting at 40 years, with a steep increase after 75 years of age. The average age of patients with hip fracture is over 80, and nearly 80% are women [44]. Although isolated hip fractures do not usually cause class III or class IV shock, long-term prognosis mostly depends on age, comorbidities, anticoagulant therapy and frailty [45]. Hip fractures are the most common cause of accident-related deaths in older people accounting for 18% deaths within 4 months of a hip fracture and 30% within a year [46]. The risk of fracture increases with the number of falls [47] and backward fall mechanism and low bone mineral density (BMD) [48]. Most hip fractures can be diagnosed by typical history and clinical presentations. The first choice

for diagnose is plain radiographs. However, it is estimated that 2–9% of fractures may be radiographically occult [49], and further imaging such as CT and MRI is required to make a definitive diagnosis. MRI has higher sensitivity than CT for detecting occult hip fractures. Additionally, nuclear medicine scintigraphy may be another choice for diagnosis due to high sensitivity. However, access to the scintigraphy usually is difficult and, it has limited capability to delineate the full nature of the fracture.

## 4. Special circumstances

### 4.1. Preexisting medical conditions

Elderly individuals are more likely to have preexisting comorbidities. The presence of a preexisting medical condition was associated with increase in mortality of elderly patients who sustained low or moderate severity trauma [50]. The most frequent preinjury comorbidities are hypertension (HT), diabetes mellitus (DM), coronary artery disease (CAD) and use of anticoagulants/antiplatelets [51]. Preinjury medical conditions usually make the management of geriatric trauma patients challenging; preexisting HT can hide the early signs of shock and cause delay or under-treatment and the presence of heart failure may cause volume overload and pulmonary edema during IV fluid therapy. ET intubation also would be challenging in the patient who has cervical or temporomandibular arthritis. Thus, early detection of preexisting medical conditions, appropriate treatment and follow-up care may improve outcomes following trauma in elderly.

### 4.2. Pre-injury medication usage

As the population ages, increasing numbers of elderly are being prescribed a medication for chronic medical conditions. It was shown that medications (especially sedatives and hypnotics, antidepressants, and benzodiazepines [52]) are particularly complex risk factors for falls and the risk of falling increases with the number of medications taken [18]. Also, polypharmacy is associated with occurrence of drug–drug interactions and adverse drug reactions which are frequently encountered in the elderly [53]. B-adrenergic blocking agents may limit the tachycardia response which can result in undesirable decreased cardiac output and reduced tissue perfusion. Calcium-channel blockers may prevent peripheral vasoconstriction and contribute to produce hypotension. Chronic diuretic use may lead to elderly patients being chronically hypovolemic, hyponatremic and hypokalemic. Additionally, declines in renal and hepatic function may alter the metabolism and clearance of these drugs. The side effects, drug interactions should always be considered and potentially nephrotoxic drugs must be given in adjusted doses based on calculated creatinine clearance.

### 4.3. Risk of bleeding

In the elderly population, both age-related structural changes and usage of some chronic medications may increase the risk of bleeding. Chronic anticoagulant therapy can increase the risk of hemorrhage, especially intracranial hemorrhage (ICH) [31]. The usage of warfarin at the time of injury also makes mortality significantly higher after the age 70 [32]. Recent

data show that Apixaban, dabigatran and rivaroxaban have lower risk of intracranial bleeding compared to warfarin [54]. However, they may potentially carry more risk of major bleeding than warfarin [55]. The influence of preinjury aspirin therapy on bleeding and the mortality is still uncertain [56]. However, the increased risk of subdural hematoma following head trauma was shown in the patients who are under preinjury aspirin plus clopidogrel therapy [57]. Hemorrhage cannot be tolerated appropriately. Therefore, the management of elderly trauma patients who are under anticoagulant therapy requires special care. Early diagnosis, close monitoring and maintaining optimal hemoglobin level are crucial. The optimal hemoglobin level for injured elderly patients is still controversial. A general suggestion is that hemoglobin concentration should be maintained over 10 g/Dl in order to maximize oxygen carrying capacity and delivery. Also, correction of coagulation defects is very important. According to the Eastern Association for the Surgery Trauma, all elderly patients with evidence of posttraumatic ICH on CT with Warfarin should have their INR be corrected toward a normal range within 2 h of admission [58]. Moreover, tranexamic acid, an antifibrinolytic agent, may reduce blood loss after traumatic injury. According to the recent data, tranexamic acid may reduce mortality without significant adverse side effects when given within 1–3 h [59]. The dose is 1 g of tranexamic acid IV bolus over 10 min, followed by 1 g IV over 8 h.

**Pain management** altered physiology changes the way analgesic drugs are distributed and metabolized therefore the pain management of geriatric trauma patient requires extra caution. The main approach should provide optimal treatment of pain while minimizing the risk of medication-related adverse effects. The standardized tools to assess the pain may be beneficial [60] (**Table 2**).

Pain type or source	Nonopioids	Opioids	Adjuvant analgesics	Other	Comments
Major trauma generalized pain	Acetaminophen, NSAIDs during posttrauma healing phase	Bolus or continuous IV opioids* during emergency phase; PO or IV opioids during healing phase	IV ketamine (very rare)	Inhaled NO	Use of ketamine is restricted to pain refractory to other treatments due to severe CNS side effects Inhaled NO is used for incident pain
Major trauma (regionalized pain)	NSAIDs (parenteral, oral) during posttrauma healing phase	Bolus or continuous IV opioids during emergency phase plus regional anesthesia	IV ketamine (very rare)	Inhaled NO	Use of ketamine is restricted to pain refractory to other treatments due to severe CNS side effects. Inhaled NO is used for incident pain

Pain type or source	Nonopioids	Opioids	Adjuvant analgesics	Other	Comments
Burns	Acetaminophen, NSAIDs, during rehabilitative phase	High dose of IV opioids ± PCA for NPO patients; oral opioids (e.g., morphine, hydromorphone) when taking PO	Parenteral ketamine (very rare) IV lidocaine (very rare)	BNZ Inhaled NO	Use of ketamine is restricted to pain refractory to other treatments due to severe CNS side effects. Inhaled NO is used for incident pain  Infusion of low-dose lidocaine is restricted to burn pain refractory to opioids.
Minor trauma	Acetaminophen, NSAIDs	Opioids for mild-to-moderate pain			
Procedural pain	NSAIDs for preemptive analgesia and postprocedural pain	IV opioids (e.g., morphine, hydromorphone, fentanyl) unless contraindicated**	Local anesthetics (e.g., EMLA®, lidocaine, bupivacaine, ropivacaine) IV ketamine	BNZ (e.g., diazepam, lorazepam, midazolam) Inhaled NO Propofol***	Local anesthetics may be applied topically (e.g., EMLA®), injected into tissue, or used for nerve blocks Use of ketamine limited by severe CNS side effects

BNZ: benzodiazepines; CNS: central nervous system; EMLA®: Eutectic Mixture of Local Anesthetics (lidocaine and prilocaine); IV: intravenous; LAs: local anesthetics; NO: nitrous oxide; NPO: nothing per os (by mouth); NSAIDs: nonsteroidal anti-inflammatory drugs, including aspirin; PO: per os (oral); PCA: patient-controlled analgesia; PRN: as needed; TD: transdermal.

Modified from American Pain Society, Section IV: Management of Acute Pain and Chronic Noncancer Pain. [http://americanpainsociety.org/uploads/education/section\\_4.pdf](http://americanpainsociety.org/uploads/education/section_4.pdf).

\*Titrate opioids carefully to maintain stable cardiovascular and respiratory status. Monitor neurological and neurovascular status continuously in patients with head injury or limb injury, respectively.

\*\*Contraindications to opioid analgesia include altered sensorium, full-term pregnancy, lung disease or inability to monitor and manage certain side effects (e.g., respiratory depression).

\*\*\*Hypnotic general anesthetic that produces good sedation.

**Table 2.** Systemic medications for acute pain management.

The search of literature mostly suggests that paracetamol should be considered as a first-line treatment for both acute and chronic pain due to its efficacy and good safety profile. NSAIDs are one of the most widely used painkillers. Clinicians must be concerned about the

Type of Abuse	Clinical Markers Indicating Abuse or Neglect
Physical abuse	<p>Abrasion and laceration in sites other than the arms and legs or multiple ones should raise suspicion.</p> <p>Bruising on face, neck, the chest wall, the abdomen, the buttocks, the palms and soles.</p> <p>Tramline bruising.</p> <p>Fractured, subluxed, or avulsed teeth</p> <p>Fractures of the zygomatic arc, mandible and maxilla</p> <p>Fractures not involving the hip, humerus, or vertebra</p> <p>A spiral fracture of a large bone with no history of gross injury or atypical site</p>
Verbal or psychological abuse	<p>Subtle signs of intimidation, such as deferring questions to a caregiver or potential abuser</p> <p>Evidence of isolation of victim from both previously trusted friends and family members</p>
Sexual abuse	<p>The majority of victims have cognitive impairment or have functional limitations.</p> <p>Bruising of the uvula and the palate may indicate forced oral copulation</p> <p>Bleeding, abrasions, lacerations in the anogenital area as well as difficulty in sitting</p> <p>New diagnosis of sexually transmitted diseases, especially in nursing home residents (and especially in cluster outbreaks)</p>
Financial exploitation	<p>Inability to pay for medicine, medical care, food, rent, or other necessities</p> <p>Failure to renew prescriptions or keep medical appointments</p> <p>Unexplained worsening of chronic medical problems that were previously controlled</p> <p>Malnutrition, weight loss, or both, without an obvious medical</p> <p>Firing of home care or other service providers by abuser</p> <p>Unpaid utility bills leading to loss of service</p>
Neglect	<p>Deep decubitus ulcers in multiple sites or foul-smelling, and necrotic ulcer may indicate neglect</p> <p>Refusal to eat may indicate improper feeding technics such as forceful assistance may lead to choking, aspiration and pneumonia.</p> <p>Need help with eating?</p> <p>Adverse side effects due to improper dosing of an indicated drug</p> <p>Overdosing patients to keep them quite and manageable</p> <p>Recent decline in personal care, dirty clothes, multiple insect bites are the signs of poor hygiene.</p>

**Table 3.** Clinical markers indicating abuse or neglect [66, 67].

potentially life-threatening side effects such as gastrointestinal hemorrhage. And, it must be given with proton-pump inhibitor (PPI) cover. In carefully selected and monitored patients, opioids usually provide fast and effective pain relief. The weak opioids including co-codamol, codeine and dihydrocodeine may elicit adverse effects such as cognitional decline and constipation. Although tramadol's GI effects lesser than other weak opioids, potential to precipitate delirium and reduced seizure threshold may limit the usage [61]. Strong opioids include morphine, oxycodone and fentanyl may also be used to treat moderate and severe pain, especially if the pain causes functional impairment. Dose titration based on patient's response is required, in order to avoid side effects such as sedation, nausea or vomiting.

#### 4.4. Elder abuse/maltreatment

Elder abuse is a global public health and human rights problem which is associated with morbidity and premature mortality. According to the latest data, the prevalence of elder abuse can vary widely. In USA, 10% of older adults have experienced some form of elder abuse [62].

Unfortunately, these statistics may represent an inaccurate underestimation because elder abuse often is not recognized and tends to be underreported. Elder abuse can be classified into five main categories and manifestations is shown in **Table 3**, but several types of abuse may occur simultaneously.

The risk factors can be stated as: shared living situation, social isolation, dementia, female gender, relationship of victim to perpetrator (spouse), personality characteristics of victim (hostility), race (black) [63].

Also it is crucial to screen for elder abuse in geriatric trauma patients, especially who have cognitive impairment or who are unwilling to report it due to fear. Health professionals are well positioned to identify elder abuse, detect vulnerabilities and evaluate interventions. If abuse or neglect is suspected or confirmed, management strategies should be applied.

## Author details

Banu Arslan

Address all correspondence to: [dr.banuarslan@hotmail.com](mailto:dr.banuarslan@hotmail.com)

Department of Emergency Medicine, Marmara University Pendik Research and Training Hospital, Istanbul

## References

- [1] U.S. Census Bureau. P23-212, 65+ in the United States: 2010. Washington, DC: U.S. Government Printing Office; 2014. <https://www.census.gov/content/dam/Census/library/publications/2014/demo/p23-212.pdf>
- [2] National Center for Health Statistics. Health, United States, 2016: With Chartbook on Long-Term Trends in Health. Hyattsville, MD; 2017. <https://www.cdc.gov/nchs/data/healthstats/16.pdf#020>
- [3] López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks of aging. *Cell*. 2013 Jun 6; **153**(6):1194-1217. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3836174/>
- [4] Johnson KN, Botros DB, Groban L, Bryan YF. Anatomic and physiopathologic changes affecting the airway of the elderly patient: Implications for geriatric-focused airway management. *Clinical Interventions in Aging*. 2015 Dec 4; **10**:1925-1934. DOI: 10.2147/CIA.S93796 eCollection 2015
- [5] D'Errico A, Scarani P, Colosimo E, Spina M, Grigioni WF, Mancini AM. Changes in the alveolar connective tissue of the ageing lung. An immunohistochemical study. *Virchows Archiv A Pathological Anatomy and Histopathology*. 1989; **415**(2):137-144



- [6] Gulshan Sharma, James Goodwin. Effect of aging on respiratory system physiology and immunology. *Clinical Interventions in Aging*. 2006;1(3):253-260
- [7] Niewoehner DE, Kleinerman J. Morphologic basis of pulmonary resistance in the human lung and effects of aging. *Journal of Applied Physiology*. 1974;36:412-418
- [8] Polkey MI, Harris ML, Hughes PD, et al. The contractile properties of the elderly human diaphragm. *American Journal of Respiratory and Critical Care Medicine*. 1997;155:1560-1564
- [9] Knudson RJ, Lebowitz MD, Holberg CJ, Burrows B. Changes in the normal maximal expiratory flow-volume curve with growth and aging. *The American Review of Respiratory Disease*. 1983 Jun;127(6):725-734
- [10] Stratton JR, Levy WC, Cerqueira MD, et al. Cardiovascular responses to exercise. Effects of aging and exercise training in healthy men. *Circulation*. 1994;89:1648
- [11] Fleg JL, O'Connor F, Gerstenblith G, Becker LC, Clulow J, Schulman SP, Lakatta EG. Impact of age on the cardiovascular response to dynamic upright exercise in healthy men and women. *Journal of Applied Physiology*. 1995;78:890-900. [PubMed: 7775334]
- [12] Tiel M, Trouwborst I, Clark BC. Skeletal muscle performance and ageing. *Journal of Cachexia, Sarcopenia and Muscle*. 2018 Feb;9(1):3-19. DOI: 10.1002/jcsm.12238. Epub 2017 Nov 19
- [13] Hickson M. Malnutrition and ageing. *Postgraduate Medical Journal*. 2006 Jan;82(963):2-8
- [14] Driscoll I, Davatzikos C, An Y, et al. Longitudinal pattern of regional brain volume change differentiates normal aging from MCI. *Neurology*. 2009;72:1906
- [15] Yamaguchi T, Kanno I, Uemura K. Reduction in regional cerebral metabolic rate of oxygen during human aging. *Stroke*. 1986;17:1220-1228
- [16] DeLatorre J, Fay L. Effects of aging on the human nervous system. In: Rosenthal R, Zenilman M, Catlic M, editors. *Principals and Practice of Geriatric Surgery*. New York: Springer-Verlag NY Inc.; 2001:926-948
- [17] Centers for Disease Control and Prevention (CDC). Fatalities and injuries from falls among older adults – United States, 1993-2003 and 2001–2005. *MMWR Morb Mortal Weekly Reports*. November 17, 2006;55(45):1221-1224
- [18] Tinetti ME, Speechley M, Ginter SFN. Risk factors for falls among elderly persons living in the community. *The New England Journal of Medicine*. 1988 Dec 29;319(26):1701-1707
- [19] Centers for Disease Control and Prevention (CDC). Self-reported falls and fall-related injuries among persons aged > or =65 years–United States, 2006. *Morbidity and Mortality Weekly Report*. 2008 Mar 7; 57(9):225-229
- [20] Rubenstein LZ. Falls in older people: Epidemiology, risk factors and strategies for prevention. *Age Ageing*. 2006; 35-S2:ii37-ii41
- [21] Tinetti ME, Williams CS. The effect of falls and fall injuries on functioning in community-dwelling older persons. *The Journals of Gerontology. Series A, Biological Sciences and Medical Sciences*. 1998 Mar; 53(2):M112-M119

- [22] National Highway Traffic Safety Administration. Traffic Safety Facts 2015 Data – Pedestrians. Washington, DC: US Department of Transportation, National Highway Traffic Safety Administration; 2017. Publication no. DOT-HS-812-375. Available at <https://crashstats.nhtsa.dot.gov/Api/Public/ViewPublication/812375>
- [23] Pham TN, Kramer CB, Wang J, Rivara FP, Heimbach DM, Gibran NS, Klein MB. Epidemiology and outcomes of older adults with burn injury: An analysis of the national Burn Repository. *Journal of Burn Care & Research*. 2009 Jan-Feb;30(1):30-36. DOI: 10.1097/BCR.0b013e3181921efc
- [24] Chang DC, Bass RR, Cornwell EE, Mackenzie EJ. Under-triage of elderly trauma patients to state-designated trauma centers. *Archives of Surgery*. 2008;143:776-781. discussion 782
- [25] Sasser SM, Hunt RC, Faul M, Sugerman D, Pearson WS, Dulski T, Wald MM, Jurkovich GJ, Newgard CD, Lerner EB. Guidelines for field triage of injured patients: recommendations of the National Expert Panel on Field Triage; 2011
- [26] Meldon SW, Reilly M, Drew B, Mancusco C, Fallon W. Trauma in the very elderly: A community based study of outcomes at trauma and non-trauma centers. *Academic Emergency Medicine*. 2000;7(10):1166
- [27] ATLS Subcommittee; American College of Surgeons' Committee on Trauma; International ATLS Working Group. Advanced trauma life support (ATLS®): The ninth edition. *The Journal of Trauma and Acute Care Surgery*. 2013 May;74(5):1363-1366. DOI: 10.1097/TA.0b013e31828b82f5
- [28] NK1 R, Medzon R, Lowery D, Pollack C, Bracken M, Barest G, Wolfson AB, Hoffman JR, Mower WR. Intracranial pathology in elders with blunt head trauma. *Academic Emergency Medicine*. 2006 Mar;13(3):302-307
- [29] Spaniolas K, Cheng JD, Gestring ML, Sangosanya A, Stassen NA, Bankey PE. Ground level falls are associated with significant mortality in elderly patients. *The Journal of Trauma*. 2010 Oct 0;69(4):821-825
- [30] Percent Distributions of TBI-related Emergency Department Visits by Age Group and Injury Mechanism—United States, 2006-2010. [https://www.cdc.gov/traumaticbraininjury/data/dist\\_ed.html](https://www.cdc.gov/traumaticbraininjury/data/dist_ed.html)
- [31] Courtney E. Collins, Elan R. Witkowski, Julie M. Flahive, Fred A. Anderson, Jr, and Heena P. Santry, Effect of preinjury warfarin use on outcomes after head trauma in Medicare beneficiaries. *The American Journal of Surgery*. 2014 Oct; 208(4):544-549.e1
- [32] Franko J, Kish KJ, O'Connell BG, Subramanian S, Yuschak JV. Advanced age and preinjury warfarin anticoagulation increase the risk of mortality after head trauma. *The Journal of Trauma*. 2006 Jul;61(1):107-110
- [33] Mina AA, Knipfer JF, Park DY, Bair HA, Howells GA, Bendick PJ. Intracranial complications of preinjury anticoagulation in trauma patients with head injury. *The Journal of Trauma* 2002 Oct;53(4):668-672
- [34] Wang H, Li C, Xiang Q, Xiong H, Zhou Y. Epidemiology of spinal fractures among the elderly in Chongqing. *China Injury*. 2012;43:2109-2116

- [35] Roche SJ, Sloane PA, McCabe JP. Epidemiology of spine trauma in an Irish regional trauma unit: A 4-year study. *Injury*. 2008;**39**:436-442
- [36] Jubert P, Lonjon G, Garreau de Loubresse C. Complications of upper cervical spine trauma in elderly subjects. A systematic review of the literature. *Orthopaedics & Traumatology: Surgery & Research*. 2013 Oct;**99**(6 Suppl):S301-S312. DOI: 10.1016/j.otsr.2013.07.007. Epub 2013 Aug 22
- [37] Hao Wanga, c, Marco Coppolaa, Richard D. Robinsona, James T. Scribnera, Veer Vithalania, Carrie E. de Moora, Raj R. Gandhib, Mandy Burtona, Kathleen A. Delaneya geriatric trauma patients with cervical spine fractures due to ground level fall: Five years experience in a level one trauma center. *Journal of Clinical Medicine Research*. 2013;**5**(2):75-83
- [38] Chang V, Holly LT. Bracing for thoracolumbar fractures. *Neurosurgical Focus* 2014; **37**(1):E3. DOI: 10.3171/2014.4.FOCUS 1477
- [39] Pitts SR, Niska RW, Xu J, Burt CW. National Hospital Ambulatory Medical Care Survey: 2006 emergency department summary. *National Health Statistics Reports*. 2008 Aug 6;**7**:1-38
- [40] Bulger EM, Arneson MA, Mock CN, Jurkovich GJ. Rib fractures in the elderly. *Journal of Trauma*. 2000 Jun; **48**(6):1040-1046; discussion 1046-7
- [41] Bergeron E, Lavoie A, Clas D, Moore L, Ratte S, Tetreault S, Lemaire J, Martin M. Elderly trauma patients with rib fractures are at greater risk of death and pneumonia. *The Journal of Trauma*. 2003 Mar;**54**(3):478-485
- [42] Stawicki SP, Grossman MD, Hoey BA, Miller DL, Reed JF 3rd. Rib fractures in the elderly: A marker of injury severity. *Journal of the American Geriatrics Society*. 2004 May;**52**(5):805-808
- [43] Nanninga GL, de Leur K, Panneman MJ, van der Elst M, Hartholt KA. Increasing rates of pelvic fractures among older adults: The Netherlands, 1986-2011. *Age Ageing*. 2014 Sep; **43**(5):648-653
- [44] Keene GS, Parker MJ, Pryor GA. Mortality and morbidity after hip fractures. *BMJ*. 1993 Nov 13; **307**(6914):1248-1250
- [45] Carpintero P, Caeiro JR, Carpintero R, Morales A, Silva S, Mesa M. Complications of hip fractures: A review. *World Journal of Orthopedics*. 2014 Sep 18; **5**(4):402-411
- [46] Roberts SE, Goldacre MJ. Time trends and demography of mortality after fractured neck of femur in an English population, 1968-98: Database study. *BMJ*. 2003;**327**:771-775
- [47] Cummings SR, Nevitt MC. Non-skeletal determinants of fractures: the potential importance of the mechanics of falls. Study of Osteoporotic Fractures Research Group. *Osteoporosis International*. 1994;**4**(Suppl 1):67-70
- [48] Nevitt MC, Cummings SR. Type of fall and risk of hip and wrist fractures: The study of osteoporotic fractures. The study of osteoporotic fractures research Group. *Journal of the American Geriatrics Society*. 1993 Nov; **41**(11):1226-1234

- [49] Diagnosis of occult fractures about the hip. Magnetic resonance imaging compared with bone-scanning. *The Journal of Bone and Joint Surgery. American Volume*. 1993 Mar;**75**(3):395-401
- [50] Hollis S, Lecky F, Yates DW, Woodford M. The effect of pre-existing medical conditions and age on mortality after injury. *Journal of Trauma*. 2006 Nov;**61**(5):1255-1260
- [51] Kirshenbom D, Ben-Zaken Z, Albilya N, Niyibizi E, Bala M. Older age, comorbid illnesses, and injury severity affect immediate outcome in elderly trauma patients. *Journal of Emergencies, Trauma and Shock*. 2017 Jul-Sep;**10**(3):146-150. DOI: 10.4103/JETS.JETS\_62\_16
- [52] Woolcott JC, Richardson KJ, Wiens MO, Patel B, Marin J, Khan KM, Marra CA. Meta-analysis of the impact of 9 medication classes on falls in elderly persons. *Archives of Internal Medicine*. 2009 Nov 23;**169**(21):1952-1960. DOI: 10.1001/archinternmed.2009.357
- [53] Field TS, Gurwitz JH, Harrold LR, Rothschild J, DeBellis KR, Seger AC, Auger JC, Garber LA, Cadoret C, Fish LS, Garber LD, Kelleher M, Bates DW. Risk factors for adverse drug events among older adults in the ambulatory setting. *Journal of the American Geriatrics Society*. 2004 Aug;**52**(8):1349-1354
- [54] Sardar P, Chatterjee S, Wu WC, Lichstein E, Ghosh J, Aikat S, Mukherjee D. New oral anticoagulants are not superior to warfarin in secondary prevention of stroke or transient ischemic attacks, but lower the risk of intracranial bleeding: Insights from a meta-analysis and indirect treatment comparisons. *PLoS One*. 2013 Oct 25;**8**(10):e77694. DOI: 10.1371/journal.pone.0077694. eCollection 2013
- [55] Kailas SD, Thambuluru SR. Efficacy and safety of direct oral anticoagulants compared to warfarin in prevention of thromboembolic events among elderly patients with atrial fibrillation. *Cureus*. 2016 Oct 18;**8**(10):e836
- [56] Batchelor JS, Grayson A. A meta-analysis to determine the effect of preinjury antiplatelet agents on mortality in patients with blunt head trauma. *British Journal of Neurosurgery*. 2013 Feb;**27**(1):12-28. DOI: 10.3109/02688697.2012.705361. Epub 2012 Aug 17
- [57] Bakheet MF, Pearce LA, Hart RG. Effect of addition of clopidogrel to aspirin on subdural hematoma: Meta-analysis of randomized clinical trials. *International Journal of Stroke*. 2015 Jun;**10**(4):501-505. DOI: 10.1111/ijss.12419. Epub 2014 Dec 3
- [58] Calland JF, Ingraham AM, Martin N, et al. Evaluation and management of geriatric trauma: An eastern Association for the Surgery of trauma practice management guideline. *Journal of Trauma and Acute Care Surgery*. 2012;**73**:S345
- [59] Roberts I, Shakur H, Coats T, Hunt B, Balogun E, Barnetson L, Cook L, Kawahara T, Perel P, Prieto-Merino D, Ramos M, Cairns J, Guerriero C. The CRASH-2 trial: A randomised controlled trial and economic evaluation of the effects of tranexamic acid on death, vascular occlusive events and transfusion requirement in bleeding trauma patients. *Health Technology Assessment*. 2013 Mar;**17**(10):1-79. DOI: 10.3310/hta17100

- [60] Kamel HK, Phlavan M, Malekgoudarzi B, Gogel P, Morley JE. Utilizing pain assessment scales increases the frequency of diagnosing pain among elderly nursing home residents. *Journal of Pain and Symptom Management*. 2001 Jun; **21**(6):450-455
- [61] Rodger KTM, Greasley-Adams C, Hodge Z, Reynish E. Expert opinion on the management of pain in hospitalised older patients with cognitive impairment: A mixed methods analysis of a national survey. *BMC Geriatrics*. 2015;**15**:56. DOI: 10.1186/s12877-015-0056-6
- [62] Institute of Medicine. *Confronting Chronic Neglect. The Education and Training of Health Professionals on Family Violence*. Washington, DC: The National Academies Press; 2002
- [63] Wallace RB, Bonnie RJ, editors. *Elder Mistreatment: Abuse, Neglect, and Exploitation in an Aging America*. Washington, DC: National Academies Press; 2003. pp. 339-381
- [64] Tintinalli JE, Stephan Stapczynski J, John ma O, Cline D, Meckler GD, Yealy DM. *Tintinallis emergency medicine: A comprehensive study guide*. New York: McGraw-Hill Education; 2016
- [65] Adapted from American College of Surgeons. *Resources for the optimal care of the injured patient*. Chicago, IL: American College of Surgeons; 2011
- [66] Dyer CB, Connolly MT, McFeeley P. The clinical and medical forensics of elder abuse and neglect. In: Wallace RB, Bonnie RJ, editors. *Elder mistreatment: Abuse, neglect, and exploitation in an aging America*. Washington, DC: National Academies Press; 2003:339-381
- [67] Lachs MS, Pillemer KA. Elder Abuse. *The New England Journal of Medicine*. 2015 Nov 12;**373**(20):1947-1956. DOI:10.1056/NEJMra1404688