

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Leishmaniases in West Africa: Past and Current

Abdoulaye Kassoum Kone, Mahamadou A. Thera,
Babacar Faye and Ogobara K. Doumbo†

Additional information is available at the end of the chapter

<http://dx.doi.org/10.5772/intechopen.77009>

Abstract

Leishmaniases are vector-borne diseases. Cutaneous leishmaniasis (CL) is endemic in West Africa. Sporadic and anecdotal cases of visceral leishmaniasis (VL) have been reported in the past. Recent data showed the changing of epidemiology of leishmaniases in West Africa, with the occurrence of outbreak of CL due to *Leishmania major* in urban and rural areas. CL is transmitted by *Phlebotomus duboscqi*. The role of *Sergentomyia (Spelaemyia) darlingi* as vector in rural areas has been evoked but not confirmed. Cases of VL due to *Leishmania spp.* have been described in West Africa; however, parasites species were not identified and dogs were suspected to be the reservoir. No humans' case of symptomatic VL due to *L. infantum* has been described in West Africa. Recent data in rural areas of Senegal confirmed dog as reservoir of *L. infantum*. In the same study in Senegal, *Sergentomyia* sandflies were found infected with *L. infantum*, indicating a possible role in leishmaniasis transmission. Coinfection leishmaniases-HIV is reported but rare. In this chapter, we included most recent publications and propose an updated landscape of CL and VL epidemiology in West Africa.

Keywords: leishmaniases, epidemiology, West Africa

1. Introduction

Leishmaniases are anthroponoses common in animals and humans. Leishmaniases are endemics in 98 countries. The annual incidence is 0.7–1.2 million cases of cutaneous leishmaniasis (CL) and 0.2–0.4 million of cases of visceral leishmaniasis (VL) causing 20,000–40,000 deaths annually [1]. In sub-Saharan African region, the estimated annual incidence of CL was between 770 and 1500 cases. *Leishmania* parasites are characterized by their enzyme electrophoretic profile that defines zymodemes. *L. major* is the main parasite causing CL and its

zymodemes MON-26, MON-25, MON-17 and MON-117 have been identified in West Africa [2–4]. CL has been cited earlier as endemic in rural areas. However, outbreaks of CL in urban areas have been recently reported, showing a change in CL epidemiology [5]. VL presents a different epidemiological pattern in West Africa. Previous data described VL as a rare condition, with occurrence of sporadic and anecdotal cases in limited areas [6–8]. Recent data on VL in West Africa showed the occurrence of several cases of canine visceral leishmaniasis and asymptomatic human infections [9–11]. We will describe an updated dynamics of CL and VL characteristics in West Africa.

2. Burden and characteristics

2.1. Cutaneous leishmaniasis

The overall burden of CL is poorly characterized, due mainly to paucity of data. The frequency of CL in suspected patients was 78.4% in Mali [12], and Niger it was 66.7% [13]. In Burkina Faso, CL was perceived as a public health problem due to occurrence of outbreaks with an average incidence of 0.1% [14] and hospital frequency of 1.1% [15]. In Senegal, the frequency of CL in hospital based study was 38 cases over 4 years (9.5 cases per year) [16].

In Mali, recent positive LST survey showed a higher prevalence (49.9%) in Diema, Kayes region [17]. LST prevalence in Kayes was 25.7% in 1969 [18]. This difference in LST prevalence in the region of Kayes over a period of more than 45 years is likely to indicate an increasing trend in CL transmission.

L. major is the main species of *Leishmania* reported in Mali and has been identified in humans as a causative agent of CL [19]. The reservoirs of *L. major* in West Africa are rodents in Senegal. *Mastomys erythrolucus*, *Tatera gambiana* and *Arvicanthus niloticus* were found infected by *L. major* [20, 21]. *Phlebotomus duboscqi* previously was cited as vector of CL. *L. major* DNA was identified recently in *P. duboscqi* confirming its role in CL transmission in Mali [22, 23]. *Sergentomyia (Spelaeomyia) darlingi* may also play a role in *L. major* transmission [22]. Other species of *Leishmania* causing CL such as *L. tropica* has been identified in *S. dissimillima*, *S. ingrami*, *S. simillima*, *S. dissimillima*, and *S. hamoni* in Ghana [24], but this species has been not identified in humans in West Africa.

Recent findings revealed *L. infantum* in a HIV-positive child suffering from CL in Senegal [25]. A new species of *Leishmania* classified as *Leishmania enrietti* complex was found in humans in Ghana [26]. These findings call to strengthen CL diagnosis and stimulate efforts to determine the causative *Leishmania* species in human infections.

The enzymes electrophoresis analysis identified several strains of *L. major* in West Africa. The most frequent strain in Mali was MON-26 and MON-74 was the more frequent in Burkina Faso [2, 4, 27]. Travelers visiting endemic areas in West Africa are at risk of getting infected with *Leishmania* [28].

In West African countries (**Figure 1**), CL outbreaks may occur in rural areas where health care centers personnel are not well trained for diagnosis or case management. Often, an

investigation post-outbreak is conducted to determine the *Leishmania spp.*, the vectors and reservoirs involved [5, 29]. In urban areas [30], outbreaks are occurring in larger population. Rapid urbanization is considered as a favoring factor that also makes uneasy outbreak control.

Coinfection CL-HIV is rare in West Africa. In Mali, the frequency of coinfection was 1% [31]. Coinfection CL-HIV has been reported in Burkina Faso and Ghana [32, 33]. In patients with coinfection, diffuse CL, mucosal involvement and bone marrow invasion have been reported [16, 34–36].

2.2. Visceral leishmaniasis

In West Africa, VL has been described, and previous data have shown the scarcity of disease (Figure 1). For several years, anecdotal and sporadic cases of VL were reported. Up to today, the data reviewed identified most of the clinical cases of VL in Ivory Coast and Niger [6–8]. In the Gambia, a case of VL has been reported in humans and in dogs [37, 38]. Underreporting of diseases is a known feature of West African health care system. This is also true for VL. The underreporting of VL in West Africa could be favored by the absence of appropriate biological diagnosis and the absence of specificity of VL clinical symptoms. VL cases may be confounded with others frequent parasitic diseases such as malaria or schistosomiasis. It is also assumed that the parasite strains found in West Africa are less virulent than those found elsewhere (Asia and East Africa). The strain identified in Senegal and likely those in West Africa are coming from Mediterranean basin [39].



Figure 1. Modified map of West Africa from map (http://d-maps.com/carte.php?num_car=36688&lang=fr) and status of leishmaniasis endemicity in West Africa (WHO weekly report 2017) [45]. ●: Cutaneous leishmaniasis endemic; ●: Cutaneous leishmaniasis previously reported; ▲: Visceral leishmaniasis endemic; ▲: Visceral leishmaniasis previously reported.

Parasite species identified in West Africa is *L. infantum* using serology method in asymptomatic Senegalese [11]. *L. donovani* has not yet been identified in West Africa. Cases of VL encountered in Niger and Ivory Coast had their parasite identified by microscopy [6–8]. It is known that microscopy cannot distinguish between *Leishmania* species. For species, diagnosis serology or molecular biology is required.

Sandflies of the genus *Sergentomyia* (*Se. dubia*, *Se. schwetzi* and *Se. magna*) have been found infected with *L. infantum* in Mont-Roland district in Senegal [40]. This raises the possibility that *Sergentomyia* spp. may be involved in VL transmission in Senegal.

Canine visceral leishmaniasis is well described in West Africa. Recent studies in domestic dogs showed that *L. infantum* was the causal pathogen in Senegal, in Burkina Faso and Nigeria [9, 10, 41].

Geographic diversity: the area of transmission could be wider, therefore underestimating the cases of leishmaniasis. In Senegal, previous studies have shown that the vectors *Phlebotomus* or *Sergentomyia* are found in many areas of the country (Kédougou, South East, Keur Moussa in Dakar region, Ferlo area [42]. Environmental changes are risk factors of explosion.

3. Management and control

VL is rarely encountered in West Africa. Most cases of human Leishmaniasis are cutaneous leishmaniasis. Often, CL cases are under-diagnosed, and their clinical management is poorly done. This is particularly true for cases encountered at peripheral health care centers with personnel poorly trained. For those encountered at referral health care facilities with good capacity for the diagnosis, treatment is available. In Mali, CL cases are referred to National Center for Diseases Control [CNAM acronym in French for Centre National d'Appui à Lutte contre la Maladie]. Treatment is done using either meglumine antimoniate locally or local thermotherapy [43]. In Burkina Faso, meglumine antimoniate is the first line treatment [15]. Treatment outcome is favorable with healing of lesions in 2–4 weeks. Treatment response in HIV coinfecting patients is also favorable but hampered by the frequency of relapses [44].

In rural areas where these treatments are not available dermatologists advice to clean skin lesions and apply tetracycline ointment until healing [43]. Meglumine antimoniate and amphotericin B have been used to treat VL in West Africa [6–8].

4. Conclusion

Compared to other endemic parts of the world, leishmaniasis are not very common in West Africa. CL is widely distributed in few West African countries such as Burkina Faso, Mali, Nigeria and Senegal. Urbanization is the main risk factor. Human VL human is sporadic in few countries. Also, VL affects more domestic dogs. Our review acknowledged the changing of CL epidemiology with more report of outbreaks and description of new parasite species in West

Africa. A surveillance system based on referral health care centers with training of health care personnel will help to better address clinical and diagnostic challenges imposed by leishmaniasis.

Conflict of interest

No conflict of interest declared.

Author details

Abdoulaye Kassoum Kone^{1*}, Mahamadou A. Thera¹, Babacar Faye² and Ogobara K. Doumbo^{1†}

*Address all correspondence to: fankone@icermali.org

1 Malaria Research and Training Center, Department of Epidemiology of Parasitic Diseases/ Faculty of Medicine, and Dentistry/ UMI-3189/ University of Science, Technique and Technology of Bamako, BP, Bamako, Mali

2 Service de Parasitologie-Mycologie, Faculté de Médecine, Université Cheikh Anta DIOP, Dakar, Sénégal

† Deceased

References

- [1] Alvar J, Velez ID, Bern C, Herrero M, Desjeux P, Cano J, et al. Leishmaniasis worldwide and global estimates of its incidence. PLoS One. 2012;**7**:e35671. DOI: 10.1371/journal.pone.0035671
- [2] Izri MA, Doumbo O, Belazzoug S, Pratlong F. Presence of *Leishmania major* MON-26 in Mali. Annales de Parasitologie Humaine et Comparée. 1989;**64**:510-511
- [3] Garin JP, Peyramond D, Piens MA, Rioux JA, Godfrey DG, Lanotte G, et al. Presence of *Leishmania major* Yakimoff and Schokhor, 1914 in Mali. Enzymatic identification of a strain of human origin. Annales de Parasitologie Humaine et Comparée. 1985;**60**:93-94
- [4] Pratlong F, Lami P, Ravel C, Balard Y, Dereure J, Serres G, et al. Geographical distribution and epidemiological features of old world *Leishmania infantum* and *Leishmania donovani* foci, based on the isoenzyme analysis of 2277 strains. Parasitology. 2013;**140**:423-434
- [5] Kone AK, Delaunay P, Djimdé AA, Thera MA, Giudice PD, Coulibaly D, Traoré K, Goita SM, Abathina A, Izri A, Marty P, Doumbo OK. Epidemiology of cutaneous leishmaniasis in five villages of Dogon country, Mali. Bulletin De La Societe De Pathologie Exotique. 2012 Feb;**105**(1):8-15

- [6] Eholié SP, Tanon AK, Folquet-Amorissani M, Doukouré B, Adoubryn KD, Yattara A, Bissagnéné E. Three new cases of visceral leishmaniasis in Côte d'Ivoire. *Bulletin De La Societe De Pathologie Exotique*. 2008 Feb;**101**(1):60-61
- [7] Kouassi B, Horo K, Achi VH, Adoubryn KD, Kakou ES, et al. Leishmaniose viscérale à Abidjan à propos de 3 observations. *La Medicina Tropical*. 2005;**65**:602-603
- [8] Djidingar D, Chippaux JP, Gragnic G, Tchani O, Meynard D, Julvez J. Visceral leishmaniasis in Niger: Six new parasitologically confirmed cases. *Bulletin De La Societe De Pathologie Exotique*. 1997;**90**(1):27-29
- [9] Faye B, Bañuls AL, Bucheton B, Dione MM, Bassanganam O, Hide M, Dereure J, Choisy M, Ndiaye JL, Konaté O, Claire M, Senghor MW, Faye MN, Sy I, Niang AA, Molez JF, Victoir K, Marty P, Delaunay P, Knecht R, Mellul S, Diedhiou S, Gaye O. Canine visceral leishmaniasis caused by *Leishmania infantum* in Senegal: Risk of emergence in humans? *Microbes and Infection*. 2010 Dec;**12**(14-15):1219-1225
- [10] Sangaré I, Djibougou Djibougou A, Yaméogo BK, Drabo F, Diabaté A, Banuls AL, Fournet F, Price H, Guiguemdé RT, Dabiré RK. First detection of *Leishmania infantum* in domestic dogs from Burkina Faso (West Africa). *Research Journal of Parasitology*. 2017;**12**(1):27-32
- [11] Faye B, Bucheton B, Bañuls AL, Senghor MW, Niang AA, Diedhiou S, Konaté O, Dione MM, Hide M, Mellul S, Knecht R, Delaunay P, Marty P, Gaye O. Seroprevalence of *Leishmania infantum* in a rural area of Senegal: Analysis of risk factors involved in transmission to humans. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2011 Jun;**105**(6):333-340
- [12] Kone AK, Niare DS, Thera MA, Kayentao K, Djimde A, Delaunay P, Kouriba B, Giudice PD, Izri A, Marty P, Doumbo OK. Epidemiology of the outbreak, vectors and reservoirs of cutaneous leishmaniasis in Mali: A systematic review and meta-analysis. *Asian Pacific Journal of Tropical Medicine*. 2016 Oct;**9**(10):985-990. DOI: 10.1016/j.apjtm.2016.07.025 Epub 2016 Aug 20
- [13] Develoux M, Blanc L, Garba S, Mamoudou HD, Warter A, Ravisse P. Cutaneous leishmaniasis in Niger. *The American Journal of Tropical Medicine and Hygiene*. 1990;**43**:29-30
- [14] Bamba S, Barro-Traoré F, Drabo MK, Gouba A, Traoré A, Guiguemdé TR. Epidemiological profile, clinical and therapeutic cutaneous leishmaniasis in the Department of Dermatology at University Hospital in Ouagadougou, Burkina Faso. *Revue Medicale De Bruxelles*. 2013 Sep–Oct;**34**(5):392-396
- [15] Bamba S, Gouba A, Drabo MK, Nezien D, Bougoum M, Guiguemdé TR. Epidemiological profile of cutaneous leishmaniasis: Retrospective analysis of 7444 cases reported from 1999 to 2005 at Ouagadougou, Burkina Faso. *The Pan African Medical Journal*. 2013 Mar **19**:14:108
- [16] Diadie S, Diatta BA, Ndiaye M, Seck NB, Diallo S, Niang SO, Dieng MT. Cutaneous leishmaniasis in Senegal: A series of 38 cases at the Aristide Le Dantec University Hospital in Dakar. *Medecine Et Sante Tropicales*. 2018 Feb **1**;**28**(1):106-108. DOI: 10.1684/mst.2017.0722

- [17] Traoré B, Oliveira F, Faye O, Dicko A, Coulibaly CA, Sissoko IM, Sibiry S, Sogoba N, Sangare MB, Coulibaly YI, Traore P, Traore SF, Anderson JM, Keita S, Valenzuela JG, Kamhawi S, Doumbia S. Prevalence of cutaneous Leishmaniasis in districts of high and low endemicity in Mali. *PLoS Neglected Tropical Diseases*. 2016 Nov 29;**10**(11):e0005141
- [18] Imperato PJ, Diakité S. Leishmaniasis in the Republic of Mali. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 1969;**63**(2):236-241
- [19] Paz C, Samake S, Anderson JM, Faye B, Traore P, Tall K, et al. *Leishmania major*, the predominant *Leishmania* species responsible for cutaneous leishmaniasis in Mali. *The American Journal of Tropical Medicine and Hygiene*. 2013;**88**:583-585
- [20] Dedet JP, Derouin F. Isolation of *Leishmania major* from *Mastomys erythroleucus* and *Tatera gambiana* in Senegal (West Africa). *Annals of Tropical Medicine and Parasitology*. 1979;**73**:433-437
- [21] Camerlynck P, Ranque P, Quilici M. Importance of systematic cultures and subcultures in research on natural viral reservoirs in cutaneous leishmaniasis. Apropos of the isolation of 5 strains of *Leishmania* in *Arvicanthus niloticus*. *Medecine Tropicale (Mars)*. 1967;**27**:89-92
- [22] Berdjane-Brouk Z, Koné AK, Djimdé AA, Charrel RN, Ravel C, Delaunay P, del Giudice P, Diarra AZ, Doumbo S, Goita S, Thera MA, Depaquit J, Marty P, Doumbo OK, Izri A. First detection of *Leishmania major* DNA in *Sergentomyia (Spelaemyia) darlingi* from cutaneous leishmaniasis foci in Mali. *PLoS One*. 2012;**7**(1):e28266
- [23] Anderson JM, Samake S, Jaramillo-Gutierrez G, Sissoko I, Coulibaly CA, Traoré B, Soucko C, Guindo B, Diarra D, Fay MP, Lawyer PG, Doumbia S, Valenzuela JG, Kamhawi S. Seasonality and prevalence of *Leishmania major* infection in *Phlebotomus duboscqi* Neveu-Lemaire from two neighboring villages in Central Mali. *PLoS Neglected Tropical Diseases*. 2011 May 10;**5**(5):e1139
- [24] Nzelu CO, Kato H, Puplampu N, Desewu K, Odoom S, Wilson MD, Sakurai T, Katakura K, Boakye DA. First detection of *Leishmania tropica* DNA and Trypanosoma species in *Sergentomyia* sand flies (Diptera: Psychodidae) from an outbreak area of cutaneous leishmaniasis in Ghana. *PLoS Neglected Tropical Diseases*. 2014 Feb 6;**8**(2):e2630
- [25] Diatta BA, Diallo M, Diadie S, Faye B, Ndiaye M, Hakim H, Diallo S, Seck B, Niang SO, Kane A, Dieng MT. Cutaneous leishmaniasis due to *Leishmania infantum* associated with HIV. *Annales de Dermatologie et de Vénérologie*. 2016 Oct;**143**(10):625-628
- [26] Kwakye-Nuako G, Mosore MT, Duplessis C, Bates MD, Puplampu N, Mensah-Attipoe I, Desewu K, Afegbe G, Asmah RH, Jamjoom MB, Ayeh-Kumi PF, Boakye DA, Bates PA. First isolation of a new species of *Leishmania* responsible for human cutaneous leishmaniasis in Ghana and classification in the *Leishmania enriettii* complex. *International Journal for Parasitology*. 2015 Sep;**45**(11):679-684. DOI: 10.1016/j.ijpara.2015.05.001. Epub 2015 Jun 19
- [27] Guiguemdé TR, Sawadogo NO, Botero S, Traore KL, Nezien D, Nikiema L, et al. *Leishmania major* and HIV coinfection in Burkina Faso. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 2003;**97**:168-169

- [28] Kelly P, Baudry T, Peyron F. Imported cutaneous leishmaniasis in a short-term traveler returning from Central Mali the role of PCR. *Travel Medicine and Infectious Disease*. 2012;**10**:97-100
- [29] Kweku MA, Odoom S, Puplampu N, Desewu K, Nuako GK, Gyan B, Raczniak G, Kronmann KC, Koram K, Botero S, Boakye D, Akuffo H. An outbreak of suspected cutaneous leishmaniasis in Ghana: Lessons learnt and preparation for future outbreaks. *Global Health Action*. 2011;**4**:5527. DOI: 10.3402/gha.v4i0.5527
- [30] Traore KS, Sawadogo NO, Traoré A, Ouedraogo JB, Traoré KL, Guiguemdé TR. Preliminary study of cutaneous leishmaniasis in the town of Ouagadougou from 1996 to 1998. *Bulletin De La Societe De Pathologie Exotique*. 2001;**94**:52-55
- [31] Mahé A, Bobin P, Coulibaly S, Tounkara A. Skin diseases disclosing human immunodeficiency virus infection in Mali *Annales de Dermatologie et de Vénérologie*. 1997;**124**:144-150
- [32] Niamba P, Traoré A, Goumbri-Lompo O, Labrèze C, Traoré-Barro F, Bonkougou M, et al. Leishmaniose cutanée chez les malades infectés par le VIH. *Annales de Dermatologie et de Vénérologie*. 2006;**133**:537-542
- [33] Lartey M, Adusei L, Hanson-Nortey L, Addy J. Coinfection of cutaneous Leishmaniasis and HIV infection. *Ghana Medical Journal*. 2006 Sep;**40**(3):110-112
- [34] Niamba P, Goumbri-Lompo O, Traoré A, Barro-Traoré F, Soudré RT. Diffuse cutaneous leishmaniasis in an HIV-positive patient in western Africa. *The Australasian Journal of Dermatology*. 2007 Feb;**48**(1):32-34
- [35] Barro-Traoré F, Preney L, Traoré A, Darie H, Tapsoba P, Bassolé A, Sawadogo S, Niamba P, Grosshans E, Geniaux M. Cutaneous leishmaniasis due to *Leishmania major* involving the bone marrow in an AIDS patient in Burkina Faso. *Annales de Dermatologie et de Vénérologie*. 2008 May;**135**(5):380-383
- [36] Ndiaye PB, Develoux M, Dieng MT, Huerre M. Diffuse cutaneous leishmaniasis and acquired immunodeficiency syndrome in a Senegalese patient. *Bulletin De La Societe De Pathologie Exotique*. 1996;**89**(4):282-286
- [37] Desjeux P, Bryan JH, Martin-Saxton P. Leishmaniasis in the Gambia. 2. A study of possible vectors and animal reservoirs, with the first report of a case of canine leishmaniasis in the Gambia. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 1983;**77**(2):143-148
- [38] Walters JH. A case of indigenous Kala-azar in the Gambia. *Transactions of the Royal Society of Tropical Medicine and Hygiene*. 1949 Nov;**43**(3):287-292. pl
- [39] Cassan C, Dione MM, Dereure J, Diedhiou S, Bucheton B, Hide M, Kako C, Gaye O, Senghor M, Niang AA, Bañuls AL, Faye B. First insights into the genetic diversity and origin of *Leishmania infantum* in Mont Rolland (Thiès region, Senegal). *Microbes and Infection*. 2016 Jun;**18**(6):412-420

- [40] Senghor MW, Niang AA, Depaquit J, Ferté H, Faye MN, Elguero E, Gaye O, Alten B, Perktas U, Cassan C, Faye B, Bañuls AL. Transmission of *Leishmania infantum* in the canine leishmaniasis focus of Mont-Rolland, Senegal: Ecological, parasitological and molecular evidence for a possible role of *Sergentomyia* sand flies. PLoS Neglected Tropical Diseases. 2016;2(10):11, e0004940
- [41] Adediran OA, Kolapo TU, Uwalaka EC. Seroprevalence of canine leishmaniasis in Kwara, Oyo and Ogun states of Nigeria. Journal of Parasitic Diseases. 2016 Jun;40(2):510-514. DOI: 10.1007/s12639-014-0535-2. Epub 2014 Sep 5
- [42] Ba Y, Trouillet J, Thonnon J, Fontenille D. Phlebotomus of Senegal: Survey of the fauna in the region of Kedougou. Isolation of arbovirus. Bulletin De La Societe De Pathologie Exotique. 1999 May;92(2):131-135
- [43] Tall K. Etude épidémiologique et prise en charge de la leishmaniose cutanée à Bamako et dans deux villages endémiques du Mali [thesis]. Faculty of Medicine and Dentistry; 2008. http://indexmedicus.afro.who.int/iah/fulltext/Thesis_Bamako/05P41.PDF
- [44] Niamba P, Traoré A, Goumbri-Lompo O, Labrèze C, Traoré-Barro F, Bonkougou M, Ilboudo L, Gaulier A, Soudré BR. Cutaneous leishmania in HIV patient in Ouagadougou: Clinical and therapeutic aspects. Annales de Dermatologie et de Vénérologie. 2006 Jun-Jul;133(6-7):537-542
- [45] WHO. Weekly Epidemiological Record. No. 38; 2017, Vol. 92. pp. 557-572. Available at: http://www.who.int/leishmaniasis/resources/REH_38_TABLEAU_S1_S2_Version_finale.pdf?ua=1

