

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Some Topological Properties of Intuitionistic Fuzzy Normed Spaces

Vakeel Ahmad Khan, Hira Fatima and Mobeen Ahmad

Abstract

In 1986, Atanassov introduced the concept of intuitionistic fuzzy set theory which is based on the extensions of definitions of fuzzy set theory given by Zadeh. This theory provides a variable model to elaborate uncertainty and vagueness involved in decision making problems. In this chapter, we concentrate our study on the ideal convergence of sequence spaces with respect to intuitionistic fuzzy norm and discussed their topological and algebraic properties.

Keywords: ideal, intuitionistic fuzzy normed spaces, Orlicz function compact operator, I-convergence

1. Introduction

In recent years, the fuzzy theory has emerged as the most active area of research in many branches of mathematics, computer and engineering [1]. After the excellent work of Zadeh [2], a large number of research work have been done on fuzzy set theory and its applications as well as fuzzy analogues of the classical theories. It has a wide number of applications in various fields such as population dynamics [3], nonlinear dynamical system [4], chaos control [5], computer programming [6], etc. In 2006, Saadati and Park [7] introduced the concept of intuitionistic fuzzy normed spaces after that the concept of statistical convergence in intuitionistic fuzzy normed space was studied for single sequence in [8]. The study of intuitionistic fuzzy topological spaces [9], intuitionistic fuzzy 2-normed space [10] and intuitionistic fuzzy Zweier ideal convergent sequence spaces [11] are the latest developments in fuzzy topology.

First, let us recall some notions, basic definitions and concepts which are used in sequel.

Definition 1.1. (See Ref. [7]). The five-tuple $(X, \mu, \nu, *, \diamond)$ is said to be an intuitionistic fuzzy normed space (for short, IFNS) if X is a vector space, $*$ is a continuous t-norm, \diamond is a continuous t-conorm, and μ and ν are fuzzy sets on $X \times (0, \infty)$ satisfying the following conditions for every $x, y \in X$ and $s, t > 0$:

- (a) $\mu(x, t) + \nu(x, t) \leq 1$,
- (b) $\mu(x, t) > 0$,
- (c) $\mu(x, t) = 1$ if and only if $x = 0$,
- (d) $\mu(\alpha x, t) = \mu\left(x, \frac{t}{|\alpha|}\right)$ for each $\alpha \neq 0$,

- (e) $\mu(x, t) * \mu(y, s) \leq \mu(x + y, t + s)$,
- (f) $\mu(x, \cdot) : (0, \infty) \rightarrow [0, 1]$ is continuous,
- (g) $\lim_{t \rightarrow \infty} \mu(x, t) = 1$ and $\lim_{t \rightarrow 0} \mu(x, t) = 0$,
- (h) $\nu(x, t) < 1$,
- (i) $\nu(x, t) = 0$ if and only if $x = 0$,
- (j) $\nu(\alpha x, t) = \nu\left(x, \frac{t}{|\alpha|}\right)$ for each $\alpha \neq 0$,
- (k) $\nu(x, t) \diamond \nu(y, s) \geq \nu(x + y, t + s)$,
- (l) $\nu(x, \cdot) : (0, \infty) \rightarrow [0, 1]$ is continuous,
- (m) $\lim_{t \rightarrow \infty} \nu(x, t) = 0$ and $\lim_{t \rightarrow 0} \nu(x, t) = 1$.

In this case (μ, ν) is called an intuitionistic fuzzy norm.

Example 1.1. Let $(X, \|\cdot\|)$ be a normed space. Denote $a * b = ab$ and $a \diamond b = \min(a + b, 1)$ for all $a, b \in [0, 1]$ and let μ_0 and ν_0 be fuzzy sets on $X \times (0, \infty)$ defined as follows:

$$\mu_0(x, t) = \frac{t}{t + \|x\|}, \quad \text{and} \quad \nu_0(x, t) = \frac{\|x\|}{t + \|x\|}$$

for all $t \in \mathbb{R}^+$. Then $(X, \mu, \nu, *, \diamond)$ is an intuitionistic fuzzy normed space.

Definition 1.2. Let $(X, \mu, \nu, *, \diamond)$ be an IFNS. Then a sequence $x = (x_k)$ is said to be convergent to $L \in X$ with respect to the intuitionistic fuzzy norm (μ, ν) if, for every $\varepsilon > 0$ and $t > 0$, there exists $k_0 \in \mathbb{N}$ such that $\mu(x_k - L, t) > 1 - \varepsilon$ and $\nu(x_k - L, t) < \varepsilon$ for all $k \geq k_0$. In this case we write $(\mu, \nu)\text{-}\lim x = L$.

In 1951, the concept of statistical convergence was introduced by Steinhaus [12] and Fast [13] in their papers “Sur la convergence ordinaire et la convergence asymptotique” and “Sur la convergence statistique,” respectively. Later on, in 1959, Schoenberg [14] reintroduced this concept. It is a very useful functional tool for studying the convergence of numerical problems through the concept of density. The concept of ideal convergence, which is a generalization of statistical convergence, was introduced by Kostyrko et al. [15] and it is based on the ideal I as a subsets of the set of positive integers and further studied in [16–20].

Let X be a non-empty set then a family $I \subset 2^X$ is said to be an **ideal** in X if $\emptyset \in I$, I is additive, i.e., for all $A, B \in I \Rightarrow A \cup B \in I$ and I is hereditary, i.e., for all $A \in I, B \subseteq A \Rightarrow B \in I$. A non empty family of sets $\mathcal{F} \subset 2^X$ is said to be a **filter** on X if for all $A, B \in \mathcal{F}$ implies $A \cap B \in \mathcal{F}$ and for all $A \in \mathcal{F}$ with $A \subseteq B$ implies $B \in \mathcal{F}$. An ideal $I \subset 2^X$ is said to be **nontrivial** if $I \neq 2^X$, this non trivial ideal is said to be admissible if $I \supseteq \{\{x\} : x \in X\}$ and is said to be **maximal** if there cannot exist any nontrivial ideal $J \neq I$ containing I as a subset. For each ideal I , there is a filter $\mathcal{F}(I)$ called as filter associate with ideal I , that is (see [15]),

$$\mathcal{F}(I) = \{K \subseteq X : K^c \in I\}, \quad \text{where } K^c = X \setminus K. \quad (1)$$

A sequence $x = (x_k) \in \omega$ is said to be **I -convergent** [21, 22] to a number L if for every $\varepsilon > 0$, we have $\{k \in \mathbb{N} : |x_k - L| \geq \varepsilon\} \in I$. In this case, we write $I\text{-}\lim_{xk} = L$.

2. IF-ideal convergent sequence spaces using compact operator

This section consists of some double sequence spaces with respect to intuitionistic fuzzy normed space and study the fuzzy topology on the said spaces. First we recall some basic definitions on compact operator.

Definition 2.1. (See [23]). Let X and Y be two normed linear spaces and $T : \mathcal{D}(T) \rightarrow Y$ be a linear operator, where $\mathcal{D} \subset X$. Then, the operator T is said to be bounded, if there exists a positive real k such that

$$\|Tx\| \leq k\|x\|, \text{ for all } x \in \mathcal{D}(T).$$

The set of all bounded linear operators $\mathcal{B}(X, Y)$ [24] is a normed linear spaces normed by

$$\|T\| = \sup_{x \in X, \|x\|=1} \|Tx\|$$

and $\mathcal{B}(X, Y)$ is a Banach space if Y is a Banach space.

Definition 2.2. (See [23]). Let X and Y be two normed linear spaces. An operator $T : X \rightarrow Y$ is said to be a compact linear operator (or completely continuous linear operator), if

(i) T is linear,

(ii) T maps every bounded sequence (x_k) in X on to a sequence $(T(x_k))$ in Y which has a convergent subsequence.

The set of all compact linear operators $\mathcal{C}(X, Y)$ is a closed subspace of $\mathcal{B}(X, Y)$ and $\mathcal{C}(X, Y)$ is Banach space, if Y is a Banach space.

In 2015, Khan et al. [11] introduced the following sequence spaces:

$$\mathcal{Z}_{(\mu, \nu)}^I = \left\{ (x_k) \in \omega : \left\{ k : \mu(x'_k - L, t) \leq 1 - \varepsilon \text{ or } \nu(x'_k - L, t) \geq \varepsilon \right\} \in I \right\},$$

$$\mathcal{Z}_{0(\mu, \nu)}^I = \left\{ (x_k) \in \omega : \left\{ k : \mu(x'_k, t) \leq 1 - \varepsilon \text{ or } \nu(x'_k, t) \geq \varepsilon \right\} \in I \right\}.$$

Motivated by this, we introduce the following sequence spaces with the help of compact operator in intuitionistic fuzzy normed spaces:

$$\mathcal{M}_{(\mu, \nu)}^I(T) = (x_k) \in \ell_\infty : \left\{ k : \mu(T(x_k) - L, t) \leq 1 - \varepsilon \text{ or } \nu(T(x_k) - L, t) \geq \varepsilon \in I \right\} \quad (2)$$

$$\mathcal{M}_{0(\mu, \nu)}^I(T) = (x_k) \in \ell_\infty : \left\{ k : \mu(T(x_k), t) \leq 1 - \varepsilon \text{ or } \nu(T(x_k), t) \geq \varepsilon \in I \right\}. \quad (3)$$

Here, we also define an open ball with center x and radius r with respect to t as follows:

$$\mathcal{B}_x(r, t)(T) = (y_k) \in \ell_\infty : \left\{ k : \mu(T(x_k) - T(y_k), t) \leq 1 - \varepsilon \text{ or } \nu(T(x_k) - T(y_k), t) \geq \varepsilon \in I \right\}. \quad (4)$$

Now, we are ready to state and prove our main results. This theorem is based on the linearity of new define sequence spaces which is stated as follows.

Theorem 2.1. The sequence spaces $\mathcal{M}_{(\mu, \nu)}^I(T)$ and $\mathcal{M}_{0(\mu, \nu)}^I(T)$ are linear spaces.

Proof. Let $x = (x_k), y = (y_k) \in \mathcal{M}_{(\mu, \nu)}^I(T)$ and α, β be scalars. Then for a given $\varepsilon > 0$, we have the sets:

$$P_1 = \left\{ k : \mu\left(T(x_k) - L_1, \frac{t}{2|\alpha|}\right) \leq 1 - \varepsilon \text{ or } \nu\left(T(x_k) - L_1, \frac{t}{2|\alpha|}\right) \geq \varepsilon \right\} \in I;$$

$$P_2 = \left\{ k : \mu\left(T(y_k) - L_2, \frac{t}{2|\beta|}\right) \leq 1 - \varepsilon \text{ or } \nu\left(T(y_k) - L_2, \frac{t}{2|\beta|}\right) \geq \varepsilon \right\} \in I.$$

This implies

$$P_1^c = \left\{ k : \mu\left(T(x_k) - L_1, \frac{t}{2|\alpha|}\right) > 1 - \varepsilon \text{ or } \nu\left(T(x_k) - L_1, \frac{t}{2|\alpha|}\right) < \varepsilon \right\} \in \mathcal{F}(I);$$

$$P_2^c = \left\{ k : \mu \left(T(y_k) - L_2, \frac{t}{2|\beta|} \right) > 1 - \varepsilon \text{ or } \nu \left(\left(T(y_k) - L_2, \frac{t}{2|\beta|} \right) < \varepsilon \right) \right\} \in \mathcal{F}(I).$$

Now, we define the set $P_3 = P_1 \cup P_2$, so that $P_3 \in I$. It shows that P_3^c is a non-empty set in $\mathcal{F}(I)$. We shall show that for each $(x_k), (y_k) \in \mathcal{M}_{(\mu, \nu)}^I(T)$.

$$P_3^c \subset \{ k : \mu((\alpha T(x_k) + \beta T(y_k)) - (\alpha L_1 + \beta L_2), t) > 1 - \varepsilon \text{ or } \nu((\alpha T(x_k) + \beta T(y_k)) - (\alpha L_1 + \beta L_2), t) < \varepsilon \}.$$

Let $m \in P_3^c$, in this case

$$\mu \left(T(x_m) - L_1, \frac{t}{2|\alpha|} \right) > 1 - \varepsilon \text{ or } \nu \left(T(x_m) - L_1, \frac{t}{2|\alpha|} \right) < \varepsilon$$

and

$$\mu \left(T(y_m) - L_2, \frac{t}{2|\beta|} \right) > 1 - \varepsilon \text{ or } \nu \left(T(y_m) - L_2, \frac{t}{2|\beta|} \right) < \varepsilon.$$

Thus, we have

$$\begin{aligned} & \mu((\alpha T(x_m) + \beta T(y_m)) - (\alpha L_1 + \beta L_2), t) \\ & \geq \mu \left(\alpha T(x_m) - \alpha L_1, \frac{t}{2} \right) * \mu \left(\beta T(y_m) - \beta L_2, \frac{t}{2} \right) \\ & = \mu \left(T(x_m) - L_1, \frac{t}{2|\alpha|} \right) * \mu \left(T(y_m) - L_2, \frac{t}{2|\beta|} \right) \\ & > (1 - \varepsilon) * (1 - \varepsilon) = 1 - \varepsilon. \end{aligned}$$

and

$$\begin{aligned} & \nu((\alpha T(x_m) + \beta T(y_m)) - (\alpha L_1 + \beta L_2), t) \\ & \leq \nu \left(\alpha T(x_m) - \alpha L_1, \frac{t}{2} \right) \diamond \nu \left(\beta T(y_m) - \beta L_2, \frac{t}{2} \right) \\ & = \nu \left(T(x_m) - L_1, \frac{t}{2|\alpha|} \right) \diamond \nu \left(T(y_m) - L_2, \frac{t}{2|\beta|} \right) \\ & < \varepsilon \diamond \varepsilon = \varepsilon. \end{aligned}$$

This implies that

$$P_3^c \subset \{ k : \mu((\alpha T(x_k) + \beta T(y_k)) - (\alpha L_1 + \beta L_2), t) > 1 - \varepsilon \text{ or } \nu((\alpha T(x_k) + \beta T(y_k)) - (\alpha L_1 + \beta L_2), t) < \varepsilon \}.$$

Therefore, the sequence space $\mathcal{M}_{(\mu, \nu)}^I(T)$ is a linear space.

Similarly, we can proof for the other space. \square

In the following theorems, we discussed the convergence problem in the said sequence spaces. For this, firstly we have to discuss about the topology of this space. Define

$$\begin{aligned} \tau_{(\mu, \nu)}^I(T) = A \subset \mathcal{M}_{(\mu, \nu)}^I(T) : & \text{ for each } x \in A \text{ there exists } t > 0 \\ & \text{ and } r \in (0, 1) \text{ such that } \mathcal{B}_x(r, t)(T) \subset A. \end{aligned}$$

Then $\tau_{(\mu, \nu)}^I(T)$ is a topology on $\mathcal{M}_{(\mu, \nu)}^I(T)$.

Theorem 2.2. Let $\mathcal{M}_{(\mu,\nu)}^I(T)$ is an IFNS and $\tau_{(\mu,\nu)}^I(T)$ is a topology on $\mathcal{M}_{(\mu,\nu)}^I(T)$. Then a sequence $(x_k) \in \mathcal{M}_{(\mu,\nu)}^I(T)$, $x_k \rightarrow x$ if and only if $\mu(T(x_k) - T(x), t) \rightarrow 1$ and $\nu(T(x_k) - T(x), t) \rightarrow 0$ as $k \rightarrow \infty$.

Proof. Fix $t_0 > 0$. Suppose $x_k \rightarrow x$. Then for $r \in (0, 1)$, there exists $n_0 \in \mathbb{N}$ such that $(x_k) \in \mathcal{B}_x(r, t_0)(T)$ for all $k \geq n_0$. So, we have

$$\mathcal{B}_x(r, t_0)(T) = \{k : \mu(T(x_k) - T(x), t) \leq 1 - r \text{ or } \nu(T(x_k) - T(x), t_0) \geq r\} \in I,$$

such that $\mathcal{B}_x^c(r, t_0)(T) \in \mathcal{F}(I)$. Then $1 - \mu(T(x_k) - T(x), t_0) < r$ and $\nu(T(x_k) - T(x), t_0) < r$. Hence $\mu(T(x_k) - T(x), t_0) \rightarrow 1$ and $\nu(T(x_k) - T(x), t_0) \rightarrow 0$ as $k \rightarrow \infty$.

Conversely, if for each $t > 0$, $\mu(T(x_k) - T(x), t) \rightarrow 1$ and $\nu(T(x_k) - T(x), t) \rightarrow 0$ as $k \rightarrow \infty$, then for $r \in (0, 1)$, there exists $n_0 \in \mathbb{N}$, such that $1 - \mu(T(x_k) - T(x), t) < r$ and $\nu(T(x_k) - T(x), t) < r$, for all $k \geq n_0$. It shows that $\mu(T(x_k) - T(x), t) > 1 - r$ and $\nu(T(x_k) - T(x), t) < r$ for all $k \geq n_0$. Therefore $(x_k) \in \mathcal{B}_x^c(r, t)(T)$ for all $k \geq n_0$ and hence $x_k \rightarrow x$.

There are some facts that arise in connection with the convergence of sequences in these spaces. Let us proceed to the next theorem on Ideal convergence of sequences in these new define spaces.

Theorem 2.3. A sequence $x = (x_k) \in \mathcal{M}_{(\mu,\nu)}^I(T)$ is I -convergent if and only if for every $\varepsilon > 0$ and $t > 0$ there exists a number $N = N(x, \varepsilon, t)$ such that

$$\left\{N : \mu\left(T(x_N) - L, \frac{t}{2}\right) > 1 - \varepsilon \text{ or } \nu\left(T(x_N) - L, \frac{t}{2}\right) < \varepsilon\right\} \in \mathcal{F}(I).$$

Proof. Suppose that $I_{(\mu,\nu)} - \lim x = L$ and let $t > 0$. For a given $\varepsilon > 0$, choose $s > 0$ such that $(1 - \varepsilon) * (1 - \varepsilon) > 1 - s$ and $\varepsilon \diamond \varepsilon < s$. Then for each $x \in \mathcal{M}_{(\mu,\nu)}^I(T)$,

$$R = \left\{k : \mu\left(T(x_k) - L, \frac{t}{2}\right) \leq 1 - \varepsilon \text{ or } \nu\left(T(x_k) - L, \frac{t}{2}\right) \geq \varepsilon\right\} \in I,$$

which implies that

$$R^c = \left\{k : \mu\left(T(x_k) - L, \frac{t}{2}\right) > 1 - \varepsilon \text{ or } \nu\left(T(x_k) - L, \frac{t}{2}\right) < \varepsilon\right\} \in \mathcal{F}(I).$$

Conversely, let us choose $N \in R^c$. Then

$$\mu\left(T(x_N) - L, \frac{t}{2}\right) > 1 - \varepsilon \text{ or } \nu\left(T(x_N) - L, \frac{t}{2}\right) < \varepsilon.$$

Now, we want to show that there exists a number $N = N(x, \varepsilon, t)$ such that

$$\{k : \mu(T(x_k) - T(x_N), t) \leq 1 - s \text{ or } \nu(T(x_k) - T(x_N), t) \geq s\} \in I.$$

For this, we define for each $x \in \mathcal{M}_{(\mu,\nu)}^I(T)$

$$S = \{k : \mu(T(x_k) - T(x_N), t) \leq 1 - s \text{ or } \nu(T(x_k) - T(x_N), t) \geq s\} \in I.$$

So, we have to show that $S \subset R$. Let us suppose that $S \not\subset R$, then there exists $n \in S$ and $n \notin R$. Therefore, we have

$$\mu(T(x_n) - T(x_N), t) \leq 1 - s \text{ or } \mu\left(T(x_n) - L, \frac{t}{2}\right) > 1 - \varepsilon.$$

In particular $\mu(T(x_N) - L, \frac{t}{2}) > 1 - \varepsilon$. Therefore, we have

$$1 - s \geq \mu(T(x_n) - T(x_N), t) \geq \mu\left(T(x_n) - L, \frac{t}{2}\right) * \mu\left(T(x_N) - L, \frac{t}{2}\right) \geq (1 - \varepsilon) * (1 - \varepsilon) > 1 - s,$$

which is not possible. On the other hand

$$\nu(T(x_n) - T(x_N), t) \geq s \text{ or } \nu\left(T(x_n) - L, \frac{t}{2}\right) < \varepsilon.$$

In particular $\nu(T(x_N) - L, \frac{t}{2}) < \varepsilon$. So, we have

$$s \leq \nu(T(x_n) - T(x_N), t) \leq \nu\left(T(x_n) - L, \frac{t}{2}\right) \diamond \nu\left(T(x_N) - L, \frac{t}{2}\right) \leq \varepsilon \diamond \varepsilon < s,$$

which is not possible. Hence SCR . $R \in I$ which implies $S \in I$. □

3. IF-ideal convergent sequence spaces using Orlicz function

In this section, we have discussed the ideal convergence of sequences in Intuitionistic fuzzy I -convergent sequence spaces defined by compact operator and Orlicz function. We shall now define the concept of Orlicz function, which is basic definition in our work.

Definition 3.1. An Orlicz function is a function $F : [0, \infty) \rightarrow [0, \infty)$, which is continuous, non-decreasing and convex with $F(0) = 0$, $F(x) > 0$ for $x > 0$ and $F(x) \rightarrow \infty$ as $x \rightarrow \infty$. If the convexity of Orlicz function F is replaced by $F(x + y) \leq F(x) + F(y)$, then this function is called modulus function.

Remark 3.1. If F is an Orlicz function, then $F(\lambda x) \leq \lambda F(x)$ for all λ with $0 < \lambda < 1$.

In 2009, Mohiuddine and Lohani [18] introduced the concept of statistical convergence in intuitionistic fuzzy normed spaces in their paper published in Chaos, Solitons and Fractals. This motivated us to introduced some sequence spaces defined by compact operator and Orlicz function which are as follows:

$$\mathcal{M}_{(\mu, \nu)}^I(T, F) = \left\{ (x_k) \in \ell_\infty : \left\{ k : F\left(\frac{\mu(T(x_k) - L, t)}{\rho}\right) \leq 1 - \varepsilon \right. \right. \\ \left. \left. \text{or } F\left(\frac{\nu(T(x_k) - L, t)}{\rho}\right) \geq \varepsilon \right\} \in I \right\}; \quad (5)$$

$$\mathcal{M}_{0(\mu, \nu)}^I(T, F) = \left\{ (x_k) \in \ell_\infty : \left\{ k : F\left(\frac{\mu(T(x_k), t)}{\rho}\right) \leq 1 - \varepsilon \right. \right. \\ \left. \left. \text{or } F\left(\frac{\nu(T(x_k), t)}{\rho}\right) \geq \varepsilon \right\} \in I \right\}. \quad (6)$$

We also define an open ball with center x and radius r with respect to t as follows:

$$\mathcal{B}_x(r, t)(T, F) = \left\{ (y_k) \in \ell_\infty : \left\{ k : F\left(\frac{\mu(T(x_k) - T(y_k), t)}{\rho}\right) \leq 1 - \varepsilon \right. \right. \\ \left. \left. \text{or } F\left(\frac{\nu(T(x_k) - T(y_k), t)}{\rho}\right) \geq \varepsilon \right\} \in I \right\}. \quad (7)$$

We shall now consider some theorems of these sequence spaces and invite the reader to verify the linearity of these sequence spaces.

Theorem 3.1. Every open ball $\mathcal{B}_x(r, t)(T, F)$ is an open set in $\mathcal{M}_{(\mu, \nu)}^I(T, F)$.

Proof. Let $\mathcal{B}_x(r, t)(T, F)$ be an open ball with center x and radius r with respect to t . That is

$$\mathcal{B}_x(r, t)(T, F) = \left\{ y = (y_k) \in \mathcal{L}_\infty : \left\{ k : F\left(\frac{\mu(T(x_k) - T(y_k), t)}{\rho}\right) \leq 1 - r \right. \right. \\ \left. \left. \text{or } F\left(\frac{\nu(T(x_k) - T(y_k), t)}{\rho}\right) \geq r \right\} \in I \right\}.$$

Let $y \in \mathcal{B}_x^c(r, t)(T, F)$, then $F\left(\frac{\mu(T(x_k) - T(y_k), t)}{\rho}\right) > 1 - r$ and $F\left(\frac{\nu(T(x_k) - T(y_k), t)}{\rho}\right) < r$. Since $F\left(\frac{\mu(T(x_k) - T(y_k), t)}{\rho}\right) > 1 - r$, there exists $t_0 \in (0, t)$ such that $F\left(\frac{\mu(T(x_k) - T(y_k), t_0)}{\rho}\right) > 1 - r$ and $F\left(\frac{\nu(T(x_k) - T(y_k), t_0)}{\rho}\right) < r$.

Putting $r_0 = F\left(\frac{\mu(T(x_k) - T(y_k), t_0)}{\rho}\right)$, so we have $r_0 > 1 - r$, there exists $s \in (0, 1)$ such that $r_0 > 1 - s > 1 - r$. For $r_0 > 1 - s$, we have $r_1, r_2 \in (0, 1)$ such that $r_0 * r_1 > 1 - s$ and $(1 - r_0) \diamond (1 - r_0) \leq s$. Putting $r_3 = \max\{r_1, r_2\}$. Now we consider a ball $\mathcal{B}_y^c(1 - r_3, t - t_0)(T, F)$. And we prove that

$$\mathcal{B}_y^c(1 - r_3, t - t_0)(T, F) \subset \mathcal{B}_x^c(r, t)(T, F).$$

Let $z = (z_k) \in \mathcal{B}_y^c(1 - r_3, t - t_0)(T, F)$, then $F\left(\frac{\mu(T(y_k) - T(z_k), t - t_0)}{\rho}\right) > r_3$ and $F\left(\frac{\nu(T(y_k) - T(z_k), t - t_0)}{\rho}\right) < 1 - r_3$. Therefore, we have

$$F\left(\frac{\mu(T(x_k) - T(z_k), t)}{\rho}\right) \geq F\left(\frac{\mu(T(x_k) - T(y_k), t_0)}{\rho}\right) * F\left(\frac{\mu(T(y_k) - T(z_k), t - t_0)}{\rho}\right) \\ \geq (r_0 * r_3) \geq (r_0 * r_1) \geq (1 - s) \geq (1 - r)$$

and

$$F\left(\frac{\nu(T(x_k) - T(z_k), t)}{\rho}\right) \leq F\left(\frac{\nu(T(x_k) - T(y_k), t_0)}{\rho}\right) \diamond F\left(\frac{\nu(T(y_k) - T(z_k), t - t_0)}{\rho}\right) \\ \leq (1 - r_0) \diamond (1 - r_3) \leq (1 - r_0) \diamond (1 - r_2) \leq s \leq r.$$

Thus $z \in \mathcal{B}_x^c(r, t)(T, F)$ and hence, we get

$$\mathcal{B}_y^c(1 - r_3, t - t_0)(T, F) \subset \mathcal{B}_x^c(r, t)(T, F).$$

Remark 3.2. $\mathcal{M}_{(\mu, \nu)}^I(T, F)$ is an IFNS.

Define

$$\tau_{(\mu, \nu)}^I(T, F) = AC\mathcal{M}_{(\mu, \nu)}^I(T, F) : \text{ for each } x \in A \text{ there exists } t > 0 \\ \text{ and } r \in (0, 1) \text{ such that } \mathcal{B}_x(r, t)(T, F) \subset A.$$

Then $\tau_{(\mu, \nu)}^I(T, F)$ is a topology on $\mathcal{M}_{(\mu, \nu)}^I(T, F)$.

In the above result we can easily verify that the open sets in these spaces are open ball in the same spaces. This theorem itself will have various applications in our future work.

Theorem 3.2. The topology $\tau_{(\mu, \nu)}^I(T, F)$ on $\mathcal{M}_{0(\mu, \nu)}^I(T, F)$ is first countable.

Proof. $\{\mathcal{B}_x(\frac{1}{n}, \frac{1}{n})(T, F) : n = 1, 2, 3, \dots\}$ is a local base at x , the topology $\tau_{(\mu, \nu)}^I(T, F)$ on $\mathcal{M}_{0(\mu, \nu)}^I(T, F)$ is first countable. \square

Theorem 3.3. $\mathcal{M}_{(\mu, \nu)}^I(T, F)$ and $\mathcal{M}_{0(\mu, \nu)}^I(T, F)$ are Hausdorff spaces.

Proof. Let $x, y \in \mathcal{M}_{(\mu, \nu)}^I(T, F)$ such that $x \neq y$. Then $0 < F\left(\frac{\mu(T(x)-T(y), t)}{\rho}\right) < 1$ and $0 < F\left(\frac{\nu(T(x)-T(y), t)}{\rho}\right) < 1$.

Putting $r_1 = F\left(\frac{\mu(T(x)-T(y), t)}{\rho}\right)$, $r_2 = F\left(\frac{\nu(T(x)-T(y), t)}{\rho}\right)$ and $r = \max\{r_1, 1 - r_2\}$. For each $r_0 \in (r, 1)$ there exists r_3 and r_4 such that $r_3 * r_4 \geq r_0$ and $(1 - r_3) \diamond (1 - r_4) \leq (1 - r_0)$.

Putting $r_5 = \max\{r_3, 1 - r_4\}$ and consider the open balls $\mathcal{B}_x(1 - r_5, \frac{t}{2})$ and $\mathcal{B}_y(1 - r_5, \frac{t}{2})$. Then clearly $\mathcal{B}_x^c(1 - r_5, \frac{t}{2}) \cap \mathcal{B}_y^c(1 - r_5, \frac{t}{2}) = \phi$. For if there exists $z \in \mathcal{B}_x^c(1 - r_5, \frac{t}{2}) \cap \mathcal{B}_y^c(1 - r_5, \frac{t}{2})$, then

$$r_1 = F\left(\frac{\mu(T(x) - T(y), t)}{\rho}\right) \geq \left(\frac{\mu(T(x) - T(z), \frac{t}{2})}{\rho}\right) * F\left(\frac{\mu(T(z) - T(y), \frac{t}{2})}{\rho}\right) \\ \geq r_5 * r_5 \geq r_3 * r_3 \geq r_0 > r_1$$

and

$$r_2 = F\left(\frac{\nu(T(x) - T(y), t)}{\rho}\right) \leq F\left(\frac{\nu(T(x) - T(z), \frac{t}{2})}{\rho}\right) \diamond F\left(\frac{\nu(T(z) - T(y), \frac{t}{2})}{\rho}\right) \\ \leq (1 - r_5) \diamond (1 - r_5) \leq (1 - r_4) \diamond (1 - r_4) \leq (1 - r_0) < r_2$$

which is a contradiction. Hence, $\mathcal{M}_{(\mu, \nu)}^I(T, F)$ is Hausdorff. Similarly the proof follows for $\mathcal{M}_{0(\mu, \nu)}^I(T, F)$. \square

4. Conclusion

The concept of defining intuitionistic fuzzy ideal convergent sequence spaces as it generalized the fuzzy set theory and give quite useful and interesting applications in many areas of mathematics and engineering. This chapter give brief introduction to intuitionistic fuzzy normed spaces with some basic definitions of convergence applicable on it. We have also summarized different types of sequence spaces with the help of ideal, Orlicz function and compact operator. At the end of this chapter some theorems and remarks based on these new defined sequence spaces are discussed for proper understanding.

Conflict of interest

The authors declare that they have no competing interests.

IntechOpen

IntechOpen

Author details

Vakeel Ahmad Khan, Hira Fatima and Mobeen Ahmad
Department of Mathematics, Aligarh Muslim University, Aligarh, India

*Address all correspondence to: vakhanmaths@gmail.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Atanassov KT. Intuitionistic fuzzy sets. *Fuzzy Sets and Systems*. 1986; **20**(1):87-96
- [2] Zadeh LA. Fuzzy sets. *Information and Control*. 1965;**8**:338-353
- [3] Barros LC, Bassanezi RC, Tonelli PA. Fuzzy modelling in population dynamics. *Ecological Modelling*. 2000; **128**:27-33
- [4] Hong L, Sun JQ. Bifurcations of fuzzy non-linear dynamical systems. *Communications in Nonlinear Science and Numerical Simulation*. 2006;**1**:1-12
- [5] Fradkov AL, Evans RJ. Control of chaos: Methods of applications in engineering. *Chaos, Solitons & Fractals*. 2005;**29**:33-56
- [6] Giles R. A computer program for fuzzy reasoning. *Fuzzy Sets and Systems*. 1980;**4**:221-234
- [7] Saddati R, Park JH. On the intuitionistic fuzzy topological spaces. *Chaos, Solution and Fractals*. 2006;**27**: 331-344
- [8] Karakus S, Demirci K, Duman O. Statistical convergence on intuitionistic fuzzy normed spaces. *Chaos, Solitons and Fractals*. 2008;**35**:763-769
- [9] Coker D. An introduction to intuitionistic fuzzy topological spaces. *Fuzzy Sets and Systems*. 1997;**88**(1): 81-89
- [10] Mursaleen M, Lohani QMD. Intuitionistic fuzzy 2-normed space and some related concepts. *Chaos, Solution and Fractals*. 2009;**42**:331-344
- [11] Khan VA, Ebadullah K, Rababah RKA. Intuitionistic fuzzy zweier I-convergent sequence spaces. *Functional Analysis: Theory, Methods and Applications*. 2015;**1**:1-7
- [12] Steinhaus H. Sur la convergence ordinaire et la convergence asymptotique. *Colloquium Mathematicum*. 1951;**2**:73-74
- [13] Fast H. Sur la convergence statistique. *Colloquium Mathematicum*. 1951;**2**:241-244
- [14] Schoenberg IJ. The integrability of certain functions and related summability methods. *American Mathematical Monthly*. 1959;**66**:361-375
- [15] Kostyrko P, Salat T, Wilczynski W. I-convergence. *Real Analysis Exchange*. 2000;**26**(2):669-686
- [16] Alotaibi A, Hazarika B, Mohiuddine SA. On the ideal convergence of double sequences in locally solid Riesz spaces. *Abstract and Applied Analysis*. 2014. 6 p. Article ID: 396254. <http://dx.doi.org/10.1155/2014/396254>
- [17] Hazarika B, Mohiuddine SA. Ideal convergence of random variables. *Journal of Function Spaces and Applications*. 2013. Article ID 148249:7
- [18] Mohiuddine SA, Lohani QMD. On generalized statistical convergence in intuitionistic fuzzy normed spaces. *Chaos, Solitons and Fractals*. 2009;**41**: 142-149
- [19] Mursaleen M, Mohiuddine SA, Edely OHH. On the ideal convergence of double sequences in intuitionistic fuzzy normed spaces. *Computers and Mathematics with Application*. 2010;**59**: 603-611
- [20] Nabiev A, Pehlivan S, Gürdal M. On I-Cauchy sequence. *Taiwanese Journal of Mathematics*. 2007;**11**(2):569-576

[21] Bromwich TJI. An Introduction to the Theory of Infinite Series. New York: MacMillan Co. Ltd; 1965

[22] Khan VA, Fatima H, Abdullaha SAA, Khan MD. On a new BV_{σ} I-convergent double sequence spaces. Theory and Application of Mathematics and Computer Science. 2016;6(2): 187-197

[23] Khan VA, Shafiq M, Guillen BL. On paranorm I -convergent sequence spaces defined by a compact operator. Afrika Matematika, Journal of the African Mathematical Union (Springer). 2014; 25(4):12. DOI: 10.1007/s13370-014-0287-2

[24] Kreyszig E. Introductory Functional Analysis with Application. New York, Chichester, Brisbane, Toronto: John Wiley and Sons, Inc; 1978