

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Chapter

Dyslipidemia in Special Populations, the Elderly, Women, HIV, Chronic Kidney Disease and ESRD, and Minority Groups

*Amarpali Brar, Jeans M. Santana, Moro O. Salifu
and Clinton D. Brown*

Abstract

This chapter discusses the management of dyslipidemia in special patient populations: the elderly, woman and pregnancy, renal disease, human immunodeficiency virus (HIV), and different racial/ethnic groups. In the elderly, dyslipidemia is often underdiagnosed and undertreated. Consideration for potential atherosclerotic risk-reduction benefits, risk of adverse effects, drug-drug interactions, and patient preferences should precede the initiation of statin therapy. Data on pregnant women are lacking and need future research. Dyslipidemia and its effects on the cardiovascular system in chronic kidney disease (CKD), end-stage renal disease (ESRD), and HIV are dynamic and multimodal. These conditions are states of chronic inflammation, where it is difficult to associate quantities of cholesterol types with outcomes. Among all racial groups, Asian Indians, Filipinos, and Hispanics are at a higher risk for dyslipidemia. Genetic differences in statin metabolism may explain this racial/ethnic difference.

Keywords: dyslipidemia, elderly, women, racial disparity, gender disparity

1. Introduction

Dyslipidemia, defined as high levels of low-density lipoprotein cholesterol (LDL-C) (≥ 130 mg/dl), total cholesterol (≥ 200 mg/dl), and triglycerides (TG) (≥ 150 mg/dl), or low levels of high-density lipoprotein cholesterol (HDL-C) [< 40 (men) and < 50 (women) mg/dl], is a major risk factor for cardiovascular disease (CVD). Significant heterogeneity in patterns of dyslipidemia exists in these special populations. There is confusion among health-care providers regarding selection and implementation of appropriate guidelines, particularly for special patient populations. Patients in special populations may not clearly fall into one of the four statin-benefit groups identified by the ACC/AHA cholesterol guidelines [1].

In this chapter, we review the evidence in patterns of dyslipidemia and management in the elderly, women and pregnancy, CKD, ESRD, HIV, and different racial/ethnic groups.

2. Dyslipidemia in the elderly

2.1 Epidemiology

According to census projections, the population age 65 and older is expected to double between 2012 and 2060, from 43.1 to 92.0 million [2]. CVD is the main cause of mortality in this age group. Lipid-lowering pharmacological intervention is one of the most successful cardiovascular preventative interventions. Concerns about its safety and efficacy in this age group have led different countries to adopt different strategies concerning the use of lipid-lowering drugs in the elderly [3]. In US population as reported in the National Health and Nutrition Examination Survey (NHANES) publications, across all age groups, triglyceride levels increase with age and reach a peak in men aged 50–59 years and in women aged 60–69 years. Apolipoprotein B (Apo B) and small dense LDLc also increase with age, while HDL-c does not seem to vary with age [4].

2.2 Management

Lifestyle modification including adhering to a heart healthy diet, regular exercise, avoidance of tobacco products, and maintenance of a healthy weight remains a critical component of ASCVD risk reduction.

2.2.1 Secondary prevention statin trials in elderly

In general, the elderly are usually underrepresented in published clinical trials. The only large randomized statin trials that focused on older patients are the Prospective Study of Pravastatin in the Elderly at Risk (PROSPER) and the Study Assessing Goals in the Elderly (SAGE) trial.

PROSPER is a randomized controlled trial in which 5804 patients aged 70–82 years with a history of, or risk factors for, vascular disease were randomized to receive pravastatin (40 mg/day; n = 2891) or placebo (n = 2913). Follow-up was 3.2 years. Pravastatin was found to reduce the incidence of coronary death, nonfatal myocardial infarction, and fatal or nonfatal stroke. Suggesting pravastatin may be utilized to manage dyslipidemia in the elderly and be prescribed to [5]. Later, the Study Assessing Goals in the Elderly (SAGE) compared the effect of intensive (atorvastatin 80 mg/day) with moderate (pravastatin 40 mg/day) cholesterol lowering with statins in a cohort of 893 patients, 65–85 years of age with coronary artery disease (CAD), and follow-up for 12 months. Atorvastatin-treated patients experienced greater LDL-C reductions than did pravastatin-treated patients, a trend toward fewer major acute cardiovascular events and a significantly greater reduction (77%) in all-cause death [6].

British Heart Foundation Heart Protection Study evaluated the role of simvastatin 40 mg/day versus placebo in 20,536 patients with coronary atherosclerosis and diabetics with coronary heart disease (CHD) risk equivalence. About 28% of randomized patients were ≥ 70 years of age. Among the 1263 individuals with 75–80 years of age at study entry, the rate of major coronary events in the simvastatin group was significantly lower than placebo group [7]. The long-term intervention with pravastatin in ischemic disease (LIPID) trial randomized patients with prior myocardial infarction or unstable angina to receive pravastatin 40 mg/day versus placebo, of which 39% (3514) were of age 65–75 years. Here, pravastatin reduced the risk for all CVD events, and similar adverse effects were observed in older and younger patients [8].

Other major secondary prevention trials, which included large numbers of elderly patients, are cholesterol and recurrent events (CARE) and Scandinavian simvastatin survival study (4S). In subgroup analyses of both studies, the absolute benefit of treatment was significantly greater in older patients as compared to younger patients for cardiovascular events [9, 10]. The 2013 ACC/AHA guidelines on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults recommended high-intensity statin therapy should be initiated or continued as first-line therapy in women and men ≤ 75 years of age who have clinical atherosclerotic cardiovascular disease (ASCVD), unless contraindicated. A larger amount of data supports the use of moderate-intensity statin therapy for secondary prevention in individuals with clinical ASCVD who are >75 years of age. However, the limited information available did not clearly support initiation of high-intensity statin therapy for secondary prevention in individuals >75 years of age [1]. Statin therapy to higher risk elderly patients is appropriate. In individuals with clinical ASCVD >75 years of age, practitioners should evaluate the potential for ASCVD risk-reduction benefits, adverse effects, drug-drug interactions and consider patient preferences when initiating a moderate- or high-intensity statin. It is reasonable to continue statin therapy in those who are tolerating it [1, 11]. Physicians treating the elderly must also consider the functional age of the patient and the impact of long-term drug therapy on safety and quality of life. Caution is recommended when statins are used in frail elderly patients, who may be more susceptible to drug-related myopathy and other side effects [12].

2.2.2 Primary prevention and statin trials

The Anglo-Scandinavian Cardiac Outcomes trial randomized 19,342 hypertensive patients (aged 40–79 years with at least three other cardiovascular risk factors) to be assigned to atorvastatin 10 mg or placebo. Nonfatal myocardial infarction and fatal cardiovascular disease were significantly lower in the statin group [13]. In a post hoc analysis, efficacy and safety of atorvastatin in 1129 patients aged 65–75 years at randomization was compared with 1709 younger patients in the Collaborative Atorvastatin Diabetes Study (CARDS). Primary end point of time to first occurrence of acute coronary heart disease events, coronary revascularizations, or stroke was similar in both groups. The overall safety profile of atorvastatin was similar between age groups [14]. Justification for the Use of statins in Prevention: An Intervention Trial Evaluating Rosuvastatin (JUPITER) is a randomized, double-blind, placebo-controlled trial with 17,802 participants of which 5695 were 70 years or older with LDL-C levels <130 mg/dl and high-sensitivity C-reactive protein levels of 2.0 mg/L or more without cardiovascular disease and were randomly assigned in a 1:1 ratio to receive 20 mg of rosuvastatin daily or placebo. In secondary analysis of this trial, no significant heterogeneity was found in treatment effects by age, absolute reductions in event rates associated with rosuvastatin were greater in elderly. The relative rate of any serious adverse event among older patients in the rosuvastatin versus placebo group was 1.05 (CI, 0.93–1.17) [15, 16].

As per 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults; four statin benefit groups are individuals with ASCVD, individuals with primary elevations of LDL-C ≥ 190 mg/day, individuals 40–75 years of age with diabetes and LDL-C 70 to 189 mg/dl without clinical ASCVD, and individuals without clinical ASCVD or diabetes who are 40–75 years of age and have LDL-C 70–189 mg/dl and an estimated 10-year ASCVD risk of $\geq 7.5\%$. This requires a clinician-patient discussion. Few data are available to indicate an ASCVD event reduction benefit in primary prevention among individuals >75 years of age who do not have clinical ASCVD. Therefore, initiation of

statins for primary prevention of ASCVD in individuals >75 years of age requires consideration of additional factors, including increasing comorbidities, safety considerations, and priorities of care. The Pooled Cohort Equations can also provide information on expected 10-year ASCVD risk for those 76–79 years of age that may inform the treatment decision. A discussion of the potential ASCVD risk-reduction benefits, risk of adverse effects, drug-drug interactions, and consideration of patient preferences should precede the initiation of statin therapy for primary prevention in older individuals [1].

2.2.3 Statin and side effects

In the PROSPER study, no rhabdomyolysis or serum CK concentration > 10 times the upper limit of normal was reported. There was no statistically significant difference in the incidence of reported myalgia between the pravastatin and placebo groups (1.2 and 1.1%, respectively). In the SAGE study, incidence of myalgia was also not different between the atorvastatin and pravastatin treatment groups (3.1% versus 2.7%) with only one individual found to have CK level > 10 times the upper limit of normal in the pravastatin treatment group but none in the atorvastatin group [5, 6]. In a retrospective study, Graham et al. reported rhabdomyolysis risk was similar and low for monotherapy with atorvastatin, pravastatin, and simvastatin. Combined statin-fibrate use was associated with increased risk, especially in older patients with diabetes mellitus [17].

All of the available statins, with the exception of pravastatin and rosuvastatin, are metabolized by the cytochrome P450 (CYP) system. Serum concentrations of these statins can potentially be increased when other medications competing for the CYP system or CYP isoenzyme inhibitors are prescribed and can lead to increased risk of myositis and rhabdomyolysis. There is an age-related decrease in glomerular filtration rate, decrease in hepatic blood flow, decrease in drug clearance, and increased expression of P-glycoproteins resulting in alterations in the rate of drug transport across cellular membranes [18–20].

2.3 Other medications

2.3.1 Ezetimibe

Improved Reduction of Outcomes: Vytorin Efficacy International Trial (IMPROVE-IT) evaluated the effect of ezetimibe combined with simvastatin, as compared to simvastatin alone in 18,144 patients with acute coronary syndrome. Subgroup analysis showed a benefit of ezetimibe in the 7971 patients' over 65 years and 2798 patients' over 75 years of age [21].

2.3.2 Proprotein convertase subtilisin/kexin type 9 (PCSK9) inhibitors

The first large cardiovascular outcome study using PCSK9 inhibitor therapy (FOURIER) was a randomized, double-blind placebo-controlled trial of 27,564 subjects with 12,254 patients over the age 65 years with ASCVD and LDL-C levels ≥ 70 mg/dl while on maximally tolerated statin therapy, randomly assigned to receive evolocumab or placebo. In these patients, there was a significant decrease in composite outcome of cardiovascular death, myocardial infarction (MI), stroke, hospitalization for unstable angina, or coronary revascularization. However, cost is prohibitive for generalized use [22].

2.3.3. Fibrates

No data are available for fibrates in patients over 75 years of age. World Health Organization Clofibrate Study, Helsinki Heart Study (HHS), Bezafibrate Infarction Prevention (BIP), and Veterans Affairs Program High-Density Lipoprotein Cholesterol Intervention Trial (VA-HIT) showed reduction in the risk of myocardial infarction. Fenofibrate is considered the preferred agent, due to once-daily dosing and a favorable adverse reactions profile. Caution is needed in elderly patients with reduced renal function [23–26].

In conclusion, cardiovascular prevention clinical trial evidence for the patients over age 75 years is very limited. Management of dyslipidemia in the elderly requires consideration of comorbidities, safety considerations, polypharmacy, and priorities of care.

3. Dyslipidemia in women

3.1 Gender and cardiovascular risk

CVD is a major cause of death in women. While the risk of CVD in men increases after age 40 years, risk develops 7–10 years later in women. Incidence of CAD in premenopausal women is 3–4 times lower than in men [27, 28]. After menopause due to loss of vasodilating property of estrogen and increased sympathetic activity, the risk of CAD is increased and is similar to men [29–31]. Risk factors unique to women are use of oral contraceptives, menopause, hormone replacement therapy, gestational hypertension, and diabetes. Obesity and metabolic syndrome are also more prevalent in women [32].

Plasma cholesterol levels and LDL-C levels are similar in both sexes during the infancy and adolescence. LDL levels increase progressively in both men and women after the age of 20, but more rapidly in men. Size of the LDL particles reduces as men age, while it remains stable in women until menopause then becomes smaller [33].

3.2 Management of dyslipidemia

In 1999, American heart association (AHA) developed the first women-specific clinical report regarding recommendations for CVD prevention [34]. In the 2011 AHA update, effectiveness-based guidelines for the prevention of CVD in women recommended lifestyle approaches for LDL-C < 100 mg/dl, HDL-C > 50 mg/dl, triglycerides < 150 mg/dl, and non-HDL-C (total cholesterol minus HDL) < 130 mg/dl (class I recommendation; level of evidence B). LDL-C-lowering drug therapy with lifestyle therapy in women with coronary heart disease (CHD) to achieve an LDL-C < 100 mg/dl (class I; level of evidence A) and is also indicated in women with other atherosclerotic CHD or diabetes mellitus or 10-year absolute risk 20% (class I; level of evidence B). A reduction to < 70 mg/dl is reasonable in very-high-risk women like those with recent acute coronary syndrome (ACS) or multiple cardiovascular risk factors with CHD and may require an LDL-lowering drug combination (class IIa; level of evidence B). In women > 60 years of age and with an estimated CHD risk > 10%, statins could be considered if hsCRP is > 2 mg/dl after lifestyle modification and no acute inflammatory process is present (class IIb; level of evidence B). Niacin or fibrate therapy can be useful when HDL-C is low (< 50 mg/dl) or non-HDL-C is elevated (> 130 mg/dl) in high-risk women after LDL-C goal is reached (class IIb; level of evidence B) [35].

In 2013 ACC/AHA guidelines added, treatment decisions for women should be based on the level of ASCVD risk. Statin treatment based on estimated 10-year ASCVD risk avoids the overtreatment of lower risk groups, such as younger, non-Hispanic white women who, despite moderate elevations in LDL-C, are typically not at significantly increased risk for ASCVD in the absence of other substantial risk factors [1]. The European Atherosclerosis Society (ESC/EAS) recommends that assessment of CV risk should be gender specific and recommended statin therapy be initiated as primary and secondary prevention for women at high risk [36].

3.3 Women in hyperlipidemia clinical trials

Women have been underrepresented in many primary and secondary prevention trials. In major primary prevention trials, the number of women varied from 15 to 49%. In AFCAPS (15%), HPS (30%), ALLHAT-LLT (49%), and in ASCOT-LLA trial (19%) were women [37, 38]. In a prospective, randomized, open-labeled, blinded Japanese primary prevention trial of patients with hypercholesterolemia, 5547 (68%) were women randomly assigned to diet alone or diet with pravastatin daily. Treatment with a low dose of pravastatin reduced the risk of CHD but subgroup analysis of CHD risk in women was not statistically significant [39]. JUPITER trial participants included 6801 women ≥ 60 years of age and 11,001 men ≥ 50 years of age with high-sensitivity C-reactive protein ≥ 2 mg/L and LDL-C < 130 mg/dl, randomized to rosuvastatin versus placebo. JUPITER demonstrated that for primary prevention, rosuvastatin reduced CVD events in women with a relative risk reduction similar to that in men, a finding supported by meta-analysis of primary prevention statin trials [40].

In major secondary prevention trials, 4S included 827 (19% woman), CARE 576 (14%), LIPID 1516 (17%), TNT 1902 (19%), and SEARCH trial included 2052 (17%) women [41–44]. Cholesterol lowering with simvastatin produced similar reductions in relative risk for major coronary events in women compared with men. There were too few female deaths to assess the effects on mortality in women [41]. A meta-analysis of 13 large trials for a total of 11,435 women in primary prevention and 8272 in secondary prevention concluded that lipid lowering does not affect CHD mortality in women for primary prevention, but it is effective for secondary prevention [27].

3.4 Gender differences in statin use

Women are more likely to have poor lipid control. There have been conflicting results regarding gender differences in benefit of statins in women with CVD. However, in a meta-analysis performed on data from 22 trials of statin therapy versus control ($n = 134,537$) and five trials of more intensive versus less intensive statin therapy, 27% of 174,149 randomized participants were women. In men and women at an equivalent risk of cardiovascular disease, statin therapy was equally effective for the prevention of major vascular events. Kostis et al. performed a meta-analysis consisting of 18 randomized clinical trials of statins with gender-specific outcomes and found statin therapy was associated with significant decreases in cardiovascular events and in all-cause mortality in women and men [45].

There is a gender-specific impact of transporter polymorphisms on statin pharmacokinetics. Estrogen-induced water and sodium retention, increased volume for lipophilic drugs, higher protein-binding globulins, higher CYP3A4 activity, lower body mass index, and lower renal clearance in women may affect absorption of different statins. Women seem to be at greater risk of statin-induced rhabdomyolysis. In a US case-control study of 252,460 users of lipid-lowering therapy, there was

trend for increased risk of rhabdomyolysis [46]. In the JUPITER study, the occurrence of serious adverse events was similar in both men and women [40].

3.5 Pregnancy

3.5.1 Normal changes in pregnancy

Metabolic adaptations during pregnancy are essential to meet the physiological demands of pregnancy as well as development of the fetus. An increase in insulin resistance results in increases in maternal glucose and free fatty acid concentrations, allowing for greater substrate availability for fetal growth. Production of progesterone leads to lipogenesis, lipids are transported across the placenta and metabolized; this signifies the essential role of lipids to normal fetal development [47, 48]. Within 6 weeks of gestation, lipid levels drop slightly followed by rise in both total cholesterol and marked increase in triglycerides (TG). HDL-C levels and apolipoprotein A-I levels also increase during normal gestation, with peak levels during the second trimester. As TG levels rise, there is a decrease in low-density lipoprotein (LDL) size with an increased proportion of atherogenic small-dense LDL. Both cholesterol and triglyceride concentrations decrease significantly within 24 h of delivery. However, while TG levels continued to decrease rapidly returning to nonpregnant levels during the puerperium, LDL-C remained elevated for at least 6–7 weeks postpartum [49, 50]. During pregnancy, lipoprotein Lp(a) levels increase with gestational age and fall to prepregnancy levels within 6 months postpartum [51, 52].

3.5.2 Lipids and complications in pregnancy

Preeclamptic women exhibit higher mean serum TG levels, elevated LDL-C fractions, increased levels of Lp(a), and lower HDL cholesterol levels compared with healthy pregnant women [49]. Amsterdam Born Children and Their Development (ABCD) cohort study showed that every unit increase in TG was linearly associated with an increased risk of preeclampsia, pregnancy-induced hypertension, and preterm delivery. Total cholesterol was not associated with any of the outcome measures [53]. Women who have higher concentrations of small-dense LDL fractions during pregnancy tend to have increased risk of cardiovascular disease later in life [53, 54].

In the multicenter, prospective Coronary Artery Risk Development in Young Adults study, 1010 women of which 49% identified as black with at least one singleton birth with 20 years of follow-up were evaluated. There was a U-shaped relationship between prepregnancy cholesterol concentrations and preterm birth risk with both low (<156 mg/dl) and high cholesterol (>195 mg/dl) related to preterm birth risk [55]. Observational and experimental evidence increasingly supports a relation between growth and development during fetal and infant life and health in later years. Additionally, preterm newborns have been found to be at an increased risk of CVD later in life [56].

3.5.2.1 Antihyperlipidemia therapy

Pregnant women are generally excluded from clinic trials, thus the data are limited. HMG-CoA reductase inhibitors have been associated with teratogenicity and congenital malformation and are not recommended. Fibrates and nicotinic acid have not been well studied and are not recommended. Omega fatty acids, on the other hand, can be safely used as monotherapy to decrease TG levels.

4. Dyslipidemia in CKD and ESRD

4.1 Epidemiology and pathogenesis

Patients with CKD and ESRD are at an increased risk of CVD, and are more likely to die because of adverse cardiovascular events [57]. In addition to traditional risk factors like hypertension, diabetes mellitus, dyslipidemia, and a family history of CAD, patients with CKD are plagued by nontraditional risk factors like homocysteinemia, mineral bone disease, carbamylation, and chronic inflammation [58, 59]). As CKD progresses, an unstable vascular environment ensues which threatens endothelial function and lipoprotein integrity and thus potentiating adverse cardiovascular outcomes. **Figure 1** shows mechanism of dyslipidemia in this population.

Dyslipidemia in these patients is undoubtedly one of the strongest risk factors for these adverse cardiovascular outcomes. Despite these data, however, CVD remains underdiagnosed and undertreated in patients with CKD. Dyslipidemia has been found to be distinctly different from the general population and variable depending on the stage of CKD [60]. The lipid profile of patients with nondialysis-dependent CKD is usually composed of low HDL-C and high triglycerides with normal to low total cholesterol and low-density lipoprotein cholesterol [61]. In fact, plasma triglycerides start to increase in early stages of CKD and show the highest concentrations in nephrotic syndrome and in dialysis patients, especially those who are treated with peritoneal dialysis [60]. In patients with nondialysis-dependent CKD, the hypertriglyceridemia has been attributed to delayed catabolism and increased hepatic production of triglyceride-rich lipoproteins, and to a smaller extent by the presence of lipase inhibitors [62]. This altered catabolism in turn results in the accumulation of triglyceride-rich lipoproteins, like IDL and small-dense LDL (sdLDL) which are highly atherogenic [63, 64].

Though elevated LDL-C is not a typical feature of these patients, sdLDL, an LDL subtype, is increased and carries the ability to penetrate the vessel wall, becomes oxidized, and triggers the atherosclerotic process. The subfractions of HDL in CKD and ESRD are also different than that of the general population. In uremia [65], LDL and HDL, their subfractions, lipidomes, and proteomes, gradually become more susceptible to structural modifications such as carbamylation, oxidation, glycation, nitration, and homocysteinylation [66, 67]. Most of these effectors are irreversible and amplify the uptake by the scavenger receptors on the surface of macrophages [60]. HDL's vital components like ApoA-I, PON-1, and LCAT are altered, ultimately attenuating HDL's cyto- and vascular-protective properties [68]. The reduced




Figure 1. Mechanism of dyslipidemia in chronic kidney disease (CKD), end stage renal disease (ESRD), and human immunodeficiency virus (HIV) patients.

PON-1 activity further predisposes LDL and HDL to more oxidation and, in turn, dysfunction and enhanced atherogenic potential [69]. In fact, it has been found that increasing serum HDL-C over time is paradoxically associated with significantly higher all-cause and cardiovascular mortality [70].

4.2 Management

Recent studies show that statin therapy can decrease cardiovascular mortality in CKD population. Heart Protection Study and Anglo-Scandinavian Cardiac Outcomes Trial (ASCOT) demonstrated that CKD patients treated with simvastatin exhibited a reduction of cardiovascular morbidity and mortality [71, 13]. Results from the Pravastatin Pooling Project showed similar cardioprotective effects [72]. Atorvastatin and rosuvastatin also reduced the relative risk of cardiovascular events in CKD patients and improved outcomes were found with higher atorvastatin doses [73–75]. Study of Heart and Renal Protection (SHARP) trial randomized 9270 patients with CKD with no prior history of CAD and found that simvastatin 20 mg plus ezetimibe 10 mg daily safely reduced the incidence of major atherosclerotic events in CKD patients [76].

KDIGO panel provides recommendations for dyslipidemia in patients at all stages of CKD based on a few large, randomized controlled trials and post hoc analyses of the subgroup of CKD patients from statin trials in the general population [76, 77]. In adult CKD patients ≥ 50 years age, statin therapy alone is recommended for those with GFR ≥ 60 ml/min and statin or statin/ezetimibe therapy is recommended for GFR < 60 ml/min. In adults aged 18–49 years with CKD, statin treatment is recommended in patients with known coronary disease, diabetes mellitus, prior ischemic stroke or estimated 10-year incidence of coronary death or nonfatal myocardial infarction $> 10\%$. Neither ACC/AHA nor KDIGO guidelines recommend initiation of statin therapy or combination treatment with statin and ezetimibe in dialysis-dependent patients on the basis of results from the AURORA (A Study to Evaluate the Use of Rosuvastatin in Subjects on Regular Hemodialysis: An Assessment of Survival and Cardiovascular Events) and Deutsche Diabetes Dialyse Studie (4-D) trials. However, patients already on lipid-lowering therapy at the time of progression to dialysis may continue treatment [76, 78–80].

Novel and innovative therapies are needed to address the multiple lipid/lipoprotein abnormalities that facilitate high cardiovascular risk and mortality in patients with dialysis-dependent CKD.

5. Dyslipidemia in renal transplant recipients

5.1 Epidemiology

Risk of cardiovascular events in kidney transplant recipients is markedly elevated as compared to people without CKD. Prevalence of hyperlipidemia ranges from 16 to 78% in kidney transplant recipients. Hypercholesterolemia has peak incidence at 12 months posttransplant and correlates with excessive body weight. After transplantation, increases in total cholesterol, LDL-C, very-low-density lipoprotein (VLDL)-cholesterol, and TG have been noted. In addition, LDL-C may be more susceptible to oxidation, making the particle more atherogenic. HDL-C levels may be normal or even high although the composition of HDL may not be normal [81].

5.2 Mechanisms of dyslipidemia

Immunosuppressive agents contribute significantly to hyperlipidemia in renal transplant recipients. Corticosteroids induce insulin resistance. Hyperinsulinemia leads to increased hepatic uptake of free fatty acids (FFA) which constitute the main substrate for VLDL cholesterol synthesis. There is increased conversion of VLDL to LDL cholesterol, leading to a rise in LDL cholesterol levels and downregulation of LDL receptors [81–83]. Calcineurin inhibitors interfere with binding of LDL-to-LDL receptor leading to increase in LDL-C, interfere with bile acid synthesis and lead to LDL receptor downregulation. Cyclosporine is highly lipophilic. Tacrolimus is associated with less hyperlipidemia [84]. Rapamycin inhibits lipoprotein lipase, associated with decrease in apolipoprotein-B 100 catabolism, and increases secretion of VLDL cholesterol [85]. Other secondary causes include nephrotic syndrome, hypothyroidism, diabetes mellitus, excessive alcohol intake, obesity, chronic liver disease, and genetic predisposition.

5.3 Management

Lipid profile evaluation including total cholesterol, LDL, HDL, and triglyceride levels is recommended. Kidney Disease Improving Global Outcomes (KDIGO) suggest treatment with statins in adult kidney transplant recipients [77]. Assessment of Lescol in Renal Transplantation (ALERT) trial is the only multicenter, randomized, double-blind, placebo-controlled trial in renal transplant recipients. In this study, 2102 renal transplant recipients with total cholesterol 4.0–9 mmol/l were randomized to receive fluvastatin (n = 1050) or placebo (n = 1052). After a mean follow-up of 5 years, fluvastatin lowered LDL cholesterol concentrations by 32%. Although cardiac deaths and nonfatal myocardial infarction seemed to be reduced, fluvastatin did not reduce rates of coronary intervention procedures or mortality [86]. Of 1787 patients who completed ALERT, 1652 (92%) were followed in the extension trial with mean follow-up of 6.7 years which showed 29% reduction in cardiac death or definite nonfatal myocardial infarction in the fluvastatin arm [87]. Due to the known interaction of calcineurin inhibitors through CYP3A isoenzyme system, lower doses of statins are generally used as compared to general population. Among fibrates, fenofibrate is less myotoxic than gemfibrozil when combined with statin but fenofibrate should be avoided in advanced CKD. Bile acid sequestrants are not widely used due to gastrointestinal side effects and can also interfere with absorption of immunosuppressants. Ezetimibe is considered safe in renal transplant recipients.

6. Dyslipidemia in HIV

Antiretroviral therapy (ART) has prolonged the survival of HIV-infected individuals, which, in effect, has increased the prevalence of comorbidities, like coronary heart disease. Studies have shown that HIV-infected individuals have higher cardiovascular disease (CVD) risk than uninfected persons in the United States [88–90]. Paisible et al. reported a 1.5–2-fold increased risk of incident myocardial infarction compared with uninfected subjects [91]. Among HIV patients, women have been found to have higher CVD risk than men [92]. Risk, however, cannot be attributed to HIV alone as traditional CVD risk factors like smoking [93], dyslipidemia, diabetes mellitus, and hypertension [92] have been found to be more common among those infected. **Table 1** summarizes the traditional and nontraditional risk factors for cardiovascular disease in special populations.

Their lipid profile is comprised of hypertriglyceridemia, increased sdLDL, and low HDL-C levels [94, 95]. Atheromas have been found to have larger lipid pools with dystrophic calcifications [96]. Consequently, the coronary arteries have a higher burden of coronary plaque and prevalence of detectable calcium [97–99]. These plaques are rupture-prone and associated with inflammation [100] and monocyte activation [101]. As a result, patients tend to have higher rates of subclinical vascular disease [102, 103]. ART-associated lipodystrophy has been linked with these cardiac and metabolic complications [104]. However, newer ART medications have had less effect on dyslipidemia and related myocardial events [105]. Still, CD4 cell depletion and immune dysfunction perpetuates HIV-related atherosclerosis, irrespective of ART [106]. Silverberg et al. reported that CD4 counts <200 cells/mm³ were significantly associated with increased risk of MI [105]. While viral load and prior ART use were not associated with MI, unsuppressed HIV viremia was associated with MI [89] and stroke [107]. Dyslipidemia and its effects on the cardiovascular system in HIV are dynamic and multimodal. In these states of chronic inflammation, it is difficult to associate quantities of cholesterol types with outcomes. That is, in these special populations, there may not be a “good” or “bad” cholesterol, but rather dysfunctional lipoprotein atherosclerosis.

Studies have shown that statins are safe and beneficial in those infected with HIV; however, the use of statins remains relatively low [89, 108, 109]. Reasons include (1) the low prevalence of elevated LDL-C, (2) their uncertain efficacy for CVD prevention, and (3) the potential to adverse side effects and negative drug-drug interactions [80]. National Lipid Association (NLA) recommends that all HIV-infected patients be first assessed for cardiovascular risk and counseled about lifestyle interventions like diet, exercise, and smoking for the prevention of atherosclerotic cardiovascular disease. In addition, a fasting lipid panel should be obtained in all newly identified HIV-infected patients before and after starting antiretroviral therapy. TG >500 mg/dl that are refractory to lifestyle modification or changes in ART (if an option) should be treated with either a fibrate (fenofibrate preferred) or prescription omega-3 fatty acids to lower TG to <500 mg/dl. Non-HDL-C and

Traditional	Non-traditional		
	CKD	HIV	Renal transplant
Diabetes mellitus and insulin resistance	Hyperhomocysteinemia	Antiretroviral therapy	Immunosuppressive therapy: Glucocorticoids Calcineurin inhibitor Rapamycin
Hypertension	Hypercalcemia and hyperphosphatemia		
Dyslipidemia	Chronic inflammation and oxidative stress		
Obesity and physical inactivity	Anemia of chronic disease		
Smoking	Microalbuminuria		
Family history of atherosclerosis	Nephrotic syndrome		
Age			
Gender			

CKD, chronic kidney disease; HIV, human immunodeficiency virus.

Table 1.
 Traditional and nontraditional risk factors for cardiovascular disease in special populations.

LDL-C should be reassessed for appropriate management with statin therapy with caution of drug-drug interactions and side effects. To this end, the NLA Expert Panel recommends clinicians to prescribe pitavastatin as the generally preferred agent in HIV-infected patients [80].

7. Racial differences in dyslipidemia

7.1 Epidemiology

National Health and Nutrition Examination Survey (NHANES) is the primary data source for national prevalence rates of dyslipidemia in the US with data on Whites, black/African Americans, and Hispanics/Latinos but is limited in data on Asian subgroups [110]. NHANES data in 2013 showed that the prevalence rate of high LDL-C was highest among Mexican men (40%) and women (30%), followed by non-Hispanic black men (33%) and women (31%). Non-Hispanic white men (30%) and women (29%) had the lowest prevalence of high LDL-C. The prevalence rates of low HDL were 20% in black men and 10% in black women as compared to prevalence rates among non-Hispanic white men (33%) and women (12%) and Hispanic American men (34%) and women (15%). NHANES data in 2008 reported prevalence of high TG in 35% of Hispanic Americans, 33% among non-Hispanic whites, and 16% among non-Hispanic blacks [111]. American Heart Association (AHA), in conjunction with the Centers for Disease Control and Prevention, and the National Institutes of Health reported prevalence of hyperlipidemia in non-Hispanic black men (32.6%) and non-Hispanic black women (36.1%), Hispanic men (43.1%), Hispanic women (41.2%), non-Hispanic white men (37%) and in women (43.4%) and for non-Hispanic Asians, prevalence was 39.9% in men and 40.5% in women [112]. Data from outpatient cohort of adults 35 years or older from 2008 to 2011 in northern California found that compared with non-Hispanic whites, every minority subgroup had an increased prevalence of high triglycerides except blacks. Most minority groups had an increased prevalence of low HDL-C, except for Japanese and blacks. The prevalence of HDL-C was increased among Asian Indians, Filipinos, Japanese, and Vietnamese compared with non-Hispanic whites [113]. **Table 2** shows cholesterol levels in minority populations compared to non-Hispanic Whites in the United States.

The Study of Health Assessment and Risk in Ethnic groups (SHARE), a prospective Canadian trial, showed that South Asians including Asian Indians had an increased prevalence of total cholesterol, high LDL-C, low HDL-C, and high TG compared to European and Chinese cohort [114]. The Multi-Ethnic Study of Atherosclerosis (MESA), a multicenter cohort study of 6814 adults aged 45 to 84 years who were free of clinical CVD at baseline were evaluated for CVD risk and self-reported use of lipid-lowering therapy. Black and Hispanic Americans had prevalence of dyslipidemia that was comparable to that of non-Hispanic whites but were less likely to be treated and controlled. Ethnic disparities were attenuated substantially by adjustment for health-care access variables [115]. Data from the Hispanic Community Health Study (HCHS)/Study of Latinos (SOL), an observational study, showed high prevalence of dyslipidemia among Central American men and Puerto Rican women [116].

7.2 Dyslipidemia and cardiovascular outcomes

In the US, cardiovascular mortality rates are highest in blacks as compared to Hispanics [117]. An early study of CHD among Japanese migrants compared with

	Percent of population	Total cholesterol	HDL	LDL	TGs	Lp(a)	ASCVD
Hispanics							
Men	17%	↑	↓	↑	↑	↔	↑
Women		↓					
African Americans							
Men	13%	↔	↑	↓	↓	↑	↑
Women		↔		↔			
Asians (<i>data limited</i>)	6%	↑	↓	↑	↑	-	↑
American Indians/ Alaska Natives (<i>data limited</i>)	2%	-	↑	-	-	-	↑

HDL, high-density lipoprotein; LDL, low-density lipoprotein; Lp(a), lipoprotein (a), ASCVD, atherosclerotic cardiovascular disease.

Table 2.
 Cholesterol levels in minority populations compared to non-Hispanic Whites in the United States.

Japanese living in Japan showed higher rates of CAD in Japanese immigrants in America [118]. In a study of immigrant Asian Indian men in US, Enas et al. reported high prevalence of CAD, low HDL-C levels, and hypertriglyceridemia. Authors suggested “insulin resistance” as a common pathogenetic mechanism [119]. Increased risk of CAD in south Asian community was seen in data from other studies as well [114, 120, 121].

The 2013 ACC-AHA guidelines for the treatment of cholesterol expand the indications for statin therapy for the prevention of CVD. Ten-year ASCVD risk assessment calculator has been added to determine statin use [122]. ASCVD risk calculator is derived from cohorts that included African-American or white participants with at least 12 years of follow-up. Data from other racial/ethnic groups were insufficient, precluding their inclusion in the final analyses. The equations were also assessed in external validation studies with data from other available cohorts [123]. For other ethnic groups, ACC/AHA recommends use of the equations for non-Hispanic whites, though it acknowledges that estimates may underestimate the risk for American Indians, some Asian Americans (e.g., of south Asian ancestry), and some Hispanics (e.g., Puerto Ricans), and may overestimate the risk for others, including some Asian Americans (e.g., of east Asian ancestry) and some Hispanics (e.g., Mexican Americans) [124].

7.3 Statin metabolism

There are several genetic variants associated with altered statin metabolism. Single-nucleotide polymorphisms in the genes that encode the organic anion-transporting polypeptide (OATP) 1B1 (521 T > C), which regulates hepatic uptake of statins, and the adenosine triphosphate-binding cassette G2 (ABCG2) transporter (421C > A), which regulates hepatic efflux, have been reported [125–127]. Plasma exposure to rosuvastatin and its metabolites was significantly higher in Chinese, Malay, and Asian-Indian subjects compared with white subjects living in Singapore [128]. Rosuvastatin, a HMG-CoA reductase inhibitor, is excreted into bile mediated by breast cancer resistance protein (BCRP). BCRP 421C > A polymorphism may play an important role in the pharmacokinetics of rosuvastatin in healthy Chinese males [129]. In a Japanese study, single-nucleotide polymorphisms in OATP-C, such as T521C (Val174Ala), has

been reported to be associated with altered pharmacokinetics of pravastatin [125]. In a US-based study cohort of 69 European-Americans and 38 African-Americans, SLCO1B1 genotype, in particular, the 521C allele, had a significant effect on the pharmacokinetics of pravastatin. European-Americans demonstrated significantly higher pravastatin levels as compared to African-Americans [130]. There is less information on statin metabolism in other racial/ethnic minority groups. In a randomized placebo-controlled trial of 25,673 patients, the absolute risk of myopathy with adding niacin-laropiprant to statin-based LDL cholesterol-lowering therapy was more than 10 times as great among Chinese as compared to their European counterparts [131]. The relative risk of myopathy with niacin-laropiprant versus placebo was also higher in Chinese patients [132].

Lifestyle risk factors include unhealthy diet, obesity, and physical inactivity. Racial/ethnic disparities also exist in these lifestyle risk factors. According to the National Health Interview Survey (NHIS) 2008–2010, Asian adults were less likely to be current smokers or to be obese. Black adults were more likely to be physically inactive, to be obese, and to get insufficient sleep. Hispanics were less likely than non-Hispanic adults to smoke cigarettes, get insufficient sleep, but were more likely to be inactive in terms of aerobic and muscle-strengthening leisure time. Immigration and acculturation have a profound impact on lifestyle in both Hispanics and Asians in the US [133]. Asian Indian, Filipino, Vietnamese women, and Asian Indian men have increased risk dyslipidemias as compared to non-Hispanic whites. Further research is needed to determine the role of dyslipidemia subtypes and other risk factors in explaining the higher risk of CVD in minority subgroups. By understanding these differences, clinicians will be able to provide more culturally competent recommendations on prevention and management of dyslipidemia.

8. Conclusion

In conclusion, differences in dyslipidemia patterns, risk factors, and management exist in the elderly, women and pregnancy, CKD, ESRD, HIV, and different racial ethnic groups. Nevertheless, they are frequently underrepresented in clinical trials, which calls for more inclusive research that will develop stronger recommendations of daily practice.

Author details

Amarpali Brar*, Jeans M. Santana, Moro O. Salifu and Clinton D. Brown
SUNY Downstate Medical Center, Brooklyn, NY, USA

*Address all correspondence to: amarpali.brar@downstate.edu

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Stone NJ, Robinson JG, Lichtenstein AH, Bairey Merz CN, Blum CB, Eckel RH, et al. 2013 ACC/AHA guideline on the treatment of blood cholesterol to reduce atherosclerotic cardiovascular risk in adults: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation*. 2014;**129** (25 Suppl 2):S1-S45 (Epub Nov 14, 2013)
- [2] Sandra LC, Ortman JM. Projections of the size and composition of the U.S. population: 2014 to 2060. *Current Population Reports*. Washington, DC: U.S. Census Bureau; 2014:25-1143
- [3] Streja D, Streja E. Management of dyslipidemia in the elderly. In: De Groot LJ, Chrousos G, Dungan K, Feingold KR, Grossman A, Hershman JM, et al., editors. *Endotext*. South Dartmouth (MA): MDText.com, Inc.; 2000
- [4] Ghandehari H, Kamal-Bahl S, Wong ND. Prevalence and extent of dyslipidemia and recommended lipid levels in US adults with and without cardiovascular comorbidities: The National Health and Nutrition Examination Survey 2003-2004. *American Heart Journal*. 2008;**156**(1):112-119 (Epub Jul 1, 2008)
- [5] Shepherd J, Blauw GJ, Murphy MB, Bollen EL, Buckley BM, Cobbe SM, et al. Pravastatin in elderly individuals at risk of vascular disease (PROSPER): A randomised controlled trial. *Lancet*. 2002;**360**(9346):1623-1630 (Epub Nov 30, 2002)
- [6] Deedwania P, Stone PH, Bairey Merz CN, Cosin-Aguilar J, Koylan N, Luo D, et al. Effects of intensive versus moderate lipid-lowering therapy on myocardial ischemia in older patients with coronary heart disease: Results of the Study Assessing Goals in the Elderly (SAGE). *Circulation*. 2007;**115**(6): 700-707 (Epub Feb 7, 2007)
- [7] Collins R, Armitage J, Parish S, Sleight P, Peto R. MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: A randomised placebo-controlled trial. *Lancet*. 2003;**361**(9374):2005-2016 (Epub Jun 20, 2003)
- [8] Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. *The New England Journal of Medicine*. 1998;**339**(19):1349-1357 (Epub Dec 5, 1998)
- [9] Sacks FM, Pfeffer MA, Moyer LA, Rouleau JL, Rutherford JD, Cole TG, et al. The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and recurrent events trial investigators. *The New England Journal of Medicine*. 1996;**335**(14):1001-1009 (Epub Oct 3, 1996)
- [10] Pedersen T et al. Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: The Scandinavian Simvastatin Survival Study (4S). *Lancet*. 1994;**344**(8934):1383-1389 (Epub Nov 19, 1994)
- [11] Jacobson TA. Overcoming 'ageism' bias in the treatment of hypercholesterolaemia: A review of safety issues with statins in the elderly. *Drug Safety*. 2006;**29**(5):421-448 (Epub May 13, 2006)
- [12] Pasternak RC, Smith SC Jr, Bairey-Merz CN, Grundy SM, Cleeman JI, Lenfant C. ACC/AHA/NHLBI clinical advisory on the use and safety of statins. *Circulation*. 2002;**106**(8):1024-1028 (Epub Aug 21, 2002)
- [13] Sever PS, Dahlof B, Poulter NR, Wedel H, Beevers G, Caulfield M, et al.

Prevention of coronary and stroke events with atorvastatin in hypertensive patients who have average or lower-than-average cholesterol concentrations, in the Anglo-Scandinavian Cardiac Outcomes Trial—Lipid Lowering Arm (ASCOT-LLA): A multicentre randomised controlled trial. *Lancet*. 2003;**361**(9364):1149-1158 (Epub Apr 11, 2003)

[14] Neil HA, DeMicco DA, Luo D, Betteridge DJ, Colhoun HM, Durrington PN, et al. Analysis of efficacy and safety in patients aged 65-75 years at randomization: Collaborative Atorvastatin Diabetes Study (CARDS). *Diabetes Care*. 2006;**29**(11):2378-2384 (Epub Oct 27, 2006)

[15] Ridker PM, Danielson E, Fonseca FA, Genest J, Gotto AM Jr, Kastelein JJ, et al. Rosuvastatin to prevent vascular events in men and women with elevated C-reactive protein. *The New England Journal of Medicine*. 2008;**359**(21):2195-2207 (Epub Nov 11, 2008)

[16] Glynn RJ, Koenig W, Nordestgaard BG, Shepherd J, Ridker PM. Rosuvastatin for primary prevention in older persons with elevated C-reactive protein and low to average low-density lipoprotein cholesterol levels: Exploratory analysis of a randomized trial. *Annals of Internal Medicine*. 2010;**152**(8):488-496, W174 (Epub Apr 21, 2010)

[17] Graham DJ, Staffa JA, Shatin D, Andrade SE, Schech SD, La Grenade L, et al. Incidence of hospitalized rhabdomyolysis in patients treated with lipid-lowering drugs. *The Journal of the American Medical Association*. 2004;**292**(21):2585-2590 (Epub Dec 2, 2004)

[18] Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron*. 1976;**16**(1):31-41 (Epub Jan 1, 1976)

[19] Wynne HA, Cope LH, Mutch E, Rawlins MD, Woodhouse KW, James OF. The effect of age upon liver volume and apparent liver blood flow in healthy man. *Hepatology*. 1989;**9**(2):297-301 (Epub Feb 1, 1989)

[20] Gupta S. P-glycoprotein expression and regulation. Age-related changes and potential effects on drug therapy. *Drugs & Aging*. 1995;**7**(1):19-29 (Epub Jul 1, 1995)

[21] Cannon CP, Blazing MA, Giugliano RP, McCagg A, White JA, Theroux P, et al. Ezetimibe added to statin therapy after acute coronary syndromes. *The New England Journal of Medicine*. 2015;**372**(25):2387-2397 (Epub Jun 4, 2015)

[22] Sabatine MS, Giugliano RP, Keech AC, Honarpour N, Wiviott SD, Murphy SA, et al. Evolocumab and clinical outcomes in patients with cardiovascular disease. *The New England Journal of Medicine*. 2017;**376**(18):1713-1722 (Epub Mar 18, 2017)

[23] Neve BP, Fruchart JC, Staels B. Role of the peroxisome proliferator-activated receptors (PPAR) in atherosclerosis. *Biochemical Pharmacology*. 2000;**60**(8):1245-1250 (Epub Sep 29, 2000)

[24] Rubins HB, Robins SJ, Collins D, Fye CL, Anderson JW, Elam MB, et al. Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. Veterans affairs high-density lipoprotein cholesterol intervention trial study group. *The New England Journal of Medicine*. 1999;**341**(6):410-418 (Epub Aug 7, 1999)

[25] Frick MH, Elo O, Haapa K, Heinonen OP, Heinsalmi P, Helo P, et al. Helsinki Heart Study: Primary-prevention trial with gemfibrozil in

- middle-aged men with dyslipidemia. Safety of treatment, changes in risk factors, and incidence of coronary heart disease. *The New England Journal of Medicine*. 1987;**317**(20):1237-1245 (Epub Nov 12, 1987)
- [26] Brunner D, Agmon J, Kaplinsky E, et al. Secondary prevention by raising HDL cholesterol and reducing triglycerides in patients with coronary artery disease. *Circulation*. 2000;**102**(1):21-27 (Epub Jul 6, 2000)
- [27] Walsh JM, Pignone M. Drug treatment of hyperlipidemia in women. *The Journal of the American Medical Association*. 2004;**291**(18):2243-2252 (Epub May 13, 2004)
- [28] Mosca L, Banka CL, Benjamin EJ, Berra K, Bushnell C, Dolor RJ, et al. Evidence-based guidelines for cardiovascular disease prevention in women: 2007 update. *Circulation*. 2007;**115**(11):1481-1501 (Epub Feb 21, 2007)
- [29] Wenger NK. Coronary disease in women. *Annual Review of Medicine*. 1985;**36**:285-294 (Epub Jan 1, 1985)
- [30] Mucha L, Stephenson J, Morandi N, Dirani R. Meta-analysis of disease risk associated with smoking, by gender and intensity of smoking. *Gender Medicine*. 2006;**3**(4):279-291 (Epub Jun 22, 2007)
- [31] Mercurio G, Longu G, Zoncu S, Cherchi A. Impaired forearm blood flow and vasodilator reserve in healthy postmenopausal women. *American Heart Journal*. 1999;**137**(4 Pt 1):692-697 (Epub Mar 31, 1999)
- [32] Pavanello C, Mombelli G. Considering gender in prescribing statins: What do physicians need to know? *Clinical Lipidology*. 2015;**10**(6):499-512
- [33] Kreisberg RA, Kasim S. Cholesterol metabolism and aging. *The American Journal of Medicine*. 1987;**82**(1B):54-60 (Epub Jan 26, 1987)
- [34] Mosca L, Grundy SM, Judelson D, King K, Limacher M, Oparil S, et al. Guide to preventive cardiology for women. AHA/ACC scientific statement consensus panel statement. *Circulation*. 1999;**99**(18):2480-2484 (Epub May 11, 1999)
- [35] Mosca L, Benjamin EJ, Berra K, Bezanson JL, Dolor RJ, Lloyd-Jones DM, et al. Effectiveness-based guidelines for the prevention of cardiovascular disease in women—2011 update: A guideline from the American Heart Association. *Circulation*. 2011;**123**(11):1243-1262 (Epub Feb 18, 2011)
- [36] Reiner Z, Catapano AL, De Backer G, Graham I, Taskinen MR, Wiklund O, et al. ESC/EAS guidelines for the management of dyslipidaemias: The task force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *European Heart Journal*. 2011;**32**(14):1769-1818 (Epub Jun 30, 2011)
- [37] Downs JR, Clearfield M, Weis S, Whitney E, Shapiro DR, Beere PA, et al. Primary prevention of acute coronary events with lovastatin in men and women with average cholesterol levels: Results of AFCAPS/TexCAPS. Air Force/Texas Coronary Atherosclerosis Prevention Study. *The Journal of the American Medical Association*. 1998;**279**(20):1615-1622 (Epub Jun 5, 1998)
- [38] ALLHAT Collaborative Research Group authors. Major outcomes in moderately hypercholesterolemic, hypertensive patients randomized to pravastatin vs. usual care: The Antihypertensive and Lipid-Lowering Treatment to Prevent Heart Attack Trial (ALLHAT-LLT). *The Journal of the American Medical Association*. 2002;**288**(23):2998-3007. (Epub Dec 20, 2002)

- [39] Nakamura H, Arakawa K, Itakura H, Kitabatake A, Goto Y, Toyota T, et al. Primary prevention of cardiovascular disease with pravastatin in Japan (MEGA Study): A prospective randomised controlled trial. *Lancet*. 2006;**368**(9542):1155-1163 (Epub Oct 3, 2006)
- [40] Mora S, Glynn RJ, Hsia J, MacFadyen JG, Genest J, Ridker PM. Statins for the primary prevention of cardiovascular events in women with elevated high-sensitivity C-reactive protein or dyslipidemia: Results from the Justification for the Use of Statins in Prevention: An Intervention Trial Evaluating Rosuvastatin (JUPITER) and meta-analysis of women from primary prevention trials. *Circulation*. 2010;**121**(9):1069-1077 (Epub Feb 24, 2010)
- [41] Miettinen TA, Pyorala K, Olsson AG, Musliner TA, Cook TJ, Faergeman O, et al. Cholesterol-lowering therapy in women and elderly patients with myocardial infarction or angina pectoris: Findings from the Scandinavian Simvastatin Survival Study (4S). *Circulation*. 1997;**96**(12):4211-4218 (Epub Jan 7, 1998)
- [42] Plehn JF, Davis BR, Sacks FM, Rouleau JL, Pfeffer MA, Bernstein V, et al. Reduction of stroke incidence after myocardial infarction with pravastatin: The Cholesterol and Recurrent Events (CARE) study. The care investigators. *Circulation*. 1999;**99**(2):216-223 (Epub Jan 20, 1999)
- [43] Hague W, Forder P, Simes J, Hunt D, Tonkin A. Effect of pravastatin on cardiovascular events and mortality in 1516 women with coronary heart disease: Results from the Long-Term Intervention with Pravastatin in Ischemic Disease (LIPID) study. *American Heart Journal*. 2003;**145**(4):643-651 (Epub Apr 8, 2003)
- [44] Link E, Parish S, Armitage J, Bowman L, Heath S, Matsuda F, et al. SLCO1B1 variants and statin-induced myopathy—A genomewide study. *The New England Journal of Medicine*. 2008;**359**(8):789-799 (Epub Jul 25, 2008)
- [45] Kostis WJ, Cheng JQ, Dobrzynski JM, Cabrera J, Kostis JB. Meta-analysis of statin effects in women versus men. *Journal of the American College of Cardiology*. 2012;**59**(6):572-582 (Epub Feb 24, 2012)
- [46] Schech S, Graham D, Staffa J, Andrade SE, La Grenade L, Burgess M, et al. Risk factors for statin-associated rhabdomyolysis. *Pharmacoepidemiology and Drug Safety*. 2007;**16**(3):352-358 (Epub Aug 8, 2006)
- [47] Hadden DR, McLaughlin C. Normal and abnormal maternal metabolism during pregnancy. *Seminars in Fetal & Neonatal Medicine*. 2009;**14**(2):66-71 (Epub Nov 7, 2008)
- [48] Lain KY, Catalano PM. Metabolic changes in pregnancy. *Clinical Obstetrics and Gynecology*. 2007;**50**(4):938-948 (Epub Nov 6, 2007)
- [49] Belo L, Caslake M, Gaffney D, Santos-Silva A, Pereira-Leite L, Quintanilha A, et al. Changes in LDL size and HDL concentration in normal and preeclamptic pregnancies. *Atherosclerosis*. 2002;**162**(2):425-432 (Epub May 9, 2002)
- [50] Potter JM, Nestel PJ. The hyperlipidemia of pregnancy in normal and complicated pregnancies. *American Journal of Obstetrics and Gynecology*. 1979;**133**(2):165-170 (Epub Jan 15, 1979)
- [51] Fanshawe AE, Ibrahim M. The current status of lipoprotein (a) in pregnancy: A literature review. *Journal of Cardiology*. 2013;**61**(2):99-106 (Epub Nov 21, 2012)
- [52] Manten GT, Franx A, van der Hoek YY, Hameeteman TM, Voorbij HA, Smolders HC, et al. Changes of plasma

lipoprotein(a) during and after normal pregnancy in Caucasians. *The Journal of Maternal-Fetal & Neonatal Medicine : The Official Journal of the European Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstet.* 2003;**14**(2):91-95 (Epub Nov 25, 2003)

[53] Vrijkotte TG, Krukziener N, Hutten BA, Vollebregt KC, van Eijnsden M, Twickler MB. Maternal lipid profile during early pregnancy and pregnancy complications and outcomes: The ABCD study. *The Journal of Clinical Endocrinology and Metabolism.* 2012;**97**(11):3917-3925 (Epub Aug 31, 2012)

[54] Enquobahrie DA, Williams MA, Butler CL, Frederick IO, Miller RS, Luthy DA. Maternal plasma lipid concentrations in early pregnancy and risk of preeclampsia. *American Journal of Hypertension.* 2004;**17**(7):574-581 (Epub Jul 6, 2004)

[55] Catov JM, Ness RB, Wellons MF, Jacobs DR, Roberts JM, Gunderson EP. Prepregnancy lipids related to preterm birth risk: The coronary artery risk development in young adults study. *The Journal of Clinical Endocrinology and Metabolism.* 2010;**95**(8):3711-3718 (Epub May 27, 2010)

[56] Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of in utero and early-life conditions on adult health and disease. *The New England Journal of Medicine.* 2008;**359**(1):61-73 (Epub Jul 4, 2008)

[57] Hemmelgarn BR, Clement F, Manns BJ, Klarenbach S, James MT, Ravani P, et al. Overview of the Alberta kidney disease network. *BMC Nephrology.* 2009;**10**:30 (Epub Oct 21, 2009)

[58] Ahmed MH, Khalil AA. Ezetimibe as a potential treatment for dyslipidemia

associated with chronic renal failure and renal transplant. *Saudi Journal of Kidney Diseases and Transplantation: An Official Publication of the Saudi Center for Organ Transplantation, Saudi Arabia.* 2010;**21**(6):1021-1029 (Epub Nov 10, 2010)

[59] Holzer M, Zangger K, El-Gamal D, Binder V, Curcic S, Konya V, et al. Myeloperoxidase-derived chlorinating species induce protein carbamylation through decomposition of thiocyanate and urea: Novel pathways generating dysfunctional high-density lipoprotein. *Antioxidants & Redox Signaling.* 2012;**17**(8):1043-1052 (Epub Apr 3, 2012)

[60] Kwan BC, Kronenberg F, Beddhu S, Cheung AK. Lipoprotein metabolism and lipid management in chronic kidney disease. *Journal of the American Society of Nephrology.* 2007;**18**(4):1246-1261 (Epub Mar 16, 2007)

[61] Mikolasevic I, Zutelija M, Mavrinac V, Orlic L. Dyslipidemia in patients with chronic kidney disease: Etiology and management. *International Journal of Nephrology and Renovascular Disease.* 2017;**10**:35-45 (Epub Feb 23, 2017)

[62] Batista MC, Welty FK, Diffenderfer MR, Sarnak MJ, Schaefer EJ, Lamon-Fava S, et al. Apolipoprotein A-I, B-100, and B-48 metabolism in subjects with chronic kidney disease, obesity, and the metabolic syndrome. *Metabolism: Clinical and Experimental.* 2004;**53**(10):1255-1261 (Epub Sep 18, 2004)

[63] McNamara JR, Shah PK, Nakajima K, Cupples LA, Wilson PW, Ordovas JM, et al. Remnant-like particle (RLP) cholesterol is an independent cardiovascular disease risk factor in women: Results from the Framingham heart study. *Atherosclerosis.* 2001;**154**(1):229-236 (Epub Jan 4, 2001)

[64] Oi K, Hirano T, Sakai S, Kawaguchi Y, Hosoya T. Role of hepatic lipase in

intermediate-density lipoprotein and small, dense low-density lipoprotein formation in hemodialysis patients. *Kidney International Supplement*. 1999;**71**:S227-S228 (Epub Jul 21, 1999)

[65] Koch M, Kutkuhn B, Trenkwalder E, Bach D, Grabensee B, Dieplinger H, et al. Apolipoprotein B, fibrinogen, HDL cholesterol, and apolipoprotein(a) phenotypes predict coronary artery disease in hemodialysis patients. *Journal of the American Society of Nephrology*. 1997;**8**(12):1889-1898 (Epub Dec 24, 1997)

[66] Ikewaki K, Schaefer JR, Frischmann ME, Okubo K, Hosoya T, Mochizuki S, et al. Delayed in vivo catabolism of intermediate-density lipoprotein and low-density lipoprotein in hemodialysis patients as potential cause of premature atherosclerosis. *Arteriosclerosis, Thrombosis, and Vascular Biology*. 2005;**25**(12):2615-2622 (Epub Oct 1, 2005)

[67] Miljkovic M, Stefanovic A, Simic-Ogrizovic S, Vekic J, Bogavac-Stanojevic N, Cerne D, et al. Association of dyslipidemia, oxidative stress, and inflammation with redox status in VLDL, LDL, and HDL lipoproteins in patients with renal disease. *Angiology*. 2018;**69**(10):861-870. (Epub Jun 19, 2018)

[68] Santana JM, Brown CD. High-density lipoprotein carbamylation and dysfunction in vascular disease. *Frontiers in Bioscience (Landmark Ed)*. 2018;**23**:2227-2234 (Epub May 18, 2018)

[69] Dantoine TF, Debord J, Charmes JP, Merle L, Marquet P, Lachatre G, et al. Decrease of serum paraoxonase activity in chronic renal failure. *Journal of the American Society of Nephrology*. 1998;**9**(11):2082-2088 (Epub Nov 10, 1998)

[70] Chang TI, Streja E, Soohoo M, Ko GJ, Rhee CM, Kovesdy CP, et al. Increments in serum high-density lipoprotein

cholesterol over time are not associated with improved outcomes in incident hemodialysis patients. *Journal of Clinical Lipidology*. 2018;**12**(2):488-497 (Epub Feb 20, 2018)

[71] Heart Protection Study Collaborative Group authors. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: A randomised placebo-controlled trial. *Lancet*. 2002;**360**(9326):7-22. (Epub Jul 13, 2002)

[72] Tonelli M, Isles C, Curhan GC, Tonkin A, Pfeffer MA, Shepherd J, et al. Effect of pravastatin on cardiovascular events in people with chronic kidney disease. *Circulation*. 2004;**110**(12):1557-1563 (Epub Sep 15, 2004)

[73] Koren MJ, Davidson MH, Wilson DJ, Fayyad RS, Zuckerman A, Reed DP. Focused atorvastatin therapy in managed-care patients with coronary heart disease and CKD. *American Journal of Kidney Diseases :The Official Journal of the National Kidney Foundation*. 2009;**53**(5):741-750 (Epub Feb 14, 2009)

[74] Shepherd J, Kastelein JJ, Bittner V, Deedwania P, Breazna A, Dobson S, et al. Effect of intensive lipid lowering with atorvastatin on renal function in patients with coronary heart disease: The treating to new targets (TNT) study. *Clinical Journal of the American Society of Nephrology*. 2007;**2**(6):1131-1139 (Epub Oct 19, 2007)

[75] Ridker PM, Macfadyen JG, Nordestgaard BG, Koenig W, Kastelein JJ, Genest J, et al. Rosuvastatin for primary prevention among individuals with elevated high-sensitivity c-reactive protein and 5% to 10% and 10% to 20% 10-year risk. Implications of the Justification for Use of Statins in Prevention: An Intervention Trial Evaluating Rosuvastatin (JUPITER)

- trial for “intermediate risk”. *Circulation Cardiovascular Quality and Outcomes*. 2010;**3**(5):447-452 (Epub Aug 26, 2010)
- [76] Baigent C, Landray MJ, Reith C, Emberson J, Wheeler DC, Tomson C, et al. The effects of lowering LDL cholesterol with simvastatin plus ezetimibe in patients with chronic kidney disease (Study of Heart and Renal Protection): A randomised placebo-controlled trial. *Lancet*. 2011;**377**(9784):2181-2192 (Epub Jun 15, 2011)
- [77] Tonelli M, Wanner C. Lipid management in chronic kidney disease: Synopsis of the kidney disease: Improving global outcomes 2013 clinical practice guideline. *Annals of Internal Medicine*. 2014;**160**(3):182 (Epub Dec 11, 2013)
- [78] Fellstrom BC, Jardine AG, Schmieder RE, Holdaas H, Bannister K, Beutler J, et al. Rosuvastatin and cardiovascular events in patients undergoing hemodialysis. *The New England Journal of Medicine*. 2009;**360**(14):1395-1407 (Epub Apr 1, 2009)
- [79] Wanner C, Krane V, Marz W, Olschewski M, Asmus HG, Kramer W, et al. Randomized controlled trial on the efficacy and safety of atorvastatin in patients with type 2 diabetes on hemodialysis (4D study): Demographic and baseline characteristics. *Kidney & Blood Pressure Research*. 2004;**27**(4):259-266 (Epub Aug 19, 2004)
- [80] Jacobson TA, Maki KC, Orringer CE, Jones PH, Kris-Etherton P, Sikand G, et al. National lipid association recommendations for patient-centered management of dyslipidemia: Part 2. *Journal of Clinical Lipidology*. 2015;**9**(6 Suppl):S1-122.e1 (Epub Dec 25, 2015)
- [81] Kobashigawa JA, Kasiske BL. Hyperlipidemia in solid organ transplantation. *Transplantation*. 1997;**63**(3):331-338 (Epub Feb 15, 1997)
- [82] Pagano G, Bruno A, Cavallo-Perin P, Cesco L, Imbimbo B. Glucose intolerance after short-term administration of corticosteroids in healthy subjects. Prednisone, deflazacort, and betamethasone. *Archives of Internal Medicine*. 1989;**149**(5):1098-1101 (Epub May 1, 1989)
- [83] Agarwal A, Prasad GV. Post-transplant dyslipidemia: Mechanisms, diagnosis and management. *World Journal of Transplantation*. 2016;**6**(1):125-134 (Epub Mar 25, 2016)
- [84] de Groen PC. Cyclosporine, low-density lipoprotein, and cholesterol. *Mayo Clinic Proceedings*. 1988;**63**(10):1012-1021 (Epub Oct 1, 1988)
- [85] Morrisett JD, Abdel-Fattah G, Hoogeveen R, Mitchell E, Ballantyne CM, Pownall HJ, et al. Effects of sirolimus on plasma lipids, lipoprotein levels, and fatty acid metabolism in renal transplant patients. *Journal of Lipid Research*. 2002;**43**(8):1170-1180 (Epub Aug 15, 2002)
- [86] Holdaas H, Fellstrom B, Jardine AG, Holme I, Nyberg G, Fauchald P, et al. Effect of fluvastatin on cardiac outcomes in renal transplant recipients: A multicentre, randomised, placebo-controlled trial. *Lancet*. 2003;**361**(9374):2024-2031 (Epub Jun 20, 2003)
- [87] Holdaas H, Fellstrom B, Cole E, Nyberg G, Olsson AG, Pedersen TR, et al. Long-term cardiac outcomes in renal transplant recipients receiving fluvastatin: The ALERT extension study. *American Journal of Transplantation* :

Official Journal of the American Society of Transplantation and the American Society of Transplant Surgeons. 2005;5(12):2929-2936 (Epub Nov 24, 2005)

[88] Klein D, Hurley LB, Quesenberry CP Jr, Sidney S. Do protease inhibitors increase the risk for coronary heart disease in patients with HIV-1 infection? *Journal of Acquired Immune Deficiency Syndromes*. 2002;30(5):471-477 (Epub Aug 3, 2002)

[89] Freiberg MS, Chang CC, Kuller LH, Skanderson M, Lowy E, Kraemer KL, et al. HIV infection and the risk of acute myocardial infarction. *JAMA Internal Medicine*. 2013;173(8):614-622 (Epub Mar 6, 2013)

[90] El-Sadr WM, Lundgren J, Neaton JD, Gordin F, Abrams D, Arduino RC, et al. CD4+ count-guided interruption of antiretroviral treatment. *The New England Journal of Medicine*. 2006;355(22):2283-2296 (Epub Dec 1, 2006)

[91] Paisible AL, Chang CC, So-Armah KA, Butt AA, Leaf DA, Budoff M, et al. HIV infection, cardiovascular disease risk factor profile, and risk for acute myocardial infarction. *Journal of Acquired Immune Deficiency Syndromes*. 2015;68(2):209-216 (Epub Jan 15, 2015)

[92] Triant VA. Epidemiology of coronary heart disease in patients with human immunodeficiency virus. *Reviews in Cardiovascular Medicine*. 2014;15(Suppl 1):S1-S8 (Epub Jul 6, 2014)

[93] Rahmanian S, Wewers ME, Koletar S, Reynolds N, Ferketich A, Diaz P. Cigarette smoking in the HIV-infected population. *Proceedings of the American Thoracic Society*. 2011;8(3):313-319 (Epub Jun 10, 2011)

[94] Grinspoon S, Carr A. Cardiovascular risk and body-fat abnormalities in HIV-infected adults. *The New England Journal of Medicine*. 2005;352(1):48-62 (Epub Jan 7, 2005)

[95] Gelpi M, Afzal S, Lundgren J, Ronit A, Roen A, Mocroft A, et al. Higher risk of abdominal obesity, elevated LDL cholesterol and hypertriglyceridemia, but not of hypertension, in people living with HIV: Results from the Copenhagen Comorbidity in HIV Infection (COCOMO) Study. *Clinical Infectious Diseases: An Official Publication of the Infectious Diseases Society of America*. 1 August 2018;67(4):579-586. (Epub Feb 23, 2018)

[96] Micheletti RG, Fishbein GA, Fishbein MC, Singer EJ, Weiss RE, Jeffries RA, et al. Coronary atherosclerotic lesions in human immunodeficiency virus-infected patients: A histopathologic study. *Cardiovascular Pathology: The Official Journal of the Society for Cardiovascular Pathology*. 2009;18(1):28-36 (Epub Apr 12, 2008)

[97] Post WS, Budoff M, Kingsley L, Palella FJ Jr, Witt MD, Li X, et al. Associations between HIV infection and subclinical coronary atherosclerosis. *Annals of Internal Medicine*. 2014;160(7):458-467 (Epub Apr 2, 2014)

[98] Burdo TH, Lo J, Abbara S, Wei J, DeLelys ME, Preffer F, et al. Soluble CD163, a novel marker of activated macrophages, is elevated and associated with noncalcified coronary plaque in HIV-infected patients. *The Journal of Infectious Diseases*. 2011;204(8):1227-1236 (Epub Sep 16, 2011)

[99] Lo J, Abbara S, Shturman L, Soni A, Wei J, Rocha-Filho JA, et al. Increased prevalence of subclinical coronary atherosclerosis detected by coronary computed tomography

angiography in HIV-infected men. *AIDS*. 2010;**24**(2):243-253 (Epub Dec 10, 2009)

[100] Tawakol A, Lo J, Zanni MV, Marmarelis E, Ihenachor EJ, MacNabb M, et al. Increased arterial inflammation relates to high-risk coronary plaque morphology in HIV-infected patients. *Journal of Acquired Immune Deficiency Syndromes*. 2014;**66**(2):164-171 (Epub May 16, 2014)

[101] Post WS, George RT, Budoff M. HIV infection and subclinical coronary atherosclerosis. *Annals of Internal Medicine*. 2014;**161**(12):923-924 (Epub Dec 17, 2014)

[102] Hulten E, Mitchell J, Scally J, Gibbs B, Villines TC. HIV positivity, protease inhibitor exposure and subclinical atherosclerosis: A systematic review and meta-analysis of observational studies. *Heart*. 2009;**95**(22):1826-1835 (Epub Jul 28, 2009)

[103] Knudsen AD, Gelpi M, Afzal S, Ronit A, Roen A, Mocroft A, et al. Prevalence of peripheral artery disease is higher in persons living with HIV compared to uninfected controls. *Journal of Acquired Immune Deficiency Syndromes*. 2018 (Epub Jul 10, 2018)

[104] Stanley TL, Grinspoon SK. Body composition and metabolic changes in HIV-infected patients. *The Journal of Infectious Diseases*. 2012;**205**(Suppl 3):S383-S390 (Epub May 18)

[105] Silverberg MJ, Leyden WA, Xu L, Horberg MA, Chao CR, Towner WJ, et al. Immunodeficiency and risk of myocardial infarction among HIV-positive individuals with access to care. *Journal of Acquired Immune Deficiency Syndromes*. 2014;**65**(2):160-166 (Epub Jan 21, 2014)

[106] Deeks SG. HIV infection, inflammation, immunosenescence,

and aging. *Annual Review of Medicine*. 2011;**62**:141-155 (Epub Nov 26, 2010)

[107] Chow FC. HIV infection, vascular disease, and stroke. *Seminars in Neurology*. 2014;**34**(1):35-46 (Epub Apr 10, 2014)

[108] Ganesan A, Crum-Cianflone N, Higgins J, Qin J, Rehm C, Metcalf J, et al. High dose atorvastatin decreases cellular markers of immune activation without affecting HIV-1 RNA levels: Results of a double-blind randomized placebo controlled clinical trial. *The Journal of Infectious Diseases*. 2011;**203**(6):756-764 (Epub Feb 18, 2011)

[109] Calza L, Trapani F, Bartoletti M, Manfredi R, Colangeli V, Borderi M, et al. Statin therapy decreases serum levels of high-sensitivity C-reactive protein and tumor necrosis factor-alpha in HIV-infected patients treated with ritonavir-boosted protease inhibitors. *HIV Clinical Trials*. 2012;**13**(3):153-161 (Epub May 18, 2012)

[110] Holland AT, Palaniappan LP. Problems with the collection and interpretation of Asian-American health data: Omission, aggregation, and extrapolation. *Annals of Epidemiology*. 2012;**22**(6):397-405 (Epub May 26, 2012)

[111] Miller M, Stone NJ, Ballantyne C, Bittner V, Criqui MH, Ginsberg HN, et al. Triglycerides and cardiovascular disease: A scientific statement from the American Heart Association. *Circulation*. 2011;**123**(20):2292-2333 (Epub Apr 20, 2011)

[112] Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, et al. Heart disease and stroke statistics—2017 update: A report from the American Heart Association. *Circulation*. 2017;**135**(10):e146-e603 (Epub Jan 27, 2017)

- [113] Frank AT, Zhao B, Jose PO, Azar KM, Fortmann SP, Palaniappan LP. Racial/ethnic differences in dyslipidemia patterns. *Circulation*. 2014;**129**(5):570-579 (Epub Nov 7, 2013)
- [114] Anand SS, Yusuf S, Vuksan V, Devanese S, Teo KK, Montague PA, et al. Differences in risk factors, atherosclerosis, and cardiovascular disease between ethnic groups in Canada: The study of health assessment and risk in ethnic groups (SHARE). *Lancet*. 2000;**356**(9226):279-284 (Epub Nov 9, 2000)
- [115] Goff DC Jr, Bertoni AG, Kramer H, Bonds D, Blumenthal RS, Tsai MY, et al. Dyslipidemia prevalence, treatment, and control in the multi-ethnic study of atherosclerosis (MESA): Gender, ethnicity, and coronary artery calcium. *Circulation*. 2006;**113**(5):647-656 (Epub Feb 8, 2006)
- [116] Rodriguez CJ, Allison M, Daviglus ML, Isasi CR, Keller C, Leira EC, et al. Status of cardiovascular disease and stroke in Hispanics/Latinos in the United States: A science advisory from the American Heart Association. *Circulation*. 2014;**130**(7):593-625 (Epub Aug 8, 2014)
- [117] Hoyert DL. 75 years of mortality in the United States, 1935-2010. *NCHS Data Brief*. 2012;**88**:1-8 (Epub May 24, 2012)
- [118] Marmot MG, Syme SL, Kagan A, Kato H, Cohen JB, Belsky J. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii and California: Prevalence of coronary and hypertensive heart disease and associated risk factors. *American Journal of Epidemiology*. 1975;**102**(6):514-525 (Epub Dec 1, 1975)
- [119] Enas EA, Garg A, Davidson MA, Nair VM, Huet BA, Yusuf S. Coronary heart disease and its risk factors in first-generation immigrant Asian Indians to the United States of America. *Indian Heart Journal*. 1996;**48**(4):343-353 (Epub Jul 1, 1996)
- [120] Balarajan R. Ethnic differences in mortality from ischaemic heart disease and cerebrovascular disease in England and Wales. *British Medical Journal*. 1991;**302**(6776):560-564 (Epub Mar 9, 1991)
- [121] Gupta M, Brister S. Is south Asian ethnicity an independent cardiovascular risk factor? *The Canadian Journal of Cardiology*. 2006;**22**(3):193-197 (Epub Mar 8, 2006)
- [122] Pencina MJ, Navar-Boggan AM, D'Agostino RB Sr, Williams K, Neely B, Sniderman AD, et al. Application of new cholesterol guidelines to a population-based sample. *The New England Journal of Medicine*. 2014;**370**(15):1422-1431 (Epub Mar 22, 2014)
- [123] Goff DC Jr, Lloyd-Jones DM, Bennett G, Coady S, D'Agostino RB Sr, Gibbons R, et al. ACC/AHA guideline on the assessment of cardiovascular risk: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology*. 2013, 2014;**63** (25 Pt B):2935-2959 (Epub Nov 19, 2013)
- [124] Wilson PW, Pencina MJ. Estimation of risk for initial atherosclerotic cardiovascular disease events: Taking stock and moving forward. *Circulation*. 2016;**134**(23):1792-1793 (Epub Dec 7, 2016)
- [125] Nishizato Y, Ieiri I, Suzuki H, Kimura M, Kawabata K, Hirota T, et al. Polymorphisms of OATP-C (SLC21A6) and OAT3 (SLC22A8) genes: Consequences for pravastatin pharmacokinetics. *Clinical Pharmacology and Therapeutics*. 2003;**73**(6):554-565 (Epub Jun 18, 2003)

[126] Niemi M. Transporter pharmacogenetics and statin toxicity. *Clinical Pharmacology and Therapeutics*. 2010;**87**(1):130-133 (Epub Nov 6, 2009)

[127] Pu J, Romanelli R, Zhao B, Azar KM, Hastings KG, Nimbal V, et al. Dyslipidemia in special ethnic populations. *Cardiology Clinics*. 2015;**33**(2):325-333 (Epub May 6, 2015)

[128] Lee E, Ryan S, Birmingham B, Zalikowski J, March R, Ambrose H, et al. Rosuvastatin pharmacokinetics and pharmacogenetics in white and Asian subjects residing in the same environment. *Clinical Pharmacology and Therapeutics*. 2005;**78**(4):330-341 (Epub Oct 4, 2005)

[129] Zhang W, Yu BN, He YJ, Fan L, Li Q, Liu ZQ, et al. Role of BCRP 421C>A polymorphism on rosuvastatin pharmacokinetics in healthy Chinese males. *Clinica Chimica Acta: International Journal of Clinical Chemistry*. 2006;**373**(1-2):99-103 (Epub Jun 21, 2006)

[130] Ho RH, Choi L, Lee W, Mayo G, Schwarz UI, Tirona RG, et al. Effect of drug transporter genotypes on pravastatin disposition in European- and African-American participants. *Pharmacogenetics and Genomics*. 2007;**17**(8):647-656 (Epub Jul 12, 2007)

[131] HPS2-THRIVE randomized placebo-controlled trial in 25 673 high-risk patients of ER niacin/laropiprant: Trial design, pre-specified muscle and liver outcomes, and reasons for stopping study treatment. *European Heart Journal*. 2013;**34**(17):1279-1291 (Epub Feb 28, 2013)

[132] Landray MJ, Haynes R, Hopewell JC, Parish S, Aung T, Tomson J, et al. Effects of extended-release niacin with laropiprant in high-risk patients. *The New England Journal of Medicine*. 2014;**371**(3):203-212 (Epub Jul 12, 2014)

[133] Schoenborn CA, Adams PF, Peregoy JA. Health behaviors of adults: United States, 2008-2010. *Vital and health statistics series 10. Data from the National Health Survey*. 2013;**257**:1-184 (Epub May 1, 2013)