

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Current Pathophysiological and Genetic Aspects of Dilated Cardiomyopathy

Deborah P. Schild, Sascha I. Ricciardi, Jens G. Hellige, Rolf Vogel and Nisha Arenja

Abstract

Dilated cardiomyopathy is the most common form of cardiomyopathy and the second leading cause of left ventricular dysfunction with highly variable clinical presentation and prognosis. The clinical courses vary and are strongly heterogeneous, ranging from asymptomatic patients to those suffering from intractable heart failure or sudden cardiac death due to arrhythmias. Previous studies have reported a 10 years cardiovascular mortality up to 40% in developed countries, due to advanced heart failure or sudden cardiac death. However, the prognosis of dilated cardiomyopathy patients is variable and depends on multiple risk factors. This chapter provides a review of dilated cardiomyopathy with specific focus on the pathophysiological aspects and genetic etiology of the disease.

Keywords: dilated cardiomyopathy, pathophysiology, etiology, diagnostics, therapy

1. Introduction

Dilated cardiomyopathy (DCM) is one of the most common cardiomyopathies causing heart failure (HF) worldwide. Although it is less common than coronary artery disease (CAD), it affects mainly young adults and presents the most frequent reason for cardiac transplantation in young age [1, 2]. According to the European Society of Cardiology (ESC), the current definition of DCM includes the presence of a dilated and poorly functioning left ventricle or of both ventricles [3]. A heterogeneous group of myocardial and systemic conditions may cause left ventricular dilatation in combination with dysfunction. In fact, identifying the etiology of DCM can be very challenging, which often leads to the common terminology of idiopathic dilated cardiomyopathy (IDC). Furthermore, DCM has a highly variable clinical presentation. While signs and symptoms of HF are most common, some patients are incidentally, for instance, by diagnosing cardiomegaly in chest X-ray. Other symptoms include arrhythmias, conduction disturbances, thromboembolic complications, or sudden cardiac death (SCD). In the last decades, major advances have been made in the understanding of molecular and genetic issues, as well as in the pathophysiology and clinical assessment of cardiomyopathies. Especially, understanding the genetic basis of DCM has improved considerably with the availability of genetic analysis. In addition, other important diagnostic approaches, particularly imaging methods are more widely available. This allows early diagnosis

and the initiation of adequate therapy, which already has led to an improvement of the prognosis in the last years. Data have shown that approximately 25% of DCM patients with recent onset of HF (<6 months) have ameliorated in symptoms or even have complete cardiac function recovery [4, 5]. However, the overall prognosis in patients with symptomatic HF and DCM is still poor. In order to further improve the prognosis of our patients, it is important to regularly gather the knowledge about the current state of DCM and adapt the appropriate diagnostic workups. Therefore, this chapter provides the reader with a comprehensive overview of the current state of DCM from definition over etiology including the genetics aspects to management for cardiovascular specialists.

2. Dilated cardiomyopathy: a frequent cardiac disease

2.1 Definition

In 1980, the World Health Organization (WHO) defined cardiomyopathies as “heart muscle diseases of unknown cause” [6]. The definition was established to distinguish cardiomyopathies from cardiac diseases with known entities such as hypertension, ischemic, or valvular heart disease. About 15 years later, the WHO/International Society and Federation of Cardiology (ISFC) Task Force classified cardiomyopathies according to anatomy and physiology into dilated cardiomyopathy (DCM), hypertrophic cardiomyopathy (HCM), restrictive cardiomyopathy (RCM), arrhythmogenic right ventricular cardiomyopathy/dysplasia (ARVC/D) and unclassified cardiomyopathies. The definitions and classifications of cardiomyopathies changed over the years. The American Heart Association (AHA) presented a new scheme in 2006 that combined genetic and clinical criteria [7]. In contrast to that, the ESC defined in 2008 cardiomyopathies in a more clinically oriented classification system according to the WHO scheme as myocardial disorders with structurally and functionally abnormalities of the heart muscle in the absence of coronary artery disease, hypertension, valvular disease, or congenital heart disease [3]. As mentioned above, right ventricular dilatation and dysfunction may be present but are not necessary for the diagnosis. Further, DCM is classified into familial/genetic and nonfamilial/nongenetic forms [3]. Recently, a revised definition of DCM was presented by the ESC working group on myocardial and pericardial diseases [8]. This position paper did not modify the above-mentioned definition, it rather added the important value of diverse etiologies and clinical manifestations with an introduction of a novel concept of a continuous phenotypic DCM spectrum, ranging from the subclinical mutation carriers to fully expressed DCM phenotypes.

2.2 Prevalence

The prevalence of DCM in the general population is unknown. The reported numbers considerably vary due to inhomogeneous study methodologies, which are mainly related to inconsistent definitions and classifications of DCM. In addition, the prevalence of DCM varies according to geographic and ethnic differences. One of the first reports was derived from an epidemiological study conducted in Olmsted County, Minnesota, from 1975 to 1984. For the diagnosis of DCM, the authors implemented echocardiography, angiography, and autopsy cases [9]. They presented a prevalence of 36.5/100,000 individuals or 1 in 2700 with a male to female ratio of 3:4 in a European-American population. A much higher prevalence with 1:250 was reported recently by a study from Hershberger et al. The authors estimated the DCM prevalence based on the known ratio of idiopathic DCM to

HCM of $\approx 2:1$. In addition, clinical data of HF patients and left ventricular dysfunction were used as a surrogate for DCM [10]. In Western countries, 25–40% of DCM patients have been described with evidence of familial DCM with predominantly autosomal dominant inheritance [4, 11–13]. In general, epidemiologic studies have stated a rate of 20 up to >50% of familial DCM in patients, who were initially diagnosed with IDC [9, 11, 14–18].

2.3 Clinical presentation

The first presentation of patients with DCM is often characterized by signs and symptoms of HF, such as dyspnea, ankle swelling, fatigue, elevated jugular venous pressure, pulmonary rales, and peripheral edema (**Table 1**). These result from reduced cardiac function with low output and/or elevated intracardiac pressures. Other clinical manifestations include chest pain caused by reduced coronary blood flow or congestion, palpitations, and syncope or sudden cardiac death (SCD). Arrhythmia results from multifactorial reasons, which are present in DCM and include structural changes with myocardial fibrosis and left ventricular dilation as well as electrophysiological changes [19]. The risk of SCD is highly heterogeneous and depends on etiologies and risk factors [20]. Some genetic constellations may be associated with arrhythmias out of proportion to the degree of left ventricular dysfunction (e.g., pathogenic variants in DES, LMNA, and SCN5A) [21]. This is described later in this chapter (see Sections 2.7.3 and 2.8). Furthermore, patients might present with pulmonary and systemic thromboembolism [22]. Possible clinical findings are summarized in **Table 1**.

	Cardiac	Pulmonary	Gastrointestinal/ urogenital	Other
Symptoms	Chest pain Palpitations Syncope	Dyspnea Wheeze Blood-tinged sputum Cough	Reduced appetite Epigastric pain Bloating Obstipation Diarrhea Swollen abdomen Nycturia Reduced libido Erectile dysfunction	Dizziness Lightheadness Concentration disturbances Fatigue Fainting Sweating Leg swelling Symptoms of multisystem disease Skin alterations Visual and hearing impairment Gait disturbance
Clinical signs	Hypotonia Tachycardia Pulsus alternans Arrhythmia Third heart sound Loud 2nd heart sound Displaced apex beat Elevated jugular venous pressure Eedema	Tachycardia Tachypnoea Low oxygen saturation Hemoptysis Rales Obstructive lung auscultation Diminished lung sounds due to pleural effusions Pulmonary edema	Hepatomegaly Ascites Jaundice in terminal liver failure Cachexia	Intellectual disability Deafness Myotonia Hyperpigmentation Palmoplantar keratoderma Woolly hair Dysmorphic appearance Polyneuropathic symptoms Carpal tunnel syndrome

Table 1.
Clinical findings in dilated cardiomyopathy.

2.4 Pathophysiological aspects and etiology

As a consequence of the definition, the etiology of left ventricular dilatation and dysfunction is heterogeneous. In developed countries, CAD is the most common cause of left ventricular dilatation and dysfunction, and responsible for approximately 50–70% of HF patients. Therefore, potentially reversible myocardial ischemia must always be excluded for the diagnosis of DCM. The following section describes in detail the possible causes of the DCM (**Figure 1**). However, the current literature underscores that the cause of DCM remains unknown, that is, IDC, in half of the patients [23].

2.4.1 Non-genetic causes of DCM

2.4.1.1 Inflammatory cardiomyopathy

DCM can occur after a cardiac infection or inflammation as an early (e.g. giant-cell myocarditis) or late stage disease. Typically, the active or fulminant myocarditis appears with preserved left ventricular size, while in contrast inflammatory DCM is defined as the presence of chronic inflammatory cells in association with left ventricular dilatation and reduced fraction [3]. The inflammatory myocarditis can result from an infection or may be mediated by autoimmune mechanisms. The infectious myocarditis is commonly caused by viral pathogens as an acute or chronic disease [24]. In the developed countries until the 1990s, the most frequently reported viruses were adenoviruses and enteroviruses. Recently, parvovirus B12 and human herpes virus-6 are increasingly reported causing DCM [25]. In the acute phase, the virus replicates actively within the myocardium. This leads to dysfunction of cardiomyocytes and endothelial cells, and thus triggering the immune response [23]. Commonly, the innate immune system clears the viral load whereas insufficient immune response results in viral persistence and progressive myocyte destruction [25]. The secondary effect is triggered by primed T-cells. In addition, some host myocardial cellular antigens may share epitopic similarities with viral antigens and induce an autoimmune response with further destruction of cardiomyocytes [23].




Figure 1.
Etiologies of the dilated cardiomyopathy.

Another important viral-related form of DCM is HIV syndrome. DCM in HIV patients is called HIV-associated cardiomyopathy. Autopsies demonstrated histological evidence of myocarditis in around 50% of patients, who died of AIDS-related illness [26]. Bacterial infections such as brucellosis, diphtheria, psittacosis, and typhoid fever are also known to cause (peri-) myocarditis [27–30]. In addition, *E. coli* bacteraemia have been described to induce myocarditis [31]. Spirochaetal myocarditis may be encountered in the setting of the Lyme disease (*Borrelia burgdorferi*), the Weil disease (Leptospirosis), and syphilis (*Treponema pallidum*) [32]. In endemic areas, the protozoan parasite *Trypanosoma cruzi* is a typical cause of DCM due to acute cardiac infection (perimyocarditis), as well as chronic myocardial fibrosis leading to DCM [33]. There are several proposed mechanisms leading to DCM in autoimmune disorders. These include immune-mediated myocarditis, progressive fibrosis, apoptosis with resultant restrictive and dilated phenotypes, and progressive atherosclerosis with subsequent ischemic cardiomyopathy [26]. Other causes for a non-infectious myocarditis are: Kawasaki disease in children with coronary vasculitis and systemic lupus erythematosus, which can affect the myocardium without involvement of the pericardium. In rare cases, connective tissue diseases such as scleroderma, rheumatoid arthritis, and polyarteritis nodosa may lead to DCM [26]. The non-infectious etiologies of myocarditis include drug-induced hypersensitivity and hypereosinophilic syndromes, as well as giant-cell myocarditis, which is one of the most aggressive non-infectious autoimmune disorders with rapid and devastating outcome if not treated appropriately [24, 26].

2.4.1.2 Peripartum cardiomyopathy

Peripartum cardiomyopathy (PPCM) affects predisposed teenagers and older women during the last month of pregnancy or within 5 months of delivery with typical signs and symptoms of HF [34, 35]. Suspected etiological factors include inflammatory myocarditis, autoimmunity caused by chimerism of hematopoietic lineage cells from the fetus, hemodynamic stress during pregnancy, and toxicity caused by an abnormal cleavage product of prolactin [3, 36]. Furthermore, genetic predisposition seems to be important [37].

2.4.1.3 Toxic cardiomyopathies

A number of chemical compounds are responsible for DCM. In western and developing countries, alcoholic cardiomyopathy (ACM) represents one of the most common forms of secondary cardiomyopathies resembling IDC with an estimated prevalence of 23–40% [24, 38]. More than women, men are affected by ACM and the occurrence correlates with a daily level and the duration of alcohol consumption [39]. However, the prevalence of ACM is variable and the mean daily amount of alcohol consumption, duration of regular intake, and patients' individual characteristics, including genetic susceptibility, are all related to the development of a respective cardiomyopathy [26]. Alcohol may result in both acute and chronic depression of myocardial contractility. Street drugs such as cocaine and methamphetamines are potent sympathomimetic drugs that induce inotropic and chronotropic effects. Mechanisms of cardiac toxicity include myocardial ischemia from increased oxygen consumption, prothrombotic effects, coronary vasospasms, and acceleration of coronary atherosclerosis [24]. Anthracycline-based chemotherapeutic agents are known to induce cardiac dysfunction. Acute or subacute injury can occur immediately after treatment with transient arrhythmias, pericarditis, and myocarditis [38]. The time course of chronic development of HF varies from an early onset (<1 year) to late onset (>1 year) or chronic progressive cardiomyopathy,

which can occur 10–30 years after exposure. Both chronic forms tend to be irreversible, are dosage dependent and are associated with ultrastructural changes in the cardiac myocytes [40]. Trastuzumab, a monoclonal antibody directed against the human epidermal growth factor receptor 2 (HER2), is widely used and very effective for the treatment of HER2-positive breast cancer. HER2 receptors are also localized on the cardiomyocyte. The inhibition of the Her2:Her4 signaling process in the myocardium is principally responsible for the cardiotoxicity [26]. In addition, a number of other, non-chemotherapeutic medications are associated with DCM, such as cyclophosphamide, phenothiazines, antidepressant drugs, carbon monoxide, lead, lithium, pseudoephedrine, ephedrine, cobalt, anabolic steroids, hydroxychloroquine, clozapine, and catecholamines [38, 41]. Possible causes of toxic cardiomyopathy are summarized in **Figure 2**.

2.4.1.4 Arrhythmia-induced DCM

Tachycardia-induced cardiomyopathy was described more than 100 years ago by Gossage et al. as a reversible form of systolic dysfunction caused by long-lasting supraventricular or ventricular arrhythmias [42]. Ongoing rapid atrial or ventricular pacing may result in systemic changes by neurohormonal activation, characterized by reduction in serum sodium, activation of the renin-angiotensin system, and an increase of plasma atrial natriuretic peptide, aldosterone, and norepinephrine. Abnormal myocardial and cellular remodeling occurs, which may result in DCM. Furthermore, epinephrine can also lead to abnormal myocardial and cellular remodeling, which further result in biventricular dilatation, decreased contractility, and elevation of left and/or biventricular filling pressure [26].

2.4.1.5 Infiltrative diseases

Several, systemic diseases may infiltrate the myocardium and result in DCM. Sarcoidosis, iron overload, and amyloidosis represent the most common clinical entities. Sarcoidosis is a multisystem inflammatory disease of unknown origin characterized by non-caseating granulomas in multiple organs. Sarcoidosis can progress to a fibrotic stage leading to DCM [26]. In the setting of iron overload, such as hereditary hemochromatosis, high blood volume, or parenteral iron infusions,




Figure 2.
Reasons for toxic cardiomyopathies.

DCM may occur due to iron-induced reactive oxygen species with cellular oxidative stress and fibrosis [43]. Cardiac amyloidosis is the most common cause of infiltrative cardiomyopathy with poor prognosis. Amyloid is an extracellular tissue deposition of misfolded amyloid protein-fibrils [44]. Cardiac amyloidosis usually presents as a cardiomyopathy with restrictive pathophysiology and can progress to severe cardiac dysfunction and to DCM in very advanced stages [26].

2.4.1.6 Endocrine/metabolic disorders

Endocrinologic disorders rarely lead to the phenotype of DCM. Especially thyroid hormones have a significant impact on cardiac function and structure via regulation and expression of key structural and regulatory genes like myosin heavy chain and phospholamban. Therefore, excess or deficiency of triiodothyronine (T3) may lead to DCM in a late stage [45]. Growth hormone disorders, pheochromocytoma, and Cushing's disease are also very rare causes of DCM [26, 46]. Furthermore, the literature names nutritional deficiencies of thiamine (beriberi disease), selenium (Keshan disease), and L-carnitine being responsible for DCM [47, 48].

2.4.1.7 Neuromuscular disorders

Neuromuscular disease like Duchenne's muscular dystrophy, Becker muscular dystrophy, and Emery-Dreifuss muscular dystrophy can affect the heart and induce DCM. Patients with DCM and muscular dystrophy commonly show X-linked mutations or autosomal recessive inheritance [24]. Further information is summarized in the following section on genetic causes of DCM.

2.4.2 Genetic causes of DCM

In up to 30% of the cases, a gene mutation may be identified as the main cause of DCM [49]. Thanks to advances in next-generation sequencing technologies more than 40 genes have already been identified causing DCM [50]. Most commonly, familial DCM is inherited as an autosomal dominant pattern. Autosomal recessive, X-linked, and mitochondrial inheritance patterns are less common [24]. Most of the genes involved in the development of DCM encode structural elements of the cardiomyocytes. Mutations in genes encoding sarcomeric, cytoskeletal, desmosomal, nuclear membrane, mitochondrial, and RNA-binding proteins have all been linked to DCM [51]. Interestingly, several of the gene mutations linked to autosomal dominant DCM encode the same contractile proteins that are also responsible for the development of HCM [38]. As well, other affected genes are described in ARVC and left ventricular non-compaction cardiomyopathy (NCCM). In the following, we will describe the most investigated gene mutations and their consequences on the cardiomyocyte functioning.

2.4.2.1 Sarcomere protein location: force generation and transmission

The sarcomere is the contracting unit of the myocyte. The thin filaments of the sarcomere emanate from the Z-disc. They consist of filamentous α -actin (gene name ACTC 1) and calcium-sensitive troponin-tropomyosin regulatory apparatus (encoded by TPM1), which includes the three troponin subunits (encoded by TNNT2, TNNC1, and TNNI3) [52, 53]. The thick filament core is formed by the β -myosin heavy chain (encoded by MYH7), the molecular motor of the thick filament, and the myosin-binding protein C (β -MYBPC3) [52, 53]. Titin (encoded by TTN), which is the largest human protein, spans the half length of the sarcomere,

where it acts as a stretch sensor and myofibril stabilizer. It limits sarcomere stretch in early diastole and restores resting sarcomere length after contraction [52, 53]. Titin (TTN) mutations are the most prevalent genetic cause of idiopathic DCM (15–20%). In fact, DCM patients with TTN mutations have a worse outcome due to a higher arrhythmic risk and progressive functional deficits [54]. Mutations in human genes encoding protein components of the sarcomere cause either HCM or DCM. Deficits of force production and transmission are the two main mechanisms that lead to DCM due to sarcomere mutations [55]. Mutations of genes encoding myosin proteins (MYH 6, MYH7, and MYBPC3), actin proteins (ACTC 1 and ACTC 2), and tropomyosin protein (TPM 1) result in alterations of coupling-uncoupling mechanisms of actin to myosin [50]. Specifically, TPM1 mutations are associated with destabilization of actin interactions and compromise force transmission to neighboring sarcomeres. ACTC mutations impair the binding of actin to the Z-disc compromising force propagation [55]. Impaired contractile force may also occur from troponin mutations. Because troponin molecules modulate calcium-stimulated actomyosin ATPase activity, the mutation causes inefficient ATP hydrolysis and decrease contractile strength [55].

2.4.2.2 Cytoskeleton protein location: force transmission and structural integrity

The Z-disc is an electron-dense structure, in which titin and the thin filaments anchor. Critical components include α -actinin, which aligns actin and titin from neighboring sarcomeres and interacts with muscle LIM protein (MLP encoded by CSRP3). Telethonin (encoded by TCAP) is another Z-disc component interacting with titin and MLP to support overall sarcomere function. In addition, Cipher/Z-band alternatively spliced PDZ-motif protein (Cipher/ZASP encoded by LDB3), which interacts with α -actinin-2 through a PDZ domain, an abundant protein interaction modules that recognize short amino acid motifs of the C-termini of target proteins, and couples to protein kinase C (PKC)-mediated signaling via its LIM domains [55, 56]. Gene mutations of multiple Z-disc proteins like MLP, cardiac ankyrin repeat protein (CARP), myopalladin, α -actinin 2, TCAP, and nexilin may result in DCM [52]. Cipher/ZASP mutations have been associated to isolated left ventricular dilatation or DCM with NCCM phenotype [50]. Metavinculin (encoded by VCL) attaches the thin filaments to the plasma membrane and plays a key role in force transmission. Gene mutations in metavinculin cause DCM by disruption of disc structure and actin-filament organization [55]. The costamere, a rib-like structure of the cytoplasmic and transmembrane proteins, interconnects the cytoskeleton to the plasma membrane and the extracellular matrix. Dystrophin and its associated proteins, sacroglycans and dystroglycan, enrich at the costamere and protect against contraction-induced injury [52]. The integrity of the dystrophin complex is critical for mechano-transduction and loss of function mutations trigger instability of the plasma membrane and myofiber loss. This mechanism leads to Duchenne and Becker muscular dystrophy [57]. Desmin, another intermediate filament in cardiomyocytes, forms a 3D scaffold that extends across the entire diameter of the cardiomyocyte, surrounds the Z-discs and interlinks them together and integrates the contractile apparatus with the sarcolemma and the nucleus. Desmin helps to sense mechanical stretch and transduces downstream signals from extracellular to the nucleus. In addition, desmin plays a crucial role during myogenesis. Inhibition of desmin expression blocks myoblast fusion and myotube formation [58]. Mutations in the desmin genes are associated with an autosomal dominant skeletal myopathy, cardiac conduction block, and DCM [59]. Prevalence of desmin mutations in familial DCM have been reported in 1–2% [60].

2.4.2.3 Desmosomes: cell-cell adhesion and intracellular signaling/mechano-signaling

Desmosomes are organized cell membrane structures that provide functional and structural contact between adjacent cells. Mutations in protein components of desmosomes like plakoglobin, desmoplakin, and plakophilin-2 can cause syndromic and non-syndromic ARVC as well as DCM due to disruption of intercellular junction [55, 61].

2.4.2.4 Sarcoplasmic reticulum: calcium homeostasis

Calcium enters the myocyte through voltage-gated L-type Ca^{2+} -channels. This triggers the release of calcium from the sarcoplasmic reticulum (SR) via the ryanodine receptor 2 (RyR2). At low intracellular calcium concentrations, troponin I and actin interactions block actomyosin ATPase activity. With increasing intracellular concentration, calcium binds to troponin C, which releases troponin I inhibition and stimulates contraction. Calcium dissociates from troponin C in cardiac relaxation. Calcium concentration decreases by calcium reuptake in the SR through the phospholamban-regulated cardiac sarcoplasmic reticulum Ca^{2+} -ATPase (SERCA2a) [55]. Mutations of phospholamban precipitate DCM by altering calcium homeostasis [54]. Specific phospholamban mutation R14del is associated with high risk of malignant ventricular arrhythmias and end-stage HF. Further, it is described in a phenotype of ARVC [62].

2.4.2.5 Nuclear envelope: maintain structural organization

The nuclear membrane protein complex contains emerin and lamin A/C (LMNA) [52, 55]. These two lamina proteins and nesprin-1 are part of the LINC complex that links the nucleus to the cytoplasm. Stress signals in the cytoplasm are hypothesized to act with the LINC complex, affecting gene expression in the nucleus. The LINC complex is crucial for an appropriate transcriptional response of the cell to mechanical stress [52]. Defects in emerin proteins can induce X-linked Emery-Dreifuss muscular dystrophy, joint contractures, conduction system disease, and DCM. Dominant lamin A/C (encoded by LMNA) mutations exhibit a more cardiac-restricted phenotype with fibrofatty degeneration of the myocardium and it is conducting system. More than 200 different lamin A/C (LMNA) mutations are associated with inherited cardiomyopathy, primarily DCM that may be associated with conduction system disease prior to the evidence of ventricular dilatation due to fibrofatty degeneration of the myocardium and conducting cells [52, 55]. Other diseases caused by lamin A/C mutations are Charcot-Marie-Tooth neuropathy, Dunningan partial familial lipodystrophy, progeria and other overlapping syndromes, all known as laminopathies [63].

2.4.2.6 Ion channel

The function of sarcolemmal transmembrane cardiac voltage-gated sodium channel is crucial in the generation of cardiac action potentials. Some mutations in the encoding gene *SCN5A* are implicated in DCM. *SCN5A* mutations causes high burden of arrhythmias. There are also many allelic variants in *SCN5A*, including those leading to Brugada syndrome, idiopathic ventricular fibrillation (VF), familial atrial fibrillation (AF), left ventricular non-compaction cardiomyopathy, and long QT syndrome type III [54, 59, 64].

2.4.2.7 Extracellular matrix-cell-adhesion and signaling

Extracellular matrix proteins such as laminin alpha-4 (LAMA4) and Fukutin (FKTN) have been described in relation to DCM. They may lead to DCM phenotype by disrupting signaling pathways and modifying cell-surface molecules [50].

The genetic evaluation of DCM is summarized in **Table 2**.

Genetic evaluation of DCM	Gene screening of DCM	Genotype correlations of DCM
Sarcomere protein related genes	<ul style="list-style-type: none"> • Titin (TNN) • α-Cardiac actin (ACTC 1 and ACTC 2) • α-Tropomyosin 1 (TPM 1) • Cardiac troponin subtypes (TNNT2, TNNC1, TNNI3) • Myosin heavy chains (MYH 6, MYH7) • Myosin-binding protein C (MYBPC) • Troponin I-interacting kinase (TNNI3K) 	<ul style="list-style-type: none"> • 12–25% of genetic related DCM are associated with titin (TTN) mutations • α-Tropomyosin 1 (TPM 1) mutations are described in 1–2% of DCM • Myosin heavy chain (MYH7)-mutations in 3–4% of DCM • 3% of DCM are linked to cardiac troponin T (TNNT2) mutations • Gene defects in sarcomere proteins are associated with defects in force generation and transmission • TNNI3K-mutations may cause conduction defects and atrial fibrillation
Cytoskeletal protein related genes: Z-disc Dystrophin complex Cytoskeleton	<ul style="list-style-type: none"> • α-Actinin 2 (ACTN 2) • Muscle LIM protein (MLP) • Cysteine- and glycine-rich protein 3 (CSRP3) • Telethonin (TCAP) • Cypher/Z-band (LDB3) • PDZ LIM domain protein 3 (PDLIM3) • Cardiac ankyrin repeat protein (CARP) • Myopalladin (MYPN) • Nexilin (NEXN) • Metavinculin (VCL) • Dystrophin (DMD) • Sacroglycan (SGCA, SGCB, SGCD, and SGCG) • Desmin (DES) 	<ul style="list-style-type: none"> • Metavinculin (CVL) mutations are related to 1% of DCM • Dystrophin (DMD) mutations are associated with Duchenne/Becker muscular dystrophy • Sacroglycan (SDC) mutations can cause Limb-girdle-muscular dystrophy • Prevalence of <i>Desmin</i> (DES) mutations in genetic related DCM is about 1–2%, the mutations are often related with myofibrillar myopathy, ARCV and cardiac conduction blocks
Desmosomal protein related genes	<ul style="list-style-type: none"> • Plakoglobin (JUP) • Desmoplakin (DSP) • Desmocollin 2 (DSC2) • Desmoglein 2 (DSG2) • Plakophilin-2 (PKP2) 	<ul style="list-style-type: none"> • Mutations in desmosomal genes are frequent in patients with advanced DCM undergoing cardiac transplantation • Desmosomal gene mutations are also linked to ARVC • Desmoplakin (DSP) causes 2% of genetic related DCM • Plakoglobin (JUP)-mutations are also associated with Naxos syndrome • Desmocollin 2 (DSC2) mutations may lead to mild palmoplantar keratoderma
Sarcoplasmic reticulum related genes	<ul style="list-style-type: none"> • Ryanodine receptor 2 (RyR2) • Phospholamban (PLN) • SR proteins, Ca-ATPase pump (SERCA2a) 	<ul style="list-style-type: none"> • Specific mutations are associated with high risk of malignant ventricular arrhythmias and end-stage heart failure • Ryanodine receptor 2 (RyR2) correlates with catecholaminergic polymorphic ventricular tachycardia and ARVC • Phospholamban (PLN) mutations can cause ARVC • Some other genes encoding for sarcoplasmic reticulum and cytoplasm related proteins like PTPN11, RAF1 and RIT1 are also associated with Noonan and Leopard Syndrome

Genetic evaluation of DCM	Gene screening of DCM	Genotype correlations of DCM
Nuclear envelope and nucleus protein related genes	<ul style="list-style-type: none"> • Emerin (EMD) • Lamin A/C (LMNA) 	<ul style="list-style-type: none"> • EMD mutations can lead to X-linked Emery-Dreyfuss muscular dystrophy, joint contractures and conduction system disease • Prevalence of genetic-related DCM due to LMNA mutation is described in 4–8% • LMNA-related heart failure is often more resistant to heart failure therapy and has a high risk for arrhythmias and sudden cardiac death • Mutations in LMNA can cause a severe and progressive DCM and can also lead to Charcot-Marie-Tooth neuropathy, Dunningan partial familial lipodystrophy, Emery-Dreyfuss muscular dystrophy and progeria
Ion channel protein related genes	<ul style="list-style-type: none"> • Sodium channel, voltage-gated, type V, alpha subunit (SCNA5) • Potassium channel (KCNQ1) 	<ul style="list-style-type: none"> • <i>SCN5A</i> mutations account for 2–3% of DCM, mutations are associated with Brugada syndrome, Long QT syndrome, atrial fibrillation and conduction defects • <i>KCNQ1</i> mutations may induce atrial fibrillation, Long QT 1, Short QT1 and Jervell and Lange-Nielsen syndrome
Extracellular matrix protein related genes	<ul style="list-style-type: none"> • Laminin alpha-4 (LAMA4) • Fukutin (FKTN) 	<ul style="list-style-type: none"> • Extracellular matrix protein relation has been described to DCM • <i>FKTN</i> and <i>LAMA2</i> mutations can also cause congenital muscular dystrophy

Table 2.
Genetic aspects of dilated cardiomyopathy.

2.5 Diagnosis

Establishing the etiology is of great importance as it may influence treatment and prognosis of patients with DCM. Beside the conventional clinical tools, modern imaging and genetic tools are available to elucidate and ensure the correct diagnosis. The recently published statement for the diagnostic workup on DCM from the ESC working group on myocardial and pericardial diseases recommend the following steps: first the diagnostic evaluation should be start with in-depth personal and family history, followed by physical examination, an electrocardiogram (ECG), and echocardiography [8]. These steps often sufficiently differentiate between acquired and familial DCM. If there is no suspicion of an acquired DCM and if ‘red flags’ are recognized, the second-level diagnostic work-up should be added. ‘Red flags’ are defined as signs and suspicion on a specific etiology. Biochemical analyses, cardiac magnetic resonance imaging (CMR), endomyocardial biopsy (EMB), and genetic testing are recommended in a second step. However, the patient’s age plays a crucial role in the decision-making during the diagnostic procedure and should be rated against the potential benefit of dedicated investigations. The detailed diagnostic workup and possible red flags are presented in **Table 3**.

2.6 Screening

In common, DCM is a slowly progressive disease and screening is essential for an early diagnosis of asymptomatic family members. Currently, screening all first-degree family members of patients with genetic proven or non-genetic forms of DCM with a positive family history is recommended. The screening comprises

Diagnostic tool	Look for	Red flags for specific disorders
Personal and familial history	Degree of symptoms Travel history Inheritance pattern Toxin exposition Involvement of other organs	Intellectual and sensorineural disabilities Muscle weakness Myotonia Gait disturbances Skin abnormalities, for example, hyperpigmentation and palmoplantar keratoderma, butterfly-shaped face rash Woolly hair
Electrocardiogram		Low P-wave amplitude Atrioventricular block Repolarization disorders with non-coronary distribution Low QRS amplitude Bundle branch block Long QTc Ventricular arrhythmias
24 h electrocardiogram	Relevant brady- and tachyarrhythmias	Relevant tachyarrhythmias
24 h ambulatory blood pressure monitoring	Exclude persistent hypertension	
Biochemistry	Blood count Electrolytes Renal function Cardiac biomarkers TSH HbA1c Serum iron, ferritin and electrophoresis Urine chemistry and proteinuria Serum free light chains HIV and hepatitis serology Other specific serology tests in accordance symptoms and clinical suspicion	High levels of creatine kinase Myoglobinuria Increased serum iron & ferritin levels Leucopenia or neutropenia Free light chains for amyloidosis Diabetes and lactacidosis Thyroid disorders Infectious etiologies
Echocardiography	Ventricular dilatation Valve diseases Right ventricular pathologies	Left ventricular hypertrabecularisation Segmental dysfunction with noncoronary distribution
Coronary angiography	Exclude coronary artery disease	
CMR	Late Gadolinium Enhancement Intramyocardial edema Intramyocardial iron deposit Right ventricular morphology Molly sequence	Patchy or inferobasal late gadolinium enhancement (LGE) distribution “Midwall sign” septal wall LGE distribution
EBM	Giant cell myocarditis	
Genetics	Screening familial DCM	

Table 3.
Diagnostic workup and possible red flags in dilated cardiomyopathy.

physical examination, 12-lead ECG and transthoracic echocardiography as well as measurement of CK levels. The CK levels may help to identify subclinical skeletal muscle abnormalities and to provide supportive evidence for the presence of an inherited myopathy. If DCM is suspected in first-degree relatives, the screening should be repeated annually. Otherwise, asymptomatic first-degree relatives should be rescreened at three- to five-year intervals because of possible late onset of DCM phenotype [21].

2.7 Therapy

Specific treatment is applicable in syndrome associated DCM, for example, infectious etiologies and infiltrative disorders. However, specific treatment is not available for most DCM patients. Therefore, the therapy focuses on improvement of clinical symptoms as well as on the control of disease progression and potential complications such as sudden cardiac arrest.

2.7.1 Heart failure treatment

Guidelines recommend the following treatment for acute HF, not including non-cardiogenic shock: oxygen, non-invasive ventilation (NIV), intravenous diuretics (20–40 mg bolus furosemide at admission), and intravenous nitrates. Intravenous nitrates have long been described to improve hemodynamic and dyspnea in HF patients by many ways: decrease in systemic blood pressure and left ventricular afterload, substantial reduction preload and therefore of in right and left ventricular filling pressure, an increase in cardiac output, and little or no change in heart rate [65]. Improvement in cardiac output by intravenous nitrates is mostly related to the reduction in left ventricular afterload, but is also influenced by a decrease in pulmonary vascular resistance, improvement in myocardial oxygenation, and a reduction of mitral regurgitation. Administration of inotropes and/or vasopressors is recommended in patients with signs of low cardiac output [66]. However, application of inotropes and/or vasopressors is associated with an increased long-term mortality risk [67–69]. Additional treatment includes the optimal dosing of angiotensin-converting enzyme inhibitors (ACEI) or angiotensin receptor blockers (ARB). ACEI have been shown to reduce cardiovascular mortality and prevent rehospitalization in patients with HF in two key randomized controlled trails (CONSENSUS and SOLVD-Treatment) [70]. Likewise, ARB improve long-term outcome in HF patients [71]. Another essential component of HF therapy is spironolactone. The RALES study has shown a reduction in mortality after addition of 25 mg of spironolactone to the standard treatment in HF patients with an LVEF <35% [72]. The international guidelines recommend spironolactone in all patients presenting with moderate to severe HF symptoms. In the PARADIGM-HF trial, the use of angiotensin receptor-neprilysin inhibitor (ARNI) (sacubitril/valsartan) showed a reduction of the composite endpoint of cardiovascular death or HF hospitalization by 20% compared with enalapril alone in symptomatic HF patients with reduced LVEF [73]. These study results are implemented in the updated ACC/AHA/HFSA guidelines on management of HF, which recommend to replace ACEI or ARB by sacubitril/valsartan in patients with reduced LVEF and ongoing symptoms [74]. Betablockers reduce mortality in HF patients even without reduced ejection fraction as has been demonstrated by multicenter placebo-controlled studies [75–77]. Because of the negative inotropic effect of betablockers, patients should not be treated in the very acute presentation with signs or symptoms of decompensation and initial doses should be low. Long-term goal is a heart rate below 70 bpm in sinus rhythm. If this is not obtainable with the maximum, or maximum tolerated dose of betablockers, the current European heart failure guidelines recommend the addition of Ivabradine [66]. In addition to pharmacological medication cardiac resynchronization therapy (CRT) has been shown to improve cardiac performance, to reduce symptoms, morbidity, and mortality [78, 79].

2.7.2 Anticoagulation

The role of anticoagulation in DCM with sinus rhythm is unclear [80]. The prospective randomized trials were either underpowered or with a too short

follow-up. At present, there are no trial data to guide anticoagulant treatment regime in DCM. Due to two studies (WATCH and WARCEF trial) showing a slight advantage of warfarin over aspirin, anticoagulation with warfarin is advised in patients with a history of thromboembolism or evidence of intracardiac thrombus [81, 82]. Current ACC/AHA HF guidelines do not recommend anticoagulation in reduced left ventricular function and sinus rhythm without prior thromboembolic events or known cardioembolic source [83]. Studies testing the non-vitamin K antagonist oral anticoagulants (NOACs) in patients with reduced left ventricular function are currently ongoing. In DCM patients with documented AF, oral anticoagulant is recommended with CHA₂DS₂-VASc score ≥ 2 , as a class I indication and in men with a CHA₂DS₂-VASc score of 1 as class IIa with level of evidence B [66, 77, 83, 84].

2.7.3 Arrhythmias

The most common arrhythmia in DCM is AF, which increases the risk of thromboembolic complications, impairs cardiac function and worsening HF symptoms. Therefore, evaluation of rate control and anticoagulation in order to preserve LV-function and prevent thromboembolic events is crucial. In the acute setting betablockers, digoxin and their combination may be used to control ventricular rate. In the chronic stage, rhythm control has been shown to be superior to rate control alone in reducing mortality [85]. Because of the commonly reduced left ventricular function, the only therapeutic option is type III antiarrhythmic drug such as amiodarone. Alternatively, electrical cardioversion can be performed. Recently, the CASTLE-AF study demonstrated the superiority of AF catheter ablation in certain patients with HF as compared to medical therapy. The ablation was associated with a significantly lower rate of death and hospitalization for worsening HF [86]. DCM patients may suffer from ventricular arrhythmias (VA), which are mainly caused by myocardial damage, fibrosis and/or loss of cell-to-cell junctions, that are described by three mechanisms: reentry, trigger activity, and automatism [87]. Monomorphic ventricular tachycardias (VTs) are frequently induced by macro-reentry mechanism, which is best treated by ablation. The main trigger mechanisms are electrolyte imbalance, mostly secondary to diuretic treatment, antiarrhythmic drugs, and bradycardia. The therapeutic options are a combination of antiarrhythmic drugs like betablockers and type III antiarrhythmics and/or implantation of an implantable cardioverter defibrillator (ICD). The indications for ICD treatment are described later [4].

2.8 Prognosis and risk stratification

Although there has been a significant improvement in prognosis of DCM patients over the last decades, mortality is still high. The prognosis is mainly influenced by HF symptoms and more relevant by the appearance of VTs. Survival data of adults with DCM have shown a one-year mortality of 25–30% and a 50% survival at 5 years. Sustained VT or VF presents the main cause for SCD, which occurs in up to 12% of DCM patients [4, 88]. In general, the prevalence of sustained VT (monomorphic or polymorphic) is estimated as less than 5% [89]. Recently, the Pediatric Cardiomyopathy Registry presented a 5-year incidence rate of SCD in children with DCM of 3% [90]. An age at diagnosis below 14 years, LV dilation, and posterior wall thinning were identified as the most important risk factors. In contrast, the mortality in adults is mostly associated with age and male gender, reduced New York Heart Association (NYHA) functional class, impaired LVEF, and the presence of specific cardiac biomarkers as well as myocardial fibrosis in CMR [91–94]. Furthermore,

genetic aspects play an important role for detection of high-risk patients. Most commonly the pathogenic mutation in the lamin A/C (LMNA) gene is associated with atrial and ventricular arrhythmias. LMNA mutation has been identified as the most malignant and penetrant condition with worse outcomes compared to other forms of DCM [95]. Beside risk models such as the Seattle Heart Failure Model for the prediction of prognosis in the general population of HF patients, there exist no specific risk tools for DCM patients [96]. For the identification of high-risk DCM patients, a personalized and precise approach is required (**Table 3**). This should include the personal and familial history, measurement of LVEF, detailed search for VA and proof of fibrosis using CMR. Other promising approaches are expected to be helpful in decision-making in high-risk DCM patients, determination of microvolt T-wave alternans analysis and detection of autonomic dysfunction using nuclear imaging. In addition, detection of LMNA gene mutation has been described to identify the high-risk DCM population. However, all these approaches need further research.

2.9 Prevention of sudden cardiac death

The most effective therapy of malignant VTs and thus prevention of SCD in DCM patients is the implantation of an ICD. Current ESC guidelines recommend the implantation of a defibrillator in patients who experienced VT or VF (secondary prevention of sudden cardiac death), as well as in high risk patients for primary prevention. The latter are patients with symptomatic HF NYHA class II–III and LVEF $\leq 35\%$ after ≥ 3 months of optimal medical therapy who are expected to survive for at least 1 year [97]. Similarly, the American College of Cardiology and American Heart Association guidelines recommend ICD therapy in patients with LVEF $\leq 35\%$ due to prior myocardial infarction (MI), at least 40 days post-MI, or non-ischemic DCM and NYHA class II or III [98]. However, existing guidelines lack sensitivity and specificity for the selection of patients with DCM for primary prevention ICD implantation. A recently presented meta-analysis by Golwala et al. has demonstrated a 23% reduction in all-cause mortality with ICD therapy compared with optimal medical therapy alone (HR, 0.77; 95% CI, 0.64–0.91) [99]. Although ICD implantation seems to be the best possible option for SCD prevention in DCM, there remain potential complications. Inappropriate shocks, risk of infection, device or lead replacement have to be considered and discussed in detail before an ICD is implanted.

3. Conclusion

DCM includes a heterogeneous group of myocardial and systemic conditions causing left ventricular dilatation and dysfunction. DCM is one of the most common cardiomyopathies worldwide. Yet, the real prevalence is unknown. The etiology contains non-genetic (e.g. myocarditis, peripartum, toxics, arrhythmia, infiltrative etiologies, endocrine, nutritional, and neuromuscular) and genetic causes. Literature on genetic mutations being responsible for DCM has increased exponentially. Today, up to 30% of the DCM cases are described to be caused by a gene mutation, the majority of which occur in autosomal genes that encode for a wide range of proteins of the cardiomyocyte's structural elements. Mutations in genes encoding sarcomeric, cytoskeletal, desmosomal, nuclear membrane, mitochondrial, and RNA-binding proteins have all been linked to DCM. However, the most common mutations occur in genes encoding sarcomeric proteins and in genes related to the nuclear envelope and the

cytoskeleton. Therefore, the diagnostic workup of DCM should involve the clinical tools as well as imaging and gen-technologies. Specific treatment is only available for syndrome-associated DCM. The majority of the DCM patients are treated for HF symptoms, prevention of thromboembolic events, and malign arrhythmias. The prognosis of DCM patients is variable and depends on multiple risk factors. Some, for example, LVEF and NYHA functional class are known for years as risk factors of SCD, others need further research before they can be established in clinical routine.

Conflict of interest

The authors declare no conflict of interest.

Acronyms and abbreviations

ACEI	angiotensin converting enzyme inhibitors
AF	atrial fibrillation
AIDS	acquired immune deficiency syndrome
AHA	American Heart Association
ARB	angiotensin receptor blockers
ARVC	arrhythmogenic right ventricular cardiomyopathy
CAD	coronary artery disease
CARP	cardiac ankyrin repeat protein
CK	creatine kinase
CMR	cardiac magnetic resonance imaging
DCM	dilated cardiomyopathy
DES	desmin
ECG	electrocardiogram
EMB	endomyocardial biopsy
ESC	European Society of Cardiology
HCM	hypertrophic cardiomyopathy
HF	heart failure
HIV	human immunodeficiency virus
ICD	implantable cardioverter defibrillator
IDC	idiopathic dilated cardiomyopathy
ISFC	International Society and Federation of Cardiology
LAMA4	laminin alpha-4
LINC	links the nucleus to the cytoplasm
LMNA	lamin A/C
LVEF	left ventricular ejection fraction
MI	myocardial infarction
MLP	muscle LIM protein
MYH	myosin proteins
NCCM	non-compaction cardiomyopathy
NIV	non-invasive ventilation
NOAC	non-vitamin K antagonist oral anticoagulant
NYHA	New York Heart Association
PPCM	peripartum cardiomyopathy
RCM	restrictive cardiomyopathy
RNA	ribonucleic acid
RyR2	ryanodine receptor 2

SCD	sudden cardiac death
TPM	troponin-tropomyosin
TNN	cardiac troponin
VA/VT/VF	ventricular arrhythmia/ventricular tachycardia/ventricular fibrillation
WHO	World Health Organization

IntechOpen

IntechOpen

Author details

Deborah P. Schild, Sascha I. Ricciardi, Jens G. Hellige, Rolf Vogel and Nisha Arenja*
Department of Cardiology, Solothurner Spitäler AG, Kantonsspital Olten and
Bürgerspital Solothurn, Switzerland

*Address all correspondence to: nisha.arenja@spital.so.ch

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Towbin JA et al. Incidence, causes, and outcomes of dilated cardiomyopathy in children. *JAMA*. 2006;**296**(15):1867-1876
- [2] Richardson P et al. Report of the 1995 World Health Organization/International Society and Federation of Cardiology Task Force on the definition and classification of cardiomyopathies. *Circulation*. 1996;**93**(5):841-842
- [3] Elliott P, Andersson B, Arbustini E, Bilinska Z, Cecchi F, Charron P, et al. Classification of the cardiomyopathies: A position statement from the European society of cardiology working group on myocardial and pericardial diseases. *European Heart Journal*. 2008;**29**:270-276
- [4] Dec GW, Fuster V. Idiopathic dilated cardiomyopathy. *The New England Journal of Medicine*. 1994;**331**(23):1564-1575
- [5] Givertz MM, Mann DL. Epidemiology and natural history of recovery of left ventricular function in recent onset dilated cardiomyopathies. *Current Heart Failure Reports*. 2013;**10**(4):321-330
- [6] Report of the WHO/ISFC task force on the definition and classification of cardiomyopathies. *British Heart Journal*. 1980;**44**(6):672-673
- [7] Maron BJ et al. Contemporary definitions and classification of the cardiomyopathies: An American Heart Association Scientific Statement from the Council on Clinical Cardiology, Heart Failure and Transplantation Committee; Quality of Care and Outcomes Research and Functional Genomics and Translational Biology Interdisciplinary Working Groups; and Council on Epidemiology and Prevention. *Circulation*. 2006;**113**(14):1807-1816
- [8] Pinto YM et al. Proposal for a revised definition of dilated cardiomyopathy, hypokinetic non-dilated cardiomyopathy, and its implications for clinical practice: A position statement of the ESC working group on myocardial and pericardial diseases. *European Heart Journal*. 2016;**37**(23):1850-1858
- [9] Codd MB et al. Epidemiology of idiopathic dilated and hypertrophic cardiomyopathy. A population-based study in Olmsted County, Minnesota, 1975-1984. *Circulation*. 1989;**80**(3):564-572
- [10] Hershberger RE, Hedges DJ, Morales A. Dilated cardiomyopathy: The complexity of a diverse genetic architecture. *Nature Reviews. Cardiology*. 2013;**10**(9):531-547
- [11] Burkett EL, Hershberger RE. Clinical and genetic issues in familial dilated cardiomyopathy. *Journal of the American College of Cardiology*. 2005;**45**(7):969-981
- [12] Ganesh SK et al. Genetics and genomics for the prevention and treatment of cardiovascular disease: Update: A scientific statement from the American Heart Association. *Circulation*. 2013;**128**(25):2813-2851
- [13] Mestroni L et al. Guidelines for the study of familial dilated cardiomyopathies. Collaborative Research Group of the European Human and Capital Mobility Project on Familial Dilated Cardiomyopathy. *European Heart Journal*. 1999;**20**(2):93-102
- [14] Gavazzi A et al. Evidence-based diagnosis of familial non-X-linked dilated cardiomyopathy. Prevalence, inheritance and characteristics. *European Heart Journal*. 2001;**22**(1):73-81
- [15] Goerss JB et al. Frequency of familial dilated cardiomyopathy.

European Heart Journal. 1995;**16**(Suppl O):2-4

[16] Grunig E et al. Frequency and phenotypes of familial dilated cardiomyopathy. *Journal of the American College of Cardiology*. 1998;**31**(1):186-194

[17] Mahon NG et al. Echocardiographic evaluation in asymptomatic relatives of patients with dilated cardiomyopathy reveals preclinical disease. *Annals of Internal Medicine*. 2005;**143**(2):108-115

[18] Mestroni L et al. Familial dilated cardiomyopathy: Evidence for genetic and phenotypic heterogeneity. *Journal of the American College of Cardiology*. 1999;**34**(1):181-190. Heart Muscle Disease Study Group

[19] Pogwizd SM, McKenzie JP, Cain ME. Mechanisms underlying spontaneous and induced ventricular arrhythmias in patients with idiopathic dilated cardiomyopathy. *Circulation*. 1998;**98**(22):2404-2414

[20] Bakalakos A, Ritsatos K, Anastasakis A. Current perspectives on the diagnosis and management of dilated cardiomyopathy beyond heart failure: A cardiomyopathy clinic doctor's point of view. *Hellenic Journal of Cardiology*. 2018. <https://doi.org/10.1016/j.hjc.2018.05.008>

[21] Hershberger RE et al. Genetic evaluation of cardiomyopathy—A Heart Failure Society of America practice guideline. *Journal of Cardiac Failure*. 2009;**15**(2):83-97

[22] Taylor MR, Carniel E, Mestroni L. Cardiomyopathy, familial dilated. *Orphanet Journal of Rare Diseases*. 2006;**1**:27

[23] Hazebroek M, Dennert R, Heymans S. Idiopathic dilated cardiomyopathy: Possible triggers and treatment strategies. *Netherlands Heart Journal*. 2012;**20**(7-8):332-335

[24] Weintraub RG, Semsarian C, Macdonald P. Dilated cardiomyopathy. *Lancet*. 2017;**390**(10092):400-414

[25] Dennert R, Crijns HJ, Heymans S. Acute viral myocarditis. *European Heart Journal*. 2008;**29**(17):2073-2082

[26] Bozkurt B et al. Current diagnostic and treatment strategies for specific dilated cardiomyopathies: A scientific statement from the American Heart Association. *Circulation*. 2016;**134**(23):e579-e646

[27] Wendt S et al. Brucella related myocarditis. *International Journal of Infectious Diseases*. 2018;**66**:126-127

[28] Villablanca P et al. Salmonella berta myocarditis: Case report and systematic review of non-typhoid Salmonella myocarditis. *World Journal of Cardiology*. 2015;**7**(12):931-937

[29] Samdani S et al. Cardiac complications in diphtheria and predictors of outcomes. *International Journal of Pediatric Otorhinolaryngology*. 2018;**104**:76-78

[30] Diaz F, Collazos J. Myopericarditis due to Chlamydia psittaci. The role of autoimmunity. *Scandinavian Journal of Infectious Diseases*. 1997;**29**(1):93-94

[31] Komuro J et al. Various cardiac abnormalities caused by bacterial myocarditis. *International Heart Journal*. 2018;**59**(1):229-232

[32] Cunha BA et al. Lyme carditis with isolated left bundle branch block and myocarditis successfully treated with oral doxycycline. *IDCases*. 2018;**11**:48-50

[33] Rassi A Jr, Rassi A, Little WC. Chagas' heart disease. *Clinical Cardiology*. 2000;**23**(12):883-889

[34] Elkayam U et al. Pregnancy-associated cardiomyopathy: Clinical

characteristics and a comparison between early and late presentation. *Circulation*. 2005;**111**(16):2050-2055

[35] Regitz-Zagrosek V et al. ESC guidelines for the management of cardiovascular diseases during pregnancy. *European Heart Journal*. 2018;**2018**, **39**(34):3165-3241

[36] Ntusi NB, Mayosi BM. Aetiology and risk factors of peripartum cardiomyopathy: A systematic review. *International Journal of Cardiology*. 2009;**131**(2):168-179

[37] van Spaendonck-Zwarts KY, van Tintelen JP, van Veldhuisen DJ, van der Werf R, Jongbloed JD, Paulus WJ, et al. Peripartum cardiomyopathy as a part of familial dilated cardiomyopathy. *Circulation*. 2010;**121**:2169-2175

[38] Sisakian H. Cardiomyopathies: Evolution of pathogenesis concepts and potential for new therapies. *World Journal of Cardiology*. 2014;**6**(6):478-494

[39] Piano MR. Alcoholic cardiomyopathy: Incidence, clinical characteristics, and pathophysiology. *Chest*. 2002;**121**(5):1638-1650

[40] Higgins AY, O'Halloran TD, Chang JD. Chemotherapy-induced cardiomyopathy. *Heart Failure Reviews*. 2015;**20**(6):721-730

[41] Figueredo VM. Chemical cardiomyopathies: The negative effects of medications and nonprescribed drugs on the heart. *The American Journal of Medicine*. 2011;**124**(6):480-488

[42] Gossage AM, Braxton Hicks JA. On auricular fibrillation. *The Quarterly Journal of Medicine*. 1913;**6**:435-440

[43] Le Lan C et al. Redox active plasma iron in C282Y/C282Y hemochromatosis. *Blood*. 2005;**105**(11):4527-4531

[44] Bhogal S et al. Cardiac amyloidosis: An updated review with emphasis on diagnosis and future directions. *Current Problems in Cardiology*. 2018;**43**(1):10-34

[45] Klein I, Danzi S. Thyroid disease and the heart. *Circulation*. 2007;**116**(15):1725-1735

[46] Rubis P. The diagnostic work up of genetic and inflammatory dilated cardiomyopathy. *e-Journal of the ESC Council for Cardiology Practice*. 2015;**13**(19)

[47] Munguti CM, Al Rifai M, Shaheen W. A rare cause of cardiomyopathy: A case of selenium deficiency causing severe cardiomyopathy that improved on supplementation. *Cureus*. 2017;**9**(8):e1627

[48] Kothari SS, Sharma M. L-carnitine in children with idiopathic dilated cardiomyopathy. *Indian Heart Journal*. 1998;**50**(1):59-61

[49] Watkins H, Ashrafian H, Redwood C. Inherited cardiomyopathies. *The New England Journal of Medicine*. 2011;**364**(17):1643-1656

[50] Mestroni L et al. Genetic causes of dilated cardiomyopathy. *Progress in Pediatric Cardiology*. 2014;**37**(1-2):13-18

[51] McNally EM, Mestroni L. Dilated cardiomyopathy: Genetic determinants and mechanisms. *Circulation Research*. 2017;**121**(7):731-748

[52] McNally EM, Golbus JR, Puckelwartz MJ. Genetic mutations and mechanisms in dilated cardiomyopathy. *The Journal of Clinical Investigation*. 2013;**123**(1):19-26

[53] Garfinkel AC, Seidman JG, Seidman CE. Genetic pathogenesis of hypertrophic and dilated cardiomyopathy. *Heart Failure Clinics*. 2018;**14**(2):139-146

- [54] Burke MA et al. Clinical and mechanistic insights into the genetics of cardiomyopathy. *Journal of the American College of Cardiology*. 2016;**68**(25):2871-2886
- [55] Morita H, Seidman J, Seidman CE. Genetic causes of human heart failure. *The Journal of Clinical Investigation*. 2005;**115**(3):518-526
- [56] Lee HJ, Zheng JJ. PDZ domains and their binding partners: Structure, specificity, and modification. *Cell Communication and Signaling: CCS*. 2010;**8**:8
- [57] Towbin JA et al. X-linked dilated cardiomyopathy. Molecular genetic evidence of linkage to the Duchenne muscular dystrophy (dystrophin) gene at the Xp21 locus. *Circulation*. 1993;**87**(6):1854-1865
- [58] Tsikitis M et al. Intermediate filaments in cardiomyopathy. *Biophysical Reviews*. 2018;**10**(4):1007-1031
- [59] Dellefave L, McNally EM. The genetics of dilated cardiomyopathy. *Current Opinion in Cardiology*. 2010;**25**(3):198-204
- [60] Taylor MR et al. Prevalence of desmin mutations in dilated cardiomyopathy. *Circulation*. 2007;**115**(10):1244-1251
- [61] Cho Y. Arrhythmogenic right ventricular cardiomyopathy. *Journal of Arrhythmia*. 2018;**34**:356-368
- [62] van Rijsingen IA et al. Outcome in phospholamban R14del carriers: Results of a large multicentre cohort study. *Circulation. Cardiovascular Genetics*. 2014;**7**(4):455-465
- [63] Goidescu CM. Dilated cardiomyopathy produced by lamin A/C gene mutations. *Clujul Medical*. 2013;**86**(4):309-312
- [64] Liu Y, Chen H, Shou W. Potential common pathogenic pathways for the left ventricular noncompaction cardiomyopathy (LVNC). *Pediatric Cardiology*. 2018;**39**(6):1099-1106
- [65] Gogia H et al. Prevention of tolerance to hemodynamic effects of nitrates with concomitant use of hydralazine in patients with chronic heart failure. *Journal of the American College of Cardiology*. 1995;**26**(7):1575-1580
- [66] Ponikowski P et al. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure: The task force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC) developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *European Heart Journal*. 2016;**2016**, **37**(27):2129-2200
- [67] Mebazaa A et al. Long-term safety of intravenous cardiovascular agents in acute heart failure: Results from the European Society of Cardiology Heart Failure Long-Term Registry. *European Journal of Heart Failure*. 2018;**20**(2):332-341
- [68] Mebazaa A et al. Short-term survival by treatment among patients hospitalized with acute heart failure: The global ALARM-HF registry using propensity scoring methods. *Intensive Care Medicine*. 2011;**37**(2):290-301
- [69] Mortara A et al. Treatment with inotropes and related prognosis in acute heart failure: Contemporary data from the Italian Network on Heart Failure (IN-HF) Outcome registry. *The Journal of Heart and Lung Transplantation*. 2014;**33**(10):1056-1065
- [70] Group, C.T.S. Effects of enalapril on mortality in severe congestive heart failure. Results of the Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS). *The New England*

Journal of Medicine. 1987;**316**(23):
1429-1435

[71] Cohn JN et al. Rationale and design of the valsartan heart failure trial: A large multinational trial to assess the effects of valsartan, an angiotensin-receptor blocker, on morbidity and mortality in chronic congestive heart failure. *Journal of Cardiac Failure*. 1999;**5**(2):155-160

[72] Pitt B et al. The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *The New England Journal of Medicine*. 1999;**341**(10):709-717

[73] McMurray JJ et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *The New England Journal of Medicine*. 2014;**371**(11):993-1004

[74] Yancy CW et al. 2016 ACC/AHA/HFSA Focused Update on New Pharmacological Therapy for Heart Failure: An Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Journal of the American College of Cardiology*. 2016;**68**(13):1476-1488

[75] Packer M et al. The effect of carvedilol on morbidity and mortality in patients with chronic heart failure. U.S. Carvedilol Heart Failure Study Group. *The New England Journal of Medicine*. 1996;**334**(21):1349-1355

[76] CIBIS-II Investigators and Committees. The cardiac insufficiency bisoprolol study II (CIBIS-II): A randomised trial. *Lancet*. 1999;**353**(9146):9-13

[77] MERIT-HF Study Group. Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL randomised

Intervention Trial in Congestive Heart Failure (MERIT-HF). *Lancet*. 1999;**353**(9169):2001-2007

[78] Sohaib SM et al. Opportunity to increase life span in narrow QRS cardiac resynchronization therapy recipients by deactivating ventricular pacing: Evidence from randomized controlled trials. *JACC: Heart Failure*. 2015;**3**(4):327-336

[79] Tang AS et al. Cardiac-resynchronization therapy for mild-to-moderate heart failure. *The New England Journal of Medicine*. 2010;**363**(25):2385-2395

[80] Mischie AN et al. Anticoagulation in patients with dilated cardiomyopathy, low ejection fraction, and sinus rhythm: Back to the drawing board. *Cardiovascular Therapeutics*. 2013;**31**(5):298-302

[81] Massie BM et al. Randomized trial of warfarin, aspirin, and clopidogrel in patients with chronic heart failure: The Warfarin and Antiplatelet Therapy in Chronic Heart Failure (WATCH) trial. *Circulation*. 2009;**119**(12):1616-1624

[82] Homma S et al. Warfarin and aspirin in patients with heart failure and sinus rhythm. *The New England Journal of Medicine*. 2012;**366**(20):1859-1869

[83] Yancy CW et al. 2013 ACCF/AHA guideline for the management of heart failure: A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology*. 2013;**62**(16):e147-e239

[84] Kirchhof P et al. ESC Guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *European Heart Journal*. 2016;**37**(38):2893-2962

[85] Roy D et al. Rhythm control versus rate control for atrial fibrillation and

heart failure. *The New England Journal of Medicine*. 2008;**358**(25):2667-2677

[86] Marrouche NF et al. Catheter ablation for atrial fibrillation with heart failure. *The New England Journal of Medicine*. 2018;**378**(5):417-427

[87] Merino JL. Mechanisms underlying ventricular arrhythmias in idiopathic dilated cardiomyopathy: Implications for management. *American Journal of Cardiovascular Drugs*. 2001;**1**(2):105-118

[88] Milner PG, Dimarco JP, Lerman BB. Electrophysiological evaluation of sustained ventricular tachyarrhythmias in idiopathic dilated cardiomyopathy. *Pacing and Clinical Electrophysiology*. 1988;**11**(5):562-568

[89] Teerlink JR et al. Ambulatory ventricular arrhythmias in patients with heart failure do not specifically predict an increased risk of sudden death. PROMISE (Prospective Randomized Milrinone Survival Evaluation) Investigators. *Circulation*. 2000;**101**(1):40-46

[90] Pahl E et al. Incidence of and risk factors for sudden cardiac death in children with dilated cardiomyopathy: A report from the Pediatric Cardiomyopathy Registry. *Journal of the American College of Cardiology*. 2012;**59**(6):607-615

[91] Michels VV et al. Progression of familial and non-familial dilated cardiomyopathy: Long term follow up. *Heart*. 2003;**89**(7):757-761

[92] Miller WL et al. Troponin, B-type natriuretic peptides and outcomes in severe heart failure: Differences between ischemic and dilated cardiomyopathies. *Clinical Cardiology*. 2007;**30**(5):245-250

[93] Gulati A et al. Association of fibrosis with mortality and sudden

cardiac death in patients with nonischemic dilated cardiomyopathy. *JAMA*. 2013;**309**(9):896-908

[94] Mahrholdt H et al. Delayed enhancement cardiovascular magnetic resonance assessment of non-ischaemic cardiomyopathies. *European Heart Journal*. 2005;**26**(15):1461-1474

[95] Pasotti M et al. Long-term outcome and risk stratification in dilated cardiomyopathies. *Journal of the American College of Cardiology*. 2008;**52**(15):1250-1260

[96] Levy WC et al. The Seattle heart failure model: Prediction of survival in heart failure. *Circulation*. 2006;**113**(11):1424-1433

[97] Priori SG et al. 2015 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: The Task Force for the Management of Patients with Ventricular Arrhythmias and the Prevention of Sudden Cardiac Death of the European Society of Cardiology (ESC). Endorsed by: Association for European Paediatric and Congenital Cardiology (AEPC). *European Heart Journal*. 2015;**36**(41):2793-2867

[98] Epstein AE et al. 2012 ACCF/AHA/HRS focused update incorporated into the ACCF/AHA/HRS 2008 guidelines for device-based therapy of cardiac rhythm abnormalities: A report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines and the Heart Rhythm Society. *Journal of the American College of Cardiology*. 2013;**61**(3):e6-e75

[99] Golwala H et al. Implantable cardioverter-defibrillator for nonischemic cardiomyopathy: An updated meta-analysis. *Circulation*. 2017;**135**(2):201-203