

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.

For more information visit www.intechopen.com



Thiosulfonates: The Prospective Substances against Fungal Infections

Vira Lubenets, Nataliya Stadnytska, Diana Baranovych, Sofiya Vasylyuk, Olena Karpenko, Viktoriya Havryliak and Volodymyr Novikov

Abstract

The synthesis of new analogs of natural biologically active substances is a promising direction for the development of effective antifungal agents. Thiosulfonic acid esters (thiosulfonates) are the structural analogs of biocidal compounds from garlic, onion, cabbage, cauliflower, etc. More than 1000 thiosulfonates of various structures of the general formula RSO_2SR' were synthesized at the Lviv Polytechnic National University, where their physicochemical properties were characterized. A high antifungal activity of the obtained substances was established in relation to the representatives of fungi of different genera. The thiosulfonates are perspective as basis for the development of effective antifungal means for the modern pharmaceutical, food industry, for the protection of various materials and agricultural products. To increase their effectiveness, antimicrobial compositions based on thiosulfonates and surfactants of microbial origin (biosurfactants) in the form of stable suspensions were developed and studied. It has been established that the use of biosurfactants in the compositions allows the enhancement of the antifungal activity of thiosulfonates and reduction of their active concentration. The possible mechanisms of the joint action of thiosulfonates and biosurfactants on fungal pathogens are proposed.

Keywords: fungal infection, thiosulfonates, biosurfactants, biological activity, pathogens

1. Introduction

Thiosulfonic acids and their esters of the general formula RSO_2SR' are close structural analogs of the natural phytoncides of garlic (*Allium sativum*), onion (*Allium cepa*), various types of cabbage, especially cauliflower [1, 2], and also deep-sea urchin *Echinocardium cordatum* [3]. It is well known that synthetic esters of thiosulfonic acids exhibit a wide range of biological activity that often exceeds the efficiency of natural analogs. Some of these esters are proposed as effective antifungal compounds [4, 5], promising substances for other applications [6–12], preservatives of fruits and vegetables, effective plant protection products, growth regulators, biocidal additives [7, 13–15], insecticides, and radioprotectors [16–19].

Esters of thiosulfonic acids are effective sulfenilating reagents in organic synthesis [19, 20] and also have valuable properties for solving complex problems of molecular biology and biochemistry [16].

Nowadays, synthesis and investigation of thiosulfoesters are carried out by Japanese [21], American [22], Italian [23], Spanish [17, 24], Korean [25], and Chinese [26] scientists from leading research centers. In Ukraine, for many years, the study on the synthesis and physicochemical and biological properties of thiosulfonic acid esters is carried out at the Lviv Polytechnic National University by the staff of the Department of Technology of Biologically Active Compounds, Pharmacy, and Biotechnology [16, 27]. At the present time, a scientific school has been formed that develops a methodology for research on the synthesis of biologically active compounds of the thiosulfonate structure. During this time more than 1000 compounds of the general structure RSO_2SR' were synthesized:



Among esters of thiosulfoacids, there are compounds with fungicidal activities against fungi of the genera *Candida*, *Fusarium*, *Mucor*, *Phragmidium*, *Ramularia*, *Penicillium*, *Aspergillus*, *Cladosporium*, *Paecilomyces*, *Phoma*, *Rhizopus*, *Saccharomyces*, *Botrytis*, *Stachybotrys*, *Alternaria*, *Aureobasidium*, *Chaetomium*, *Myrothecium*, *Epidermophyton*, *Trichophyton*, *Microsporum*, *Sclerotinia*, *Monilia*, *Trichoderma*, *Verticillium*, *Pullularia*, *Cryptococcus*, *Trichosporon*, and *Geotrichum* [4–16].

Investigation of esters of thiosulfonic acids began after the isolation of natural antibiotic allicin from the garlic juice, which manifests the antimicrobial activities. Allicin is a low stable allyl ester of allylthiosulfine acid [17]. Esters of thiosulfonic acid, in comparison with esters of thiosulfine acid, are stable compounds; the effectiveness of its antimicrobial activity, in particular, antifungal, is equal to or higher than the activity of thiosulfonates [17].

The antimicrobial activity of esters of thiosulfoacids is closely related to their ability to block the normal metabolism of microorganisms through sulfenylation of thiol groups of their enzymes [28]. It is known that these esters are highly reactive compounds that interact with nucleophiles, electrophiles, and radicals. Nucleophilic substitution reactions occur with breaking of -S-S-bond due to the redistribution of electron density in thiosulfogroup that determines the direction of nucleophile attack [9, 16, 29].

The information in the review about the possibility of practical application of thiosulfonates as antifungal substances is based on published data on its use in the modern pharmaceutical, food, other industries, and agriculture.

2. Protection of agricultural products

Plant-fungal infections cause great economic losses due to the reduction in crop yields during its growing and storage. The loss of crops at all stages of production ranges from 15 to 40%. To solve these problems, esters of thiosulfonic acid can be

used because they are characterized by low toxicity and are permitted as preservatives for the food industry [30–34]. The products of their decomposition are environmentally safe, so these substances do not harm the environment [29, 35].

S-esters of thiosulfonic acid are proposed for the prevention and control of agricultural products damage, in particular, fruit and vegetable, as these compounds are capable of inhibiting or eliminating rotting processes [36].

2.1 Effect of thiosulfoesters on microorganisms

The minimal inhibitory concentrations (MIC) against such fungi (*Penicillium* sp., *Aspergillus* sp., *Fusarium* sp., *Rhizopus* sp., *Mucor* sp., *Saccharomyces ellipsoideus*, *Candida albicans*) were determined for alkyl, cycloalkyl, trichloromethyl, aryl, and alkyl-functionalized esters of alkane and cycloalkane thiosulfonic acids (**Table 1**).

#	Substances		MIC, µg/ml						
	R	R'	A	B	C	D	E	F	G
1	2	3	4	5	6	7	8	9	10
1	CH ₃	CH ₃	1.0	4.0	2.0	20.0	1.0	1.0	0.4
2		C ₂ H ₅	40.0	20.0	20.0	20.0	20.0	20.0	- ^a
3		C ₄ H ₉	10.0	4.0	4.0	20.0	20.0	20.0	- ^a
4		i-C ₄ H ₉	4.0	4.0	4.0	10.0	0	4.0	- ^a
5		CCl ₃	2.0	20.0	2.0	20.0	20.0	2.0	2.0
6		C ₂ H ₅ Cl	20.0	100.0	2.0	100.0	100.0	2.0	- ^a
7		C ₂ H ₄ OH	4.0	100.0	4.0	100.0	100.0	4.0	20.0
8		CH ₂ COOH	20.0	20.0	2.0	100.0	100.0	2.0	- ^a
9		<i>cycl</i> - C ₅ H ₉	- ^a	- ^a	2.0	- ^a	10.0	- ^a	0.5
10		<i>cycl</i> - C ₆ H ₁₁	- ^a	- ^a	2.0	- ^a	5.0	- ^a	1.0
11	C ₂ H ₅	CH ₃	20.0	100.0	4.0	100.0	20.0	4.0	4.0
12		C ₂ H ₅	10.0	10.0	10.0	10.0	20.0	- ^a	- ^a
13		C ₄ H ₉	10.0	10.0	10.0	100.0	100.0	- ^a	- ^a
14		C ₄ H ₉ - <i>iso</i>	10.0	10.0	10.0	100.0	0	- ^a	- ^a
15		CCl ₃	4.0	2.0	2.0	4.0	2.0	2.0	- ^a
16		C ₂ H ₅ Cl	4.0	20.0	2.0	100.0	100.0	2.0	- ^a
17		C ₂ H ₅ OH	1.0	20.0	1.0	100.0	20.0	2.0	- ^a
18		CH ₂ NH ₂ *HBr	0	0	0	0	0	0	- ^a
19		CH ₂ COOH	0	0	0	0	0	- ^a	- ^a
20		<i>cycl</i> - C ₅ H ₉	- ^a	- ^a	2.0	- ^a	100.0	- ^a	50.0
21	<i>cycl</i> - C ₆ H ₁₁	- ^a	- ^a	0.4	- ^a	20.0	- ^a	0.2	
22	C ₃ H ₇	<i>cycl</i> - C ₅ H ₉	- ^a	- ^a	5.0	- ^a	10.0	- ^a	1.0
23		<i>cycl</i> - C ₆ H ₁₁	- ^a	- ^a	0.4	- ^a	100.0	- ^a	0.2
24	C ₄ H ₉	<i>cycl</i> - C ₅ H ₉	- ^a	- ^a	5.0	- ^a	50.0	- ^a	1.0
25		<i>cycl</i> - C ₆ H ₁₁	- ^a	- ^a	1.0	- ^a	- ^a	- ^a	0.2
26	C ₆ H ₅ CH ₂	C ₂ H ₅	4.0	10.0	4.0	40.0	- ^a	10.0	- ^a
27		CH ₂ CH(CH ₃) ₂	4.0	4.0	4.0	40.0	- ^a	4.0	- ^a
28		CH(CH ₃)C ₂ H ₅	4.0	100.0	2.0	100.0	100.0	4.0	- ^a
29		C ₂ H ₅ Cl	20.0	20.0	20.0	0	0	- ^a	4.0
30		C ₂ H ₅ OH	0	0	100.0	0	0	- ^a	0
31	CH ₂ COOH	0	0	0	0	0	- ^a	0	

Note. # — number of compounds
 A: *Penicillium* sp.; B: *Aspergillus* sp.; C: *Fusarium* sp.; D: *Rhizopus* sp.;
 E: *Mucor* sp.; F: *Saccharomyces ellipsoideus*; G: *Candida albicans*
 0 Not activity -^a Not tested

Table 1.
 Minimal inhibitory concentrations of AlkSO₂SR' [37].

It has been shown that the fungicidal activity of cyclopentyl (## 9, 20, 22, 24, **Table 1**) and cyclohexyl S-esters of alkanthiosulfonic acids (## 10, 21, 23, 25, **Table 1**) is similar or higher than the activity of alkyl S-esters of alkanthiosulfonic acid (## 1–4, 11–14, **Table 1**), especially for *Fusarium* sp., *Mucor* sp., and *Candida albicans*.

The minimal inhibitory concentration for cyclopentyl S-esters (## 9, 20, 22, 24, **Table 1**) varied from 0.5 to 50.0 µg/ml, and for cyclohexyl S-esters, it was in the range of 0.2–5.0 µg/ml. For cyclohexyl S-esters of alkanthiosulfonic acid (## 10, 21, 23, 25, **Table 1**), the inhibitory concentration is from 0.2 to 2.0 µg/ml relative to *Fusarium* sp., and *Candida albicans* [28, 29]. For cyclohexyl S-esters of alkanthiosulfonic acid (## 10, 21, 23, 25, **Table 1**), the minimal inhibitory concentration ranged from 0.2 to 2.0 µg/ml against *Fusarium* sp. and *Candida albicans* [37, 38].

Among the S-esters of cyclopentane-hexanethiosulfonic acid (## 1–19, **Table 2**) and cyclohexanethiosulfonic acid (## 10–19, **Table 2**), trichloromethyl S-esters (## 4, 13, **Table 2**) are most effective against *Fusarium* sp. and *Candida albicans* but less active against *Mucor* sp. [38].

It has been found that cyclopentyl C₆H₅-SO₂-S-C₆H₁₁-cycl (# 6, **Table 3**) and cyclohexyl C₆H₅-SO₂-S-C₅H₉-cycl (# 3, **Table 3**) S-esters of benzenethiosulfonic acid C₆H₅-SO₂-S-C₆H₁₁-cycl (# 6, **Table 3**) exhibit high activity against *Candida albicans* (MIC 2.0 µg/ml and 1 µg/ml, respectively), *Fusarium avenaceum* (MIC 5.0 µg/ml and 2.0 µg/ml, respectively), and a lower activity against *Mucor* sp. [38].

#	Substances		MIC, µg/ml						
	R	R'	A	B	C	D	E	F	G
1	2	3	4	5	6	7	8	9	10
1	cycl-C ₅ H ₉	CH ₃	- ^a	- ^a	1.0	- ^a	4.0	- ^a	4.0
2		C ₂ H ₅	- ^a	- ^a	2.0	- ^a	20.0	- ^a	0.2
3		C ₃ H ₇	- ^a	- ^a	1.0	- ^a	20.0	- ^a	0.4
4		CCl ₃	- ^a	- ^a	0.05	- ^a	5.0	- ^a	0.02
5		CH ₂ CH ₂ Cl	- ^a	- ^a	5.0	- ^a	50.0	- ^a	1.0
6		CH ₂ CH ₂ OH	- ^a	- ^a	20.0	- ^a	200.0	- ^a	20.0
7		CH ₂ COOH	- ^a	- ^a	50.0	- ^a	50.0	- ^a	5.0
8		C ₆ H ₅	- ^a	- ^a	10.0	- ^a	100.0	- ^a	10.0
9		4-NO ₂ C ₆ H ₄	- ^a	- ^a	50.0	- ^a	200.0	- ^a	20.0
10	cycl-C ₆ H ₁₁	CH ₃	- ^a	- ^a	2.0	- ^a	5.0	- ^a	2.0
11		C ₂ H ₅	- ^a	- ^a	5.0	- ^a	10.0	- ^a	2.0
12		C ₃ H ₇	- ^a	- ^a	0.4	- ^a	100.0	- ^a	2.0
13		CCl ₃	- ^a	- ^a	0.1	- ^a	100.0	- ^a	0.05
14		CH ₂ CH ₂ Cl	- ^a	- ^a	1.0	- ^a	20.0	- ^a	0.2
15		CH ₂ CH ₂ OH	- ^a	- ^a	20.0	- ^a	100.0	- ^a	20.0
16		CH ₂ COOH	- ^a	- ^a	4.0	- ^a	100.0	- ^a	50.0
17		C ₆ H ₅	- ^a	- ^a	20.0	- ^a	100.0	- ^a	10.0
18		4-NO ₂ C ₆ H ₄	- ^a	- ^a	100.0	- ^a	200.0	- ^a	20.0
19	4-CH ₃ OC ₆ H ₄	- ^a	- ^a	50.0	- ^a	100.0	- ^a	50.0	
Note. # – number of compounds A: <i>Penicillium</i> sp.; B: <i>Aspergillus</i> sp.; C: <i>Fusarium</i> sp.; D: <i>Rhizopus</i> sp.; E: <i>Mucor</i> sp., F: <i>Saccharomyces ellipsoideus</i> , G: <i>Candida albicans</i> 0 – Not activity; - ^a Not tested									

Table 2.
Minimal inhibitory concentrations of cycl-AlkSO₂SR' [38].

#	Substances		MIC, µg/ml		
	R-	Alk-cycl	<i>Fusarium avenaceum</i>	<i>Mucor sp.</i>	<i>Candida albicans</i>
1	2	3	4	5	6
1	4-CH ₃ O	cycl-C ₅ H ₉	20.0	100.0	10.0
2	4-CH ₃ CONH	cycl-C ₅ H ₉	5.0	50.0	2.0
3	H	cycl-C ₅ H ₉	5.0	100.0	2.0
4	4-CH ₃ O	cycl-C ₆ H ₁₁	5.0	200.0	5.0
5	4-CH ₃ CONH	cycl-C ₆ H ₁₁	20.0	- ^a	4.0
6	H	cycl-C ₆ H ₁₁	2.0	50.0	1.0

Note. # - number of compounds
^a. Not tested

Table 3.
 Minimal inhibitory concentrations of 4-R-C₆H₄-SO₂-S-Alk-cycl [38].

2.2 Anti-phytopathogenic activities of esters of thiosulfoacids

The prospects for the use of esters of thiosulfonic acid for the control of clamp rot of beet have revealed. The causative agents of clamp rot of beet are phytopathogens *Botrytis cinerea*, *Fusarium betae*, and *Phoma betae*, which cause the loss of root resource during its storage.

These pathogens damage the roots of beet seedlings, cause spotty leaves, and dry rot of root crops, which lead to a decrease in the taste and commodity performance of products. In laboratory conditions, 18 esters of thiosulfoacid were investigated against these phytopathogens (**Table 4**) [39].

#	Substances		MFC, µg/ml		
	R	R'	<i>Botrytis cinerea</i>	<i>Fusarium betae</i>	<i>Phoma betae</i>
1	2	3	4	5	6
1	CH ₃	CH ₃	200.0	40.0	40.0
2		C ₂ H ₄ OH	400.0 mycelium growth	-	-
3		C ₂ H ₄ Cl	100.0	200.0	100.0
4	C ₂ H ₅	CH ₃	40.0	100.0	400.0
5		C ₂ H ₅	200.0	100.0	40.0
6		C ₄ H ₉	100.0	200.0	40.0
7		CCl ₃	4.0	4.0	200.0
8		C ₂ H ₄ OH	400.0 mycelium growth		
9	C ₃ H ₇	CH ₃	100.0	40.0	200.0
10		C ₃ H ₇	100.0	100.0	40.0
11		CCl ₃	4.0	10.0	400.0
12		CH ₂ CH ₂ Cl	400.0	—	—
13	C ₄ H ₉	CH ₃	200.0	—	—
14	C ₆ H ₅	CH ₃	100.0	40.0	200.0
15	C ₆ H ₅	i-C ₃ H ₇	400.0	—	—
16	4-CH ₃ CONHC ₆ H ₄	C ₂ H ₅	400.0 mycelium growth	—	—
17	NH ₂ C ₆ H ₄	C ₂ H ₅	400.0 mycelium growth	—	—
18	β-C ₁₀ H ₇	CCl ₃	40.0	—	—

Note. # –number of compounds

Table 4.
 Minimal fungicidal concentrations of RSO₂SR' against pathogens of clamp rot [39].

The highest efficacy against *Botrytis cinerea* and *Fusarium betae* was observed for trichloromethyl S-esters of ethane and propanethiosulfonic acid (# 7, 11, **Table 4**). Trichloromethyl S-ester β -naphthalene thiosulfonic acid is less active among the synthesized trichloromethyl esters. The most effective against *Phoma betae* are methyl S-ester of methanethiosulfonic acid (# 1, **Table 4**), ethyl and butyl S-esters of ethanethiosulfonic acid (# 5, 6, **Table 4**), and propyl ester of propanethiosulfonic acid [10]. It has been shown that thiosulfoesters (# 1, 7, 11, **Table 4**) at a concentration of 200 $\mu\text{g/ml}$ are toxic to sugar beet (*Beta vulgaris var. saccharifera*) but have fungicidal activity against these phytopathogens.

It is interesting that trichloromethyl S-esters of methane- and propanethiosulfonic acids, as well as methyl S-esters of methanesulfonic acid at a concentration of 200.0 $\mu\text{g/ml}$, are nontoxic to sugar beet but exhibit a fungicidal effect on the abovementioned phytopathogenic fungi.

Approval in the farm conditions of the trichloromethyl S-ester of propanethiosulfonic acid (# 11 **Table 4**), as well as its mixtures with methyl-S-ester of methanesulfonic acid, revealed its effectiveness for the treatment of sugar beet root crops for prolonged storage. Improvement of the quality parameters of sugar beet after treatment with the synthesized preparations was established.

Thus, the objects of study were the roots of beet of different qualities. Wilted roots on 15%, damaged ones near the head and tail, roots with 3–4% of green mass and 10% of the earth, and healthy clean roots were used in experiments. After 104 days of storage, all samples of beet treated with solutions of these substances had rotten roots in 2.5–4.0 times less than in untreated control.

The effectiveness of the preparations based on the esters of thiosulfonic acid against the causative agents of clamp rot is confirmed by comparative studies of healthy sugar beet with control, where the amount of rotten mass is greater by 9.3 times [39].

The high antifungal activity of alkyl and trichloromethyl esters of methane-, ethane-, and propanethiosulfonic acids in experiments in vitro was discovered. These data indicate that these compounds can be used for the prevention of fruits and vegetables against fungal damage during prolonged storage. Effective fungicidal concentrations of synthesized nine esters of alkanthiosulfonic acid were found for 13 genera of fungi (**Table 5**):



The highest fungicidal activity of trichloromethyl esters was found at concentrations of 40–1.25 $\mu\text{g/ml}$, while alkyl esters of alkanthiosulfonic acids exhibit fungicidal activity at concentrations of 5–100 $\mu\text{g/ml}$ against fungi of the genera *Fusarium*, *Rhizopus*, and *Mucor*, which in some cases is higher than for trichloromethyl esters [40].

The antifungal activity of some alkyl and trichloromethyl esters of aromatic thiosulfonic acids as preservatives for protecting fruits and vegetables during storage against fungal damage was also studied (**Table 6**) [40].

The positive effect was observed after the use of 4-aminophenyl ester of 4-aminobenzenesulfonic acid (# 2a, **Table 6**) to prevent potato rot during storage. It is significant that this ester is less toxic than alkyl esters of alkanthiosulfonic acid. The potato was treated with a solution of this substance at concentrations of 40 and 20 $\mu\text{g/ml}$ (at a rate of 100 ml or 4–8 mg of dry substance per 1 ton of potatoes). After 1.5 months storage, the amount of potato waste decreased by 2.5 times. The ability of some esters of thiosulfonic acids to protect tomatoes from fungal damage during storage was also studied. The most active and low toxic were the ethyl

#	Fungi	MFC of substances, µg/ml								
		1a	1b	2a	2b	2c	2d	2e	3a	3b
1	2	3	4	5	6	7	8	9	10	11
1	<i>Penicillium glaucum</i>	40.0	10.0	40.0	10.0	10.0	10.0	20.0	10.0	1.67
2	<i>Penicillium expansum</i>	40.0	1.25	40.0	5.0	5.0	1.25	—	10.0	1.67
3	<i>Penicillium digitatum</i>	20.0	5.0	20.0	10.0	5.0	5.0	10.0	20.0	2.5
4	<i>Penicillium italicum</i>	20.0	10.0	40.0	10.0	10.0	40.0	10.0	20.0	1.67
5	<i>Aspergillus niger</i>	20.0	2.5	40.0	10.0	10.0	20.0	—	40.0	2.5
6	<i>Fusarium solani</i>	20.0	80.0	20.0	10.0	10.0	40.0	20.0	40.0	20.0
7	<i>Fusarium moniliforme</i>	20.0	40.0	20.0	10.0	10.0	40.0	10.0	20.0	40.0
8	<i>Rhizopus nigricans</i>	80.0	100.0	80.0	5.0	100.0	180.0	10.0	80.0	20.0
9	<i>Mucor racemosus</i>	20.0	40.0	20.0	10.0	10.0	20.0	20.0	40.0	20.0
10	<i>Botrytis cinerea</i>	40.0	1.67	20.0	20.0	20.0	5.0	40.0	40.0	1.67
11	<i>Botrytis allii</i>	20.0	20.0	50.0	10.0	20.0	20.0	40.0	20.0	2.5
12	<i>Sclerotinia sp.</i>	40.0	10.0	10.0	40.0	20.0	10.0	20.0	20.0	20.0
13	<i>Monilia fructigena</i>	40.0	20.0	80.0	20.0	50.0	20.0	40.0	40.0	10.0

Note. # – number of compounds
 1a: R=CH₃, R'=CH₃; 1b: R=CH₃, R'=CCl₃;
 2a: R=C₂H₅, R'=CH₃; 2b: R=C₂H₅, R'=CCl₃; 2c: R=C₂H₅, R'=C₄H₉; 2d: R=C₂H₅, R'=CCl₃;
 2e: R=C₂H₅, R'=C₆H₅
 3a: R=C₃H₇, R'=CH₃; 3b: R=C₃H₇, R'=CCl₃

Table 5.
 Minimal fungicidal concentrations of RSO₂SR' for the prevention of fruits and vegetables [40].

#	Funges	MFC of substances, µg/ml		
		1a	1b	2a
1	2	3	4	5
1	<i>Penicillium glaucum</i>	10.0	2.5	10.0
2	<i>Penicillium expansum</i>	5.0	1.25	-
3	<i>Penicillium digitatum</i>	5.0	5.0	10.0
4	<i>Penicillium italicum</i>	20.0	10.0	5.0
5	<i>Aspergillus niger</i>	40.0	2.5	-
6	<i>Fusarium solani</i>	20.0	2.5	10.0
7	<i>Fusarium moniliforme</i>	20.0	20.0	10.0
8	<i>Rhizopus nigricans</i>	180.0	80.0	2.5
9	<i>Mucor racemosus</i>	20.0	10.0	5.0
10	<i>Botrytis cinerea</i>	40.0	1.67	5.0
11	<i>Botrytis allii</i>	20.0	20.0	20.0
12	<i>Sclerotinia sp.</i>	40.0	10.0	5.0
13	<i>Monilia fructigena</i>	40.0	20.0	20.0

Note. # — number of compounds
 1a: R=HR'=CH₃; 1b: R=H, R'=CCl₃; 2a: R=NH₂, R'=4-NH₂C₆H₄

Table 6.
 Minimal fungicidal concentrations of 4-RC₆H₄SO₂SR' for the prevention of fruits and vegetables [40].

esters of 4-acetylamino- and 4-aminobenzenethiosulfonic acids. It was established that the treatment of tomatoes by these preparations in laboratory conditions was accompanied by a significant decrease in the amount of waste after 10–12 days of storage.

The effectiveness of alkyl esters of 4-aminobenzenethiosulfonic acids against phytopathogens—the causative agents of citrus fruits' damage—was studied. Citrus fruits were purchased in the Ukrainian trade networks (Lviv). Mycelial fungi *Ramularia chelidonii* and *Phragmidium fragariae* were isolated and identified as pathogens, which cause the damage of lemons and mandarins. It has been shown that isolated cultures exhibit middle and high sensitivity to all studied esters (**Table 7**). Growth inhibition zones of isolated microorganisms at concentrations of active substances of 0.5% and 1% were on average 25–30 mm [41].

In experiments, the ability of S-ethyl ester of 4-aminobenzenethiosulfoacid (ETS) to elicit an antifungal effect against pathogens of fruit and vegetable damage was tested. Minimal inhibitory and minimal fungicidal concentrations against four genera of fungi *Aspergillus*, *Penicillium*, *Paecilomyces*, and *Cladosporium* by methods of diffusion and serial dilutions at a microbial load of 5×10^5 CFU/ml were determined (**Table 8**) [41].

2.3 Effect of Propyl propanethiosulfinate and Propyl propanethiosulfonate on phytopathogens

Propyl propanethiosulfinate (PTS), $C_3H_7SOSC_3H_7$, and propylpropylthiosulfonate (PTSO), $C_3H_7SO_2SC_3H_7$, isolated from garlic (*Allium sativum* L.) and onion (*Allium cepa* L.) were proposed by the Spanish researches for the prevention and control of fungal diseases of plants, for crop storage and as disinfectants for food industry, and for the sanitary treatment of cold rooms, equipment, fruit packaging, and vegetables. PTS and PTSO can be used both in pure form and in the form of aqueous mixtures or suspensions with other synthetic or natural antifungal agents, fertilizers, antioxidants, and plant growth regulators. These compounds can be used in different ways such as immersion, wetting, spraying, applying to soil, etc. [17].

The results, presented in **Table 9**, indicate that PTSO is more active compound against *Penicillium solitum* (wild-type strain) than less stable PTS.

These substances are effective protection agents of tomatoes, peppers, cucumbers, melons, lettuce, stone fruits, citrus fruits, strawberries, tropical fruits like avocado, and mango against *Pseudoperonospora cubensis*, *Phytophthora infestans*, *Erysiphe* sp., *Sphaerotheca* sp., *Leveillula taurica*, *Botrytis cinerea* Pers., *Alternaria dauci*, *Alternaria citri*, *Venturia inaequalis*, *Monilia fructicola*, *Monilia laxa*,

#	Compound	Concentration, %	Zones of growth inhibition, mm	
			<i>Phragmidium fragariae</i>	<i>Ramularia chelidonii</i>
1	2	3	4	5
1	$NH_2C_6H_4SO_2SC_3H_5$	0.1	21	10
		0.5	27	20
		1.0	30	23
2	$NH_2C_6H_4SO_2SCH_3$	0.1	21	0
		0.5	25	17
		1.0	28	24
3	$NH_2C_6H_4SO_2SC_2H_5$	0.1	19	11
		0.5	24	23
		1.0	26	30

Table 7.
Antifungal activity of $NH_2C_6H_4SO_2Salk$ [41].

#	Fungi	Method of diffusion, mm		Method of serial dilution	
		Concentration, 0.1 %	Concentration, 0.3 %	MIC, µg/ml	MFC, µg/ml
1	2	3	4	5	6
1	<i>Aspergillus niger</i>	0	25.0	125.0	250.0
2	<i>Aspergillus terreus</i>	15.0	20.0	62.5	125.0
3	<i>Aspergillus fungus</i>	0	22.0	- ^a	- ^a
4	<i>Penicillium chrysogenum</i>	20.0	25.0	62.5	62.5
5	<i>Penicillium sp.</i>	18.0	20.0	- ^a	- ^a
6	<i>Paecilomyces variotti</i>	20.0	20.0	31.2	62.5
7	<i>Cladosporium resinae</i>	- ^a	- ^a	15.6	31.2
- ^a Not tested					

Table 8.
 Antifungal activity of $\text{NH}_2\text{C}_6\text{H}_4\text{SO}_2\text{SC}_2\text{H}_5$ [41].

Microorganisms	PTS, µg/ml					PTSO, µg/ml				
	1000	500	250	125	60	1000	500	250	125	60
1	2	3	4	5	6	7	8	9	10	11
<i>Penicillium solitum</i> wild strain	35	28	16	11	0	40	30	20	12	0
<i>Enterococcus faecalis</i> ATCC 29212	30	30	26	22	15	45	45	45	33	26
<i>Listeria innocua</i> CECT 4030	40	37	32	23	13	52	43	35	20	13

Table 9.
 Comparative activities of PTS and PTSO against test-microorganisms [17].

Taphrina deformans, *Phytophthora* spp., *Phytophthora infestans*, *Oidium fragariae*, and *Colletotrichum gloeosporioides* [17], which cause downy mildew, *Botrytis* (gray mold), alternariosis, spotted, powdery mildew, *Monilia*, bruised, *Alternaria* of citrus fruits, and gummosis.

Pathogens *Penicillium expansum*, *Botrytis cinerea*, *Physalospora obtusa*, *Glomerella cingulata*, and *Botryosphaeria ribis* can cause blue mold, gray mold, black rot, sour rot, and white rot in postharvested apples, pears, and quince [17].

3. Protection of materials against biodamage

A serious problem is the biodamage to various materials and products, which is accompanied by annual economic losses. Therefore, there is a need to find ways to solve problems related to the protection of raw materials and products from the action of causative agents during its prolonged storage, transportation, and operation.

In view of this, S-esters of thiosulfonic acid RSO_2SR can be promising biocides to protect materials and products since they have a wide range of antifungal and antibacterial effects. The sulfonyl and thiol component of S-esters determines their biocidal action spectrum; therefore, it is important to expand the information on the correlation of the dependence of the structure of S-esters of thiosulfonic acids with its useful characteristics [14, 42].

Ninety-three esters of thiosulfonic acids of various structures in relation to the protection of industrial materials (adhesives, wood, paper, textiles, leather, lubricating liquids, paints, and polymer products) against the fungi, yeasts, bacteria,

algae, and mucilages were studied by researchers of the concern “Bayer.” These compounds are also proposed as biocidal additives in circulating water circuits at industrial plants, in particular, oil refineries [43].

The effectiveness of the synthesized compounds was studied on the following species of fungi: *Alternaria tenuis*, *Aspergillus niger*, *Chaetomium globosum*, *Coniophora puteana*, *Lentinus tigrinus*, *Penicillium glaucum*, *Polyporus versicolor*, *Aureobasidium pullulans*, *Sclerophoma pithyophila*, and *Trichoderma viride*. These thiosulfoesters can be used in various forms: powders, wet powders, suspensions, pastes, soluble powders, dust, and granules [43].

The optimal active concentrations of esters of thiosulfoacids were investigated in the range of 0.001–5.0% by weight. Effective fungicidal concentrations to protection of the materials are ranged within 0.05–1.0% by weight [43].

The minimal inhibitory concentration of some esters against *Penicillium brevicaulis* (200 µg/ml), *Chaetomium globosum* (300 µg/ml), and *Aspergillus niger* (400 µg/ml) were determined.

The antifungal effect of S-esters of thiosulfonic acids for the protection of various materials (lubricating liquids, products of the oil refining industry and equipment at profile enterprises) is determined [44, 45].

These substances can be used for biocidal protection and conservation of works of art, library, and archive funds for long-term storage [46].

3.1 Effect of esters of thiosulfonic acids on paper protection from biodamage.

The effectiveness of the paper protection with S-esters of thiosulfonic acids against test-cultures—*Aspergillus niger*, *Penicillium chrysogenum*, and *Candida tenuis*—compared with reference nipagin (methyl ester of 4-hydroxybenzoic acid) was established [47].

Ethyl S-ester 4'-nitrobenzylidene-4-aminobenzenesulfonic acid in a concentration of 0.01% inhibits the growth of test-cultures of mold fungi by 20%, more than nipagin at a concentration of 0.1% [47].

A comparative study of the resistance of the paper treated with solutions of synthesized compounds (S-ethyl-4'-nitrobenzylidene-4-aminobenzenethiosulfonate, a mixture of ethylthiosulfanilate and polyvinylpyrrolidone in a ratio of 1:2, nipagin) was performed at 100% humidity and 28°C for 36 days against test-cultures (*Penicillium chrysogenum* BKMF-245, *Aspergillus niger* BKMF-1119, *Mucor plumbeus* BKPMF-520) and evaluated by conditional score (Table 10) [48].

It is interesting that the reference preparation nipagin was less effective against fungal action than thiosulfonates (Table 10) [48]. Acetone solutions based on thiosulfonates and their formulations with polyvinylpyrrolidone were developed. Adding polyvinylpyrrolidone improves the physico-mechanical and fungal resistant indices of paper [46, 49].

3.2 Influence of esters of thiosulfonic acids on the protection of the oil and oil refining industry equipment and lubricants from biodegradation

Oil, refined products, and equipment for the oil, oil refining, and petrochemical industries can also be subjected to biological damage. Therefore, the main way to fight the harmful microflora is to use environmentally friendly biocides that violate the enzymatic systems of microorganisms and inhibit its activity [50, 51]. To solve this problem, a nontoxic (LD₅₀ = 2000 mg/kg) ethyl S-ester 4-aminobenzenethiosulfonic acid (ETC) biocide was proposed [44]. Comparing the antifungal activity of the ETS with the reference industrial biocide on the basis of water-soluble 1,3,5-tris-(2-hydroxyethyl)-perhydro-1,3,5-triazine, it was found that the reference biocide

Compound	Concentration, %	<i>Aspergillus niger</i> BKMF – 1119			<i>Penicillium chrysogenum</i> BKMF – 245			<i>Mucor plumbeus</i> BKIMF – 520		
		Duration of experiment, day								
		7	12	36	7	12	36	7	12	36
		Degree of lesion, conditional score								
1	2	3								
<i>n</i> -nitrophenyl-ethylsulfanilate	0.1	0.5	1	1	0	0	0	0	0	0
	0.05	1	2	2	0	0.5	1	0	0.5	0
	0.01	2	3	3	1	1	1	0	1	1
Ethylsulfanilate: polyvinylpyrrolidone 1:2	0.1	1	1	1	0	0.5	1	0	0.5	0
	0.05	1.5	2	2.5	0.5	1	1.5	0	1	0.5
	0.01	2	3	3	1	2	2	0.5	2	1
Nipagin	0.1	1	2	3	1	1	2	0.5	1	0.5
	0.05	1.5	3	4	2	2	3	1	2	1
	0.01	2	3	4	3	4	4	1	4	2
control	—	4	5	5	4	5	5	1	5	2

0 – no growth; 0.5 – 5 colonies; 1 – 10 colonies; 2 – 20 colonies; 3 – 40 colonies; 4 – 80 colonies; 5 – more than 100 colonies.

Table 10.
 Fungal resistance of paper, treated by biocides $RC_6H_4SO_2SC_2H_5$ [48].

even at a concentration of 0.15% by weight is noneffective, whereas the effective fungicidal concentration of the ETS is 0.01% by weight (Table 11). The MIC and MFC of the ethyl S-ester of 4-aminobenzenethiosulfonic acid are, respectively, 40–160 and 2–160 times lower than those of the reference preparation. The advantage of the ETC is its solubility in organic hydrocarbon compounds, which provides its effect in the total volume of the organic phase of petroleum products.

In addition, the introduction of the ETC into the emulsion reduces the damage to the equipment of the oil and oil refining industry as a result of increasing the corrosion resistance of materials [44].

Experimental and industrial studies have shown the stabilization of the lubricating liquids by the addition of ETC, which increases the usage period of ones. In the control cycle during cold rolling of the metal, the number of microorganisms in lubricating liquids after 6 days was 60 million/ml, that is, 30 times more than in the experimental cycle after using the ETS (2 million/ml).

The ETC improves the physicochemical parameters of lubricating liquids and reduces emulsion consumption and the time to replace the spent emulsion. As a

Fungi	Zones of growth inhibition, mm			MIC, $\mu\text{g/ml}$	MFC, $\mu\text{g/ml}$	MIC, $\mu\text{g/ml}$	MFC, $\mu\text{g/ml}$
	A		B				
	0.01%	0.1%	0.15%				
1	2	3	4	5		6	
<i>Aspergillus niger</i>	14	20	0	31.2	62.5	1250	1250
<i>Cladosporium resinae</i>	18	30	0	15.6	31.0	2500	5000
<i>Paecilomyces variotii</i>	17	23	0	62.5	125	5000	5000
<i>Trichoderma viride</i>	15	25	0	62.5	125	2500	5000

A — S-ethyl of 4-amino-benzenethiosulfonic acid;
 B — 1,3,5-tris-(2-hydroxyethyl)-perhydro-1,3,5-triazine

Table 11.
 Antifungal activity of ethyl ester of 4-amino-benzenethiosulfonic acid and 1,3,5-tris-(2-hydroxyethyl)-perhydro-1,3,5-triazine [44].

whole, productive time of the equipment of enterprises significantly increases. The use of ETC as an antifungal agent for lubricating liquids and emulsion in technological processes creates favorable sanitary and hygienic conditions of work and solves a number of environmental issues [44].

4. The effect of ethylthiosulfanilate on the fungal infections

According to the WHO, more than 20% of the world's population of different age groups is affected by mycoses, especially mycoses of the feet and hands with the damage to the nail plate. The resistance of fungal pathogens to known drugs causes increased mycosis diseases and their complications (secondary infections, allergic reactions, eczema, etc.).

It is well known that ETS exhibits a broad spectrum of antifungal activity against pathogenic fungi [4, 5, 14]. The antifungal effects of ETS against 17 strains of various fungi were studied. It has been found that the MIC of the ETC varies from 3.6 to 500 µg/ml (**Table 12**) and depends on the genus and strain [5, 52].

The causative agents that cause skin and systemic diseases are yeast of the genus *Candida*. The antifungal activity of ETS was determined for the most virulent representatives of *Candida albicans*, *Candida tropicalis*, and *Candida stellatoidea*. The MIC of ETS for *C. albicans* is 30 µg/ml, *C. tropicalis* 250 µg/ml, and for *C. stellatoidea* 500 µg/ml. ETS is highly effective against *Aspergillus foetidus* and *Acremonium chrysogenum*, MIC of which is 3.6 µg/ml and 62.5 µg/ml, respectively (**Table 12**) [5, 52, 53].

ETC was proposed as an active substance of the antifungal 1% Esulanum ointment after a detailed study of the antimicrobial effects of a number of esters of thiosulfonic acids in S. Ordzhonikidze All-Union Scientific-Research Institute

Fungi	MIC, µg/ml	Citation
1	2	3
<i>Aspergillus terreus</i>	250	53
<i>Aspergillus foetidus</i>	3.6	53
<i>Aspergillus niger</i>	500	53
<i>Aspergillus awamory</i>	250	53
<i>Aspergillus niger mold</i>	60.0	5
<i>Penicillium canescens</i>	250	53
<i>Acremonium chrysogenum</i>	62.5	53
<i>Trichoderma viride</i>	250	53
<i>Trichoderma terricola</i>	500	53
<i>Candida albicans</i>	30.0	5
<i>Candida tropicalis</i>	250	53
<i>Candida stellatoidea</i>	500	53
<i>Trichophyton gypseum</i>	60.0	5
<i>Microsporum lanosum</i>	15.0	5
<i>Achorion schoenleinii</i>	30.0	5
<i>Actinomyces sp.</i>	15.0	5
<i>Rhizopus nigricans</i>	50.0	54

Table 12.
Antifungal activity of $\text{NH}_2\text{C}_6\text{H}_4\text{SO}_2\text{SC}_2\text{H}_5$ [5, 52, 53].

of Pharmaceutical Chemistry (Moscow). This ointment was developed based on doegling oil and intended for the treatment of tinea pedis and other fungal skin diseases [4, 5, 54].

The advantage of the Esulanum ointment is its keratolytic properties, which promotes rapid penetration of the drug in the deep tissue and provides an effective long-term therapeutic effect [5, 54]. The study of antimicrobial activity of 1% Esulanum emulsion ointment in vitro against a number of microorganisms (*Staphylococcus aureus*, *Streptococcus haemolyticus*, *Escherichia coli*, *Salmonella typhosa*, *Flexner's Bacillus dysenteriae*, *Diphtheria bacillus* (strain PW3), *Bacillus pyocyaneus*, *Proteus vulgaris*, *Anthrax spores*, *Human tubercle bacillus* (H37), *Avian tubercle bacillus*, *Mycobacterium B5*, *Microsporium lanosum*, *Trichophyton gypseum*, *Achorion schoenleinii*, *Actinomycetes*, and fungi *Candida albicans*, *Aspergillus niger*) showed fungicidal properties of the active substance of the ETS. In this case, the MIC and MFC of ETS are practically similar [5].

Clinical studies of the therapeutic effect of 1% ETS ointment in patients with various forms of fungal skin diseases (epidermophytosis, rubrophytosis, microsporia, and trichophytosis) and different clinical manifestations have proven their effectiveness. Based on these results, an instruction for the application of 1% ETC ointment (Esulanum) was developed by the Pharmacological Committee of the Ministry of Health of the former USSR [5].

Experimental part production of 1% Esulanum was introduced into medical practice, but taking into account that the basis of the dosage form was doegling oil, which resources were limited, industrial production was not realized.

The study of the qualitative and quantitative composition of the dosage form based on the ETS and its therapeutic effects is ongoing. The dynamics of the death of *C. tropicalis* cells affected by fungicidal concentrations of the ETS on the model system, similar to human organism processes (37°C), using a microorganism suspension with a cell load of 4×10^5 cells/ml, were detected. It has been found that the minimum fungicidal concentration for cells is 250 µg/ml [52]. Thus, the fungicidal effect of the ETS on *C. tropicalis* is founded after 30 min of exposure (9.09%) and after an hour by 98.48%, and the full therapeutic effect is achieved after 6 h of exposure.

Adding ETS in fungicidal concentrations to the growth medium changes the morphogenesis of *C. tropicalis*. Bumps and fractures appear on the surface of yeast cells, and their manifestation depends on the time of exposure [52]. ETC affects the metabolism of *C. tropicalis*, suppressing endogenous respiration by 87% and the decrease in nucleic acid pool in pathogen cells to 27.52% for DNA and up to 39.13% for RNA. Significant differences were observed in *C. tropicalis* lipogenesis under the influence of this biocide. Thus, subfungicidal concentration of ETS reduces the concentration of almost all classes of phospholipids in cells but increases in the content of lysophosphatidylcholine by 16.25% and phosphatidylcholine by 11.11% [52].

The ETC at fungistatic and subfungicidal concentrations exhibits a membranotropic effect and provides a high degree of co-operability of the membrane structural transitions of cells [55, 56]. The functional state of cell membranes of *C. tropicalis* was assessed by the release of low molecular nucleotide component from cells (pyrimidine and purine bases) under the action of various concentrations of the ETS. Detection of substances at $\lambda = 260$ nm is a sensitive test that characterizes the state of the barrier permeability of the membrane and can be used to study the kinetics of this process.

The obtained results showed that the release of low molecular nucleotide components from *C. tropicalis* begins immediately after the adding of ETS into the medium (*C. tropicalis* cells concentration, 10^6 cells/ml). The increase of output of purine and pyrimidine-containing compounds from the cell in 4.8 times compared

to reference cells was observed, changing the concentration of ETS from 0 to 62.5 µg/ml. The concentration of ETS in the range of 62.5–125 µg/ml causes almost complete loss of these compounds' pool.

The increase in the permeability of *C. tropicalis* membranes under the influence of different concentrations of the ETS or different time of its exposure can be related to high saturation of cell membranes by lipids. The change in permeability of membranes under the action of ETS is probably due to their dynamic structure.

Interaction of ETS and the surface structures of the cell initiate deep structural rearrangements of membranes, which results in increased permeability and, possibly, inhibition of one's physiological functions.

The obtained data suggest that the mechanism of action of the ETS may be related to the disruption of the cytoplasmic membrane, which will lead to significant defects in the delivery of nutrient components to cells and the removal of vital metabolites from them.

So, summing up our results, it can be considered that the mechanism of ETS effects can be related to the disruption of the cytoplasmic membrane that leads to significant defects in the flow of nutrients into the cells and the removal of metabolites from ones [55].

5. Compositions of esters of thiosulfoacids and biosurfactants and their effects on microorganisms

Alkyl esters of alkane- and arene-thiosulfonic acids are hydrophobic compounds, since they are poorly soluble in water, which limits their use as antimicrobial agents. In addition, the surfaces of microbial cells provide the protective barrier to antimicrobials. To increase the solubility and bioavailability of the thiosulfonates, substances that are capable of their solubilization are used, in particular, surface-active substances (surfactants), which can be used as components of ointments, gels, and creams. The most promising ones are biogenic surfactants—products of microbial synthesis (biosurfactants) [57, 58]. Vasileva-Tonkova et al. [59] and Sotirova (2012) described the permeabilizing effect of biosurfactants. They can interact with membrane phospholipids, influence cell hydrophobicity, and have emulsifying ability [53, 59].

The formation of the supramolecular complexes of rhamnolipids with model membrane phospholipids is considered as a possible molecular mechanism of membranotropic action of the biosurfactant, which was shown by electrospray ionization mass spectrometry [60]. These complexes can affect the liquid-crystalline state of the lipid matrix of microbial membranes and change some membrane processes such as cells transport [61].

The permeabilization of cell membranes with surfactants can overcome the barriers and increase the efficacy of various antimicrobial agents. In this regard, biosurfactants can play a significant role as additives in the development of pharmaceuticals due to their ability to enhance solubility or bioavailability of poorly soluble substances [58].

A new promising approach to the creation of effective antimicrobials is proposed, which consists in a synergistic combination of thiosulfonic esters, ethylthiosulfanilate (ETS) or methylthiosulfanilate (MTS), with the biosurfactants of *Pseudomonas* sp. PS-17 strain—rhamnolipid biocomplex (RBC) and rhamnolipids (RL). The introduction of biosurfactants into the compositions allowed the reduction of the active concentrations of thiosulfonates in the resulting products. The antimicrobial potential of compositions based on ETS and MTS with RL was investigated against model strains of various genera and taxonomic groups,

Microorganisms	Zones of growth inhibition, mm					
	0,1 %		0,5 %		1,0 %	
	ETS	ETS + RBC	ETS	ETS + RBC	ETS	ETS + RBC
1	2	3	4	5	6	7
<i>Aspergillus niger</i>	40	50	40	55	48	65
<i>Candida lipolytica</i>	18	30	23	35	33	38
<i>Bacillus mesentericus</i>	17	20	20	23	22	24
<i>Escherichia coli</i>	20	24	25	28	28	31

Note: ETS — ethylthiosulfanilate, RBC — rhamnolipid biocomplex.
 RBC was used in concentration 0,01 %; the mixture of polyethylene glycols (PEG 400 and PEG 500) was used as a base for the ointment compositions

Table 13.
 Influence of the contents of ointment compositions on the growth of microorganisms [54].

capable of causing damages to human health, agriculture, industrial products, as well as phytopathogens. So, for culture *Rhizopus nigricans*, MFCs for MTS, 30 µg/ml, and for ETS, 50 µg/ml, and for their compositions RL were, respectively, 10 and 20 µg/ml [53, 60].

The explanation of these results could be related to permeabilization of the microbial cell membranes by rhamnolipids (increase in levels of extracellular proteins). Biosurfactants provoke changes in cell surface and affect different components of the membranes [62]. Biosurfactant also can damage the surface structure of the spores [53, 63].

The developed ointment composition exhibits high antifungal and antibacterial activity in comparison with known biocidal agents. The addition of rhamnolipid biosurfactants into thiosulfonate compositions contributed to the decrease of minimal fungicidal and bactericidal concentrations [53, 64]. The criteria for selecting the ratios of ETS and RBC in the compositions were the formation of stable emulsions and the antifungal and antibacterial activity of the compositions. Comparative studies of the effectiveness of ointment preparations based on ETC and the composition of ETS + RBC with respect to microorganisms of various taxonomic groups were carried out (Table 13).

Thus, the presence of the biosurfactant enhanced the biocidal effect of MTS and ETS. A possible explanation could be related to the higher protein leakage as a result of permeabilization with the rhamnolipid biosurfactant. Probably, the decrease in concentrations of the studied substances, which are able to suppress the microbial growth completely, is due to the increased access of inhibitors to the bacterial cell. The use of rhamnolipids or other biosurfactants active against variety of microorganisms, in combination with antibiotic treatment, or antimicrobials, may represent new productive antimicrobial strategy [65].

6. Antifungal activity of various esters of carboxylic and heterocyclic thiosulfonic acids

The series of alkyl-, cycloalkyl- and aryl-esters of arylthiosulfonic acids $\text{RNHC}_6\text{H}_4\text{SO}_2\text{SR}'$ was synthesized, and the spectrum of their antifungal action was studied (Table 14) [27].

For $\text{RNHC}_6\text{H}_4\text{SO}_2\text{SR}'$ (Table 14), MIC was determined for *Candida albicans* ATCC 90028, *Candida glabrata* ATCC 90030, *Aspergillus fumigatus* IHEM 13934, and MFC for *Candida albicans*, *Verticillium dahliae*, *Trichophyton gypseum*. All synthesized esters are characterized by rather high fungicidal activity. The compounds

#	Compound		MIC, µg/ml			MFC, µg/ml		
	R	R'	A.	B	C	D	E	F
1	2	3	4	5	6	7	8	9
1	CH ₃ CO	CH ₃	50.0	- ^a	25.0	100.0	40.0	20.0
2		C ₂ H ₅	25.0	- ^a	12.5	40.0	40.0	10.0
3		C ₃ H ₅	100.0	- ^a	12.5	- ^a	- ^a	- ^a
1		3	4	5	6	7	8	9
4	CH ₃ CO	C ₄ H ₉	- ^a	- ^a	- ^a	20.0	10.0	10.0
5		cycl-C ₅ H ₉	- ^a	- ^a	- ^a	2.0	- ^a	- ^a
6		cycl-C ₆ H ₁₁	- ^a	- ^a	- ^a	4.0	- ^a	- ^a
7		C ₆ H ₅	- ^a	- ^a	- ^a	10.0	2.0	4.0
8		C ₆ H ₄ Cl-p	- ^a	- ^a	- ^a	10.0	4.0	20.0
9		C ₆ H ₄ NO ₂ -o	- ^a	- ^a	- ^a	- ^a	400.0	40.0
10	F ₃ CCO	CH ₃	50.0	100.0	25.0	- ^a	- ^a	- ^a
11		C ₂ H ₅	50.0	50.0	25.0	- ^a	- ^a	- ^a
12	ClC ₂ H ₄ CO	CH ₃	- ^a	- ^a	- ^a	40.0	40.0	- ^a
13		C ₃ H ₇	- ^a	- ^a	- ^a	400.0	200.0	200.0
14		i-C ₃ H ₇	- ^a	- ^a	- ^a	200.0	- ^a	- ^a
15		C ₄ H ₉	- ^a	- ^a	- ^a	400.0	- ^a	- ^a
16		C ₆ H ₅	- ^a	- ^a	- ^a	40.0	10.0	10.0
17		C ₆ H ₄ Cl-p	- ^a	- ^a	- ^a	200.0	200.0	100.0
18		C ₆ H ₄ NO ₂ -p	- ^a	- ^a	- ^a	200.0	100.0	200.0
Note. # - number of compounds								
^a Not tested								
A: <i>Candida albicans</i> ATCC 90028; B: <i>Candida glabrata</i> ATCC90030;								
C: <i>Aspergillus fumigatus</i> IHEM 13934; D: <i>Candida albicans</i> ;								
E: <i>Verticillium dahliae</i> ; F: <i>Trichophyton gypseum</i>								

Table 14.

Minimal inhibitory and fungicidal concentrations of substances of structure RNHC₆H₄SO₂SR' [27].

with an acetyl fragment exhibit a higher fungicidal activity than compounds with a 3-chloropropionyl or trifluoroacetyl fragments. The MIC of ethyl ester of 4-acetylaminobenzenethiosulfonic acid relative to *C. albicans* is 25 µg/ml and for fungi *A. fumigatus*, 12.5 µg/ml, while the MIC of the ethyl ester of 4-trifluoroacetylaminobenzenethiosulfonic acid was, respectively, 50 and 25 µg/ml. Regarding the influence of substituents (R') from the side of sulfide sulfur, cycloalkyl- and aryl thiosulfone esters were more active than alkylic. The MFC values of phenyl esters of 4-acetyl and 3-chloropropionyl aminobenzenethiosulfonic acids were determined between 2–10 and 10–40 µg/ml, respectively.

The evaluation of the fungistatic effect of methyl, ethyl, and allyl esters of 3-acetyl-amino-4-methoxybenzenethiosulfonic on test-cultures of *C. albicans* and *Penicillium* sp. with an exposure time of 24–120 h was conducted [64]. At a concentration of 200 µg/ml methyl and allyl esters partially delay growth at exposure for 24 h. With an exposure of 48 h at a concentration of 100 µg/ml, allyl ester completely inhibits the growth of *C. albicans*, which is associated with a change in the antifungal mechanism [66].

People with reduced immunity are often exposed to disease, provoked by fungal infections. It was established that alkyl esters of alkane and arylthiosulfonic acids, alkyl esters of heterocyclic thiosulfonic acids have high antifungal activity. MIC of alkyl esters of 2,3-dioxo-1,2,3,4-tetrahydroquinoxaline-6-thiosulfonic acid was established for various fungal strains of *Candida*, *Cryptococcus*, *Trichosporon*, and *Geotrichum* [67–71]. For fungi of the genus *Candida*, the most effective is the allyl ester (C₃H₅) (Table 15).


Fungi	MIC [$\mu\text{g/ml}$]					
						
	CH ₃		C ₂ H ₅		C ₃ H ₅	
	24h	48h	24h	48h	24h	48h
1	2	3	4	5	6	7
<i>C. albicans</i> ATCC 90028	25.0	25.0	50.0	50.0	10.0	10.0
<i>C. tropicalis</i> ATCC 750	25.0	50.0	12.5	50.0	25.0	50.0
<i>C. parapsilosis</i> ATCC 90018	100.0	100.0	<6.25	25.0	6.25	25.0
<i>C. krusei</i> ATCC 6258	25.0	50.0	25.0	50.0	10.0	10.0
<i>C. inconspicua</i> IHEM 15763	25.0	25.0	12.5	12.5	25.0	25.0
<i>C. norvegensis</i> IHEM 19639	25.0	50.0	25.0	50.0	12.5	25.0
<i>C. kefyr</i> IHEM 15765	25.0	50.0	50.0	100.0	50.0	100.0
<i>C. lusitaniae</i> IHEM 3979	-	12.5	-	100.0	-	25.0
<i>C. glabrata</i> ATCC 90030	100.0	>100.0	25.0	100.0	25.0	100.0

Table 15.
 Fungicidal action of alkyl esters of 2,3-dioxo-1,2,3,4-tetrahydroquinoxaline-6-thiosulfonic acid against *Candida* sp. [67].


Strain	MIC [$\mu\text{g/ml}$]								
									
	CH ₃			C ₂ H ₅			C ₃ H ₅		
	48 h	72 h	96 h	48 h	72 h	96 h	48 h	72 h	96 h
1	2	3	4	5	6	7	8	9	10
<i>Cryptococcus neoformans</i> IHEM 14227	<6.25	12.5	^a	25.0	50.0	^a	6.25	12.5	^a
<i>Cryptococcus neoformans</i> IHEM 3969	^a	<6.25	6.25	^a	25.0	50.0	^a	<6.25	12.5
<i>Trichosporon asahii</i> IHEM 17912	<6.25	<6.25	^a	25.0	25.0	^a	<6.25	12.5	^a
<i>Geotrichum capitatum</i> Lab1969	12.5	25.0	^a	12.5	25.0	^a	<6.25	6.25	^a
<i>Geotrichum capitatum</i> Lab1967	^a	^a	<6.25	^a	^a	12.5	^a	^a	<6.25
^a Not tested									

Table 16.
 Fungicidal action of alkyl esters of 2,3-dioxo-1,2,3,4-tetrahydroquinoxaline-6-thiosulfonic acid against *Cryptococcus*, *Trichosporon*, and *Geotrichum* [67].

This regularity is characteristic to fungi of genera *Cryptococcus*, *Trichosporon*, and *Geotrichum* (Table 16).

The fungistatic effect of methyl, ethyl, and allyl esters of 8-quinolinethiosulfonic acid on the test-culture of *C. albicans* exposed during 24–120 h was evaluated. For all

investigated esters, the inhibitory concentration is 200 µg/ml. Ethyl and allyl esters are more active and their MIC is 100 µg/ml. At a concentration of 1.0 µg/ml, the ethyl ester partially delay the growth of the test-culture exposed for 24–48 h [70, 71].

Summing up our results of the antimicrobial activity of esters of thiosulfonic acids, various ways of their practical application can be proposed. It has been shown that there is a correlation between the structure of thiosulfonate esters, their reactivity in chemical and biochemical reactions, and biological activity. This is confirmed by the specificity of the effects of thiosulfonates against various pathogenic fungi, depending on the structure of their sulfenyl and thiol components.

Our finding showed the advantages of synthetic esters of thiosulfonic acids in comparison with their natural analogs by the biological activity, as well as their competitiveness compared with commercial antifungal substances. The competitiveness of thiosulfonates is determined by their low fungicidal and bactericidal concentrations and low resistance of microorganisms to their action.

Thus, new thiosulfoacid esters as biologically active compounds are suitable for the development of more efficient and safe medicines, biocides, remedies, and growth stimulators.

Author details

Vira Lubenets^{1*}, Nataliya Stadnytska¹, Diana Baranovych¹, Sofiya Vasylyuk¹, Olena Karpenko², Viktoriya Havryliak¹ and Volodymyr Novikov¹

1 Lviv Polytechnic National University, Lviv, Ukraine

2 Department of Physical Chemistry of Fossil Fuels InPOCC, L. M. Lytvynenko InPOCC, NAS of Ukraine, Lviv, Ukraine

*Address all correspondence to: vlubenets@gmail.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Nakamura Y, Matsuo T, Shimoi K, Nakamura Y, Tomita I. S-methyl methanethiosulfonate, a new antimutagenic compound isolated from *Brassica oleracea* L. var. *Botrytis*. *Biological & Pharmaceutical Bulletin*. 1993;**16**:207-209. DOI: 10.1248/bpb.16.207
- [2] Benkeblia N, Lanzotti V. Allium thiosulfinates: Chemistry, biological properties and their potential utilization in food preservation. *Food - Global Science Book*. 2007;**1**(2):193-201
- [3] Takada N, Watanabe M, Suenaga K, Yamada K, Kita M, Uemura D. Isolation and structures of hedathiosulfoniscasids A and B, novel thiosulfonic acids from the deep-sea urchin *Echinocardium cordatum*. *Tetrahedron Letters*. 2001;**42**:6557-6560. DOI: 10.1016/S0040-4039(01)01314-4
- [4] Boldyirev BG, Kolmakova LE, Pershin GM, Milovanova SN, Pozharskaya AM. i dr: Esulan – novoe sredstvo dlya lecheniya epidermofitii stop. *Khymyko-farmatsevtichesky zhurnal*. 1968;**2**(4):12-16. (In Russian)
- [5] Pershin GN, Kukushkina TS, Guskova TA. Esulan. *Khimiko-Farmatsevticheskii Zhurnal*. 1976;**10**(11):146-150. (In Russian)
- [6] Bernard S, Yves B, Henri D (Frantslya). MKI S 07 D 235/28; A 61K 31/47/Nouveaux derives thiosulfonates, leur procede de preparation ainsi que les compositions pharmaceutiques les contenant; № 8417286; Zayavlen. November 13, 84; Opubl. May 16, 86; RZhH. 9O138P. Patent 2573077 Frantslya
- [7] Benkeblia N, Lanzotti V. Allium thiosulfinates: Chemistry, biological properties and their potential utilization in food preservation. *Food - Global Science Books*. 2007;**1**(2):193-201
- [8] Gabriele E, Ricci CM, Eneghetti F, Ferri N, Asai A, Sparatore A. Methanethiosulfonate derivatives as ligands of the STAT3-SH2 domain. *Journal of Enzyme Inhibition and Medicinal Chemistry*. 2017;**32**(1):337-344. DOI: 10.1080/14756366.2016.1252757
- [9] Smith M, Hunter R, Stellenboom N, Kusza DA, Parker MI, Hammouda ANH, et al. The cytotoxicity of garlic-related disulphides and thiosulfonates in WHCO1 oesophageal cancer cells is dependent on S-thiolation and not production of ROS. *Biochimica et Biophysica Acta*. 2016;**1860**:1439-1449. DOI: 10.1016/j.bbagen.2016.03.032
- [10] Foskolos A, Siurana A, Rodriguez-Prado M, Ferret A, Bravo D, Calsamiglia S. The effects of a garlic oil chemical compound, propyl-propane thiosulfonate, on ruminal fermentation and fatty acid outflow in a dual-flow continuous culture system. *Journal of Dairy Science*. 2015;**98**(8):5482-5549. DOI: 10.3168/jds.2014-8674
- [11] Fleischauer AT, Arab L. Garlic and cancer: A critical review of the epidemiologic literature. *The Journal of Nutrition*. 2001;**3S**:1032S-1040S. DOI: 10.1093/jn/131.3.1032S
- [12] Perrino E, Cappelletti G, Tazzari V, Giavini E, Del Soldato P, Sparatore A. New sulfurated derivatives of valproic acid with enhanced histone deacetylase inhibitory activity. *Bioorganic & Medicinal Chemistry Letters*. 2008;**18**:1893-1897. DOI: 10.1016/j.bmcl.2008.02.007
- [13] Shvets V, Karpenko O, Karpenko I, Novikov V, Lubenets V. Antimicrobial action of compositions based on thiosulfonates and biosurfactants on phytopathogens. *Innovative Biosystems and Bioengineering*. 2017;**1**(1):43-48

- [14] Boldarev BG, Bilozor TK, Vlyazlo RI, Kozharskiy AI, Lubenets VI, Luzhetskaya OV, et al. Protivomikrobnaya i fiziologicheskaya aktivnost efirov tiosulfo-kislot i vozmozhnyie puti ih prakticheskogo ispolzovaniya v razlichnyih otraslyah narodnogo hozyaystva. Biopovrezhdeniya v promyshlennosti. Mezhvuzovskiy sbornik. — Gorkiy, izdatelstvo Gorkovskogo gosudarstvennogo universiteta, 1983:44-51. (In Russian)
- [15] Peinado MJ, Ruiz R, Echávarri A, Rubio LA. Garlic: Derivative propyl propane thiosulfonate is effective against broiler enteropathogens in vivo. Poultry Science. 2012;**91**(9):2148-2157. DOI: 10.3382/ps.2012-02280
- [16] Lubenets VI. Tiosulfonaty: syntezi i vlastyvosti. Ukrainskii Khimicheskii Zhurnal. 2003;**69**(3):109-117. (in Ukrainian)
- [17] Garcia-Pareja MP, Sanchez-Vaquero E, Ayala EG, Lopez FM, Alhendin (Granada). Use of antimicrobial agents derived from alliaceous plants for the prevention and control of crop diseases, post-harvest rot and as environmental disinfectant products. Appl. No.: 12/219,136; Filed: July 11, 2008; Pub. Date: January 15, 2009. Patent 2009/0018194A1 US
- [18] Dos Santos EA, Gonçalves FM, Prado PC, Sasaki DY, de Lima DP, Macedo ML. Synthesis method for thiosulfonate and report of its insecticidal activity in *Anagasta kuehniella* (Lepidoptera: Pyralidae). International Journal of Molecular Sciences. 2012;**13**(11):15241-15251. DOI: 10.3390/ijms131115241
- [19] Reeves B, Hilmer J, Mellmann L, Hartzheim M, Poffenberger K, Johnson K, et al. Selective trapping of SNO-BSA and GSNO by benzenesulfinic acid sodium salt: Mechanistic study of thiosulfonate formation and feasibility as a protein S-nitrosothiol detection strategy. Tetrahedron Letters. 2013;**54**:5707-5710. DOI: 10.1016/j.tetlet.2013.08.011
- [20] Trost BM. Alpha-sulphenylated carbonyl compounds in organic synthesis. Chemical Reviews. 1978;**78**:363-382. DOI: 10.1021/cr60314a002
- [21] Taniguchi N. Copper-catalyzed synthesis of thiosulfonates by oxidative coupling of thiols with sodium sulfonates. European Journal of Organic Chemistry. 2014;**26**:5691-5694. DOI: 10.1002/ejoc.201402847
- [22] Rathnayaka H, Mozziconacci O, Sreedhara A, Schoneich C. Fragmentation of a monoclonal antibody by peroxotungstate. Pharmaceutical Research. 2018;**35**(11):1-15
- [23] Gabriele E, Brambilla D, Ricci C, Regazzoni L, Taguchi K, Ferri N, et al. New sulfurated derivatives of cinnamic acids and rosmarinic acid as inhibitors of STAT3 and NF- β transcription factors. Journal of Enzyme Inhibition and Medicinal Chemistry. 2017;**32**(1):1012-1028. DOI: 10.1080/14756366.2017.1350658
- [24] Abad P, Arroyo-Manzanares N, Garcia-Campana AM. A rapid and simple UHPLC-ESI-MS/MS method for the screening of propyl propane thiosulfonate, a new additive for animal feed. Analytical Methods. 2016;**8**:3730-3739. DOI: 10.1039/c6ay00219f
- [25] Hwang SJ, Shyam PK, Jang H-Y. Synthesis of vinyl sulfones via I₂-mediated alkene sulfonylations with thiosulfonates. Bulletin of the Korean Chemical Society. 2018;**39**(4):535-539
- [26] Liang C, Luo S-H, Kai J, Wang B-W, Chu-Ming P, Wang Z-Y, et al.

Disproportionate coupling reaction of sodium sulfinates mediated by BF₃·OEt₂: An approach to symmetrical/unsymmetrical thiosulfonates. *Organic Letters*. 2018;**20**(16):4754-4758

[27] Lubenets V, Vasylyuk S, Monka N, Bolibrukh K, Komarovska-Porokhnyavets O, Baranovych D, et al. Synthesis and antimicrobial properties of 4-acylaminobenzenethiosulfoacid S-esters. *Saudi Pharmaceutical Journal*. 2017;**25**:266-274

[28] Pylypets R, Iskra Y, Havryliak VV, Nakonechna AV, Novikov VP, Lubenets VI. Effects of thiosulfonates on the lipid composition of rat tissues. *Ukrainian Biochemical Journal*. 2017;**89**(6):58-64. <http://ukrbiochemjournal.org/2017/12/effects-of-thiosulfonates-on-the-lipid-composition-of-rat-tissues.html>

[29] Banya AR, Karpenko OY, Lubenets VI, Baranov VI, Novikov VP, Karpenko OV. Influence of surface-active rhamnolipid biocomplex and ethylthiosulfanilate on growth and biochemical parameters of plants in oil contaminated soil. *Biotechnologia Acta*. 2015;**8**(5):71-77. DOI: 10.15407/biotech8.05.071

[30] Boldyirev BG, Grim AI, Nikitina KV. Sposob predokhraneniya plodov ot porchi pri hranenii B.I. (SSSR) A.s. № 186,274 SSSR; 1966. p. 18. (In Russian)

[31] Allyak IV, Boldyirev BG, Melnichenko EL. Sposob predokhraneniya ovoschey ot porchi pri hranenii. (SSSR) A.s. № 186,849 SSSR; 1966. p. 19. (In Russian)

[32] Beltyukova KI, Boldyirev BG. Efiry ti sulfokislota—Novyye sredstva dlya borby s bakterialnyimi boleznyami rasteniy. *DAN SSSR*. 1957;**117**(3):500-503. (In Russian)

[33] Boldyirev BG, Ayzeman BE, Beltyukova KI, Zelepuha SI. Sposob borby s gribkovyimi i bakterialnyimi

boleznyami selskohozyaystvennyih kultur i stimulyatsii razvitiya rasteniy. *Izobreteniya. Prom. obraztsyi. Tovarnyye znaki. (SSSR) A.s. № 103382 MSH SSSR*; 1956. p. 12. (In Russian)

[34] Boldyirev BG, Gar KA, Evteeva NV. Efiry ti sulfokislota novyye fungitsyidnyie sredstva. *DAN SSSR* 1960;**132**(2):346-348. (In Russian)

[35] Lubenets VI, Vasyliuk SV, Nakonechna AV, Shvets VV, Bania AR, Karpenko OV, et al. Zastosuvannya S-etyl-4-aminobenzentiosul-fonatu yak rehuliatora rostu ta biotsydu dlia borotby z fitopatohennoiu mikrofloroiu pry fitorekultyvatsii naftozabrudnenykh gruntiv. zaiavnyk i vlasnyk NU “Lvivska politekhnika”. – № u201707007; zaiavl. 03.07.2017; opubl. 10.04.2018, Biul. №7. Patent UA 124324 S07S381/04 (in Ukrainian)

[36] Lubenets V, Vasylyuk S, Baranovych D, Komarovska O, Rayevska K, Zaichenko O, et al. The plant protection remedies of thiosulfonate type. *Chemicals in Agriculture and Environment*. 2007;**8**:163-167

[37] Boldarev B, Ayzeman B, Zelepuha S, Shvayger M, Mandrik T. O deystvii efirov ti sulfokislota na bakterii i gribyi. *Fiziologicheskii aktivnyie veschestva. Respublikanskiy Mezhhvuzovski Sbornik*. 1973;**5**:24-32. (In Russian)

[38] Ayzeman B, Skorobogatko T, Boldarev B, Aristahova L. O protivomikrobnoy aktivnosti efirov ti sulfokislota - proizvodnyih tsiklopentana i tsiklogeksana. *Fiziologicheskii aktivnyie veschestva. Respublikanskiy Mezhhvuzovski Sbornik*. 1975;**7**:113-116. (In Russian)

[39] Helemskiy MZ, Chepegina FD, Boldarev BG. Primenenie efirov ti sulfokislota dlya borby s kagatnoy gnilyu. *Fiziologicheskii aktivnyie veschestva. Respublikanskiy*

- Mezhvuzovskiy Sbornik. 1972;4:110-113. (In Russian)
- [40] Boldarev B, Hrymm A, Nykytyna K. Efiru tyosulfokyslot—Novi konservanty dlia predokhranenyia plodov y ovoshchei ot porchy pry khranennyi Fyzyolohycheskyi aktyvni veshchestva. Respublykanskyyi Mezhvuzovskyyi Sbornyk. 1971;3:148-154. (In Russian)
- [41] Stadnytska N, Shved O, Chervetsova V, Monka N, Lubenets V, Novikov V. Doslidzhennia in vitro tiosulfonativ dlia zakhystu ahroproduktii vid fitopatohehiv. Zbirka naukovykh prats IX zizdu Ukrainskoho tovarystva henetykiv i selektsioneriv im. M.I. Vavylova “Dosiahnennia i problemy henetyky, selektsii ta biotekhnolohii”. Kyiv. Lohos. 2012;4:609-612. (in Ukrainian)
- [42] Lechado CN, Arjona AB, Ayala EG, Lopez VA, Moll CN, Rus AS. Use of propyl propane thiosulfinate and thiosulfonate for the prevention and reduction of parasites in aquatic animal. Appl. No.: 15/713,840; Filed: August 25, 2017; Date: January 18, 2018. Patent 2018/0014538 A1 US
- [43] Kretschik O, Koln DE, Kugler M, Leichlingen DE, Jaetsch T, Koln DE. Thiosulfonic acid S-esters as agents for protecting material. Appl. No.: 10/015,321; Filed: December 12, 2001; Date: October 17, 2002. Patent 2002/0151570 A1 US
- [44] Shved OV, Lubenets VI, Baranovych DB, Novikov VP, Yarish M Ie. Zastosuvannia S-etyl-4-aminobenzentiosulfonatu yak biotsydu dlia zakhystu naftoproduktiv, metaliv ta obladnannia. № a 2005 04194 zaiav. May 04, 2005; opublik. December 15, 2006. - Biul. № 12. Patent UA 77586. MPK 7 A 01N 29/08, 31/14, 33/06, 33/08, S07S.381/04
- [45] Novikov VP, Lubenets VI, Baranovych DB. Biotsyd dlia zakhystu naftoproduktiv ta obladnannia naftopererobnykh pidpryiemstv vid bioposhkodzhen. № 2003042917; Zaiavl. April 03, 2003; Opubl. January 15, 2004.- Biul. № 1. Patent UA 63325 A. UA. MPK 7 A01N29/08. 31/14. 33.00
- [46] Kadyliak MS, Komarovska-Porokhniavets OZ, Onyshchenko TI, Shved OV, Novikov VP. Sirkovmisna kompozytsiia dlia zakhystu paperu. MPK (2007) D21H 25/00, A01N 31/00, A01N 25/02, C07C 309/00. U 20031110091 opubl. January 15, 2007, Biul.№2. Patent UA 67615
- [47] Komarovska-Porokhniavets O, Hubrii Z, Shved O, Moskalenko N, Iskiv O, Novikov V. Antymikrobnii vlastyvosti N-zamishchenykh tiosulfonativ. Visnyk Natsionalnoho Universytetu “Lvivska politekhnika”. Khimiia, Tekhnolohiia Rechovyn ta Yikh Zastosuvannia. 2008;622:44-46
- [48] Kadyliak M, Komarovska-Porokhniavets O, Shved O, Onyshchenko T, Novikov V. Doslidzhennia biotsydiv dlia zakhystu paperu vid bioposhkodzhen. Visnyk Natsionalnoho Universytetu “Lvivska politekhnika”. Khimiia, Tekhnolohiia Rechovyn ta Yikh Zastosuvannia. 2004;497:71-73
- [49] Kadyliak MS, Komarovska-Porokhniavets OZ, Onyshchenko TI, Skorokhoda VI, Suberliak OV, Shved OV, et al. Biotsydna kompozytsiia dlia zakhystu paperu. MPK (2006) D21H 25/00. U 20040503904 opubl. July 17, 2006, Biul.№7. Patent UA 76229
- [50] Koval E, Sydorenko L. Mykodestruktory nefteproduktov. Mikodestruktoryi promyishlennykh materialov: nauchnoe izdanie inctituta mikrobiologii i virusologii im. D.K. Zabolotnogo AN USSR. - Kiev: Naukova dumka. 1989:142-171: il. - ISBN 5-12-000779-1. (In Russian)
- [51] Lozynskyyi M, Kachan V, Hevaza Y, Maskaiev AK. Stan i osnovni

napriamy robit po zakhystu mastylno-okholodzhuiuchykh ridyn vid bioushkodzhzen. Visnyk - AN URSR. 1983;8:55-58

[52] Starovoitova S, Oriabinska L, Lubenets V. Spectrum antimicrobial action of the original antifungal drug Esulanum. Naukovi visti Natsionalnoho tekhnichnoho universytetu Ukrainy "Kyivskyi politekhnichnyi instytut. 2015;3:68-76. (In Ukrainian)

[53] Sotirova A, Avramova T, Stoitsova S, Lazarkevich I, Lubenets V, Karpenko E, et al. The importance of rhamnolipid-biosurfactant-induced changes in bacterial membrane lipids of *Bacillus subtilis* for the antimicrobial activity of thiosulfonates. Current Microbiology. 2012;65(19):534-541

[54] Lubenets V, Karpenko O, Ponomarenko M, Zahoriy G, Krychkovska A, Novikov V. Development of new antimicrobial compositions of thiosulfonate structure. Chemistry & Chemical Technology. 2013;7:119-124

[55] Oriabinska, Starovoitova S, Vasylyuk S, Novikov V, Lubenets V. Ethylthiosulfanilate effect on *Candida tropicalis*. Ukrainian Biochemical Journal. 2017;89(5):70-76

[56] Starovoitova S, Oriabinska L, Lubenets V. Influence of antimycotic esulanum on lipid composition and functions of cytoplasmic membrane of *Candida tropicalis* yeast. Biotechnology and Microbiology. Scientific Works of NUFT. 2015;21(3):31-37

[57] Rodrigues L, Teixeira J. Chapter 6: Biomedical and therapeutic applications of biosurfactants. In: Sen R, editor. Biosurfactants. Advances in Experimental Medicine and Biology. Vol. 672. New York, NY: Springer; 2010. pp. 75-78

[58] Bhadoriya S, Madoriya N, Shukla K, Parihar MS. Biosurfactants: A new

pharmaceutical additive for solubility enhancement and pharmaceutical development. Biochemical Pharmacology. 2013;2:113. DOI: 10.4172/2167-0501.100011

[59] Vasileva-Tonkova E, Galabova D, Karpenko E, Shulga A. Biosurfactant-rhamnolipid effects on yeast cells. Letters in Applied Microbiology. 2001;33:280-284

[60] Pashynska V. Mass spectrometric study of rhamnolipid biosurfactants and their interactions with cell membrane phospholipids. Biopolymers and Cell. 2009;25(6):504-508

[61] Pashinskaya V, Kosevich M, Gomory A, Lisetski L. Mechanistic investigation of the interaction between bisquaternary antimicrobial agents and phospholipids by liquid secondary ion mass spectrometry and differential scanning calorimetry. Rapid Communications in Mass Spectrometry. 2002;16(18):1706-1713

[62] Sotirova A, Spasova D, Vasileva-Tonkova E, Galabova D. Effects of rhamnolipid-biosurfactant on cell surface of *Pseudomonas aeruginosa*. Microbiological Research. 2009;164:297-303

[63] Banat I, Franzetti A, Gandolfi I, Bestetti G, Martinotti M, Fracchia L, et al. Microbial biosurfactants production, applications and future potential. Applied Microbiology and Biotechnology. 2010;87:427-444. DOI: 10.1007/s00253-010-2589-0

[64] Futami J, Yamada H, Kyuragi G, Yagi K. Thiosulfonate compound, reversible cationization agent for protein and/or peptide, and method for solubilization. Appl. No.: 13/636,789; Filed: March 24, 2011; Date: February 18, 2014. Patent No US 8,653,240 B2

[65] Sotirova A, Avramova T, Lazarkevich I, Lubenets V, Karpenko E, Galabova D. Antimicrobial potential

of selected thiosulfonates-based biocides and biosurfactants against bacteria and fungi. *Comptes Rendus de L'Academie Bulgare des Sciences*. 2010;**63**(9):1307-1314

[66] Lubenets V, Baranovich D, Lisitsa F, Mel'nik O, Vovk I, Gudz O, et al. Synthesis and antimicrobial activity of 3,4-disubstituted benzenethiosulfonic acid esters. *Pharmaceutical Chemistry Journal*. 2000;**34**(3):118-121

[67] Nawrot U, Zaczyńska E, Czarny A, Lubenets V, Karpenko E. Aktywność przeciw grzybic zasyntheticznych pochodnych allicyny—Kontynuacja badań. *Mikologia Lekarska*. 2012;**4**(19):143-146

[68] Lubenets VI, Vasylyuk SV, Novikov VP. Synthesis of S-(3-chloroquinoxalin-2-yl) esters of aliphatic and aromatic thiosulfonic acids. *Chemistry of Heterocyclic Compounds*. 2005;**41**(12):1547-1548

[69] Lubenets V, Vasyliuk S, Hoi O, But S, Cherneha O, Novikov V. Syntez ta vlastyvoli S-esteriv 2,3-diokso-1,2,3,4-tetrahidrokhinoksalin-6-tiosulfokysloty. *Zhurnal Orhanichnoi ta Farmatsevychnoi Khimii*. 2007;**53**(19):56-63

[70] Stadnytska N, Lubenets V, Novikov V, Komarovska O, Vovk N, Kryvoruchko O. Syntez ta biolohichna aktyvnist S-alkil(-8-khinolin)tiosulfonativ. *Zhurnal Fiziolohichno Aktyvnykh Rechovyn*. 2000;**2**(30):27-30

[71] Baranovych D, Stadnytska N, Komarovska O, Hoi O, Lubenets V, Novikov V. Vyvchennia zalezhnosti biolohichnoi aktyvnosti tiosulfonativ vid kyslotnoi i tiolnoi skladovoi. *Visnyk DU "Lvivska politekhnika". Khimii, Tekhnolohii Rechovyn ta Yikh Zastosuvannia*. 2000;**395**:96-99