

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



---

# Introductory Chapter: The Overview of Phosphorous Recovery

---

Huanhuan Li and Tao Zhang

Additional information is available at the end of the chapter


<http://dx.doi.org/10.5772/intechopen.85967>

---

## 1. Introduction

Phosphorus (P) is a finite, non-substitutable, non-renewable, and geographically restricted resource. Substantial interest in P availability was sparked [1]. Some researchers demonstrated sufficient availability to sustain production beyond the twenty-first century, or a maximum occurring late twenty-first century [2–4]. But there are no financial incentives to support mineral resource inspections worldwide without the exploitation of phosphate deposits. The anthropogenic influences on this critical resource are likely to bring about a number of challenges to P sustainability. Advances in technology, public health, and food production over the last couple of centuries have fundamentally interrupted the natural global P cycle. Phosphate deposits have been mined to supply human production, which generated a mostly one-way flow of P from mines to farms to surface waters, ultimately impairing freshwater and coastal waters environment and function [5]. Rapid increases in human population and the subsequent need for high agricultural productivity have led to substantial increases in fertilizer use [6]. The P used as fertilizer consumes more than 80% of the P resources [7]. P is simultaneously an important non-renewable agricultural nutrient and an environmental pollutant [8].

The use of P resources in different countries or regions is different. But the basic P flow is as follows: phosphates are extracted from phosphate rock, passing through crops, animals and human, and ending up in landfill or emission into rivers and the sea (if sewage or animal wastes are not adequately treated) (**Figure 1**). The P cycle is extremely inefficient and wasteful. P loss from wastewater into natural water bodies accounts for about 10% of inorganic P source fertilizers in arable soils globally [9]. Eutrophication is the consequence of human behavior changing the natural P cycle. Researchers and practitioners in multiple fields have responded with numerous strategies to reduce P loading to aquatic ecosystems in an era of eutrophication control [10]. P has become the principal contributor and limiting factor to water eutrophication.



**Figure 1.** Diagram of key global P flows (the widths of the arrows semiquantitatively represents figures reported by Cordell [1] in million tons (Mt) of P per year. Point “A” denotes the flow of P contained in treated or untreated sewage to natural water bodies and represents approximately 1.5 Mt P/year. Point “B” denotes the flow of P contained in erosion losses and is about 8 Mt P/year. \*Other uses include industrial uses such as the production of some detergents) [9].

P, as a resource in fertilizer production or as a pollutant in wastewater treatment, is often simultaneous. The removed P has the potential to be reused as a substitute for mined P fertilizer [11]. With different fractions of P in different water or waste, direct use as fertilizer could not be the sustainable and efficient way and an increasing number of physical, chemical, or biological methods with various functions were designed to recover and reuse available P in the past few years: struvite crystallization, P adsorption, anaerobic digestion, membrane concentration technology, and integration technologies. Most of the P recovery processes are developed to be applied on industrial and municipal wastewater and only a few techniques are developed for P recovery from manure and digestate [12]. More sustainable techniques, such as P recovery techniques for both solid and liquid wastes, are important to maintain the P cycle in modern human society. Sustainable P use has been largely driven by pollution concerns and the shortage of P resources over the past few decades, which P will need to be recovered for productive reuse as a fertilizer to replace increasingly scarce P resource. But for P recycling and reusing, technological advances alone are not enough. There is no single solution to achieving a P-secure future. Partnerships and strategic frameworks to stimulate and support the development of renewable P is an urgent need to establish.

## Acknowledgements

The work was supported by a grant from the National Key Technology Research and Development Program of China (Nos. 2017YFD0800202 and 2016YFD0501404-6).

## Author details

Huanhuan Li and Tao Zhang\*

\*Address all correspondence to: taozhang@cau.edu.cn

Biomass Engineering Center, Beijing Key Laboratory of Farmland Soil Pollution Prevention and Remediation, Key Laboratory of Plant-Soil Interactions of Ministry of Education, College of Resources and Environmental Sciences, China Agricultural University, Beijing, China

## References

- [1] Koppelaar RHEM, Weikard HP. Assessing phosphate rock depletion and phosphorus recycling options. *Global Environmental Change*. 2013;**23**(6):1454-1466
- [2] Cooper J, Lombardi R, Boardman D, Carliell-Marquett C. The future distribution and production of global phosphate rock reserves. *Resources, Conservation and Recycling*. 2011;**57**:78-86
- [3] van Vuuren DP, Bouwman AF, Beusen AHW. Phosphorus demand for the 1970-2100 period: A scenario analysis of resource depletion. *Global Environmental Change*. 2010;**20**:428-439
- [4] Sverdrup HU, Ragnarsdottir KV. Challenging the planetary boundaries. II: Assessing the sustainable global population and phosphate supply, using a system dynamics assessment model. *Applied Geochemistry*. 2011;**26**(Supplement):307-310
- [5] Chowdhury RB, Moore GA, Weatherley AJ, et al. Key sustainability challenges for the global phosphorus resource, their implications for global food security, and options for mitigation. *Journal of Cleaner Production*. 2017;**140**:945-963
- [6] Kaushik V, Mcnamara PJ, Mayer BK. Meta-analysis of non-reactive phosphorus in water, wastewater, and sludge, and strategies to convert it for enhanced phosphorus removal and recovery. *Science of the Total Environment*. 2018;**644**:661-674
- [7] Jeong YS, Matsubae-Yokoyama K, Kubo H, et al. Substance flow analysis of phosphorus and manganese correlated with South Korean steel industry. *Resources, Conservation and Recycling*. 2009;**53**(9):479-489
- [8] Cordell D, White S. Life's bottleneck: Sustaining the World's phosphorus for a food secure future. *Annual Review of Environment and Resources*. 2014;**39**(1):161-188
- [9] Melia PM, Cundy AB, Sohi SP, et al. Trends in the recovery of phosphorus in bioavailable forms from wastewater. *Chemosphere*. 2017;**186**:381-395
- [10] Roy ED. Phosphorus recovery and recycling with ecological engineering: A review. *Ecological Engineering*. 2017;**98**:213-227

- [11] Cornel P, Schaum C. Phosphorus recovery from wastewater: Needs, technologies and costs. *Water Science and Technology*. 2009;**59**(6):1069
- [12] Desmidt E, Ghyselbrecht K, Zhang Y, et al. Global phosphorus scarcity and full-scale P-recovery techniques: A review. *Critical Reviews in Environmental Science and Technology*. 2015;**45**(4):336-384

IntechOpen

IntechOpen