

# We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

**4,800**

Open access books available

**122,000**

International authors and editors

**135M**

Downloads

Our authors are among the

**154**

Countries delivered to

**TOP 1%**

most cited scientists

**12.2%**

Contributors from top 500 universities



**WEB OF SCIENCE™**

Selection of our books indexed in the Book Citation Index  
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?  
Contact [book.department@intechopen.com](mailto:book.department@intechopen.com)

Numbers displayed above are based on latest data collected.

For more information visit [www.intechopen.com](http://www.intechopen.com)



# Applications of Genomics in Regulatory Food Safety Testing in Canada

Catherine D. Carrillo, Adam Koziol, Neil Vary  
and Burton W. Blais

## Abstract

Recent developments in the field of pathogen genomics herald a new paradigm for analytical food microbiology in which pathogenic bacteria will be characterized on the basis of their genetic profile rather than traditional approaches relying on phenotypic properties. The ability to identify gene markers associated with virulence, antimicrobial resistance, and other properties relevant to the identification, risk profiling, and typing of foodborne bacterial isolates will play a critical role in informing regulatory decisions and tracing sources of food contamination. Here we present several scenarios illustrating current and prospective roles for pathogen genomics in food inspection.

**Keywords:** pathogen genomics, virulence, foodborne pathogens, whole-genome sequencing, typing, bioinformatics, food inspection, *Escherichia coli*, *Salmonella*, *Listeria*

## 1. Introduction

The food microbiology testing laboratory has a key role in supporting regulatory food safety investigations, whether stemming from a contamination incident identified through routine monitoring food inspection programs or a foodborne illness outbreak event where human lives and well-being are at risk. While such investigations typically involve the concerted actions of food inspection and public health authorities at different levels of government, the main role of the regulatory testing laboratory is to confirm the presence of a specified hazard in a food vehicle and provide data indicating the scope and source of a contamination event. The extent to which the laboratory can contribute critical information to an investigation will to a large degree depend on the application of leading-edge technologies for detection and characterization of foodborne pathogens. Approaches capable of maximizing the amount of information obtained in the course of conducting laboratory testing of inspection samples will foster the most appropriate regulatory responses, for example, by informing the health risk assessment process undertaken to categorize the degree of risk attending a contamination incident.

The impact of analytical service delivery on public health outcomes depends on the ability to process large numbers of investigative samples and produce accurate test results in the shortest timeframe possible. While cultural enrichment of food samples to amplify pathogens to detectable levels is generally necessary for their

recovery, current approaches often use protracted identification processes relying on phenotypic characteristics elucidated by time-consuming cultivation, biochemical and serological techniques. While effective under certain circumstances, there are shortcomings to such a limited approach, when dealing with novel pathogens for which analytical parameters may not have been comprehensively worked out or in trying to attribute contamination sources.

The exploitation of the genetic blueprint, or known parts thereof, associated with a targeted bacterial pathogen is now widely accepted as an effective means for the identification of food pathogens. Polymerase chain reaction (PCR) technology is now well established as an analytical tool in the regulatory food laboratory [1–4]. Its implementation in regulatory testing programs underscores the growing acceptance of redefining the terms for the identification and characterization of bacteria from a phenotypic to a genotypic basis. Indeed, the advent of leading-edge genomics technologies opens new possibilities for comprehensive analyses of microbial isolates recovered from food inspection samples; for example, next-generation sequencing technologies (whole-genome sequencing, WGS) can now render a bacterial genome much faster (i.e., within 1–3 days) and at a significantly lower cost (about 100 dollars) than previously possible, making it feasible to sequence foodborne isolates within the timeframes of food safety investigations [5–7].

Currently available bioinformatics tools are sufficiently advanced to enable the rapid processing of raw sequence data into a usable form for many purposes. Sequencing pathogenic bacteria, whether in the context of outbreak investigations or information gathering in the course of research, can yield an unprecedented level of information regarding the presence of virulence and other marker genes relevant to the identification and risk characterization of food isolates [6–10]. WGS data can provide an exquisite degree of resolution capable of ascertaining differences between strains and determining phylogenetic relationships among different bacterial isolates for pinpoint precision in the attribution of contamination sources [6, 11]. Finally, the identification of strain-specific features such as unique DNA sequences, metabolic properties, and antimicrobial resistance will enable testing labs to implement customized tests addressing specific strains of interest in determining the scope of contamination events.

While genomics, including WGS technology, already plays a significant role in the clinical sciences, its role in regulatory food microbiology inspection programs remains to be fully delineated. Currently, methods used to characterize foodborne pathogens recovered in regulatory food testing programs aim to answer three questions: (1) what is it? (2) have we seen it before? and (3) is it dangerous? This work describes some of the ways in which characterization of bacterial pathogens using genomics technologies has provided or may contribute to faster, more reliable and cost-effective results addressing these questions. Our purpose is to present different scenarios to illustrate impacts of leading-edge genomics technologies, some imagined, and others already achieved, on food inspection programs.

## **2. Impacts of the implementation of genomics in regulatory food microbiology**

### **2.1 What is it? Definitive identification of pathogenic bacteria based on genomics techniques**

#### *2.1.1 In the beginning: detection of genomic markers by PCR*

*Escherichia coli* O157:H7 have been implicated in outbreaks of foodborne illness associated with the consumption of contaminated foods such as ground beef and

produce [12, 13]. In the event of an outbreak, it is imperative that production lots associated with the primary food vehicle are identified as quickly as possible in order to recall all contaminated products from the marketplace [14]. Traditional techniques for the detection of *E. coli* O157:H7 in foods rely on a multistep process involving pre-enrichment in a selective broth followed by plating to reveal the presence of sorbitol-negative colonies, which are then purified and subjected to a battery of biochemical and serological tests to confirm their identity [15, 16]. This process can take up to 1 week to complete before the contaminant can be definitively identified because of the requirement for growth and phenotypic expression of the organism.

As an alternative to classic phenotypic techniques, the identification of foodborne colony isolates can be achieved on the basis of detection of defining gene markers. Detection platforms incorporating PCR techniques are particularly well suited for same-day analysis of a primary colony isolate. The CFIA microbiology laboratory network has undertaken a program of method development aimed at the rapid identification of colonies isolated on plating media at an early stage during the enrichment process. A key technology platform adopted by CFIA for this purpose is the cloth-based hybridization array system (CHAS) providing for identification of pathogens through amplification of key target genes by multiplex PCR, followed by rapid hybridization of the amplicons with an array of immobilized capture probes on a polyester cloth support [2–4, 17]. This approach enables facile detection of many gene markers in a single reaction, with specificity assured through the hybridization process.

A CHAS method for the identification of *E. coli* O157:H7 [17] has been validated following the guidelines of the Health Canada *Compendium of Analytical Methods* (CAM) (available at [18]). This method has been published in the CAM (MFLP-22: Characterization of verotoxigenic *Escherichia coli* O157:H7 colonies by polymerase chain reaction and cloth-based hybridization array system, [2]), enabling its implementation for regulatory testing purposes in Canada. It is notable as the first instance of a genetic marker-based approach for the definitive identification of a foodborne pathogen isolate in our laboratories. The *E. coli* O157:H7 CHAS was used by CFIA laboratories on two separate occasions in 2013 to provide critical evidence supporting health risk assessments in connection with foodborne illness outbreaks implicating ground beef distributed in Canada. This method enabled the testing laboratories to issue official results of analysis a full 2 days ahead of the traditional approach, leading to more timely interventions minimizing public exposure to the contaminated product. The CHAS technology has now been incorporated into routine diagnostic testing schemes in CFIA food testing laboratories.

### *2.1.2 Enter the next generation: whole-genome sequencing*

Non-O157 STEC, particularly strains bearing certain O antigenic determinants (O26, O45, O103, O111, O121, O145, and O157), are emerging as a serious foodborne public health concern [19]. Unlike *E. coli* O157:H7, there are no biochemical features by which these so-called priority STEC strains can be differentiated from commensal *E. coli* or other STEC which are not a public health concern. However, it is universally recognized that foodborne STEC posing a public health risk can be defined on the basis of possession of certain gene markers, including the Shiga toxin genes *st1* or *st2*, the intimin-coding gene *eae*, and markers for the specific serogroups of concern [1, 19, 20]. Thus, priority STEC constitute a striking example illustrating the benefit of the use of gene markers for pinpointing pathogens otherwise not readily amenable to identification by classic means. This in turn has enabled practical strategies for multiplex gene detection methods for detection of such pathogens

during routine monitoring of the food supply to verify industry compliance with food safety regulations. In fact, such an approach is the basis for the Canadian STEC Method which was developed jointly by CFIA and Health Canada for the detection of this family of pathogens in meats and produce [3, 4].

The potential of PCR technology to provide informative test results is limited by its rather fragmentary nature, that is, the fact that only a relatively small number of different DNA markers can be assessed in a single analytical procedure. A considerable effort is required to optimize and validate the performance of PCR systems, particularly those in which multiple primer pairs are combined (e.g., identification of priority Shiga toxigenic *E. coli* (STEC) using a combination of 11 primer pairs; [1]). For all intents and purposes, validated assays are locked into their original configurations, and it is not possible to modify them (e.g., add or alter primers) on an ad hoc basis without first having to undertake a laborious and time-consuming re-evaluation of assay performance characteristics. Thus, novel queries cannot be undertaken if, in the course of an active food safety investigation, new questions arise regarding the occurrence of an unanticipated assay target. This was the case in a 2011 German outbreak involving STEC with an unusual virulence profile [21, 22].

The use of WGS technologies for STEC characterization can provide a more complete picture [8]; however, completion of WGS analyses typically requires 3–5 days. As this timeline may be too long in an outbreak situation, we have developed practical processes in which genomic DNA isolated from a single STEC colony is sequenced using the Illumina MiSeq platform, followed by analysis of the sequence data during the course of the sequencing run for the determination of genomic markers for phylogenetic identity, virulence profile, serotyping, as well as biological metrics serving as quality control features supporting the validity of the process [5]. Identification and characterization of isolates can be completed within 9 h, comparable to the current method used for characterization of STEC. This real-time WGS approach produces high-resolution characterization of bacterial pathogens at a cost and within a time-frame that are similar to standard microbiological techniques and has the potential to replace lengthy biochemical characterization and molecular and serological typing procedures widely used in food testing laboratories. Our laboratory is currently studying strategies for broad implementation of WGS technology in support of regulatory food inspection objectives through the detection, identification, and characterization of priority bacterial pathogens such as STEC, *Salmonella enterica*, and *Listeria monocytogenes* [5, 23–25]. We have developed guidelines and implemented validated methods and bioinformatics tools for automated analyses of sequence data to ensure reliability and reproducibility of WGS-based analyses [24, 26].

## **2.2 Have we seen it before? Impact of high-resolution molecular typing by WGS for distinguishing new isolates from control and historical laboratory isolates**

Following detection of a pathogen in foods, (sub)typing methods are often used to generate a profile of the isolated organism to determine similarity to previously characterized isolates from clinical or food sources (reviewed in [27–29]). Typing of isolates recovered from food samples can provide important information regarding the complexity and source(s) of a given contamination incident, enables tracking of foodborne bacterial strains, and is frequently used to support regulatory decisions.

Early methods of typing were based on phenotypic properties, for example, serotyping based on proteins expressed by the organism on surface structures [30–32] and phage typing based on susceptibility to reference panels of bacteriophage for distinguishing closely related isolates [33]. The development of methods based on DNA sequences followed [29]. Methods such as multilocus sequence typing

(MLST), involving sequencing of PCR-amplified fragments of a small number (i.e., 6–9) of housekeeping genes, can be used to infer evolutionary relationships among organisms [34, 35]. This portable and highly reproducible method of typing has been widely deployed, and MLST schema have been developed for all of the priority foodborne pathogens [34]. For public health surveillance in Canada, as in other jurisdictions, the standard approach for typing of foodborne pathogens for cluster identification has been pulsed-field gel electrophoresis (PFGE) and multilocus variable-number tandem repeat analysis (MLVA) [27, 28, 36, 37]. In North America, data is shared among public health agencies through PulseNet [27, 38]. The selection of a typing method depends on a number of factors, including proven utility of the method for the pathogen being investigated. Each method requires costly training of lab personnel and in many cases the purchase of specialized equipment [39]. Furthermore, comparisons of typing data among different strains can only be done in cases where the same method has been applied. In some cases, variability in the execution of methods by different analysts or different labs significantly impacts the comparability of molecular typing data [36].

These typing methods are based on a limited subset of genomic sequences and often lack the discriminatory power to differentiate among closely related organisms [39]. DNA typing profiles from two isolates appearing indistinguishable might be interpreted as evidence that the bacteria have a common source. However, the strength of this type of evidence rests on the extent to which the DNA profile consists of a combination of rare traits. When the traits defining a DNA profile are not rare, there is the possibility that two isolates are in fact unrelated and that matches are mere chance occurrences. In highly clonal strains (e.g., *Salmonella*) serovar Enteritidis, where only a few single-nucleotide changes may be observed among epidemiologically unrelated strains, most methods are not sufficiently discriminatory [40].

WGS provides a high-resolution molecular typing platform that can be universally applied to bacterial pathogens [6, 38, 41]. In principle, strains differing by even a single nucleotide can be distinguished [42, 43]. Furthermore, WGS can now be done more cheaply than lower resolution methods such as MLST and is backward-compatible with previous methods since, in some cases, typing data can be generated from minimally processed genomic data in silico [8, 9, 44]. Strains characterized by WGS can be compared to strains characterized by any other DNA-based subtyping method, enabling optimal use of historical data. Molecular typing data have generally been developed as a surrogate measure of the genetic similarity between bacterial strains. Using databases of WGS information, the utility of existing subtyping methods can be rigorously assessed, and improved subtyping schemes that reflect true strain relationships can be developed [45].

One advantage of the availability of WGS for typing bacterial isolates is the ability to evaluate datasets at different levels of resolution as needed to resolve biological questions. In this regard, MLST continues to be a valuable approach for tracking foodborne pathogens in the genomic era [41, 44]. WGS data can be matched to current and historical databases at different levels of resolution including pathogen-specific MLST (described above), core genome MLST (cgMLST) which uses hundreds of genes that are conserved within a species, and whole-genome MLST (wgMLST) which considers all genes within a species [38]. Similarly, ribosomal MLST (rMLST) is a 53-gene scheme with the advantage of being universally applicable to bacteria typically encountered in a food testing laboratory [46].

The highest-resolution WGS-based analyses such as wgMLST and single-nucleotide variant (SNV) analyses provide unrivaled DNA fingerprinting capability and offer tremendous potential for food safety applications [41, 42, 47]. Still, the use of these analyses in a food safety context is in its infancy, and the interpretation of genomic data from foodborne pathogens in support of regulatory interventions

remains challenging [48]. For example, how many SNVs are required to exclude a sample from a food safety investigation? This question remains difficult to answer, in part because of differences in rates at which DNA accumulates changes within a species or among strains within a species. Bacterial strains with a mutator phenotype have an elevated mutation rate, typically due to mutations in genes encoding components of DNA replication and repair pathways [49]. Mutator phenotypes are commonly found in clinical bacterial populations and may contribute significantly to the acquisition of antimicrobial resistance [50–52]. For example, the mutator phenotype has been attributed to the development of multidrug-resistant *S. typhimurium* [50]. In *E. coli*, this phenotype has recently been shown to be induced in response to stress conditions [53]. The possibility that environmental stress could lead to the development of the mutator phenotype may be highly relevant to analyses of foodborne pathogens.

### *2.2.1 Identification of persistent contamination*

Persistent contamination of food manufacturing environments with bacteria such as *Listeria monocytogenes* poses significant public health risks as these events serve as a source for the continual contamination of food products, primarily because (by definition) persistent contaminants defy attempts to sanitize the manufacturing environment using standard protocols [54–56]. *Listeria monocytogenes* is the causative agent of listeriosis, a potentially fatal foodborne illness in susceptible populations such as the very young, the elderly, and the immunocompromised [57]. The ability of this organism to contaminate food contact surfaces (e.g., conveyors, saws, etc.) and survive in the manufacturing environment increases the risk of food product contamination, with possible serious public health consequences. Routinely monitoring the food manufacturing environment for the presence of pathogenic bacteria is a key to preventing contamination of food products [56]. The presence of environmental contamination is usually ascertained through swabbing and testing of food manufacturing environment surfaces using standard microbiology techniques. Transient contamination is of a sporadic nature and may be effectively managed through the application of standard sanitation regimens followed by testing to ensure treatment efficacy. Persistent contamination occurs when a specific strain becomes a permanent resident of specific niches in the manufacturing environment. Persistence may be attributable to the incorporation of *L. monocytogenes* into biofilms occurring on equipment surfaces, resistance to commonly used sanitizers such as benzalkonium chloride, or through mechanical sequestration in hard-to-clean areas (e.g., meat cutting saw arbors, as was the case in the 2008 Canadian listeriosis outbreak resulting in many fatalities associated with the consumption of contaminated ready-to-eat meats [58]).

The ability to distinguish the two modes of contamination in the analysis of environmental isolates recovered during routine monitoring activities would constitute an important element to inform the best approach for the management of microbial hazards in food manufacturing plants [54, 55]. For example, while regular sanitation procedures may be effective in dealing with removal of sporadic surface contaminants, a more comprehensive approach requiring equipment tear-down and aggressive sanitation would be required to deal with persistent contaminants, which are by nature highly resistant to sanitizers and cleaning procedures. The traditional approach to identify the occurrence of persistent contamination in food manufacturing environments involves the characterization of successive isolates using typing procedures such as PFGE to determine their relatedness [54]. However, PFGE is of limited value for this purpose because it is not sufficiently discriminatory to unequivocally establish whether two strains are clonally related

(i.e., one being descendant from the other). Depending on the scope of the contamination, there may be multiple related populations within the food production environment.

Whole-genome sequencing approaches offer the prospect of determining the degree of relatedness among isolates on the basis of very fine base sequence differences, because more closely or clonally related isolates have fewer SNV differences. Therefore, it should be possible to compare two isolates (e.g., recovered on successive sampling incursions in the same plant) using high-resolution WGS typing methods to ascertain whether they are clonally related or different [55]. In the former case, this would be a strong indication that there is either an unresolved source of contamination in the plant, or more likely, a case of persistent contamination, whereas the latter case would suggest two independent contamination incidents. Each scenario would warrant a different approach to decontamination, and the ability to differentiate persistent and sporadic strains on the basis of the relatedness of successive isolates would constitute a powerful risk assessment and risk management tool for the use in the most highly proactive food safety programs. There is one caveat in the use of SNV-based typing, and that is the temporal drift which naturally occurs in bacteria, resulting in the accumulation of SNVs among the progeny derived from a single parent. The question remains under which conditions this occurs for bacterial strains in a food manufacturing environment and how many SNV differences constitute a real difference in terms of the provenance of isolates under comparison.

One cause of *L. monocytogenes* persistence has been identified as resistance to sanitizers such as quaternary ammonium compounds (QACs) [59–61]. An important determinant of QAC resistance is efflux systems such as those encoded by the *bcrABC* gene cassette [62] or *emrE* [63]. In a recent study to characterize the genomes of *L. monocytogenes* isolates collected at a pig slaughterhouse to determine the molecular basis for their persistence, we found that successive environmental isolates (persistent types) linked on the basis of SNV analyses all harbored the *bcrABC* and concluded that high-resolution typing and determination of the cassette may serve to distinguish between persistent and sporadic *L. monocytogenes* isolates [59]. This in turn may have important ramifications for risk management actions when *L. monocytogenes* is recovered from a food manufacturing environment, since the ability of a strain to persist casts doubt on the efficacy of standard sanitization protocols, and more intensive cleaning procedures (e.g., equipment teardown, use of alternative sanitizers) may be warranted. These types of analyses of food inspection isolates are greatly facilitated by WGS technology [55, 59].

### 2.2.2 Contribution to surveillance programs and outbreak investigations

Although numerous methods are used by food safety and public health agencies to support regulatory decisions during outbreak investigations, demonstrating that food and clinical isolates originated from the same source can be challenging. As the results generated by WGS make their way into situation rooms to guide decision-makers, concise metrics for the interpretation and contextualization of genomics-derived data will be required to achieve more precise assessments [64]. The value of WGS-based typing for cluster identification has already been demonstrated through global surveillance networks such as GenomeTrakr and PulseNet [6, 38, 41]. Smaller clusters of cases can be linked through WGS analyses and investigated [65], and conversely, unrelated cases can be excluded from an epidemiological investigation leading to improved outcomes for rapid identification of foods implicated in outbreaks [66]. Nonetheless, WGS results should not be interpreted in the absence of epidemiological context as some lineage rates of mutation are low and strains from different sources may appear to be linked [67].



The concepts of “match probability” and “likelihood ratios” are well known in human forensic sciences where they facilitate the interpretation of DNA profiles in matching individuals to a crime scene [68]. For example, when the DNA profile found on a crime scene matches that of a suspect and there is only a one-in-one million probability that this DNA profile might be found in another individual, there is a strong case linking the suspect to the crime scene. Food inspectors face a similar situation during outbreak investigations when trying to establish causal links between isolates from different sources [69]. Bacteria may undergo subtle changes in their genomes during the course of a foodborne illness outbreak event, with possible impacts on the typing profiles of clonally related isolates recovered over time. The question arises as to how much change in a genome constitutes a significant difference between individual isolates (i.e., different origins or strains). Through statistical analyses of comprehensive pathogen genome databases, it should be possible to develop a likelihood ratio approach to determine the probability of finding a given profile in a defined population and, hence, develop criteria to measure sequence diversity between isolates with different degrees of relatedness and even among clonally related isolates recovered over the course of an outbreak event [69]. This in turn would provide a greater degree of confidence in attributing the origins of isolates, identifying clusters of foodborne illness and their food vehicles, and the scope of contamination. This information can also be used to revise and adjust detection tools (e.g., PCR primers) to ensure their effectiveness in identifying “moving” genomic targets. The development of a forensic likelihood ratio approach would provide a valuable tool to assess the reliability of genomic information underlying regulatory decision-making.

### *2.2.3 Attribution of food vehicles through genomic surveillance*

The advent of genomic typing augurs well for the creation of highly refined databases of bacterial isolates from various sources (foods, production facilities, farms, environmental and clinical strains) providing high-resolution characterization of individual strains with established linkages to their geographic and temporal origins. Historically, the use of low-resolution typing approaches such as MLST profiles or serotypes has been valuable for the association of specific lineages with a given food type, production environment, or country [70, 71]. Initiatives such as the GenomeTrakr and the PulseNet WGS networks represent rich resources from which to draw valuable information linking isolates to their origin in the food production continuum [6, 38, 72]. For example, an analysis of *E. coli* O157:H7 identified SNVs associated with country of origin [73]. Similarly, an outbreak involving *S. Bareilly* in the United States was tracked to a food originating in India based on high-resolution SNV typing [74].

With the aid of bioinformatics tools, databases can be queried to identify genomic signatures that are overrepresented in particular host sources for bacterial isolates. For example, a study by Thépault et al. [75] identified 15 host-associated *C. jejuni* markers and demonstrated utility of these markers for identifying host association of strains with 80% accuracy. The ability to discern this type of information would be a tremendous boon for foodborne illness investigations: WGS data could be used to determine the “source signature” of clinical isolates, enabling a highly proactive approach in rapidly narrowing the field for the attribution of food vehicles. Regulatory food inspection agencies such as the CFIA would have an important role to play in such a scheme. Ongoing, extensive sampling plans will be required to ensure adequate representation of different food production elements, such as food types and geographic provenance. Given that most cases of foodborne illness occur sporadically, this approach would enable public health authorities to track the causes

of a larger proportion of cases of foodborne illness. This would ultimately lead to a better understanding of foods commonly implicated in disease and the implementation of more effective interventions to reduce the burden of foodborne illness.

### 2.3 Is it dangerous? Rapid identification of virulence, antimicrobial resistance, and epidemiological markers through WGS

Genomic information is highly complex, and there are many knowledge gaps with respect to the significance of various marker genes to public health [76]. Nonetheless, there is a growing body of evidence linking certain well-defined gene markers to virulence characteristics of bacteria, for example, the role of intimin (coded by the *eae* gene) in the pathogenesis of STEC, epidemiological associations between certain serotype features and outbreaks of serious foodborne illness (e.g., *L. monocytogenes* serogroups 1/2a, 1/2b, and 4b, STEC serogroups O26, O45, O103, O111, O121, O145, and O157), and even the type of toxin secreted (e.g., Shiga toxin 2) and the attendant severity of illness [77, 78].

In the case of STEC, regulatory food testing programs currently define priority target strains as bearing markers for Shiga toxin genes and intimin, in addition to markers associated with a narrow family of O serogroups [2]. However, the question arises whether in the course of conducting routine monitoring of food inspection samples the occurrence of an isolate bearing markers for Shiga toxin and intimin, but none of the so-called priority serogroups would be actionable. There are varying subjective opinions on the matter, ranging from a narrow interpretation of test results in which only isolates bearing all the designated factors are considered hazardous to the more precautionary approach whereby any isolate bearing both Shiga toxin and intimin factors, regardless of O serogroup, constitutes a public health risk. There is also evidence suggesting that severity or likelihood of foodborne illness varies with Shiga toxin type and subtype (e.g., STEC strains possessing st2a tend to be more frequently implicated in cases of severe foodborne illness [79]) and that this should be a factor in determining the appropriate response to the presence of a food contaminant. These properties are readily discoverable through the analysis of WGS data. For instance, the Shiga toxin subtype can be reliably determined using the V-typer tool, which is an automated assembly-independent subtyping module that can be integrated in a bioinformatics pipeline for the analysis of foodborne STEC isolates [23]. Yet another possibility would be to define priority STEC on the basis of contemporary public health data (reviewed periodically) identifying STEC serogroups most frequently associated with illness in a given jurisdiction. The serogroup of an *E. coli* isolate can be determined using tools such as SeroTypeFinder [8] which can be freely accessed through the Center for Genomic Epidemiology website [80].

Such considerations raise problems for health risk assessment specialists who must interpret laboratory results (among other factors) to determine the degree of risk informing the course of regulatory interventions [76]. It is tempting to speculate that it may be possible to devise an objective scheme for rating the degree of hazard associated with a given isolate on the basis of genomic analyses. For instance, the public health and food inspection communities could agree on a list of key factors relevant to the characterization of a given pathogen (**Table 1**). For organisms such as *E. coli*, acquired virulence gene databases have been established, and isolates may be profiled using tools such as VirulenceFinder [7], accessible through the Center for Genomic Epidemiology website [80]. Since not all factors have the same impact, it should be possible to develop a weighted index approach in which each constituent factor determined by genomic analysis makes up a fraction of a final index value which is proportional to the degree of hazard. Such an index value (hazard characterization or HazChar Score), used in conjunction with

Key factors	Element	Relevance
Primary virulence	Toxin	Presence or absence
	Attachment and colonization	<i>eae</i> , enteroaggregative factors
	Pathogenicity	Pathogenesis mechanisms (e.g., hemolysin)
Severity modulator	Type	<i>st1</i> vs. <i>st2</i>
	Subtype	<i>st2<sub>a</sub></i> vs. <i>st2<sub>f</sub></i>
Accessory functions	Antibiotic resistance	Therapeutic impact
	Antimicrobial resistance	Sanitizer efficacy
	Biofilm formation	Persistence
	Pathogenicity islands	Signatures for novel pathogens
Epidemiological markers	Serotype	Outbreak vs. sporadic vs. nil association
	Phage type	Reservoirs, illness outbreaks
	Molecular type	PFGE/SNV cluster
Phylogenetic markers	Genus/species	Identity (e.g., <i>Salmonella</i> , <i>E. coli</i> )
	Family or group	STEC

<sup>a</sup>A list of key factors is developed for a given pathogen, and each element is assigned a weighted value based on its significance in human illness. Genomic analysis of a foodborne isolate by whole-genome sequencing with the application of bioinformatics tools to determine the presence of targeted features. The individual weighted values are summed giving the HazChar Score, which is then compared against a set of predetermined criteria to categorize the degree of risk.

**Table 1.**  
Proposed concept for hazard characterization: HazChar Score<sup>a</sup>.

numerical criteria derived from historical data, could form a basis for attributing the degree of hazard associated with a particular isolate, which in turn would enable an objective categorization of risk to inform the appropriate regulatory response.

The antimicrobial resistance profile of pathogenic bacteria, while not a virulence attribute per se, remains an important factor in the ultimate public health outcomes of foodborne illness events, since a significant fraction of the affected population (e.g., the elderly and the immunocompromised) may critically require antibiotic therapy to recover. Furthermore, antimicrobial use at sub-therapeutic levels for growth promotion in food animal production has been implicated in the development of antimicrobial resistance (AMR) in animals and humans [81, 82], though there is paucity of data to support this claim. Food testing laboratories can play an important role in contributing data on the occurrence of AMR bacteria to national and international surveillance initiatives seeking to understand the role of production practices in the emergence of these microorganisms. As an alternative to labor- and time-consuming phenotypic testing, AMR profiles can be predicted from WGS data through the identification of genetic markers by querying the subject genome using DNA sequence information deposited in curated AMR gene databases, such as well-cataloged AMR gene markers. A number of tools are currently available to predict AMR from bacterial WGS data (e.g., ResFinder [10, 83], SEAR [84], Resistance Gene Identifier [85], and ARMI [86]). These AMR marker prediction tools rely on curated international AMR gene databases such as CARD [85], ARDB [87], and ARG-ANNOT [88]. WGS-based methods for prediction of AMR phenotype have been shown to be highly accurate [86, 89–91].

### 3. Future applications

#### 3.1 Deployment of ad hoc methods in support of outbreak investigations

Despite recent efforts of regulatory food safety agencies to implement test methods targeting defined serogroups of so-called priority STEC, the history of foodborne disease outbreaks is rife with examples of causative strains with unexpected characteristics (e.g., the 2011 German outbreak in which the etiologic agent belonged to serogroup O104, not a designated priority serogroup, and lacked the definitive virulence marker *eae* [21]), making it difficult to anticipate detection methods suiting all contingencies. Detection is further complicated by variability among non-O157 STEC strains in resistance to selective agents commonly used in enrichment culture techniques, hindering their recovery from foods bearing high levels of background microbiota [92–94]. Genomic information garnered from clinical bacterial isolates implicated in outbreaks of foodborne illness will be very useful in the customization of selective recovery and identification procedures to facilitate their detection in food samples during outbreak investigations [86].

The state of the art in WGS technology has reached the point where clinical isolates implicated in foodborne disease outbreaks are routinely sequenced in public health laboratories at an early stage during these events [38]. With the application of appropriate bioinformatics tools to analyze the ensuing data, it should be possible to develop customized strain-specific test methods that can be rapidly deployed to food testing labs conducting analyses in support of outbreak investigations. The availability of WGS information for these strains should make it possible to ascertain the presence of traits conferring resistance to antimicrobial agents such as antibiotics, quaternary ammonium compounds, and tellurite, suggesting an avenue for the formulation of customized selective enrichment media enabling recovery of specific outbreak strains [86]. This would be a particular advantage in instances where a food matrix (e.g., meats, sprouts, etc.) contains high levels of background microbiota, which might otherwise interfere with recovery of the target organism. Genomic AMR prediction tools can be used to discern the AMR marker profile of a strain of interest (e.g., outbreak strain) to identify an antibiotic resistance trait which can be exploited for customization of selective enrichment media favoring its recovery from samples with high background bacteria loads [86, 95]. In addition, WGS data can be analyzed using a pipeline such as SigSeekr [96] designed to identify DNA sequences associated with a particular strain for its rapid identification by PCR. By combining strain-specific selective enrichment and PCR detection tools, it should be possible to deploy custom recovery and identification tools for the efficient detection of STEC outbreak strains within the timeframe of an active investigation. The feasibility of such an approach has been demonstrated using laboratory STEC strains as models, with resistance to a variety of antibiotic classes used as the basis for their selective recovery against high backgrounds of commensal *E. coli* bacteria in ground beef samples [86].

#### 3.2 Characterization of food microbiomes in support of improved method development

Metagenomic analysis of enrichment dynamics can be used to inform the development of improved methods for cultural enrichment of pathogens [97–99]. A practical approach for this is to selectively amplify hypervariable regions within the 16S rDNA and to sequence amplicons using WGS technologies. Sequences are then mapped to databases to determine composition of microbial communities using bioinformatics tools such as QIIME [100] or mothur [101]. Samples from

enrichment cultures can be used to evaluate growth of target pathogens relative to background food microbiota over time [97, 99]. Such studies can provide valuable insight into species that could potentially interfere with target pathogens that can be applied to the development of improved methodology.

A modern concept in the study of pathogenic bacteria is the emergence of novel pathogens among commensals in a given environment through the acquisition of virulence factors by horizontal gene transfers from other bacteria [102–104]. The evolutionary trail of the STEC family suggests a priori transformations of benign *E. coli* strains into virulent STEC having acquired exogenous DNA segments such as bacteriophage carrying Shiga toxin genes and pathogenicity islands harboring host colonization factors [105]. There is evidence that other food pathogens such as *L. monocytogenes* strains implicated in serious outbreaks of foodborne illness may have acquired enhanced virulence characteristics through horizontal gene transfer processes [106]. This is believed to occur on a relatively short time scale, perhaps on the order of weeks or months, making the emergence of novel pathogens in food production environments or animal reservoirs in near real time a significant possibility. Furthermore, food-acquired coinfections may arise in which two or more bacterial strains complement each another, for example, a toxigenic strain lacking adherence factors might colonize a host by cross-utilizing a factor secreted by another strain [107]. It may be possible to predict the probability of emergence of novel pathogens with enhanced virulence or antibiotic resistance characteristics in the food supply through periodic microbial metagenomic analyses to ascertain the presence of key indicators in the background microflora of food commodities (e.g., ground beef, trim), food manufacturing environments (e.g., floors, food contact surfaces), and animal reservoirs (e.g., cattle, poultry) [108, 109]. A weighted index approach much like that described for the HazChar Score above could be employed here, with possible inclusion of a more comprehensive catalog of known virulence, AMR, and other critical factors relevant to public health.

#### **4. Conclusions**

Modern food microbiology research has generated a critical understanding of the epidemiology, pathogenic mechanisms, virulence factors, and other salient characteristics of the major food pathogens. The convergence of expanded scientific knowledge and sophisticated technological capability create exciting new opportunities for the refinement of food microbiology testing programs to meet the needs of a comprehensive risk-based inspection approach. Advances in next-generation sequencing technologies have made it possible for investigators to carry out sequencing and processing of bacterial genomes within the time course of a typical foodborne illness outbreak investigation. It may reasonably be expected that in the near future, analysts will be moving from traditional DNA hybridization approaches (e.g., PCR and microarrays) toward rapid whole-genome sequencing allowing a much more comprehensive examination of the isolate at hand. This new approach will require broad access to leading-edge bioinformatics capability for analysis of complex genomics data in silico to ascertain the presence of key genetic markers (e.g., presence of virulence genes in bacterial pathogens, completeness and functionality of gene products, markers for molecular typing, etc.). The generation and analysis of WGS information requires the migration of large packets of information between laboratory sites involved in the exploitation of this information, remote computing sites, and internet databases for data manipulation and comparative analyses. There are many ways in which the high-tech needs of the future can be met, even for relatively small laboratories with low operating budgets.

Opportunities abound for food inspection, public health, and academic laboratories to pool their resources and serve one another in the common purpose of safeguarding citizens from preventable food-acquired illness.

## **Acknowledgements**

The authors thank Paul Manninger, Mylène Deschênes, and Martine Dixon for providing laboratory support throughout many years in developing the food pathogen genomics program at the Canadian Food Inspection Agency and Andrew Low for providing bioinformatics support and for critical review of this manuscript.

IntechOpen

## **Author details**

Catherine D. Carrillo, Adam Koziol, Neil Vary and Burton W. Blais\*  
Research and Development, Ottawa Laboratory (Carling), Canadian Food  
Inspection Agency, Ottawa, Ontario, Canada

\*Address all correspondence to: [burton.blais@canada.ca](mailto:burton.blais@canada.ca)

## **IntechOpen**

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

## References

- [1] Blais BW, Gauthier M, Deschênes M, Huszczyński G. Polyester cloth-based hybridization array system for identification of enterohemorrhagic *Escherichia coli* serogroups O26, O45, O103, O111, O121, O145, and O157. *Journal of Food Protection*. 2012;75:1691-1697. DOI: 10.4315/0362-028X.JFP-12-116
- [2] Blais BW, Gauthier M, Deschenes M, Huszczyński G. Characterization of verotoxigenic *Escherichia coli* O157:H7 colonies by polymerase chain reaction (PCR) and cloth-based hybridization array system (CHAS) (MFLP-22). In: *Compendium of Analytical Methods, Laboratory Procedures for the Microbiological Analysis of Foods*. Vol. 3. Ottawa, Canada: Health Canada; 2013. <https://www.canada.ca/en/health-canada/services/food-nutrition/research-programs-analytical-methods/analytical-methods/compendium-methods.html> [Accessed: February 20, 2019]
- [3] Blais BW, Martinez A, Gill A, McIlwham S, Mohajer S, Gauthier M. Isolation and identification of priority verotoxigenic *Escherichia coli* (VTEC) in meat and vegetable products (MFLP-52). In: *Compendium of Methods, Laboratory Procedures for the Microbiological Analysis of Foods*. Vol. 3. Ottawa, Canada: Health Canada; 2014. Available from: <https://www.canada.ca/en/health-canada/services/food-nutrition/research-programs-analytical-methods/analytical-methods/compendium-methods.html> [Accessed: February 20, 2019]
- [4] Huszczyński G, Gauthier M, Mohajer S, Gill A, Blais B. Method for the detection of priority Shiga toxin-producing *Escherichia coli* in beef trim. *Journal of Food Protection*. 2013;76:1689-1696. DOI: 10.4315/0362-028X.JFP-13-059
- [5] Lambert D, Carrillo CD, Koziol AG, Manninger P, Blais BW. GeneSippr: A rapid whole-genome approach for the identification and characterization of foodborne pathogens such as priority Shiga toxigenic *Escherichia coli*. *PLoS One*. 2015;10:e0122928. DOI: 10.1371/journal.pone.0122928
- [6] Allard MW, Strain E, Melka D, Bunning K, Musser SM, Brown EW, et al. Practical value of food pathogen traceability through building a whole-genome sequencing network and database. *Journal of Clinical Microbiology*. 2016;54:1975-1983. DOI: 10.1128/JCM.00081-16
- [7] Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, et al. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *Journal of Clinical Microbiology*. 2014;52:1501-1510. DOI: 10.1128/JCM.03617-13
- [8] Joensen KG, Tetzschner AMM, Iguchi A, Aarestrup FM, Scheutz F. Rapid and easy *in silico* serotyping of *Escherichia coli* isolates by use of whole-genome sequencing data. *Journal of Clinical Microbiology*. 2015;53:2410-2426. DOI: 10.1128/JCM.00008-15
- [9] Yoshida CE, Kruczkiewicz P, Laing CR, Lingohr EJ, Gannon VPJ, Nash JHE, et al. The *Salmonella* In Silico Typing Resource (SISTR): An open web-accessible tool for rapidly typing and subtyping draft *Salmonella* genome assemblies. *PLoS One*. 2016;11:e0147101. DOI: 10.1371/journal.pone.0147101
- [10] Kleinheinz KA, Joensen KG, Larsen MV. Applying the ResFinder and VirulenceFinder web-services for easy identification of acquired antibiotic resistance and *E. coli* virulence genes in bacteriophage and prophage

nucleotide sequences. Bacteriophage. 2014;**4**:e27943. DOI: 10.4161/bact.27943

[11] Ronholm J, Nasheri N, Petronella N, Pagotto F. Navigating microbiological food safety in the era of whole-genome sequencing. *Clinical Microbiology Reviews*. 2016;**29**:837-857. DOI: 10.1128/CMR.00056-16

[12] Sharapov UM, Wendel AM, Davis JP, Keene WE, Farrar J, Sodha S, et al. Multistate outbreak of *Escherichia coli* O157:H7 infections associated with consumption of fresh spinach: United States, 2006. *Journal of Food Protection*. 2016;**79**:2024-2030. DOI: 10.4315/0362-028X.JFP-15-556

[13] Berenger BM, Berry C, Peterson T, Fach P, Delannoy S, Li V, et al. The utility of multiple molecular methods including whole genome sequencing as tools to differentiate *Escherichia coli* O157:H7 outbreaks. *Eurosurveillance*. 2015;**20**:30073. DOI: 10.2807/1560-7917.ES.2015.20.47.30073

[14] Seys SA, Sampedro F, Hedberg CW. Assessment of Shiga toxin-producing *Escherichia coli* O157 illnesses prevented by recalls of beef products. *Foodborne Pathogens and Disease*. 2015;**12**:800-805. DOI: 10.1089/fpd.2015.1968

[15] Microbiological Methods Committee. Isolation of *Escherichia coli* O157:H7/NM from foods and environmental surface samples (MFHPB-10). In: *Compend. Anal Methods*. Ottawa, Canada: Health Canada. Vol. 2. Ottawa, Canada: Health Canada; 2017. Available from: <https://www.canada.ca/en/health-canada/services/food-nutrition/research-programs-analytical-methods/analytical-methods/compendium-methods.html> [Accessed: February 20, 2019]

[16] USDA-FSIS. Detection, isolation and identification of top seven

Shiga toxin-producing *Escherichia coli* (STECs) from meat products and carcass and environmental sponges. United States Department of Agriculture, Athens, GA, 2019. Available from: <https://www.fsis.usda.gov/wps/wcm/connect/fsis-content/internet/main/topics/science/laboratories-and-procedures/guidebooks-and-methods/microbiology-laboratory-guidebook/microbiology-laboratory-guidebook> [Accessed April 18, 2019]

[17] Martinez-Perez A, Blais BW. Cloth-based hybridization array system for the identification of *Escherichia coli* O157:H7. *Food Control*. 2010;**21**:1354-1359. DOI: 10.1016/j.foodcont.2010.04.009

[18] Health Canada. The Compendium of Analytical Methods. n.d. Available from: <https://www.canada.ca/en/health-canada/services/food-nutrition/research-programs-analytical-methods/analytical-methods/compendium-methods.html> [Accessed: February 20, 2019]

[19] Catford A, Kouamé V, Martinez-Perez A, Gill A, Buenaventura E, Couture H, et al. Risk profile on non-O157 verotoxin-producing *Escherichia coli* in produce, beef, milk and dairy products in Canada. *International Food Risk Analysis Journal*. 2014;**4**:1-25. DOI: 10.5772/59208

[20] EFSA Panel on Biological Hazards (BIOHAZ). Scientific opinion on VTEC-seropathotype and scientific criteria regarding pathogenicity assessment. *EFSA Journal*. 2013;**11**:3138. DOI: 10.2903/j.efsa.2013.3138

[21] Burger R. EHEC O104:H4 in Germany 2011: Large Outbreak of Bloody Diarrhea and Haemolytic Uraemic Syndrome by Shiga Toxin-producing *E. coli* via Contaminated Food. In: Institute of Medicine (US). *Improving Food Safety Through a One Health Approach: Workshop Summary*. Washington DC: National Academies Press (US); 2012.



Available from: <https://www.ncbi.nlm.nih.gov/books/NBK114499/> [Accessed February 20, 2019]

[22] Frank C, Werber D, Cramer JP, Askar M, Faber M, an der Heiden M, et al. Epidemic profile of Shiga-toxin-producing *Escherichia coli* O104:H4 outbreak in Germany. The New England Journal of Medicine. 2011;**365**:1771-1780. DOI: 10.1056/NEJMoa1106483

[23] Carrillo CD, Koziol AG, Mathews A, Goji N, Lambert D, Huszczyński G, et al. Comparative evaluation of genomic and laboratory approaches for determination of Shiga toxin subtypes in *Escherichia coli*. Journal of Food Protection. 2016;**79**:2078-2085. DOI: 10.4315/0362-028X.JFP-16-228

[24] Lambert D, Pightling A, Griffiths E, Van Domselaar G, Evans P, Berthelet S, et al. Baseline practices for the application of genomic data supporting regulatory food safety. Journal of AOAC International. 2017;**100**:721-731. DOI: 10.5740/jaoacint.16-0269

[25] Low AJ, Koziol AG, Manninger PA, Blais BW, Carrillo CD. ConFindr: Rapid detection of intraspecies and cross-species contamination in bacterial whole-genome sequence data. 2019;**7**:e27499v1. DOI: 10.7287/peerj.preprints.27499v1

[26] Angers-Loustau A, Petrillo M, Bengtsson-Palme J, Berendonk T, Blais B, Chan K-G, et al. The challenges of designing a benchmark strategy for bioinformatics pipelines in the identification of antimicrobial resistance determinants using next generation sequencing technologies. F1000 Research. 2018;**7**:459. DOI: 10.12688/f1000research.14509.2

[27] Ribot EM, Hise KB. Future challenges for tracking foodborne diseases: PulseNet, a 20-year-old US surveillance system for foodborne

diseases, is expanding both globally and technologically. EMBO Reports. 2016;**17**:1499-1505. DOI: 10.15252/embr.201643128

[28] Swaminathan B, Barrett TJ, Hunter SB, Tauxe RV, CDC PulseNet Task Force. PulseNet: The molecular subtyping network for foodborne bacterial disease surveillance, United States. Emerging Infectious Diseases. 2001;**7**:382-389. DOI: 10.3201/eid0703.010303

[29] Taboada EN, Clark CG, Sproston EL, Carrillo CD. Current methods for molecular typing of *Campylobacter* species. Journal of Microbiological Methods. 2013;**95**:24-31. DOI: 10.1016/j.mimet.2013.07.007

[30] Herikstad H, Motarjemi Y, Tauxe RV. *Salmonella* surveillance: A global survey of public health serotyping. Epidemiology and Infection. 2002;**129**:1-8

[31] Penner JL, Hennessy JN. Passive hemagglutination technique for serotyping *Campylobacter fetus* subsp. *jejuni* on the basis of soluble heat-stable antigens. Journal of Clinical Microbiology. 1980;**12**:732-737

[32] Evans DJ, Evans DG. Classification of pathogenic *Escherichia coli* according to serotype and the production of virulence factors, with special reference to colonization-factor antigens. Reviews of Infectious Diseases. 1983;**5**(Suppl 4):S692-S701

[33] Hickman-Brenner FW, Stubbs AD, Farmer JJ. Phage typing of *Salmonella enteritidis* in the United States. Journal of Clinical Microbiology. 1991;**29**:2817-2823

[34] Maiden MCJ. Multilocus sequence typing of bacteria. Annual Review of Microbiology. 2006;**60**:561-588. DOI: 10.1146/annurev.micro.59.030804.121325

- [35] Maiden MCJ, Bygraves JA, Feil E, Morelli G, Russell JE, Urwin R, et al. Multilocus sequence typing: A portable approach to the identification of clones within populations of pathogenic microorganisms. *Proceedings of the National Academy of Sciences of the United States of America*. 1998;**95**:3140-3145
- [36] Schjørring S, Niskanen T, Torpdahl M, Björkman JT, Nielsen EM. Evaluation of molecular typing of foodborne pathogens in European reference laboratories from 2012 to 2013. *Euro Surveillance*. 2016;**21**:pii=30429. DOI: 10.2807/1560-7917.ES.2016.21.50.30429
- [37] Barrett TJ, Lior H, Green JH, Khakhria R, Wells JG, Bell BP, et al. Laboratory investigation of a multistate food-borne outbreak of *Escherichia coli* O157:H7 by using pulsed-field gel electrophoresis and phage typing. *Journal of Clinical Microbiology*. 1994;**32**:3013-3017
- [38] Nadon C, Van Walle I, Gerner-Smidt P, Campos J, Chinen I, Concepcion-Acevedo J, et al. PulseNet international: Vision for the implementation of whole genome sequencing (WGS) for global food-borne disease surveillance. *Eurosurveillance*. 2017;**22**:pii=30544. DOI: 10.2807/1560-7917.ES.2017.22.23.30544
- [39] Grant K, Jenkins C, Arnold C, Green J, Zambon M. Implementing Pathogen Genomics: A Case Study. *Public Health England*. 2018. Available from: <https://www.gov.uk/government/publications/implementing-pathogen-genomics-a-case-study> [Accessed February 23, 2019]
- [40] Deng X, Desai PT, den Bakker HC, Mikoleit M, Tolar B, Trees E, et al. Genomic epidemiology of *Salmonella enterica* serotype Enteritidis based on population structure of prevalent lineages. *Emerging Infectious Diseases*. 2014;**20**:1481-1489. DOI: 10.3201/eid2009.131095
- [41] Jagadeesan B, Gerner-Smidt P, Allard MW, Leuillet S, Winkler A, Xiao Y, et al. The use of next generation sequencing for improving food safety: Translation into practice. *Food Microbiology*. 2019;**79**:96-115. DOI: 10.1016/j.fm.2018.11.005
- [42] Petkau A, Mabon P, Sieffert C, Knox NC, Cabral J, Iskander M, et al. SNVPhyl: A single nucleotide variant phylogenomics pipeline for microbial genomic epidemiology. *Microbial Genomics*. 2017;**3**:e000116. DOI: 10.1099/mgen.0.000116
- [43] Markell JA, Koziol AG, Lambert D. Draft genome sequence of *Escherichia coli* O157:H7 ATCC 35150 and a nalidixic acid-resistant mutant derivative. *Genome Announcements*. 2015;**3**:e00734-15. DOI: 10.1128/genomeA.00734-15
- [44] Maiden MCJ, Jansen van Rensburg MJ, Bray JE, Earle SG, Ford SA, Jolley KA, et al. MLST revisited: The gene-by-gene approach to bacterial genomics. *Nature Reviews. Microbiology*. 2013;**11**:728-736. DOI: 10.1038/nrmicro3093
- [45] Carrillo CD, Kruczkiewicz P, Mutschall S, Tudor A, Clark C, Taboada EN. A framework for assessing the concordance of molecular typing methods and the true strain phylogeny of *Campylobacter jejuni* and *C. coli* using draft genome sequence data. *Frontiers in Cellular and Infection Microbiology*. 2012;**2**:57. DOI: 10.3389/fcimb.2012.00057
- [46] Jolley KA, Bliss CM, Bennett JS, Bratcher HB, Brehony C, Colles FM, et al. Ribosomal multilocus sequence typing: Universal characterization of bacteria from domain to strain. *Microbiology*. 2012;**158**:1005-1015. DOI: 10.1099/mic.0.055459-0

- [47] Chen Y, Luo Y, Carleton H, Timme R, Melka D, Muruvanda T, et al. Whole genome and core genome multilocus sequence typing and single nucleotide polymorphism analyses of *Listeria monocytogenes* associated with an outbreak linked to cheese, United States, 2013. *Applied and Environmental Microbiology*. 2017;**83**:pii: e00633-17. DOI: 10.1128/AEM.00633-17
- [48] Gilmour MW, Graham M, Reimer A, Van Domselaar G. Public health genomics and the new molecular epidemiology of bacterial pathogens. *Public Health Genomics*. 2013;**16**:25-30. DOI: 10.1159/000342709
- [49] LeClerc JE, Li B, Payne WL, Cebula TA. High mutation frequencies among *Escherichia coli* and *Salmonella* pathogens. *Science*. 1996;**274**:1208-1211. DOI: 10.1126/science.274.5290.1208
- [50] Kozyreva VK, Ilina EN, Malakhova MV, Carattoli A, Azizov IS, Tapalski DV, et al. Long-term dissemination of CTX-M-5-producing hypermutable *Salmonella enterica* serovar typhimurium sequence type 328 strains in Russia, Belarus, and Kazakhstan. *Antimicrobial Agents and Chemotherapy*. 2014;**58**:5202-5210. DOI: 10.1128/AAC.02506-14
- [51] Pope CF, O'Sullivan DM, McHugh TD, Gillespie SH. A practical guide to measuring mutation rates in antibiotic resistance. *Antimicrobial Agents and Chemotherapy*. 2008;**52**:1209-1214. DOI: 10.1128/AAC.01152-07
- [52] Eliopoulos GM, Blázquez J. Hypermutation as a factor contributing to the acquisition of antimicrobial resistance. *Clinical Infectious Diseases*. 2003;**37**:1201-1209. DOI: 10.1086/378810
- [53] Swings T, Van den Bergh B, Wuyts S, Oeyen E, Voordeckers K, Verstrepen KJ, et al. Adaptive tuning of mutation rates allows fast response to lethal stress in *Escherichia coli*. *eLife*. 2017;**6**:e22939. DOI: 10.7554/eLife.22939
- [54] Leong D, Alvarez-Ordóñez A, Jordan K. Monitoring occurrence and persistence of *Listeria monocytogenes* in foods and food processing environments in the Republic of Ireland. *Frontiers in Microbiology*. 2014;**5**:436. DOI: 10.3389/fmicb.2014.00436
- [55] Nastasijevic I, Milanov D, Velebit B, Djordjevic V, Swift C, Painset A, et al. Tracking of *Listeria monocytogenes* in meat establishment using whole genome sequencing as a food safety management tool: A proof of concept. *International Journal of Food Microbiology*. 2017;**257**:157-164. DOI: 10.1016/j.ijfoodmicro.2017.06.015
- [56] Malley TJV, Butts J, Wiedmann M. Seek and destroy process: *Listeria monocytogenes* process controls in the ready-to-eat meat and poultry industry. *Journal of Food Protection*. 2015;**78**:436-445. DOI: 10.4315/0362-028X.JFP-13-507
- [57] Thomas MK, Murray R, Flockhart L, Pintar K, Fazil A, Nesbitt A, et al. Estimates of foodborne illness-related hospitalizations and deaths in Canada for 30 specified pathogens and unspecified agents. *Foodborne Pathogens and Disease*. 2015;**12**:820-827. DOI: 10.1089/fpd.2015.1966
- [58] Government of Canada. Report of the Independent Investigator into the 2008 listeriosis outbreak 2009. Available from: <http://publications.gc.ca/site/eng/361474/publication.html> [Accessed: February 18, 2019]
- [59] Cherifi T, Carrillo C, Lambert D, Miniaï I, Quessy S, Larivière-Gauthier G, et al. Genomic characterization of *Listeria monocytogenes* isolates

reveals that their persistence in a pig slaughterhouse is linked to the presence of benzalkonium chloride resistance genes. *BMC Microbiology*. 2018;**18**:220. DOI: 10.1186/s12866-018-1363-9

[60] Martínez-Suárez JV, Ortiz S, López-Alonso V. Potential impact of the resistance to quaternary ammonium disinfectants on the persistence of *Listeria monocytogenes* in food processing environments. *Frontiers in Microbiology*. 2016;**7**:638. DOI: 10.3389/fmicb.2016.00638

[61] Yu T, Jiang X, Zhang Y, Ji S, Gao W, Shi L. Effect of benzalkonium chloride adaptation on sensitivity to antimicrobial agents and tolerance to environmental stresses in *Listeria monocytogenes*. *Frontiers in Microbiology*. 2018;**9**:2906. DOI: 10.3389/fmicb.2018.02906

[62] Elhanafi D, Dutta V, Kathariou S. Genetic characterization of plasmid-associated benzalkonium chloride resistance determinants in a *Listeria monocytogenes* strain from the 1998-1999 outbreak. *Applied and Environmental Microbiology*. 2010;**76**:8231-8238. DOI: 10.1128/AEM.02056-10

[63] Kovacevic J, Ziegler J, Wałęcka-Zacharska E, Reimer A, Kitts DD, Gilmour MW. Tolerance of *Listeria monocytogenes* to quaternary ammonium sanitizers is mediated by a novel efflux pump encoded by *emrE*. *Applied and Environmental Microbiology*. 2016;**82**:939-953. DOI: 10.1128/AEM.03741-15

[64] Franz E, Gras LM, Dallman T. Significance of whole genome sequencing for surveillance, source attribution and microbial risk assessment of foodborne pathogens. *Current Opinion in Food Science*. 2016;**8**:74-79. DOI: 10.1016/j.cofs.2016.04.004

[65] Schjørring S, Lassen SG, Jensen T, Moura A, Kjeldgaard JS, Müller L, et al. Cross-border outbreak of listeriosis caused by cold-smoked salmon, revealed by integrated surveillance and whole genome sequencing (WGS), Denmark and France, 2015 to 2017. *Eurosurveillance*. 2017;**22**:17-00762. DOI: 10.2807/1560-7917.ES.2017.22.50.17-00762

[66] Lienau EK, Strain E, Wang C, Zheng J, Ottesen AR, Keys CE, et al. Identification of a salmonellosis outbreak by means of molecular sequencing. *The New England Journal of Medicine*. 2011;**364**:981-982. DOI: 10.1056/NEJMc1100443

[67] Rantsiou K, Kathariou S, Winkler A, Skandamis P, Saint-Cyr MJ, Rouzeau-Szynalski K, et al. Next generation microbiological risk assessment: Opportunities of whole genome sequencing (WGS) for foodborne pathogen surveillance, source tracking and risk assessment. *International Journal of Food Microbiology*. 2018;**287**:3-9. DOI: 10.1016/j.ijfoodmicro.2017.11.007

[68] National Research Council (US) Committee on DNA Forensic Science: An Update. *The Evaluation of Forensic DNA Evidence.5. Statistical Issues*. Washington DC: National Academies Press (US); 1996. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK232615/> [Accessed February 24 2019]

[69] Wilson MR, Allard MW, Brown EW. The forensic analysis of foodborne bacterial pathogens in the age of whole-genome sequencing. *Cladistics*. 2013;**29**:449-461. DOI: 10.1111/cla.12012

[70] Rosner BM, Schielke A, Didelot X, Kops F, Breidenbach J, Willrich N, et al. A combined case-control and molecular source attribution study of human *Campylobacter* infections

in Germany, 2011-2014. Scientific Reports. 2017;7:5139. DOI: 10.1038/s41598-017-05227-x

[71] Pires SM, Vieira AR, Hald T, Cole D. Source attribution of human salmonellosis: An overview of methods and estimates. Foodborne Pathogens and Disease. 2014;11:667-676. DOI: 10.1089/fpd.2014.1744

[72] Yachison CA, Yoshida C, Robertson J, Nash JHE, Kruczkiewicz P, Taboada EN, et al. The validation and implications of using whole genome sequencing as a replacement for traditional serotyping for a National *Salmonella* Reference Laboratory. Frontiers in Microbiology. 2017;8:1044. DOI: 10.3389/fmicb.2017.01044

[73] Mellor GE, Fegan N, Gobius KS, Smith HV, Jennison AV, D'Astek BA, et al. Geographically distinct *Escherichia coli* O157 isolates differ by lineage, Shiga toxin genotype, and total Shiga toxin production. Journal of Clinical Microbiology. 2015;53:579-586. DOI: 10.1128/JCM.01532-14

[74] Hoffmann M, Luo Y, Monday SR, Gonzalez-Escalona N, Ottesen AR, Muruvanda T, et al. Tracing origins of the *Salmonella* Bareilly Strain causing a food-borne outbreak in the United States. The Journal of Infectious Diseases. 2016;213:502-508. DOI: 10.1093/infdis/jiv297

[75] Thépault A, Méric G, Rivoal K, Pascoe B, Mageiros L, Touzain F, et al. Genome-wide identification of host-segregating epidemiological markers for source attribution in *Campylobacter jejuni*. Applied and Environmental Microbiology. 2017;83. DOI: 10.1128/AEM.03085-16

[76] Cocolin L, Membré J-M, Zwietering MH. Editorial: Integration of omics into MRA. International Journal of Food Microbiology. 2018;287:1-2. DOI: 10.1016/j.ijfoodmicro.2017.11.006

[77] Farber JM, Peterkin PI. *Listeria monocytogenes*, a food-borne pathogen. Microbiological Reviews. 1991;55:476-511

[78] de Boer RF, Ferdous M, Ott A, Scheper HR, Wisselink GJ, Heck ME, et al. Assessing the public health risk of Shiga toxin-producing *Escherichia coli* by use of a rapid diagnostic screening algorithm. Journal of Clinical Microbiology. 2015;53:1588-1598. DOI: 10.1128/JCM.03590-14

[79] Ashton PM, Perry N, Ellis R, Petrovska L, Wain J, Grant KA, et al. Insight into Shiga toxin genes encoded by *Escherichia coli* O157 from whole genome sequencing. PeerJ. 2015;3:e739. DOI: 10.7717/peerj.739

[80] Center for Genomic Epidemiology. n.d. Available from: <http://www.genomicepidemiology.org/> [Accessed: February 20, 2019]

[81] Dutil L, Irwin R, Finley R, Ng LK, Avery B, Boerlin P, et al. Ceftiofur resistance in *Salmonella enterica* serovar Heidelberg from chicken meat and humans, Canada. Emerging Infectious Diseases. 2010;16:48-54. DOI: 10.3201/eid1601.090729

[82] Tang KL, Caffrey NP, Nóbrega DB, Cork SC, Ronksley PE, Barkema HW, et al. Restricting the use of antibiotics in food-producing animals and its associations with antibiotic resistance in food-producing animals and human beings: A systematic review and meta-analysis. Lancet Planetary Health. 2017;1:e316-e327. DOI: 10.1016/S2542-5196(17)30141-9

[83] Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, et al. Identification of acquired antimicrobial resistance genes. The Journal of Antimicrobial Chemotherapy. 2012;67:2640-2644. DOI: 10.1093/jac/dks261

- [84] Rowe W, Baker KS, Verner-Jeffreys D, Baker-Austin C, Ryan JJ, Maskell D, et al. Search engine for antimicrobial resistance: A cloud compatible pipeline and web interface for rapidly detecting antimicrobial resistance genes directly from sequence data. *PLoS One*. 2015;**10**:e0133492. DOI: 10.1371/journal.pone.0133492
- [85] McArthur AG, Wagglechner N, Nizam F, Yan A, Azad MA, Baylay AJ, et al. The comprehensive antibiotic resistance database. *Antimicrobial Agents and Chemotherapy*. 2013;**57**:3348-3357. DOI: 10.1128/AAC.00419-13
- [86] Knowles M, Stinson S, Lambert D, Carrillo C, Koziol A, Gauthier M, et al. Genomic tools for customized recovery and detection of foodborne Shiga toxigenic *Escherichia coli*. *Journal of Food Protection*. 2016;**79**: 2066-2077. DOI: 10.4315/0362-028X.JFP-16-220
- [87] Liu B, Pop M. ARDB-antibiotic resistance genes database. *Nucleic Acids Research*. 2009;**37**:D443-D447. DOI: 10.1093/nar/gkn656
- [88] Gupta SK, Padmanabhan BR, Diene SM, Lopez-Rojas R, Kempf M, Landraud L, et al. ARG-ANNOT, a new bioinformatic tool to discover antibiotic resistance genes in bacterial genomes. *Antimicrobial Agents and Chemotherapy*. 2014;**58**:212-220. DOI: 10.1128/AAC.01310-13
- [89] Tyson GH, McDermott PF, Li C, Chen Y, Tadesse DA, Mukherjee S, et al. WGS accurately predicts antimicrobial resistance in *Escherichia coli*. *The Journal of Antimicrobial Chemotherapy*. 2015;**70**:2763-2769. DOI: 10.1093/jac/dkv186
- [90] McDermott PF, Tyson GH, Kabera C, Chen Y, Li C, Folster JP, et al. Whole-genome sequencing for detecting antimicrobial resistance in nontyphoidal *Salmonella*. *Antimicrobial Agents and Chemotherapy*. 2016;**60**:5515-5520. DOI: 10.1128/AAC.01030-16
- [91] Zhao S, Tyson GH, Chen Y, Li C, Mukherjee S, Young S, et al. Whole genome sequencing analysis accurately predicts antimicrobial resistance phenotypes in *Campylobacter*. *Applied and Environmental Microbiology*. 2015;**82**:459-466. DOI:10.1128/AEM.02873-15
- [92] Gill A, Huszczyński G, Gauthier M, Blais B. Evaluation of eight agar media for the isolation of Shiga toxin-producing *Escherichia coli*. *Journal of Microbiological Methods*. 2014;**96**:6-11. DOI: 10.1016/j.mimet.2013.10.022
- [93] Gill A, Martinez-Perez A, McIlwham S, Blais B. Development of a method for the detection of Verotoxin-producing *Escherichia coli* in food. *Journal of Food Protection*. 2012;**75**: 827-837. DOI: 10.4315/0362-028X.JFP-11-395
- [94] Vimont A, Delignette-Muller M-L, Vernozy-Rozand C. Supplementation of enrichment broths by novobiocin for detecting Shiga toxin-producing *Escherichia coli* from food: A controversial use. *Letters in Applied Microbiology*. 2007;**44**:326-331. DOI: 10.1111/j.1472-765X.2006.02059.x
- [95] Blais BW, Tapp K, Dixon M, Carrillo CD. Genomically informed strain-specific recovery of Shiga toxin-producing *Escherichia coli* during foodborne illness outbreak investigations. *Journal of Food Protection*. 2019;**82**:39-44. DOI: 10.4315/0362-028X.JFP-18-340
- [96] Knowles M, Lambert D, Huszczyński G, Gauthier M, Blais BW. PCR for the specific selection of an *Escherichia coli* O157:H7 laboratory control strain. *Journal of Food*

Protection. 2015;78:1738-1744. DOI: 10.4315/0362-028X.JFP-15-147

[97] Margot H, Tasara T, Zwietering MH, Joosten H, Stephan R. Effects of different media on the enrichment of low numbers of Shiga toxin-producing *Escherichia coli* in mung bean sprouts and on the development of the sprout microbiome. International Journal of Food Microbiology. 2016;232:26-34. DOI: 10.1016/j.ijfoodmicro.2016.05.005

[98] Jarvis KG, Daquigan N, White JR, Morin PM, Howard LM, Manetas JE, et al. Microbiomes associated with foods from plant and animal sources. Frontiers in Microbiology. 2018;9:2540. DOI: 10.3389/fmicb.2018.02540

[99] Ottesen A, Ramachandran P, Reed E, White JR, Hasan N, Subramanian P, et al. Enrichment dynamics of *Listeria monocytogenes* and the associated microbiome from naturally contaminated ice cream linked to a listeriosis outbreak. BMC Microbiology. 2016;16:275. DOI: 10.1186/s12866-016-0894-1

[100] Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, et al. QIIME allows analysis of high-throughput community sequencing data. Nature Methods. 2010;7:335-336. DOI: 10.1038/nmeth.f.303

[101] Schloss PD, Westcott SL, Ryabin T, Hall JR, Hartmann M, Hollister EB, et al. Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. Applied and Environmental Microbiology. 2009;75:7537-7541. DOI: 10.1128/AEM.01541-09

[102] Morschhäuser J, Köhler G, Ziebuhr W, Blum-Oehler G, Dobrindt U, Hacker J. Evolution of microbial pathogens. Philosophical Transactions of the Royal Society of London. Series B, Biological

Sciences. 2000;355:695-704. DOI: 10.1098/rstb.2000.0609

[103] Iraola G, Pérez R, Naya H, Paolicchi F, Pastor E, Valenzuela S, et al. Genomic evidence for the emergence and evolution of pathogenicity and niche preferences in the genus *Campylobacter*. Genome Biology and Evolution. 2014;6:2392-2405. DOI: 10.1093/gbe/evu195

[104] Wirth T, Falush D, Lan R, Colles F, Mensa P, Wieler LH, et al. Sex and virulence in *Escherichia coli*: An evolutionary perspective. Molecular Microbiology. 2006;60:1136-1151. DOI: 10.1111/j.1365-2958.2006.05172.x

[105] Lacher DW. The evolutionary model of *Escherichia coli* O157:H7. In: Walk S, Feng P, editors. Population Genetics of Bacteria. Washington, DC: ASM Press; 2011:227-239. DOI: 10.1128/9781555817114.ch13

[106] Kelly BG, Vespermann A, Bolton DJ. Horizontal gene transfer of virulence determinants in selected bacterial foodborne pathogens. Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association. 2009;47:969-977. DOI: 10.1016/j.fct.2008.02.007

[107] Gilmour MW, Tabor H, Wang G, Clark CG, Tracz DM, Olson AB, et al. Isolation and genetic characterization of a coinfection of non-O157 Shiga toxin-producing *Escherichia coli*. Journal of Clinical Microbiology. 2007;45:3771-3773. DOI: 10.1128/JCM.01125-07

[108] Oniciuc EA, Likotrafiti E, Alvarez-Molina A, Prieto M, Santos JA, Alvarez-Ordóñez A. The present and future of whole genome sequencing (WGS) and whole metagenome sequencing (WMS) for surveillance of antimicrobial resistant microorganisms and antimicrobial resistance genes across the food chain. Genes. 2018;9:268. DOI: 10.3390/genes9050268

[109] Auffret MD, Dewhurst RJ, Duthie C-A, Rooke JA, John Wallace R, Freeman TC, et al. The rumen microbiome as a reservoir of antimicrobial resistance and pathogenicity genes is directly affected by diet in beef cattle. *Microbiome*. 2017;5:159. DOI: 10.1186/s40168-017-0378-z

IntechOpen

IntechOpen