

We are IntechOpen, the world's leading publisher of Open Access books Built by scientists, for scientists

4,800

Open access books available

122,000

International authors and editors

135M

Downloads

Our authors are among the

154

Countries delivered to

TOP 1%

most cited scientists

12.2%

Contributors from top 500 universities



WEB OF SCIENCE™

Selection of our books indexed in the Book Citation Index
in Web of Science™ Core Collection (BKCI)

Interested in publishing with us?
Contact book.department@intechopen.com

Numbers displayed above are based on latest data collected.
For more information visit www.intechopen.com



Biotransformation of Steroids Using Different Microorganisms

Arturo Cano-Flores, Javier Gómez and Rigoberto Ramos

Abstract

The introduction of a hydroxyl group “biohydroxylation” in the steroid skeleton is an important step in the synthesis of new steroids used physiologically as hormones and active drugs. There are currently about 300 known steroid drugs whose production constitutes the second category within the pharmaceutical market after antibiotics. Several biotransformations at industrial scale have been applied in the production of steroid hormones and drugs, which have functionalized different types of raw materials by means of *chemo-*, *regio-*, and *stereoselective* reactions (hydroxylation, Baeyer-Villiger oxidation, oxidation reactions, reduction of group carbonyl, isomerization, and Michael additions, condensation reactions, among others). In Green Chemistry, biotransformations are an important chemical methodology toward more sustainable industrial processes.

Keywords: biotransformation, steroid compounds, biological transformation, bioconversions, microorganisms

1. Introduction

Steroids (stereos = solids) are organic compounds derived from alcohols, which are widely distributed in the animal and plant kingdoms. Their base skeleton has 17 carbon atoms in a tetracyclic ring system known as cyclopentanoperhydrophenanthrenes (gonane and estrane). In this group of substances, life-vital compounds are categorized, such as cholesterol, bile acids, sex hormones, vitamin D, corticosteroids, cardiac aglycones, and antibiotics, among others.

Some of the most potent toxins are steroid alkaloids. Steroids are responsible for important biological functions in the cell; for example, the steroids derived from androstane, pregnane, and estrane have hormonal activity [1–5]; bile acids are important for the digestion and absorption of fats; and cardiotonic aglycones are used for the treatment of heart disease. Sterols are constituents of the cell membrane, essential for cell stability and development; also, they are precursors of bile acids and steroid hormones.

A large number of steroids are used as anti-inflammatory agents [6], immunosuppressants, progestational agents, diuretics, anabolics, and contraceptives [7–9]. Some are used for the treatment of prostate and breast cancer [10, 11], for adrenal insufficiency [12], for prevention of heart disease [13], as antifungal agents [14], and as active ingredients used for the treatment of obesity [15] and AIDS [16]. Recently, the antiviral activity against the herpes simplex virus type I of some steroid glycosides was determined [17].

The therapeutic action of some steroid hormones has been associated with their interaction with intracellular receptors, which act as transcription factors in the regulation of gene expression [18]. It has been reported that some steroids, such as dehydroepiandrosterone (DHEA), progesterone, pregnenolone and its sulfated derivatives [19, 20], as well as, 17β -estradiol, allopregnanolone and its synthetic derivatives (afoxolaner and ganaxolone) are considered neurosteroids, due to their action at the level of the CNS [19].

The physiological activity of steroids depends on their structure, the type, number, spatial orientation, and reactivity of the different functional groups present in the tetracyclic core as well as the oxidation state of the rings. For example, the presence of an oxygenated function in C-11 β is crucial for the anti-inflammatory activity; the hydroxyl function in C-17 β determines androgenic properties; the aromatization of ring A confers estrogenic effect; and corticosteroids have the 3-keto-4-ene group and the pregnane side chain at C-17 [21, 22].

Currently, about 300 steroid drugs are known, and this number tends to grow. Their production represents the second category in the pharmaceutical market after antibiotics [24, 25]. Nowadays, steroids represent one of the largest sectors in pharmaceutical industry with world markets in the region of US\$ 10 billion and the production exceeding 1,000,000 tons per year [23].

The production of steroid drugs and hormones is one of the best examples of the applications that biotransformations have on an industrial scale [3, 21]. Microbiological transformations are an effective tool for the preparation of various compounds [26], which can be difficult to obtain by conventional chemical methods and have been widely used in the bioconversion of steroids [25]. In 1950, the pharmacological effects of cortisol and progesterone were reported, in addition to the hydroxylation of the latter in C-11 α using *Rhizopus* species. This began a very important stage in the development of the synthesis of steroids with biological activity [4, 5].

Currently, a great versatility of microbial systems in the pharmaceutical industry for the commercial production of steroids and other drugs is recognized [27, 28]. Several hundreds of microbiological transformations of steroids have been reported in the literature; also, many bioconversions have been incorporated into numerous partial syntheses of new compounds for their evaluation such as hormones or drugs [21, 29–32]. Chemical derivatives of some steroids are reported to have better therapeutic advantages than the starting materials.

However, the main objectives in the research and development of the steroid drug industry currently consist of the detection and isolation of microbial strains with novel activity or more efficient transformation capacity, where genetic engineering and metabolic engineering can play a prominent role in the metabolism of bacteria, fungi, and plants [33–36].

The aim of the present review is to emphasize the importance of biotransformation using microorganisms to obtain steroid compounds with pharmaceutical interest, as a chemical-biological strategy that alternates with the chemical synthesis, and to highlight the chemical reaction made by different types of microorganisms in the functionalization of the steroid skeleton.

2. Microbiological transformations of steroids

In Green Chemistry, biotransformations constitute an important methodology in organic chemistry [37]. The microbiological transformations of steroids have been an essential chemical tool used for the preparation of many intermediaries and in the generation of new drugs, where chemical functionalization-hydroxylation, Baeyer-Villiger oxidation, reduction, isomerization, Michael additions, and condensation

reactions can be carried out in different positions of the steroid skeleton in *chemo*-, *regio*-, and *stereoselective* ways, being very complicated or even impossible by the classic chemical methods. Currently, any stereogenic center of the steroid skeleton can be specifically hydroxylated stereoselectively. Nowadays, *biohydroxylations* in C-11 α , 11 β , 15 α , and 16 α are industrially carried out via a microbial hydroxylation with good yields and enantiomeric excess (ee). Below are some of the microbiological transformations performed on different natural and synthetic steroids [25].

In the literature, it is the well-documented *regio*- and *stereoselective* hydroxylation in C-14 with α orientation in progesterone (**1**) and other steroids by well-functioning fungi, such as *Thamnostylum piriforme* (ATCC 8992), *Mucor griseocyanus* (ATCC 1207a), *Actinomucor elegans* (MMP 3132), and *Zygodessmus* sp. (ATCC 14716).

From the incubation of **1** with *T. piriforme*, 14 α -hydroxyprogesterone (**2**, 32%) and 9 α -hydroxyprogesterone (**3**, 1.4%) were obtained; whereas in the incubation of **1** with *M. griseocyanus*, **2** (13.4%), 7 α ,14 α -dihydroxyprogesterone (**4**, 6.5%) and 6 β ,14 α -dihydroxyprogesterone (**5**, 2.8%) were obtained. In the biotransformation of **1** using *A. fumigatus* after 24 h of incubation, different mono- and dihydroxylated products were obtained: 11 α -hydroxyprogesterone (**6**, 33%), 11 α ,15 β -dihydroxyprogesterone (**7**, 17%), 7 β ,15 β -dihydroxyprogesterone (**8**, 14%), 15 β -hydroxyprogesterone (**9**), 7 β -hydroxyprogesterone (**10**), where **9** and **10** were detected in minimal quantity. Finally, at 72 h, the main products were **7** (48%) and **8** (25%), with the positions 11 α and 15 β being hydroxylated more easily than the position 7 β in **1** [38, 39].

In the incubation of **1** with *Saprolegnia hypogyna*, 4-androstene-3,17-dione (**11**), testosterone (**12**), and testolactone (**13**) were obtained [40]. The compounds **13** (98%) were also obtained from the bioconversion of **1** using *A. sojae* (PTCC 5196). The biotransformation pathway indicating the presence of Baeyer-Villiger monooxygenase (BVMO) can carry out both oxygenative esterification of 20-ketosteroids and oxygenative lactonization of 17-ketosteroids [41]. The compounds 15 α -hydroxyprogesterone (**14**, 47%) and 12 β ,15 α -dihydroxyprogesterone (**15**, 25%) were isolated in the biotransformation of **1** using *Fusarium culmorum* [42]. In the biotransformation of **1** using the bacterium, thermophilic *Bacillus stearothermophilus*, four products of monohydroxylation, 20 α -hydroxyprogesterone (**16**, 61%), 6 β -hydroxyprogesterone (**17**, 21%) and 6 α -hydroxyprogesterone (**18**, 14%), and 9,10-seco-pregnen-3,9,20-trione (**19**, 4%), were isolated [43].

An efficient *regio*- and stereoselectivity was observed in the biotransformation of **1** on a large scale by the system *Mucor* 881 (M881) to give the hydroxylated derivatives **6**, 6 β ,11 α -dihydroxyprogesterone (**20**), and 6 β -hydroxypregn-4-ene-3,11,20-trione (**21**). In the literature, it is described that species of the genus *Mucor* and *Rhizopus* can hydroxylate said positions but with lower yields. The fungal system M881 showed the ability to carry out hydroxylation at 6 β and 11 α positions of 4-ene-3-one steroids (**1**, **11**, **12** and **211**) [44].

Recently, it was reported that in the biotransformation of **1** using *Penicillium aurantiogriseum* for 10 days, **11** and androsta-1,4-dien-3,17-dione (**22**) were obtained. These products were observed in the biotransformation of **1** using *Bacillus sphaericus*; the hydroxylation in C-17 was mainly observed [45, 46]. Biotransformation of **1** using *Geobacillus gargensis* (DSM 15378) has resulted in the production of secoderivatives: **19** and **23** (9,10-seco-4-pregnene-20 α -hydroxy-3,9-dione), which are produced by the rupture of the ring B of **1** (**Figure 1**) [47]. Secosteroids are an important group, which exhibits a variety of different biological activities [48, 49].

In the biotransformation of 5 β -dihydroprogesterone (**24**) using *T. piriformis*, 14 α -hydroxy-5 β -pregnan-3,20-dione (**25**, 11.8%), 3 β ,14 α -dihydroxy-5 β -pregnan-20-one (**26**, 0.5%), and 14 α ,15 β -dihydroxy-5 β -pregnan-3,20-dione (**27**, 0.4%) were characterized, while in the biotransformation of 3 β -hydroxy-5 β -pregnan-20-one (**28**), **26**




Figure 1.
 Biotransformation products of progesterone (1).

(0.6%) and $3\beta,9\alpha,14\alpha$ -trihydroxy- 5β -pregnan-20-one (29, 16%) were isolated, after being incubated for 96 h. The microbiological transformation of 28 using *Actinomucor elegans* produced the compounds 25 and 28 in lower yield than *T. piriforme* and a minor product identified as $3\beta,9\alpha$ -dihydroxy- 5α -pregnan-20-one (30) (Figure 2) [38].

The biotransformation of 16-dehydropregesterone (4,16-pregnadien-3,20-dione, 31) using *Mucor piriformis* has been reported to give different hydroxylation products: 14α -hydroxypregna-4,16-dien-3,20-dione (32, 1%), $7\alpha,14\alpha$ -dihydroxypregna-4,16-dien-3,20-dione (33, 78%), $3\beta,7\alpha,14\alpha$ -trihydroxy- 5α -pregna-16-en-20-one (34, 3%), and $3\alpha,7\alpha,14\alpha$ -trihydroxy- 5α -pregna-16-en-20-one (35, 2%); while the microsomes




Figure 2.
 Biotransformation products of 5β -dihydroprogesterone (24).

prepared from **31** transformed the hydroxylate to 14α -hydroxy derivative (**32**). Incubation of **32** with *M. piriformis* resulted in the formation of **33–35** (**Figure 3**) [50].

In contrast, in the biotransformation of 17α -hydroxyprogesterone (**36**) using *M. piriformis*, after 48 h of incubation, four compounds were obtained: $17\alpha,20\alpha$ -dihydroxypregn-4-en-3-one (**37**, 19%), $7\alpha,17\alpha$ -dihydroxypregn-4-en-3,20-dione (**38**, 25%), $6\beta,17\alpha,20\alpha$ -trihydroxy-pregn-4-en-3-one (**39**, 18%), and $11\alpha,17\alpha,20\alpha$ -trihydroxypregn-4-en-3-one (**40**, 25%); it was observed that *M. piriformis* was able to hydroxylate the C-6, C-7, C-11, and C-14 positions stereospecifically, in addition to reducing the 4-en-3-one system in ring A and the keto group of C-20 (**Figure 4**) [50]. The biotransformation of **36** using *Fusarium culmorum* led to the formation of **14** (47%) and **15** (25%) [42].

Pregnenolone (3β -hydroxypregn-5-en-20-one, **41**), the precursor of many steroid hormones, was biotransformed by *Mucor piriformis* to obtain two metabolites, $3\beta,7\alpha$ -dihydroxypregn-5-en-20-one (**42**) and $3\beta,7\alpha,11\alpha$ -trihydroxypregn-5-en-20-one (**43**) [51], where **43** (46.4%) was also a bioconversion product of **41** using *Mucor circinelloides* var. *lusitanicus* [52]. Two metabolites of pregnenolone (**41**) obtained from biotransformation of *B. cinereae* were characterized as $3\beta,11\alpha,16\beta$ -trihydroxypregn-5-en-20-one (**44**, 39%) and $11\alpha,16\beta$ -dihydroxypregn-4-en-3,20-dione (**45**, 6%). The formation of the hydroxylation products in C-11 and C-16 by *B. cinereae* can be determined by the presence of the acetyl group in C-20 [53]. The biotransformation of **41** using different microorganisms (*Cunninghamella elegans*, *R. stolonifer*, and *G. fujikuroi*) was reported by Choudhary et al. [54]. Incubation of **41** with *C. elegans* produced $3\beta,7\beta,11\alpha$ -trihydroxypregn-5-en-20-one (**46**, 28%), $3\beta,6\alpha,11\alpha,12\beta,15\beta$ -pentahydroxypregn-4-en-20-one (**47**, 4%), and $3\beta,6\beta,11\alpha$ -trihydroxypregn-4-en-20-one (**48**, 2%), while incubation with *G. fujikuroi*, two products $3\beta,7\beta$ -dihydroxypregn-5-en-20-one (**49**, 3%) and $6\beta,15\beta$ -dihydroxypregn-4-en-3,20-dione (**50**, 2%) were obtained. In the microbiological transformation of **41** using different *Bacillus* strains, **42**, **49**, and 7-oxo-pregnenolone (**51**) were the major products obtained [55], while by using *Fusarium oxysporum* var. *cubense*, **42** was the only product obtained [56]. The biotransformation of pregnenolone acetate (**52**) using *C. elegans* generated **41**, **22**, $6\beta,15\beta$ -dihydroxyandrosta-4-en-3,17-dione (**53**), and $11\alpha,15\beta$ -dihydroxypregn-4-en-3,20-dione (**54**), while by using *R. stolonifer*, 11α -hydroxypregn-4-en-3,20-dione (**55**) and **53** were obtained (**Figure 5**) [54].

The microbiological transformation of the racemic mixture of 13-ethyl-17 β -hydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one (**56**) was tested with different fungi *Rhizopus nigricans*, *R. arrhizus*, *Aspergillus niger*, *A. ochraceus*, and *Curvularia lunata*. The bioconversion of the racemic mixture of **56** by *R. arrhizus* produced only one major product, (\pm)-13-ethyl-10 $\beta,17\beta$ -dihydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one




Figure 3.
Biotransformation products of 16-dehydroprogesterone (**31**).




Figure 4.
Biotransformation products of 17 α -hydroxyprogesterone (36).




Figure 5.
Biotransformation products of pregnenolone (41) and acetyl derivative (52).

(57, 28.4%), whereas *R. nigricans*, *A. niger*, and *C. lunata* biotransformed 56 to 57 more slowly and inefficiently [57].

The racemic mixture (\pm)-13-ethyl-7 β ,17 β -dihydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one (58, 4.3%) was obtained as product of incubating mixture 56 with *A. ochraceus*; none of the fungi tested were able to differentiate the two enantiomers of 56 in the course of the hydroxylation reaction; in addition, the absence of the hydroxylated derivative in C-11 is due to the presence of the ethyl group in C-13 or the ethynyl group in C-17 [57]. The microbiological transformation of the racemic mixture and the dextro enantiomer of compound 56 has been described using different species of *Cunninghamella* [58]. For example, the transformation of the racemic mixture of 56 by *C. blakesleeana* (AS 3.910) produced 57 (5.3%), 13-ethyl-6 β ,17 β -dihydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one (59, 3.6%), 13-ethyl-15 α ,17 β -dihydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one (60, 3.0%), and 13-ethyl-6 β ,10 β ,17 β -trihydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one (61, 3.6%), while by using *C. echinulata* (AS 3.1990), 61 (3.2%), 57 (1.2%), and enantiomer dextro of 58 (2.9%) were obtained. The transformation of the enantiomer dextro of 56 using *C. blakesleeana* produced 57 (1.2%), 58 (2.9%), and 61 (3.2%), by using *C. echinulata*, the same compounds were obtained but in lower yield. Therefore, the microbial transformation of the racemic mixture and the *d*-enantiomer of 56 using different *Cunninghamella* species gave poor yields and poor resolutions, which were obtained for the hydroxylation reaction (Figure 6) [58].




Figure 6.
Biotransformation products of (+)-13-ethyl-17 β -hydroxy-18, 19-dinor-17 α -pregn-4-en-20-yn-3-one (56).

The biotransformation of danazol (17 β -hydroxy-17 α -pregna-2,4-dien-20-yno-[2,3-d]-isoxazole, **62**), a heterocyclic steroid drug in which an isoxazole ring is fused with ring-A of a steroid nucleus, using *Fusarium lini*, *A. niger* and *Cephalosporium aphidicola* yielded 17 β -hydroxy-2-(hydroxymethyl)-17 α -pregn-4-en-20-yn-3-one (**63**) and 17 β -hydroxy-2-(hydroxymethyl)-17 α -pregn-1,4-dien-20-yn-3-one (**64**); while *Bacillus cereus* afforded **64**, as the only product [59]. Microbial transformation of danazol (**62**) using *C. blakesleeana* yielded four compounds: 14 β ,17 β -dihydroxy-2-(hydroxymethyl)-17 α -pregn-4-en-20-yn-3-one (**65**, 1.2%), 1 α ,17 β -dihydroxy-17 α -pregna-2,4-dien-20-yno-[2,3-d]-isoxazole (**66**, 1.2%), and 6 β ,7 β -dihydroxy-17 α -pregna-2,4-dien-20-yno-[2,3-d]-isoxazole (**67**, 0.8%) and **64** (1.2%). This involves hydroxylations at C-1, C-6 and C-15, whereas oxidation at C-3, and N-O bond cleavage has also occurred (**Figure 7**) [60].

Norethisterone (17 α -ethynyl-19-nortesterone, **68**) is a potent progestin used as a contraceptive agent; its biotransformation with *Cephalosporium aphidicola* (IMI 68689) produced the aromatization of ring A that yielded 17 α -ethynylestradiol (**69**), whereas **69** was biotransformed by *Cunninghamella elegans* (NRRL 1392) producing the compounds 19-nor-17 α -pregna-1,3,5(10)-trien-20-yn-3,4,17 β -triol (**70**), 19-nor-17 α -pregna-1,3,5(10)-trien-20-yn-3,7 α ,17 β -triol (**71**), 19-nor-17 α -pregna-1,3,5(10)-trien-20-yn-3,11 α ,17 β -triol (**72**), 19-nor-17 α -pregna-1,3,5(10)-trien-20-yn-3,6 β ,17 β -triol (**73**), and 19-nor-17 α -pregna-1,3,5(10)-trien-20-yn-3,17 β -diol-6 β -methoxy (**74**) (**Figure 8**) [61].

Mestranol (**75**) and 17 β -methoxymestranol (**76**) are the mono- and dialkylated derivatives of **69**, respectively. In incubating **75** with *C. elegans*, two hydroxylated compounds were obtained: 6 β -hydroxymestranol (**77**, 2.8%) and 6 β ,12 β -dihydroxymestranol (**78**, 3.6%), inferring that the presence of the methoxyl group in C-3 reduces the number of biotransformation products and introduces hydroxyl groups in C-6 and C-12 with β orientation, while **76** was not biotransformed due to the presence of the methoxyl group in C-17 (**Figure 9**) [62].

Microbial transformation of 6-dehydropregesterone (**79**) using *A. niger* yielded five metabolites: 6 β -chloro-7 α ,11 α -dihydroxypregna-4-en-3,20-dione (**80**, 1.0%), 7 α -chloro-6 β ,11 α -dihydroxypregna-4-en-3,20-dione (**81**, 1.33%), 6 α ,7 α ,-epoxy-11 α -hydroxypregna-4-en-3,20-dione (**82**, 1.33%), 6 α ,7 α ,-epoxy-pregna-4-en-3,20-dione (**83**, 2.0%), and 11 α -hydroxypregna-4,6-dien-3,20-dione (**84**, 2.33%). Compound 11 α -hydroxyandrost-4,6-dien-3-one (**85**, 15.4%) was obtained through whole cell biotransformation of **79** by *G. fujikuroi* (ATCC 10704). The formation of **80** and **81** is an interesting finding. This route provides an efficient method for the obtention of chlorohydrins from alkene functionality [63]. The compound **84** was obtained through the microbial transformation of **79** using *R. nigricans* [64], *Nigrospora sphaerica*, *Mucor racemosus*, and *Botryosphaeria obtusa*. 6-dehydropregesterone (**79**)




Figure 7.
 Biotransformation products of danazol (62)




Figure 8.
 Biotransformation products of norethisterone (68) and 17 α -ethinylestradiol (69).




Figure 9.
 Biotransformation of products of mestranol (75).

is a synthetic derivate of progesterone. *Botryodiplodia theobromae* was used for the synthesis of 6-DPH from progesterone (**Figure 10**) [65].

Incubation of melengestrol acetate (**86**) with *C. blakesleeana*, which provides an route for the monohydroxylation of the (**86**) at C-11, yielded a 17 α -acetoxy-11 β -hydroxy-6-methylenepregna-4,6-diene-3,20-dione (**87**) (**Figure 11**) [66].

Biotransformation of 3 β -hydroxy-17 β -carboxyethyl-5 β -androsthenol (**88**) using *T. pyriformis* resulted in the mixture of




Figure 10.
Biotransformation products of 6-dehydropregesterone (79).




Figure 11.
Biotransformation products of melengestrol acetate (86).




Figure 12.
 Biotransformation products of 3 β -17 β -carboxyethyl-5 β -androsteno (88).

3 β ,14 α -dihydroxy-17 β -carboxyethyl-5 β -androsteno (89, 9%) with 9 α ,14 α -dihydroxy derivative (90, 12%) and two minor products 14 α ,15 α -dihydroxy (91) and 15 β -hydroxy (92). Compound 92 was identified as a product of biotransformation using *A. elegans*, *M. griseocyamus*, and *Zygodemus sp.* (Figure 12) [38].

Androst-4-en-3,17-dione (11), which plays an important role in the metabolism of drugs, among many other functions, was biotransformed using *M. piriformis* to give one main product, 6 β -hydroxyandrost-4-en-3,17-dione (93, 13%), and four minor products, 14 α -hydroxyandrost-4-en-3,17-dione (94, 2%), 7 α -hydroxyandrost-4-en-3,17-dione (95, 2%), testosterone (12, 3%), and 6 β -hydroxytestosterone (96, 1%). In the biotransformation of 11 using *M. griseocyamus* 94 (9%), 95 (4%) and 14 α -hydroxytestosterone (97, 9%) were the major products obtained; likewise, 11 and 93 were identified in the mixture of biotransformation products [67]. From the incubation of 11 with *M. piriformis*, 94–97 and 7 α ,14 α -dihydroxytestosterone (98) were obtained [38]. Hydroxylated steroids in C-9 are important intermediaries in the synthesis of highly effective anti-inflammatory drugs. The microbiological transformation of 11 to 9 α -hydroxyandrost-4-en-3,17-dione (99) was studied using *Rhodococcus sp.* in a low-nutrient culture medium at a fixed pH (Figure 13) [68]. When 11 was incubated with *Bacillus* strain HA-V6–3, the metabolites 12, 93–97,




Figure 13.
 Biotransformation products of androst-4-en-3,17-dione (11).

6 β ,14 α -dihydroxyandrost-4-en-3,17-dione (**100**), 11 α -hydroxyandrost-4-en-3,17-dione (**101**), androst-4-en-3,6,17-trione (**102**), and 5 α -androst-3,6,17-trione (**103**) were produced as described by Schaaaf and Dettner [69].

In the bioconversion of **11** using *C. aphidicola*, **93** and **94** were obtained [70], while in the fermentation of **11** using *Curvularia lunata*, the products **101** (4%), 17 β -hydroxyandrost-1,4-dien-3-one (**104**, 4.4%), androsta-1,4-dien-3,17-dione (**105**, 3%), 11 α ,17 β -dihydroxyandrost-4-en-3-one (**106**, 4%), and **107** (15 α -hydroxyandrost-1,4-dien-3,17-dione, 2.8%) were obtained (**Figure 13**) [71]. Biotransformation of **11** using *Beauveria bassiana* was studied in times and with culture media at different pH (pH 6 and 7) [72]. At pH 6, two products were obtained: **106** and 6 β ,11 α -dihydroxyandrost-4-en-3,17-dione (**108**), where the stereoselective hydroxylation was observed at C-11 α and C-6 β ; while at pH 7, the compounds **12**, **106**, 3 α ,11 α ,17 β -trihydroxy-5 α -androstane (**109**), and 6 β ,11 α ,17 β -trihydroxyandrost-4-en-3-one (**110**) were obtained. Products **93** (14%) and **94** (75%) were isolated from the biotransformation of **11** using *Chaetomium* sp. (**Figure 13**) [73].

Obtaining hydroxylated derivatives in a specific position is one of the objectives of the steroid industry; for example, 14 α -hydroxysteroids are shown to have anti-inflammatory, contraceptive, and antitumor activities. With the biotransformation of **11** and **105** using different strains of the fungus, *C. lunata* allowed in the case of **11**, the production of a major product, **94**; while with **105**, 14 α -hydroxyandrost-1,4-dien-3,17-dione (**111**, 70%) was obtained (**Figure 13**) [74].

Androsta-1,4-dien-3,17-dione (**105**) is a useful precursor in the chemical or microbiological preparation of other steroid hormones and pharmaceutical. Transformation of **105** by *Colletotrichum lini* (As3.486) produced the hydroxylated compounds at C-11 α and C-15 α : 15 α -hydroxyandrost-1,4-dien-3,17-dione (**107**), 11 α ,15 α -dihydroxyandrost-1,4-dien-3,17-dione (**112**), and 15 α ,17 β -dihydroxyandrost-1,4-dien-3-one (**113**) (**Figure 14**) [75].

Testosterone (**12**) was metabolized by *M. griseocyamus* and *T. piriforme*. In the biotransformation of **12** using *M. griseocyamus*, **97** (35%) and other products were obtained, where **94** was identified as the major product. Conversely, the microbiological transformation of **12** using *T. piriforme* produced **97** (10%), as the main product at 24 h; after 72 h of biotransformation, four products were obtained: **93** (13%), **96** (7%), **97** (13%), and **111** (5%). It was discovered that *T. piriforme* produced smaller quantity of 14 α -hydroxy derivatives (**Figure 15**) [38].

In the biotransformation of **12** using *Nectria haematococca*, four substances were isolated, whose performance was dependent on the incubation time; majority of the products were produced at 72 h. The hydroxylated derivatives in C-11 with α




Figure 14.
Biotransformation products of androsta-1,4-dien-3,17-dione (**105**).




Figure 15.
Biotransformation products of testosterone (**12**).

orientation and dehydrogenation in C₁-C₂ resulted in the following compounds: 11 α -hydroxyandrost-1,4-dien-3,17-dione (**114**, 8.0%), 11 α ,17 β -dihydroxyandrost-1,4-dien-3-one (**115**, 4.3%), **101** (1.9%), and **104** (2.3%) [76]. Incubation of **12** with *Fusarium culmorum* produced **93** (10%) and **96** (32%) with hydroxylated derivatives at C-6 β , including the products, 15 α ,17 β -dihydroxyandrost-4-en-3-one (**116**, 22%) and 15 α -hydroxyandrost-4-en-3,17-dione (**117**). Selective hydroxylation of **103** at C-6 with a β orientation and allylic position at the unsaturated 3-keto-system is favored by the system π and the presence of the hydroxyl group at C-17, while hydroxylation at C-15 is a very frequent process carried out by fungi of the genus *Fusarium* [42]. Metabolites **11**, **85**, **105**, and **115** were obtained as oxidation and hydroxylation products of **12** using the fungus *F. oxysporum* var. *cubense* [56]. The fungus, *Cephalosporium aphidicola*, was hydroxylated with **12** to give the products **96** (47%) and **97** (3%), with hydroxylated derivatives in C-6 β and C-14 α , respectively [70]. Incubation of **12** with *C. lunata* and *Pleurotus ostreatus* yielded compounds **11** (17%) and **115** (13%), respectively [77]. The phytopathogenic fungus, *Botrytis cinerea*, produced 7 β ,17 β -dihydroxyandrost-3-one (**118**, 73%), as the only biotransformation product of **12**. It seems that the presence of the hydroxyl group in C-17 in the androstane skeleton directed the hydroxylation at C-7 with a β orientation (**Figure 15**) [53].

In the biotransformation of **12** using *Bacillus stearothermophilus*, thermophilic bacterium, the major product obtained was **11** (90.2%); it was generated by the oxidation of C-17, and the hydroxylated derivatives of **11** in C-6 (**93**, C-6 β , 1.1%) and (**119**, C-6 α , 0.9%) include two monohydroxy derivatives of **12**, **96** (C-6 β , 3.9%) and **120** (C-6 α , 3.9%). This indicates that hydroxylation with α orientation in C-6 may be a common action of some thermophilic bacteria [78]. Biotransformation of **11** using *B. stearothermophilus* in the presence of hydrolase inducers—salicylic acid, chloramphenicol, cyclodextrin, dexamethasone, riboflavin, and rifampicin—resulted in obtaining a higher concentration of the compounds: 9,10-seco-4-androst-3,9,17-trione (**121**), 5 α -androst-3,6,17-trione (**103**), 17 β -hydroxy-5 α -androst-3,6-dione (**122**), 3 β ,17 β -dihydroxyandrost-4-en-6-one (**123**), and 17 β -hydroxyandrost-4,6-dien-3-one

(124). For example, the presence of glucose and cycloheximide favored the obtaining of 123, while the production of 124 was achieved in the presence of rifampicin [79]. The products isolated from the biotransformation of 12 using *Chaetomium sp.* were 93 (21%), 94 (39%), and 99 (19%); after 24 h of incubation, the presence of 11 was detected. Janeczko et al. [73] concluded that the steric factors associated with the substrate determine the location and orientation of the hydroxyl group. For example, the carbonyl group in C-17 at 11 directs the entry of the hydroxyl group at C-14 with α orientation, while the hydroxylation in C-6 β is favored by the presence of the hydroxyl group in C-17, as in 12. In the case of progesterone (1), which has an acyl group, dihydroxylated derivatives were observed in C-6 and C-14 (Figure 15) [73].

Incubation of 11 and 12 with *C. lini* ST-1 displayed different catalytic characteristics. Biotransformation of 11 afforded two products: 15 α -hydroxyandrost-4-en-3,17-dione (117, 5%) and 11 α ,15 α -dihydroxyandrost-4-en-3,17-dione (125, 64%), while 12 yielded 15 α -hydroxyandrost-4-en-3,17-dione (117, 60%). Incubation of 1 resulted in the isolation of 14. Wu et al. [80] concluded that the different hydroxylation sites between 11 and 12 suggested that the hydroxyl group or carbonyl group on the substrate at C-17 had influence on the location of introduced hydroxyl groups (Figure 15).

Dehydroepiandrosterone (3 β -hydroxyandrost-5-en-17-one, 126) endogenous prohormone secreted by the adrenal glands is a precursor of androgens and estrogens. Incubation with *M. piriformis* allowed the isolation of five compounds: 3 β ,17 β -dihydroxyandrost-5-ene (127), 3 β ,7 α -dihydroxyandrost-5-en-17-one (128), 3 β -hydroxyandrost-5-en-7,17-dione (129), 3 β ,17 β -dihydroxyandrost-5-en-7-one (130), and 3 β ,7 α ,17 β -trihydroxyandrost-5-ene (131). The action of the fungus was the stereospecific hydroxylated products at C-7 α (128 and 131) and the reduction of the carbonyl group at C-17 [51]. From the microbiological transformation of 126 using *Rhizopus stolonifer*, six products were isolated: 127 (20%), 128 (12%), 129 (20%), 3 β ,17 β -dihydroxyandrost-4-ene (132, 12%), 17 β -hydroxyandrost-4-en-3-one (133, 34%), and 3 β ,11 β -dihydroxyandrost-4-en-17-one (134, 15%) [81]. *Fusarium oxysporum* biotransformed to 126 in a mixture of four hydroxylated derivatives (127–129 and 130), which were characterized as their acetylated derivatives; the hydroxylation was favorably in C-7 stereospecifically (α orientation) in the 3 β -hydroxy- Δ^5 -steroids, while *Colletotrichum musae* biotransformed to 126–127 by reducing the carbonyl group in C-17 (Figure 16) [56].




Figure 16.
Biotransformation products of dehydroepiandrosterone (126).

In the biotransformation of **126** using *Penicillium griseopurpureum* and *P. glabrum*, the following was produced; hydroxylated derivatives in C-7 α (**95**), C-14 α (**94**) and C-15 α (**117**), with **11** being the main product. In addition, *P. griseopurpureum* generated products for the Baeyer Villiger oxidation to give the lactone D ring (testolactone, **13**) and its hydroxylated derivative at C-15 α (15 α -hydroxy-17 α -oxa-D-homo-androst-4-en-3,17-dione, **135**); while *P. glabrum* generated the compounds, 3 β -hydroxy-17 α -oxa-D-homo-androst-5-en-17-one (**136**) and 3 β -hydroxy-17 α -oxa-D-homo-5 α -androstan-17-one (**137**) (**Figure 16**) [82].

The biotransformation of 17 α -ethynyl-17 β -hydroxyandrost-4-en-3-one (ethisterone, **138**) and 17 α -ethyl-17 β -hydroxyandrost-4-en-3-one (**139**) was described using the fungi *Cephalosporium aphidicola* and *Cunninghamella elegans*. The bioconversion of **138** using *C. aphidicola* yielded 17 α -ethynyl-17 β -hydroxyandrost-1,4-dien-3-one (**140**, 5.5%), while by using *C. elegans*, 17 α -ethynyl-11 α ,17 β -dihydroxyandrost-4-en-3-one (**141**, 3.4%) was obtained. The biotransformation of **138** using *C. aphidicola* generated 17 α -ethyl-17 β -hydroxyandrost-1,4-dien-3-one (**142**, 2.2%). In contrast, when incubating **139** with *C. elegans*, two new products were obtained: 17 α -ethyl-11 α ,17 β -dihydroxyandrost-4-en-3-one (**143**, 2.8%) and 17 α -ethyl-6 α ,17 β -dihydroxy-5 α -androstan-3-one (**144**, 1.6%) (**Figure 17**) [83].

Adrenosterone (**145**) is an inhibitor of the enzyme estrogen synthetase responsible for the formation of estrogen, and it has a great clinical application. Biotransformation of **145** using *C. aphidicola* produced androst-1,4-dien-3,11,17-trione (**146**, 3%), 17 β -hydroxyandrost-4-en-3,11-dione (**147**, 2%), and 17 β -hydroxyandrost-1,4-dien-3,11-dione (**148**, 17%). **145** (11.2%) and **12** (8.1%) were obtained from the biotransformation of **145** using *Fusarium lini*, while **147** (36.8%) was obtained from the biotransformation of **145** using *Trichothecium roseum* (**Figure 18**) [84].

The biotransformation of mesterolone (1 α -methyl-17 β -hydroxy-5 α -androst-3-one, **149**), a synthetic androgenic steroid, was performed using different fungi as described by Choudhary et al. [85]. From the biotransformation of **149** using *C. aphidicola*, the compounds 1 α -methyl-5 α -androst-3,17-dione (**150**), 1 α -methyl-5 α -androst-3,17-diol (**151**), and 1 α -methyl-15 α -hydroxy-5 α -androst-3,17-dione (**152**) were obtained. Incubation of **149** with *Fusarium lini* produced the compounds **152**, 1-methyl-5 α -androst-1-en-3,17-dione (**153**), 1 α -methyl-6 α ,17 β -dihydroxy-5 α -androst-3-one (**154**), 1 α -methyl-15 α ,17 β -dihydroxy-5 α -androst-3-one (**155**), and 1-methyl-15 α ,17 β -dihydroxy-5 α -androst-1-en-3-one (**156**). The products obtained from the biotransformation of **149** using *R. stolonifer* were **150**, **154**, **156**, 1 α -methyl-7 α ,17 β -dihydroxy-5 α -androst-3-one (**157**), and 1 α -methyl-11 α ,17 β -dihydroxy-5 α -androst-3-one (**158**) [85]. Bioconversion of **149** using *C. blakesleeana* produced




Figure 17.

Biotransformation products of 17 α -ethynyl-17 β -hydroxyandrost-4-en-3-one (**138**) and 17 α -ethyl-17 β -hydroxyandrost-4-en-3-one (**139**).




Figure 18.
 Biotransformation products of androsterone (145).

seven biotransformation products, such as 154, 157, 158, in addition to 1 α -methyl-1 β ,11 β ,17 β -trihydroxy-5 α -androst-3-one (159), 1 α -methyl-7 α ,11 β ,17 β -trihydroxy-5 α -androst-3-one (160), 1 α -methyl-1 β ,6 α ,17 β -trihydroxy-5 α -androst-3-one (161), and 1 α -methyl-1 β ,11 α ,17 β -trihydroxy-5 α -androst-3-one (162). *Macrophomina phaseolina* biotransformed 149 to obtain 1 α -methyl-17 β -hydroxy-5 α -androst-3,6-dione (155) [86]. Additionally, the biotransformation of 141 using *C. blakesleeana* (ATCC 8688A) yielded three metabolites: 1 α -methyl-11 β ,14 α ,17 β -trihydroxy-5 α -androst-3-one (163, 0.4%), 1 α -methyl-7 β ,17 β -dihydroxy-5 α -androst-3-one (164, 0.47%), and 1 α -methyl-17 β -hydroxy-5 α -androst-3,7-dione (165, 0.67%). *C. blakesleeana* catalyzed the β -hydroxylation in C-11, and dihydroxylation and oxidations at various positions of steroid skeleton (Figure 19) [87].

In the microbiological transformation of 3-hydroxyestra-1,3,5-(10)-trien-17-one (166) using *Fusarium oxysporum* var. *cubense*, the compounds, reduced in C-17 (3,17-dihydroxyestra-1,3,5-(10)-triene, 167) and hydroxylated in C-15 (3,15 α -dihydroxyestra-1,3,5-(10)-triene, 168), were isolated (Figure 20) [56].

Prednisone (169) is a synthetic corticosteroid (prodrug) used for the treatment of autoimmune, inflammatory and kidney diseases, among others.




Figure 19.
 Biotransformation products of mesterolone (149).




Figure 20.
Biotransformation products of 3-hydroxy-1,3,5-(10)-trien-17-one (**166**).




Figure 21.
Biotransformation products of prednisone (**169**).

Biotransformation of **169** using *C. elegans* occurred by hydrogenation of the $\Delta^{4(5)}$ and reduction of C-20, to produce the compounds 17 α ,21-dihydroxy-5 α -pregn-1-en-3,11,20-trione (**170**, 15.6%) and 17 α , (20S),21-trihydroxy-5 α -pregn-1-en-3,11-dione (**171**, 6.5%); whereas as the only biotransformation product, **169** using *F. lini* (5.2%), *R. stolonifer* (5.5%) and *C. lunata* (6.2%), was 1,4-pregnadien-17 α , (20S),21-trihydroxy-3,11-dione (**172**) (**Figure 21**) [88].

The main chemical transformation carried out by different *Acremonium* species in various steroid compounds have been oxidations, reductions, hydroxylations in different positions, isomerizations, and hydrolysis of the chain in C-17. Hydrocortisone (**173**) is an important anabolic, used clinically as anti-inflammatory and antiallergic drug, besides being a raw material for the synthesis of many steroid hormones. Biotransformation of **173** using *Acremonium strictum* generated the products 11 β ,17 β -dihydroxyandrost-4-en-3-one (**174**, 8%), 11 β ,17 α ,20 β ,21-tetrahydropregn-4-en-3-one (**175**, 11.2%), and 21-acetoxy-17 β ,17 α ,20-trihydroypregn-4-en-3-one (**176**, 7.6%); it was observed that the actions of the said species were as the reduction, acetylation and degradation of the chain in C-17, without modification of the unsaturated ketone- α,β [89]. Biotransformation of **173** using *Gibberella fujikuroi* yielded 11 β -hydroxyandrost-4-en-3,17-dione (**177**, 41%), while *B. subtilis* and *R. stolonifer* yielded **175** (15%). The products **173** (45%) and 3 β ,11 β ,17 α ,21-tetrahydroxy-5 α -pregnan-20-one (**178**, 31%) were obtained from the bioconversion of **173** using *Bacillus cereus* (**Figure 22**) [90].




Figure 22.
 Biotransformation products of hydrocortisone (173).




Figure 23.
 Biotransformation products of 17β-methoxy-5α-androst-3-one (179).

Incubation of 17β-methoxy-5α-androst-3-one (179) with *Cephalosporium aphidicola* produced 17β-methoxy-5α-androst-3β-ol (180) and 6β,11α-dihydroxy-17β-methoxy-5α-androst-3-one (181); while the biotransformation of 17β-methoxyestra-4-en-3-one (182) using *C. aphidicola* produced a major metabolite 6β-hydroxy-17β-methoxyestra-4-en-3-one (183). Similarly, the microbiological transformation of 3β-methoxyandrost-5-en-17-one (184) gave a mixture of products: 7α-hydroxy-3β-methoxyandrost-5-en-17-one (185) and 7β-hydroxy-3β-methoxyandrost-5-en-17-one (186) (Figure 23) [91].

In the literature, several species of fungi belonging to the genera *Aspergillus*, *Fusarium*, *Mortierella*, and *Penicillium* and capable of hydroxylating various steroids in C-15 have been described. For example, Jekkel et al. [92] described that more than 3000 fungi hydroxylate 13β-ethyl-4-gonene-3,17-dione (187) in C-15 position, the genus being *Fusarium*, particularly *F. nivale*; the fungus preferentially hydroxylated 187 with an α orientation in C-15 (15α-hydroxy-13β-ethyl-4-gonene-3,17-dione, 188, 77%) and C-7β (7β,15α-dihydroxy-13β-ethyl-4-gonene-3,17-dione, 189). On the other hand, the biotransformation of 187 using *Mortierella pusilla* produced 188, 190 (10β-hydroxy-13β-ethyl-4-gonene-3,17-dione) and 191 (6β-hydroxy-13β-ethyl-4-gonene-3,17-dione) (Figure 24).

The ethynodiol diacetate (192) is a synthetic derivative 1, used as an oral contraceptive because it inhibits the ovulation process. The microbiological transformation of 192 using *Cunninghamella elegans* produced four hydroxylated compounds




Figure 24.
Biotransformation products of 13 β -ethyl-4-gonene-3, 17-dione (**187**).

characterized as: 17 α -ethynylestr-4-en-3 β ,17 β -diacetoxy-6 α -ol (**193**, 0.5%), 17 α -ethynylestr-4-en-3 β ,17 β -diacetoxy-6 β -ol (**194**, 1.0%), 17 α -ethynylestr-4-en-3 β ,17 β -diacetoxy-10 β -ol (**195**, 0.5%), and 17 α -ethynyl-17 β -acetoxiestr-4-en-3-one (**196**, 1.4%) (**Figure 25**) [93].

Desogestrel (13-ethyl-17-methylene-18,19-dinor-17 α -pregn-4-en-20-yn-17-ol, **197**) is an orally active third-generation contraceptive steroid drug. Conversion of **197** by *C. blackesleeana* (ATCC 8688 A) yielded four metabolites: 13-ethyl-11-methylene-18,19-dinor-17 α -pregn-4-en-20-yn-6 β ,15 β ,17 β -triol (**198**), 13-ethyl-11-methylene-18,19-dinor-17 α -pregn-4-en-20-yn-3 β ,6 β ,17 β -triol (**199**), 13-ethyl-11-methylene-18,19-dinor-17 α -pregn-4-en-20-yn-3 α ,5 α ,6 β ,17 β -tetraol (**200**), and 13-ethyl-11-methylene-18,19-dinor-17 α -pregn-4-en-20-yn-6 β ,17 β -dihydroxy-3-one (**201**). Compounds **197** and **198** showed a potent growth inhibition against drug-resistant strains of *S. aureus* (**Figure 26**) [94].

The drugs mexrenone (**202**) and canrenone (**203**) are steroids with a spironolactone in C-17 and are potent antagonists of mineralocorticoids [95]. The biotransformation of **202** and **203** using a wide variety of microorganisms resulted in the production of monohydroxylated products in different positions, where *Beauveria bassiana* generated 11 α -hydroxymexrenone (**204**, 67%) as the major product, while 12 β -hydroxymexrenone (**205**, 50%) and 6 β -hydroxymexrenone (**206**, 33%) were obtained using *Mortierella isabellina*. The dehydrogenation product ($\Delta^{1(2)}$ -mexrenone, **207**, 15%) was favored with *Bacterium* cyclooxygenases as well as *Rhodococcus equi*, *Nocardia aurentia*, and *Comamonas testosteroni*. From the biotransformation of **203** using *Corynespora cassiicola*, 9 α -hydroxycanrenone




Figure 25.
Biotransformation products of ethynodiol diacetate (**192**).




Figure 28.

Biotransformation products of exemestane (211).

products in the C-7 (7 α -hydroxymethyltestosterone, 223, 35%) and C-15 (15 α -hydroxymethyltestosterone, 224, 21%) positions, plus a dihydroxylated product (12,15 α -dihydroxymethyltestosterone, 225, 22%) [100]. Recently, three additional products were identified: 11 α -hydroxy-17 α -methyltestosterone (226), 6 β -hydroxy-17 α -methyltestosterone (227), and 6 β ,11 α -dihydroxy-17 α -methyltestosterone (228). Isolation of hydroxylation products have been reported in different carbons from 222 with different orientations, C-6 β , C-7 β , C-9 α , C-11 α , C-12 β , and C-15 α (Figure 30).

Dianabol (methandrostenolone, 17 α -methyl-17 β -hydroxyl-androst-1,4-dien-3-one, 229) is an oral anabolic steroid that promotes the synthesis of proteins (increasing the muscle tissue). From the biotransformation of 229 using *Cunninghamella elegans*, five bioconversion products were obtained: 6 β -hydroxydianabol (230), 15 α -hydroxydianabol (231), 11 α -hydroxydianabol (232), 6 β ,12 β -dihydroxydianabol (233), and 6 β ,15 α -dihydroxydianabol (234). The products 17 β -hydroxy-17 α -methyl-5 α -androst-1,4-dien-3,6-dione (235), 7 β -hydroxydianabol (236), 15 β -hydroxydianabol (237), 17 β -hydroxy-17 α -methyl-5 α -androst-1,4-dien-3,11-dione






Figure 29.

Biotransformation products of formestane (216).




- 222) $R_1 = R_2 = R_3 = R_4 = R_5 = H$
 223) $R_1 = OH, R_2 = R_3 = R_4 = R_5 = H$
 224) $R_1 = R_3 = R_4 = R_5 = H, R_2 = OH$
 225) $R_1 = R_2 = R_5 = H, R_2 = R_3 = OH$
 226) $R_1 = R_2 = R_3 = R_5 = H, R_4 = OH$
 227) $R_1 = R_2 = R_3 = R_4 = H, R_5 = OH$
 228) $R_1 = R_2 = R_3 = H, R_4 = R_5 = OH$

Figure 30.
 Biotransformation products of methyltestosterone (222).




- 229) $R_1 = R_2 = R_3 = R_4 = R_5 = H$
 230) $R_1 = OH, R_2 = R_3 = R_4 = R_5 = H$
 231) $R_1 = R_3 = R_4 = R_5 = H, R_2 = \alpha OH$
 232) $R_1 = R_2 = R_4 = R_5 = H, R_3 = \alpha OH$
 233) $R_1 = R_4 = OH, R_2 = R_3 = R_5 = H$
 234) $R_1 = OH, R_2 = \alpha OH, R_3 = R_4 = R_5 = H$
 236) $R_1 = R_2 = R_3 = R_4 = H, R_5 = \beta OH$
 237) $R_1 = R_3 = R_4 = R_5 = H, R_2 = \beta OH$
 239) $R_1 = R_2 = R_4 = R_5 = H, R_3 = \beta OH$
 240) $R_1 = R_2 = R_3 = R_4 = H, R_5 = \alpha OH$




- 235) $R_1 = O, R_2 = H$
 238) $R_1 = H, R_2 = O$


Figure 31.
 Biotransformation products of dianabol (229).



- 241) $R_1 = R_2 = R_3 = R_4 = R_5 = R_6 = R_7 = H$
 242) $R_1 = O, R_2 = R_3 = R_4 = R_5 = R_6 = R_7 = H$
 243) $R_1 = R_2 = R_3 = R_4 = R_6 = H, R_5 = OH$
 244) $R_1 = R_2 = R_4 = R_6 = R_7 = H, R_3 = R_5 = OH$
 245) $R_1 = R_2 = R_3 = R_5 = R_6 = H, R_4 = R_7 = OH$
 246) $R_1 = R_2 = R_3 = R_6 = R_7 = H, R_4 = R_5 = OH$
 247) $R_1 = R_2 = R_3 = R_4 = R_5 = H, R_6 = R_7 = OH$



- 248) $R_1 = R_2 = OH, \Delta^{14(15)}$
 249) $R_1 = H, R_2 = OH$
 252) $R_1 = O, R_2 = H$
 253) $R_1 = R_2 = H$



251

Figure 32.
 Biotransformation products of masthasterone (241).

(238), and 11 β -hydroxydianabol (239) were obtained from the biotransformation of 229 using *Macrophomina phaseolina* [101]. Biotransformation of 229 using several microorganisms has been reported, for example, *Penicillium notatum* [102] transformed 229 into 230 and 231, while *Trichoderma hamatum* produced 232 [103].

Similarly, *B. bassiana*, *A. ochraceus*, *Colletotrichum lagenarium*, and *Sporotrichum sulfurreducens* gave a biotransformed product **232** [104]. *Absidia glauca* metabolized **229** in compounds **230**, and **236–237** [105]. In contrast, the biotransformation of **229** using *A. coerulea* yielded **239** along with 7 α -hydroxydianabol (**240**) [106], while by using *B. cinerea*, **237** was obtained as the only product (**Figure 31**) [107].

Methasterone (**241**) is a synthetic anabolic steroid, known to gain muscle mass. Microbial transformation of **241** using *M. phaseolina* yielded 17 β -hydroxy-17 α -(hydroxymethyl)-2 α -methyl-5 α -androstane-3,6-dione (**242**), while by using *C. blakesleeana*, 7 α -hydroxymethasterone (**243**, 2.0%), 7 α ,16 β -dihydroxymethasterone (**244**, 0.7%), 5 α ,12 β -dihydroxymethasterone (**245**, 1.0%), 7 α ,12 β -dihydroxymethasterone (**246**, 1.5%), and 7 α ,9 α -dihydroxy-methasterone (**247**, 0.5%) were obtained. Incubation of **241** with *Fusarium lini* yielded different metabolites with dehydrogenation in ring A and D: 6 β ,17 β -dihydroxy-2,17 α -dimethyl-5 α -androst-1,4-diene-3-one (**248**, 1.0%), 15 α ,17 β -dihydroxy-2 α ,17 α -dimethyl-5 α -androst-1,4-diene-3-one (**249**, 0.6%), 6 β ,17 β -dihydroxy-2,17 α -dimethylandrost-1,4-diene-3-one (**250**, 0.4%), 14 α ,15 α -dihydroxy-2,17-dimethyl-5 α -androst-1,4,16-trien-3-one (**251**, 0.3%), 17 β -hydroxy-2,17 α -dimethyl-5 α -androst-5 α -1,4-dien-3,6-dione (**252**, 0.3%), and 17 β -hydroxy-2,17 α -dimethyl-5 α -androst-1,4-dien-3-one (**253**, 1.0%) (**Figure 32**) [108].

3. Conclusions

The biotransformation processes of different steroid compounds described in this review, although not exhaustive, aim to highlight the importance of biotransformation through different microorganisms, as a useful chemical-biological tool for obtaining novel derivatives for research purpose and as industrial applications. An example includes obtaining steroid compounds for the pharmaceutical industry.

Biotransformation of steroids has been implemented in an important way in the partial synthesis of new steroids, for their evaluation as hormones and drugs. Currently, there is a wide variety of steroids used as diuretics, anabolic, anti-inflammatory, antiandrogenic, anticontraceptive, antitumor, among other applications. Chemical functionalization in different carbon atoms of the steroid skeleton is related to the biological activity of the molecule. This is why microbiological transformations play an important role in obtaining these compounds through chemical transformations, such as the oxidation of hydroxyl group at C-3 and C-17, isomerization of the double bond $\Delta^{5(6)}$ to $\Delta^{4(5)}$, hydrogenation of double bonds $\Delta^{1(2)}$ and $\Delta^{4(5)}$, and reduction of the carbonyl group at C-17 and C-20 with β orientation. Biohydroxylations performed in different positions of the steroid skeleton—C-11 α , C-11 β , C-15 β , and C-16 α —using different species of fungi of the genera *Rhizopus*, *Aspergillus*, *Curvularia*, *Cunninghamella*, and *Streptomyces* with high yields are an important chemical transformation in many synthesis schemes of new steroids with a determined biological activity.

Hydroxylation of steroids—progesterone, testosterone, 17 α -methyltestosterone, and 4-androsten-3,17-dione—presenting the 4-en-3-one system, proceeds with a high stereo- and regioselectivity in the C-6 and C-11 positions, with a β orientation in C-6 and α orientation in C-11. The presence of the methyl group in C-10 is necessary for the hydroxylation in C-11, as can be seen in the derivatives of 19-nortestosterone.

The interest in the biotransformation of steroid compounds has been increasing in recent years, due to the obtaining of new and useful pharmacologically active compounds. In addition to the development of new genetically modified strains, there is an increase in the availability of immobilized enzymes and the manipulation of culture media.

Biotransformation of steroids proceeds with low to moderate yields in general. One of the main causes is their low solubility in water. Currently, methodologies are developed that allow the incorporation of chemicals—surfactants, ionic liquids, cyclodextrins, liposomes, among others—that contribute to improve the yields of each biotransformation process and the processes friendly to the environment.

Acknowledgements


A la Carrera de Biología, FES-Zaragoza, UNAM. Ms. Fabiola Cano thank for her suggestions and comments toward improving the manuscript.

Author details

Arturo Cano-Flores*, Javier Gómez and Rigoberto Ramos
Laboratorio de Biotransformaciones y Química de Productos Naturales, L-314,
Facultad de Estudios Superiores Zaragoza, UNAM, México, D.F., Mexico

*Address all correspondence to: acano1750@gmail.com

IntechOpen

© 2019 The Author(s). Licensee IntechOpen. This chapter is distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/3.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. 

References

- [1] Mensah-Nyagan AG, Meyer L, Schaeffer V, Kibaly C, Patten-Mensah C. Evidence for a key role of steroids in the modulation of pain. *Psychoneuroendocrinology*. 2009;**34**:169-177. DOI: 10.1016/j.psyneuen.2009.06.004
- [2] Funder JW. Minireview: Aldosterone and mineralocorticoid receptors: Past, present, and future. *Endocrinology*. 2010;**151**:5098-5102. DOI: 10.1210/en.2010-0465
- [3] Fernandez P, Cruz A, Angelova B, Pinheiro HM, Cabral JMS. Microbial conversion of steroid compounds: Recent developments. *Enzyme and Microbial Technology*. 2003;**32**:688-705. DOI: 10.1016/S0141-0229(03)00029-2
- [4] Hogg JA. Steroids, steroids community and Upjohn in perspective: A profile of innovation. *Steroids*. 1992;**257**:593-616
- [5] Sedlaczek L. Biotransformations of steroids. *Critical Reviews in Biotechnology*. 1988;**7**:187-236. DOI: 10.3109/07388558809146602
- [6] Ko D-H, Heiman AS, Chen M, Lee HJ. New steroidal anti-inflammatory antedugs: Methyl 21-desoxy-21-chloro-11 β ,17 α -dihydroxy-3,20-dioxo-1,4-pregnadiene-16 α -carboxylate, methyl 21-desoxy-21-chloro-11 β -hydroxy-3,20-dioxo-1,4-pregnadiene-16 α -carboxylate, and their 9 α -fluoro derivatives. *Steroids*. 2000;**65**:210-218
- [7] Fragkaki AG, Angelis YS, Koupparis M, Tsantili-Kakoulidou A, Kokotos G, Georgakopoulos C. Structural characteristics of anabolic androgenic steroids contributing to binding to the androgen receptor and to their anabolic and androgenic activities: Applied modifications in the steroidal structure. *Steroids*. 2009;**74**:172-197. DOI: 10.1016/j.steroids.2008.10.016
- [8] Agoston ES, Hastcher MA, Kensler TW, Posner GH. Vitamin D analogs as anti-carcinogenic agents. *Anti-Cancer Agents in Medicinal Chemistry*. 2006;**6**:53-71. DOI: 10.2174/187152006784111369
- [9] Tuba Z, Barden CW, Dancsi A, Francsics-Czinege E, Molnar EC, Csorgei J, et al. Synthesis and biological activity of a new progestogen, 16-methylene-17 α -hydroxy-18-methyl-19-norpregn-4-ene-3,20-dione acetate. *Steroids*. 2000;**65**:266-274. DOI: 10.1016/S0039-128X(99)00109-9
- [10] Siemes C, Visser LE, De-Jong FH, Jan-Willem Coebergh W, Uitterlinden AG. Cytochrome P450 3A gene variation, steroid hormone serum levels and prostate cancer. *Steroids*. 2010;**75**:1024-1032. DOI: 10.1016/j.steroids.2010.07.001
- [11] Diaz-Chico BN, Rodriguez FG, Gonzalez A, Ramirez R, Bilbao C, Leon AC, et al. Androgens and androgen receptors in breast cancer. *Journal of Steroid Biochemistry and Molecular Biology*. 2007;**105**:1-15. DOI: 10.1016/j.jsbmb.2006.11.019
- [12] Johnston JO. Aromatase inhibitors. In: Parish EJ, Nes WD, editors. *Biochemistry and Function of Sterols*. Boca Raton, FL: CRC Press; 1987. pp. 23-53
- [13] Craigie E, Mullins JJ, Bailey MA. Glucocorticoids and mineralocorticoids. In: Bader M, editor. *Cardiovascular Hormone Systems: From Molecular Mechanisms to Novel Therapeutics*. Weinheim: Wiley-VCH Verlag GmbH & Co; 2009. pp. 1-64
- [14] Chung SK, Ryoo CH, Yang HW, Shim JY, Kang MG, Lee KW, et al.

- Synthesis and bioactivities of steroid derivatives as antifungal agents. *Tetrahedron*. 1998;**52**:15899-15914. DOI: 10.1016/S0040-4020(98)01000-X
- [15] Suzuki K, Nakata T, Shimizu T, Enomoto K. Anti-obesity agents. US Patent 5,846,962; 1998
- [16] Dombrowski AW, Hazuda DJ, Polishook JD, Felock PJ, Singh SB, Zink DL. HIV-integrase-inhibitors. WO Patent 0036132; 2000
- [17] Athan D, Svasti J, Kittakoop P, Pittayakhachonwut D, Tanticharpen M, Thebtaranonth Y. Antiviral isoflavonoid sulfate and steroidal glycoside from the fruits of *Solanum torvum*. *Phytochemistry*. 2002;**59**:459-463. DOI: 10.1016/S0031-9422(01)00417-4
- [18] Rupprecht R, Holsboer F. Neuroactive steroids: Mechanisms of action and neuropsychopharmacology perspectives. *Trends in Neurosciences*. 1999;**22**:410-416. DOI: 10.1016/S0166-2236(99)01399-5
- [19] Mellon SH, Griffin LD. Neurosteroids: Biochemistry and clinical significance. *Trends in Endocrinology and Metabolism*. 2002;**13**:35-43. DOI: 10.1016/S1043-2760(01)00503-3
- [20] Baulieu E-E, Schumacher M. Progesterone as a neuroactive neurosteroid, with special reference to the effect of progesterone on myelination. *Steroids*. 2000;**65**:605-612
- [21] Fernandez P, Cabral JMS. Steroid bioconversion. In: Flickinger M, editor. *Encyclopedia of Industrial Biotechnology: Bioprocess, Bioseparation, and Cell Technology*. New York: John Wiley & Sons, Inc.; 2010. pp. 1-32
- [22] Donova MV, Ergova OV. Microbial steroid transformations: Current state and prospect. *Applied Microbiology and Biotechnology*. 2012;**94**:1423-1447. DOI: 10.1007/978-1-4939-7183-1
- [23] Sultana N. Microbial biotransformation of bioactive and clinically useful steroids and some salient features of steroids and biotransformation. *Steroids*. 2018;**136**:76-92. DOI: 10.1016/j.steroids.2018.01.007
- [24] Tong W-Y, Dong X. Microbial biotransformation: Recent developments on steroid drugs. *Recent Patents on Biotechnology*. 2009;**3**:141-153
- [25] Bureik M, Bernhardt R. Steroid hydroxylation: Microbial steroid biotransformation using cytochrome P450 enzymes. In: Schmid RD, Urlacher VB, editors. *Modern Biooxidation: Enzymes, Reactions and Applications*. Weinheim: Wiley-VCH Verlag GmbH & Co. KGaA; 2007. pp. 155-176
- [26] Lehman LR, Stewart JD. Filamentous fungi: Potentially useful for the biohydroxylation of non-activated carbon centres. *Current Organic Chemistry*. 2001;**5**:439-470. DOI: 10.2174/1385272013375490
- [27] Pollard DJ, Woodley JM. Biocatalysis for pharmaceutical intermediates: The future is now. *Trends in Biotechnology*. 2006;**25**:66-72. DOI: 10.1016/j.tibtech.2006.12.005
- [28] Straathof AJJ. The production of fine chemicals by biotransformation. *Current Opinion in Biotechnology*. 2002;**13**:548-546. DOI: 10.1016/S0958-1669(02)00360-9
- [29] Mahato SB, Majumdar I. Current trends in microbial steroid biotransformation. *Phytochemistry*. 1993;**34**:883-898. DOI: 10.1016/S0031-9422(00)90685-X

- [30] Mahato SB, Garai S. Advances in microbial steroid biotransformation. *Steroids*. 1997;**62**:332-345. DOI: 10.1016/S0039-128X(96)00251-6
- [31] Donova MV. Transformation of steroids by actinobacteria. *Applied Biochemistry and Microbiology*. 2007;**43**:1-14
- [32] Prabha V, Ohri M. Review: Bacterial transformations of bile acids. *World Journal of Microbiology and Biotechnology*. 2006;**22**:191-196. DOI: 10.1007/s11274-005-9019-y
- [33] Huttel W, Hoffmeister D. Fungal Biotransformation in pharmaceutical sciences. In: Hofrichter M, editor. *Industrial Applications, the Mycota*. Berlin: Springer; 2010. pp. 293-317
- [34] Petric S, Hakki T, Bernhardt R, Zigon D, Cresnar B. Discovery of a steroid 11 α -hydroxylase from *Rhizopus oryzae* and its biotechnological application. *Journal of Biotechnology*. 2010;**150**:428-437. DOI: 10.1016/j.jbiotec.2010.09.928
- [35] Rodina NV, Andryushina VA, Stysenko TS, Turova TP, Baslerov RV, Panteleeva AN, et al. The introduction of the 9 α -hydroxy group into andros-4-en-3,17-dione using a new actinobacterium strain. *Applied Biochemistry and Microbiology*. 2009;**45**:395-400
- [36] Faber K, Kroutil W. New enzymes for biotransformation. *Current Opinion in Chemical Biology*. 2005;**9**:181-187. DOI: DOI 10.1016/j.cbpa.2005.01.0
- [37] Ran N, Zhao L, Chen Z, Tao J. Recent applications of biocatalysis in developing green chemistry for chemical synthesis at the industrial scale. *Green Chemistry*. 2008;**10**:361-372. DOI: 10.1039/b716045c
- [38] Hu SH, Genain G, Azerad R. Microbial transformation of steroids: Contribution to 14 α -hydroxylations. *Steroids*. 1995;**60**:337-352
- [39] Smith KE, Ahmed F, Williams RAD, Kelly SL. Microbial transformations of steroids-VIII. Transformation of progesterone by whole cells and microsomes of *Aspergillus fumigatus*. *The Journal of Steroid Biochemistry and Molecular Biology*. 1994;**49**:93-100
- [40] Khalil AM, Eman MM. Microbiological transformation of progesterone by some zoosporic fungi. *Journal of Basic Microbiology*. 1996;**36**:255-259
- [41] Javid M, Nickavar B, Vahidi H, Faramarzi MA. Baeyer-Villiger oxidation of progesterone by *Aspergillus sojae* PTCC 5196. *Steroids*. 2018;**140**:52-57. DOI: 10.1016/j.steroids.2018.07.008]
- [42] Kolek T, Swizzer A. Biotransformation XLV. Transformations of 4-ene-3-oxosteroids in *Fusarium culmorum* culture. *The Journal of Steroid Biochemistry and Molecular Biology*. 1998;**67**:63-69
- [43] Al-Awadi S, Afzal M, Oommen S. Studies on *Bacillus stearothermophilus*. Part I. Transformation of progesterone to a new metabolite 9,10-seco-4-pregnene-3,9,20-trione. *The Journal of Steroid Biochemistry and Molecular Biology*. 2001;**78**:493-498
- [44] Kolet SP, Niloferjahan S, Haldar S, Gonnade R, Thulasiram HV. Biocatalyst mediated production of 6 β ,11 α -dihydroxy derivatives of 4-ene-3-one steroids. *Steroids*. 2013;**78**:1152-1158. DOI: 10.116/j.steroids.2013.08.00
- [45] Eshrat G-F, Aroona C. Biotransformation of progesterone by *Penicillium aurantiogriseum*. *Research Journal of Microbiology*. 2011;**6**:98-104. DOI: 10.3923/jm.2011.98.104
- [46] Gopal SKV, Naik S, Somal P, Sharma P, Arjuna A, Hassam R, et al. Production

of 17-ketoandrostene steroids by the side chain cleavage of progesterone with *B. sphaericus*. Biocatalysis and Biotransformation. 2008;**26**:272-279. DOI: 10.1080/10242420701707637

[47] Al-Beloshei NE, Al-Awadhi HA, Al-Khalaf RAR, Oommen S, Afzal M. Biocatalyzed transformation of progesterone by *Geobacillus gargensis* DSM 15378. Applied Biochemistry and Microbiology. 2015;**51**:321-328

[48] Gas KP, Sakac M, Jovanovic-Santa S, Djurendic E. An overview of partial synthesis and transformations of secosteroids. Current Organic Chemistry. 2014:216-259. DOI: 10.2174/138527281131176660147.]

[49] Lin Z, Marepally SR, Ma D, Kim T-K, Oak ASW, Myers LK, et al. Synthesis and biological evaluation of vitamin D3 metabolite 20S,23S-dihydroxyvitamin D3 and its 23R epimer. Journal of Medicinal Chemistry. 2016:136, 5102-5108. DOI: 10.1021/acs.jmedchem.6b00182

[50] Madyastha KM, Joseph T. Transformation of 16-dehydroprogesterone and 17 α -hydroxyprogesterone by *Mucor piriformis*. Applied Microbiology and Biotechnology. 1994;**41**:170-177

[51] Madyastha KM, Joseph T. Transformation of dehydroepiandrosterone and pregnenolone by *Mucor piriformis*. Applied Microbiology and Biotechnology. 1995;**44**:339-343

[52] Shan L-H, Liu H-M, Huang K-H, Dai G-F, Cao C, Dong R-J. Synthesis of 3 β ,7 α ,11 α -trihydroxy-pregn-21-benzylidene-5-en-20-one derivatives and their cytotoxic activities. Bioorganic & Medicinal Chemistry Letters. 2009;**19**:6637-6639. DOI: 10.1016/j.bmcl.2009.10.019

[53] Farooq A, Tahara S. Biotransformation of testosterone and

pregnenolone catalyzed by the fungus *Botrytis cinerea*. Journal of Natural Products. 2000;**63**:489-491. DOI: 10.1021/np990520b

[54] Choudhary MI, Batool I, SAA S, Nawaz SA, Ali Shah SA, Atta-ur-Rahman. Microbial hydroxylation of pregnenolone derivatives. Chemical & Pharmaceutical Bulletin. 2005;**55**:1455-1459

[55] Schaaaf O, Dettner K. Transformation of steroids by *Bacillus* strains isolated from the foregut of water beetles (Coleoptera: Dytiscidae): II. Metabolism of 3 β -hydroxypregn-5-en-20-one (pregnenolone). The Journal of Steroid Biochemistry and Molecular Biology. 2000;**75**:187-199

[56] Wilson MR, Gallimore WA, Reese PB. Steroid transformations with *Fusarium oxysporum* var. cubense and *Colletotrichum musae*. Steroids. 1999;**64**:834-843. DOI: 10.1016/S0039-128X(99)00067-7

[57] Hu SH, Tian XF, Sun YH, Han GD. Microbial hydroxylation of 13-ethyl-17 β -hydroxy-18,19-dinor-17 α -pregn-4-en-20-in-17-ona. Steroids. 1996;**61**:407-410

[58] Hu SH, Tian XF, Han GD. Novel microbial hydroxylation of 13-ethyl-17 β -hydroxy-18,19-dinor-17 α -pregn-4-en-20-yn-3-one. Steroids. 1998;**63**:88-92

[59] Choudhary MI, Azizuddin, Atta-ur-Rehman. Microbial transformation of danazol. Natural Product Letters. 2002;**16**:101-106. DOI: 10.1080/10575630290019994

[60] Baydoun E, Atia-tul-Wahab, Mehmood H, Ahmad MS, Malik R, Smith C, et al. Microbial transformation of danazol with *Cunninghamella blakesleeana* and anti-cancer activity of danazol and its transformed products. Steroids. 2016;**105**:121-127. DOI: 10.1016/j.steroids.2015.11.010

- [61] Choudhary MI, Musharraf SG, Ali RA, Atif M, Atta-ur-Rahman. Microbial transformation of antifertility agents. Norethisterone and 17 α -ethynylestradiol. *Zeitschrift für Naturforschung*. 2004;**59B**:319-323
- [62] Choudhary MI, Musharraf SG, Ali-Siddiqui Z, Khan NT, Alli RA, Atta-ur-Rahman. Microbial transformation of mestranol by *Cunninghamella elegans*. *Chemical & Pharmaceutical Bulletin*. 2005;**53**:1011-1013
- [63] Ahmad MS, Zafar S, Yousuf S, Atta-tul-Wahab, Atta-ur-Rahman, Choudhary MI. Biotransformation of 6-dehydroprogesterone with *Aspergillus niger* and *Gibberella fujikuroi*. *Steroids*. 2016;**112**:62-67. DOI: 10.1016/j.steroids.2016.04.008
- [64] Peterson DH, Nathan AH, Meister PD, Eppstein SH, Murray HC, Weintraub A, et al. Microbiological transformation of steroids VI. Preparation of 11 α -hydroxy-6-dehydroprogesterone. *Journal of the American Chemical Society*. 1953;**75**:419-421
- [65] Smith KE, Latif S, Kirk DN, White KA. Microbial transformations of steroids IV. 6,7-dehydrogenation; a new class of fungal steroids transformation product. *Journal of Steroid Biochemistry*. 1989;**33**:271-276. DOI: 10.1016/0022-4731(89)90304-X
- [66] Baydoun E, Bano S, Atia-tu-Wahab, Jabeen A, Yousuf S, Mesaik A, et al. Fungal transformation and T-cell proliferation inhibitory activity of melengestrol acetate and its metabolite. *Steroids*. 2014;**86**:56-61. DOI: 10.1016/j.steroids.2014.04.012
- [67] Madyasth KM. Novel microbial transformation of steroids. *Advances in Experimental Medicine and Biology*. 1995;**405**:259-270
- [68] Angelova B, Mutafov S, Avramova T, Dimova I, Boyadjieva L. 9 α -Hydroxylation of 4-androstene-3,17-dione by resting *Rhodococcus* sp. cells. *Process Biochemistry*. 1996;**31**:179-184
- [69] Schaaaf O, Dettner K. Transformation of steroids by *Bacillus* strains isolated from the foregut of water beetles (Coleoptera: Dytiscidae): I. Metabolism of androst-4-en-3,17-dione (AD). *The Journal of Steroid Biochemistry and Molecular Biology*. 1998;**67**:451-465
- [70] Hanson JR, Nasir H, Pervez A. The hydroxylation of testosterone and some derivatives by *Cephaslosporium aphidicola*. *Phytochemistry*. 1996;**42**:411-415
- [71] Choudhary MI, Sultan S, MTH K, Yasin A, Shaheen F, Atta-ur-Rehman. Biotransformation of (+)-androst-4-ene-3,17-dione. *Natural Product Research*. 2004;**18**:529-535. DOI: 10.1080/14786410310001620628
- [72] Xiong Z, Wei Q, Chen H, Chen S, Xu W, Qiu G, et al. Microbial transformation of androst-4-en-3,17-dione by *Beauveria bassiana*. *Steroids*. 2006;**71**:979-983. DOI: 10.1016/j.steroids.2006.07.007
- [73] Janeczko T, Dmochowska-Gładysz J, Kostrzewa-Susłow E, Białonska A, Ciunik Z. Biotransformations of steroid compounds by *Chaetomium* sp. KCH 6651. *Steroids*. 2009;**74**:657-661. DOI: 10.1016/j.steroids.2009.02.006
- [74] Andryushina VA, Voishvillo NE, Druzhinina AV, Stytsenko TS, Yaderets VV, Petrosyan MA, et al. 14 α -hydroxylation of steroids by mycelium of the mold fungus *Curvularia lunata* (VKPM F-981) to produce precursors for synthesizing new steroidal drugs. *Pharmaceutical and Chemical Journal*. 2013;**47**:103-108
- [75] Qiao Y, Shen Y, Huang W, Wang Y, Ren J, Xia T, et al. Biocatalyst-mediated production of 11,15-dihydroxy derivatives

- of androst-1,4-dien-3,17-dione. Journal of Bioscience and Bioengineering. 2017;123:692-697. DOI: 10.1016/j.jbiosc.2017.01.008.], 10.1016/j.jbiosc.2017.01.008.]
- [76] Ahmed F, Williams RAD, Smith KE. Microbial transformations of steroids-X. Cytochromes P-450 11 α -hydroxylase and C17-C20 lyase and a 1-ene-dehydrogenase transform steroids in *Nectria haematococca*. The Journal of Steroid Biochemistry and Molecular Biology. 1996;58:337-349. DOI: 10.1016/0960-0760(96)00032-5
- [77] Atta-ur-Rahman, Choudhary MI, Asif F, Farooq A, Yaqoob M. Microbial transformations of testosterone. Natural Product Letters. 1998;12:255-261. DOI: 10.1080/10575639808048299
- [78] Al-Awadi S, Afzal M, Oommen S. Studies of *Bacillus stearothermophilus*. Part III. Transformation of testosterone. Applied Microbiology and Biotechnology. 2003;62:48-52. DOI: 10.1007/s00253-003-1269-8
- [79] Al-Awadi S, Afzal M, Oommen S. Studies on *Bacillus stearothermophilus*. Part IV. Influence of enhancers on biotransformation of testosterone. Steroids. 2005;70:323-327
- [80] Wu Y, Li H, Zhang XM, Gong JS, Ra ZM, Shi JS, et al. Efficient hydroxylation of functionalized steroids by *Colletotrichum lini* ST-1. Journal of Molecular Catalysis B: Enzymatic. 2015;120:111-118. DOI: 10.1016/j.molcatb.2015.07.003
- [81] Choudhary MI, Ali Shah SA, Musharraf SG, Shaheen F, Atta-ur-Rahman. Microbial transformation of dehydroepiandrosterone. Natural Product Research. 2003;17:215-220. DOI: 10.1080/1057563021000040835
- [82] Huang L-H, Li J, Xu G, Zhang X-H, Wang Y-G, Yin Y-L, et al. Biotransformación of dehydroepiandrosterone (DHEA) with *Penicillium griseopurpureum* Smith and *Penicillium glabrum* (Wehmer) Westling. Steroids. 2010;75:1039-1046. DOI: 10.1016/j.steroids.2010.06.008
- [83] Choudhary MI, Sultan S, Khan MTH, Atta-ur-Rehman. Microbial transformation of 17 α -ethynyl- and 17 α -ethylsteroids, and tyrosinase inhibitory activity of transformed products. Steroids. 2005;70:798-802. DOI: 10.1016/j.steroids.2005.05.002
- [84] Musharraf SG, Atta-ur-Rahman, Choudhary MI, Sultan S. Microbial transformation of (+)-adrenosterone. Natural Product Letters. 2002;16:345-349. DOI: 10.1080/10575630290033105
- [85] Choudhary MI, Sultan S, Jalil S, Anjum S, Rahman AA, Fun HK, et al. Microbial transformation of mesterolone. Chemistry & Biodiversity. 2005;2:392-400. DOI: 10.1002/cbdv.200590019
- [86] Ahmad MS, Zafar S, Bibi M, Bano S, A-tul W, Atta-ur-Rahman, et al. Biotransformation of androgenic steroid mesterolone with *Cunninghamella blakesleeana* and *Macrophomina phaseolina*. Steroids. 2014;82:53-59. DOI: 10.1016/j.steroids.2014.01.001
- [87] Ahmada MS, Farooqa R, Hussaina N, Atia-tul-Wahabb, Atta-ur-Rahmana, Choudhary MI. Three new analogues of androgenic drug mesterolone through biotransformation with *Cunninghamella blakesleeana*. Journal of Molecular Catalysis B: Enzymatic. 2016;133:S395-S399. DOI: 10.1016/j.molcatb.2017.03.001
- [88] Choudhary MI, Siddiqui ZA, Musharraf SG, Nawaz SA, Atta-ur-Rahman. Microbial transformation of prednisone. Natural Product Research. 2005;19:311-317. DOI: 10.1080/14786410410001729168

- [89] Faramarzi MA, Yazdi MT, Shafiee A, Zarrini G. Microbial transformation of hydrocortisone by *Acremonium strictum* PTCC 5282. *Steroids*. 2002;**67**:869-872. DOI: 10.1016/S0039-128X(02)00049-1
- [90] Choudhary MI, Sultan S, Yaqoob M, Musharraf SG, Yasin A, Shaheen F, et al. Microbial transformation of cortisol and prolyl endopeptidase inhibitory activities of its transformed products. *Natural Product Research*. 2003;**17**:389-395. DOI: 10.1080/14786410310001617640
- [91] Kiran I, Hanson JR, Hunter AC. The microbiological hydroxylation of some methoxysteroids by *Cephalosporium aphidicola*. *Journal of Chemical Research*. 2004;**5**:362-363. DOI: 10.3184/0308234041639827
- [92] Jekkel A, Ilkoy E, Horvath G, Pallagi I, Süto J, Ambrus G. Microbial hydroxylation of 13 β -ethyl-4-gonene-3, 17-dione. *Journal of Molecular Catalysis B: Enzymatic*. 1998;**5**:385-387. DOI: 10.1016/S1381-1177(98)00097-6
- [93] Zafar S, Yousuf S, Kayani HA, Saifullah S, Khan S, Al-Majid AM, et al. Biotransformation of oral contraceptive ethynodiol diacetate with microbial and plant cell cultures. *Chemistry Central Journal*. 2012;**6**:109-116. DOI: 10.1186/1752-153X-6-109
- [94] Atia-tul-Wahab, Siddiqui M, Ibrahim I, Hussain A, Ajandouz EH, Hijazi A, et al. *Cunninghamella blakesleeana* mediated biotransformation of a contraceptive drug, desogestrel, and anti-MDR-*Staphylococcus aureus* activity of its metabolites. *Bioorganic & Medicinal Chemistry*. 2018;**77**:152-158. DOI: 10.1016/j.bioorg.2017.12.027
- [95] Huyet J, Pinon GM, Fay MR, Fagart J, Rafestin-Oblin M-E. Structural basis of spiro lactone recognition by the mineralocorticoid receptor. *Molecular Pharmacology*. 2007;**72**:563-571. DOI: 10.1124/mol.107.036459
- [96] Preisig CL, Laakson JA, Mocek UM, Wang PT, Baez J, Byng G. Biotransformation of the cardiovascular drugs mexrenone and canrenone. *Journal of Natural Products*. 2003;**66**:350-356. DOI: 10.1021/np020347a
- [97] Baydoun E, Bibi M, Iqbal MA, Wanab A, Farran D, Smith C, et al. Microbial transformation of anti-cancer steroid exemestane and cytotoxicity of its metabolites against cancer cell lines. *Chemistry Central Journal*. 2013;**7**:57-63. DOI: 10.1186/1752-153X-7-57
- [98] Martin GDA, Narvaez J, Bulmer R, Durrant MC. Biotransformation and molecular docking studies of aromatase inhibitor. *Steroids*. 2016;**113**:95-102. DOI: 10.1016/j.steroids.2016.07.003
- [99] Martin GDA, Narvaez J, Marti A. Synthesis and bioconversions of formestane. *Journal of Natural Products*. 2013;**76**:1966-1969. DOI: 10.1021/np400585t
- [100] Torshabi M, Badiee M, Faramarzi MA, Rastegar H, Forootanfar H, Mohit E. Biotransformation of methyltestosterone by the filamentous fungus *Mucor racemosus*. *Chemistry of Natural Compounds*. 2011;**47**:59-63
- [101] Khan NT, Zafar S, Noreen S, Al Majid AM, Al Othman ZA, Al-Rasayes SI, et al. Biotransformation of dianabol with the filamentous fungi and β -glucuronidase inhibitory activity of resulting metabolites. *Steroids*. 2014;**85**:65-72. DOI: 10.1016/j.steroids.2014.04.004
- [102] Bartmanska A, Dmochowska-Gladysz J, Huszcza E. Steroids transformation in *Penicillium notatum* culture. *Steroids*. 2005;**70**:193-198. DOI: 10.1016/j.steroids.2004.11.011

[103] Bartmanska A, Dmochowska-Gladysz J. Transformation of steroids by *Trichoderma hamatum*. *Enzyme and Microbial Technology*. 2007;**40**:1651-1621

[104] Huszcza E, Dmochowska-Gladysz J, Bartmanska A. Transformation of steroids by *Beauveria bassiana*. *Zeitschrift für Naturforschung*. 2005;**60c**:103-108

[105] Huszcza E, Dmochowska-Gladysz J. Transformation of testosterone and related steroids in *Absidia glauca* culture. *Journal of Basic Microbiology*. 2003;**43**:113-120

[106] Brezezowska E, Dmochowska-Gladysz J, Kolek T. Biotransformation XXXIX. Metabolism of testosterone, androstenedione, progesterone and testosterone derivatives in *Absidia coerulea* culture. *The Journal of Steroid Biochemistry and Molecular Biology*. 1996;**57**:357-362

[107] Huszcza E, Dmochowska-Gladysz J. Transformation of testosterone and related steroids by *Botrytis cinerea*. *Phytochemistry*. 2003;**62**:155-158

[108] Ahmad MS, Yousuf S, Atia-tul-Wahab, Jabeen A, Atta-ur-Rahman, Choudhary MI. Biotransformation of anabolic compound methasterone with *Macrophomina phaseolina*, *Cunninghamella blackesleeana*, and *Fusarium lini*, and TNF- α inhibitory effect of transformed products. *Steroids*. 2017;**128**:75-84. DOI: 10.1016/j.steroids.2017.04.001