



Some Remarks on Lukasiwicz disjunction and conjunction operators On Intuitionistic Fuzzy Matrices

T.Muthuraji and S.Sriram

Mathematics Section, Faculty of Engg and Tech,
Annamalai University, Annamalai nagar,
Chidambaram, TamilNadu, India-608002.

ABSTRACT

Some properties of two operations - conjunction and disjunction from Lukasiwicz type – over Intuitionistic Fuzzy Matrices are studied.

Keywords and Phrases: Intuitionistic Fuzzy Set (IFS); Intuitionistic Fuzzy Matrix (IFM).



Council for Innovative Research

Peer Review Research Publishing System

Journal: JOURNAL OF ADVANCES IN MATHEMATICS

Vol .11, No.3

www.cirjam.com , editorjam@gmail.com

1 .INTRODUCTION:

After the introduction of fuzzy theory in [1], intuitionistic fuzzy theory [2] has been found to be highly meaningful to consider vagueness. In [3] Krassimir T. Atanassov defined so many new operations over intuitionistic fuzzy sets and some operations deal with uncertainty case of the element also. Using the theory of IFS, Im et al [9] defined the notion of IFMs. X.Zhang [7] studied Intuitionistic Fuzzy Value and introduced the concept of composition two Intuitionistic Fuzzy matrices. Z.Xu [6], Ronald R.Yager [8], Meenakshi A.R and Gandhimathi [5] and several authors discussed IFMs. Here we give some inequalities contains Lukasiewicz fuzzy conjunction and disjunction operators and some properties in IFMs. The above said operators are introduced by Atanassov in [4] over IFSs and so many operations were defined by him in [3] also. In [10] we define the above operations over intuitionistic fuzzy matrix and this work is the extension of our previous research deals with conjunction and disjunction operators in [10] and [11].

2. PRELIMINARIES:

Definition 2.1[2,6,7]:

Let a set $X = \{x_1, x_2, \dots, x_n\}$ be fixed, then an intuitionistic fuzzy set (IFS) can be defined as $A = \{(x_i, \mu_A(x_i), \nu_A(x_i)) / x_i \in X\}$ which assigns to each element x_i a membership degree $\mu_A(x_i)$ and a non membership degree $\nu_A(x_i)$ with the condition $0 \leq \mu_A(x_i) + \nu_A(x_i) \leq 1$ for all $x_i \in X$.

Definition 2.2.[6][7]:

The 2-tuple $\alpha(x_i) = (\mu_\alpha(x_i), \nu_\alpha(x_i))$ is called an intuitionistic fuzzy value (IFV), if $\mu_\alpha(x_i) \in [0,1]$, $\nu_\alpha(x_i) \in [0,1]$ and $\mu_\alpha(x_i) + \nu_\alpha(x_i) \leq 1$.

Definition 2.3[6]:

Let $A = (a_{ij})$ be a matrix of order $m \times n$, if all a_{ij} ($i = 1,2, \dots, m, j = 1,2, \dots, n$) are IFVs, then A is called an intuitionistic fuzzy matrix (IFM).

Definition 2.4[3]:

Let A and B be two IFSs and $\alpha \in [0,1]$ then

(i) $A \subseteq B$ if and only if for all $x_i \in X$, $\mu_A(x_i) \leq \mu_B(x_i)$ and $\nu_A(x_i) \geq \nu_B(x_i)$.

(ii) $\square A = \{(x_i, \mu_A(x_i), 1 - \mu_A(x_i)) / x_i \in X\}$

(iii) $C(A) = \{(x_i, K, L) / x_i \in X\}$ where $K = \max. \mu_A(x_i)$ and $L = \nu_A(x_i)$.

(iv) $D_\alpha(A) = \{(x_i, \mu_A(x_i) + \alpha \pi_A(x_i), \nu_A(x_i) + (1 - \alpha) \pi_A(x_i)) / x_i \in X\}$, where $\pi_A(x_i) = 1 - \mu_A(x_i) - \nu_A(x_i)$.

Definition 2.5[12,13,14]:

In [12] and [13] $Norm(A)$ was defined as follows. Let X be a nonempty universal set. The normalization of an IFS A denoted by $Norm(A)$ and defined by

$$Norm(A) = \left\{ \left\langle x, \frac{\mu_A(x)}{\sup \mu_A(x)}, \frac{\gamma_A(x) - \inf \gamma_A(x)}{1 - \inf \gamma_A(x)} \right\rangle \right\}.$$

Some remarks were given about the definition of $Norm(A)$ in [14] such that the above definition will be valid only if $\pi_A(x)$ is negligible. Here we consider that case only.

Definition 2.6[10,11]:

Consider that two elements in IFS $\langle x, x' \rangle$ and $\langle y, y' \rangle$ such that $0 \leq x+x' \leq 1$ and $0 \leq y+y' \leq 1$

Now the Lukasiewicz conjunction and disjunction operators are defined as follows

$$\langle x, x' \rangle \oplus \langle y, y' \rangle = \langle (x+y) \wedge 1, (x'+y'-1) \vee 0 \rangle$$

$$\langle x, x' \rangle \odot \langle y, y' \rangle = \langle (x+y-1) \vee 0, (x'+y') \wedge 1 \rangle$$

Now we define all the previous operations on IFMs as follows.

Let $A = [\langle a_{ij}, a'_{ij} \rangle]$ and $B = [\langle b_{ij}, b'_{ij} \rangle]$ be two IFMs of order $m \times n$. Then the ij^{th} element of all the operations are given below.

i. $A \oplus B = \langle (a_{ij} + b_{ij}) \wedge 1, (a'_{ij} + b'_{ij} - 1) \vee 0 \rangle$

ii. $A \odot B = \langle (a_{ij} + b_{ij} - 1) \vee 0, (a'_{ij} + b'_{ij}) \wedge 1 \rangle$

iii. $A \leq B \Rightarrow a_{ij} \leq b_{ij}$ and $a'_{ij} \geq b'_{ij}$

iv. $\square A = \langle a_{ij}, 1 - a_{ij} \rangle$

v. $D_\alpha(A) = \langle a_{ij} + \alpha a''_{ij}, a'_{ij} + (1 - \alpha) a''_{ij} \rangle$ Where $a''_{ij} = 1 - a_{ij} - a'_{ij}$, the value of indetermination.



- vi. An IFM $J = [(1,0)]$ for all entries is known as Universal Matrix and $O = [(0,1)]$ for all entries is known as Zero matrix.
- vii. $A^{[k+1]} = A^{[k]} \oplus A$ For all $k=1, 2, \dots$
- viii. $A^{(k+1)} = A^k \odot A$ For all $k=1, 2, \dots$
- ix. $Norm(A) = \langle \frac{a_{ij}}{\max a_{ij}}, \frac{a'_{ij} - \min a'_{ij}}{1 - \min a'_{ij}} \rangle$ for all i, j .
- x. If A is reflexive then $A \geq I_n$ where I_n is the identity IFM contains $\langle 1,0 \rangle$ when $i=j$ otherwise $\langle 0,1 \rangle$.
- xi. If A is irreflexive then $\langle a_{ij}, a'_{ij} \rangle = \langle 1,0 \rangle$ when $i = j$.
- xii. $c[A] = \langle K, L \rangle$ where $K = \max_{i,j} a_{ij}$ and $L = \min_{i,j} a'_{ij}$.

Here after F_{mn} means set of all Intuitionistic Fuzzy Matrices of order $m \times n$.

3. MAIN RESULTS:

Theorem 3.1:

For any two IFMs $A, B \in F_{mn}$, the following inequalities are true.

- i. $D_\alpha(A \oplus B) \leq D_\alpha(A) \oplus D_\alpha(B)$
- ii. $D_\alpha(A \odot B) \geq D_\alpha(A) \odot D_\alpha(B)$ for some $\alpha \in [0,1]$

Proof :

(i) From the Definition 2.5, we have

$$A \oplus B = \langle (a_{ij} + b_{ij}) \wedge 1, (a'_{ij} + b'_{ij} - 1) \vee 0 \rangle \text{ and}$$

$$D_\alpha[A \oplus B] = [(a_{ij} + b_{ij}) \wedge 1 + \alpha \{1 - (a_{ij} + b_{ij}) \wedge 1 - (a'_{ij} + b'_{ij} - 1) \vee 0\}, \\ (a'_{ij} + b'_{ij} - 1) \vee 0 + (1 - \alpha) \{1 - (a_{ij} + b_{ij}) \wedge 1 - (a'_{ij} + b'_{ij} - 1) \vee 0\}] \text{ ---3.1}$$

Now consider $D_\alpha(A) = [(a_{ij} + \alpha a''_{ij}, a'_{ij} + (1 - \alpha) a'_{ij})]$, where $a''_{ij} = 1 - a_{ij} - a'_{ij}$

$$D_\alpha(B) = [(b_{ij} + \alpha b''_{ij}, b'_{ij} + (1 - \alpha) b'_{ij})], \text{ where } b''_{ij} = 1 - b_{ij} - b'_{ij}$$

Using the above two equations we have

$$D_\alpha(A) \oplus D_\alpha(B) = [\{a_{ij} + b_{ij} + \alpha(a''_{ij} + b''_{ij})\} \wedge 1, \{a'_{ij} + b'_{ij} + (1 - \alpha)(a''_{ij} + b''_{ij}) - 1\} \vee 0] \\ \text{-----3.2}$$

Case (i)

If $a_{ij} + b_{ij} \geq 1$ and $a'_{ij} + b'_{ij} - 1 \leq 0$ then

$$3.1 \text{ becomes } D_\alpha[A \oplus B] = [1, 0]$$

$$3.2 \text{ becomes } D_\alpha(A) \oplus D_\alpha(B) = [1, 0]$$

Hence in this case $D_\alpha[A \oplus B] = D_\alpha(A) \oplus D_\alpha(B)$

Case (ii)

If $a_{ij} + b_{ij} \leq 1$ and $a'_{ij} + b'_{ij} - 1 \geq 0$ then

$$D_\alpha[A \oplus B] = [a_{ij} + b_{ij} + \alpha\{1 - (a_{ij} + b_{ij}) - (a'_{ij} + b'_{ij} - 1)\}, (a'_{ij} + b'_{ij} - 1) \\ + (1 - \alpha)\{1 - (a_{ij} + b_{ij}) - (a'_{ij} + b'_{ij} - 1)\}] \\ = [a_{ij} + b_{ij} + \alpha(a''_{ij} + b''_{ij}), (a'_{ij} + b'_{ij} - 1) + (1 - \alpha)(a''_{ij} + b''_{ij})] \\ = D_\alpha(A) \oplus D_\alpha(B)$$

Case (iii)

If $a_{ij} + b_{ij} \leq 1$ and $(a'_{ij} + b'_{ij} - 1) \leq 0$ then

$$D_\alpha[A \oplus B] = [a_{ij} + b_{ij} + \alpha\{1 - (a_{ij} + b_{ij})\}, (1 - \alpha)\{1 - (a_{ij} + b_{ij})\}]$$

And the membership value of

$$D_\alpha(A) \oplus D_\alpha(B) = [a_{ij} + b_{ij} + \alpha\{1 - (a_{ij} + b_{ij}) + 1 - (a'_{ij} + b'_{ij})\}] \wedge 1$$

Since $a_{ij}' + b_{ij}' \leq 1 \Rightarrow 1 - (a_{ij}' + b_{ij}')$ is positive which means the following

$$[1 - (a_{ij} + b_{ij})] + [1 - (a_{ij}' + b_{ij}')] \geq [1 - (a_{ij} + b_{ij})]$$

Gives membership value of $D_\alpha[A \oplus B] \leq$ membership value of $D_\alpha(A) \oplus D_\alpha(B)$

Now consider the nonmembership value of $D_\alpha(A) \oplus D_\alpha(B)$ as follows

$[(a_{ij}' + b_{ij}' - 1) + (1 - \alpha)\{1 - (a_{ij} + b_{ij}) + 1 - (a_{ij}' + b_{ij}')\}] \vee 0$ which can be written as

$$[1 - (a_{ij} + b_{ij}) - \alpha\{1 - (a_{ij} + b_{ij}) + 1 - (a_{ij}' + b_{ij}')\}] \vee 0 \leq [1 - (a_{ij} + b_{ij}) - \alpha\{1 - (a_{ij} + b_{ij})\}]$$

Which is the nonmembership value of $D_\alpha[A \oplus B]$

Therefore in this case $D_\alpha[A \oplus B] \leq D_\alpha(A) \oplus D_\alpha(B)$.

Case (iv)

If $a_{ij} + b_{ij} \geq 1$ and $(a_{ij}' + b_{ij}' - 1) \geq 0$ then A and B are not an IFMs.

Hence from all the above cases we conclude $D_\alpha[A \oplus B] \leq D_\alpha(A) \oplus D_\alpha(B)$.

(ii) From the Definition 2.5

$$A \odot B = [(a_{ij} + b_{ij} - 1) \vee 0, (a_{ij}' + b_{ij}') \wedge 1]$$

$$\text{Now } [A \odot B] = [(a_{ij} + b_{ij} - 1) \vee 0 + \alpha\{1 - (a_{ij} + b_{ij} - 1) \vee 0 - (a_{ij}' + b_{ij}') \wedge 1\}, (a_{ij}' + b_{ij}') \wedge 1 + (1 - \alpha)\{1 - (a_{ij} + b_{ij} - 1) \vee 0 - (a_{ij}' + b_{ij}') \wedge 1\}]$$

Case (i)

If $(a_{ij} + b_{ij} - 1) \geq 0$ and $(a_{ij}' + b_{ij}') \leq 1$ then

$$\begin{aligned} D_\alpha[A \odot B] &= [(a_{ij} + b_{ij} - 1) + \alpha\{a_{ij}'' + b_{ij}''\}, (a_{ij}' + b_{ij}') + (1 - \alpha)\{a_{ij}'' + b_{ij}''\}] \\ &\text{Where } a_{ij}'' = 1 - a_{ij} - a_{ij}' \text{ and } b_{ij}'' = 1 - b_{ij} - b_{ij}' \\ &= [a_{ij} + \alpha a_{ij}'' + b_{ij} + b_{ij}'' - 1, a_{ij}' + (1 - \alpha)a_{ij}'' + b_{ij}' + (1 - \alpha)b_{ij}''] \\ &= D_\alpha[A] \odot D_\alpha(B). \end{aligned}$$

Case (ii)

If $(a_{ij} + b_{ij} - 1) \leq 0$ and $(a_{ij}' + b_{ij}') \geq 1$ then $D_\alpha[A \odot B] = [0, 1]$ and

$$D_\alpha[A] \odot D_\alpha(B) = [(a_{ij} + \alpha a_{ij}'' + b_{ij} + \alpha b_{ij}'' - 1) \vee 0, \{a_{ij}' + (1 - \alpha)a_{ij}'' + b_{ij}' + (1 - \alpha)b_{ij}''\} \wedge 1]$$

Now consider the membership value of $D_\alpha[A] \odot D_\alpha(B)$ which can be written as

$$[(a_{ij} + b_{ij} - 1) + \alpha\{1 - (a_{ij} + b_{ij} - 1) - (a_{ij}' + b_{ij}')\}] \vee 0 = [(1 - \alpha)(a_{ij} + b_{ij} - 1) + \alpha - \alpha(a_{ij}' + b_{ij}')] \vee 0 = 0$$

Since $(1 - \alpha)(a_{ij} + b_{ij} - 1) \leq 0$ and $\alpha - \alpha(a_{ij}' + b_{ij}') \leq 0$

Similarly the nonmembership value $D_\alpha[A] \odot D_\alpha(B)$ can be considered as

$$a_{ij} + b_{ij}' + (1 - \alpha)(a_{ij}'' + b_{ij}'') \wedge 1 = 1 \text{ since } (1 - \alpha)(a_{ij}'' + b_{ij}'') \geq 0$$

Hence $D_\alpha[A] \odot D_\alpha(B) = [0, 1] = D_\alpha[A \odot B]$.

Case (iii)

If $(a_{ij} + b_{ij} - 1) \leq 0$ and $(a_{ij}' + b_{ij}') \leq 1$ then we have $D_\alpha[A \odot B] = [\alpha\{1 - (a_{ij}' + b_{ij}')\}, (a_{ij}' + b_{ij}') + (1 - \alpha)(1 - a_{ij}' - b_{ij}')]]$

$$\text{and } D_\alpha[A] \odot D_\alpha[B] = [\{(a_{ij} + b_{ij} - 1) + \alpha(a_{ij}'' + b_{ij}'')\} \vee 0, \{(a_{ij}' + b_{ij}') + (1 - \alpha)(a_{ij}'' + b_{ij}'')\} \wedge 1]$$

Now rewrite the membership value of the above as follows

$$[(a_{ij} + b_{ij} - 1)(1 - \alpha) + \alpha\{1 - (a_{ij}' + b_{ij}')\}] \vee 0 \leq \alpha\{1 - (a_{ij}' + b_{ij}')\} \text{ Which is the membership value of } D_\alpha[A \odot B].$$

Similarly we can prove the nonmembership value of $D_\alpha[A] \odot D_\alpha(B) \geq$ Nonmembership of $D_\alpha[A \odot B]$

Hence from all above cases $D_\alpha[A \odot B] \geq D_\alpha[A] \odot D_\alpha[B]$

Theorem 4.2:

For any two IFMs $A, B \in F_{mn}$, the following inequalities are valid

- (i) $c[A] \oplus c[B] \geq c[A \oplus B]$
- (ii) $c[A] \odot c[B] \geq c[A \odot B]$
- (iii) $c[D_\alpha(A)] \leq D_\alpha[c(A)]$

Proof :

(i) Let $A = [a_{ij}, a'_{ij}]$ and $B = [b_{ij}, b'_{ij}]$ be two IFMs of order $m \times n$.

From the Definition 2.6, $c[A] = [\max_{i,j} a_{ij}, \min_{i,j} a'_{ij}]$, $c[B] = [\max_{i,j} b_{ij}, \min_{i,j} b'_{ij}]$

Now $c[A] \oplus c[B] = [\{\max_{i,j} a_{ij} + \max_{i,j} b_{ij}\} \wedge 1, \{\min_{i,j} a'_{ij} + \min_{i,j} b'_{ij} - 1\} \vee 0]$

We know that $A \oplus B = [(a_{ij} + b_{ij}) \wedge 1, (a'_{ij} + b'_{ij} - 1) \vee 0]$

$c[A \oplus B] = [\max_{i,j} (a_{ij} + b_{ij}) \wedge 1, \min_{i,j} (a'_{ij} + b'_{ij} - 1) \vee 0]$

For all i, j it is clear that $a_{ij} + b_{ij} \leq \max_{i,j} a_{ij} + \max_{i,j} b_{ij}$

$$\Rightarrow \max_{i,j} (a_{ij} + b_{ij}) \wedge 1 \leq \{\max_{i,j} a_{ij} + \max_{i,j} b_{ij}\} \wedge 1 \dots \dots \dots 3.3$$

Similarly for all i, j we have $a'_{ij} + b'_{ij} \geq \min_{i,j} a'_{ij} + \min_{i,j} b'_{ij}$

$$\Rightarrow \min_{i,j} (a'_{ij} + b'_{ij} - 1) \vee 0 \geq \{\min_{i,j} a'_{ij} + \min_{i,j} b'_{ij} - 1\} \vee 0 \dots \dots \dots 3.4$$

From 3.3 and 3.4 we have the inequality $c[A] \oplus c[B] \geq c[A \oplus B]$

(ii) It is trivial from the above.

(iii) $D_\alpha[A] = [a_{ij} + \alpha a''_{ij}, a'_{ij} + (1 - \alpha) a'_{ij}]$

Now $c[D_\alpha(A)] = [\max_{i,j} (a_{ij} + \alpha a''_{ij}), \min_{i,j} (a'_{ij} + (1 - \alpha) a'_{ij})]$

And $D_\alpha[c(A)] = [\max_{i,j} a_{ij} + \alpha \{1 - \max_{i,j} a_{ij} - \min_{i,j} a'_{ij}\}, \min_{i,j} a'_{ij} + (1 - \alpha) \{1 - \max_{i,j} a_{ij} - \min_{i,j} a'_{ij}\}]$

for all i, j it is clear that $(a_{ij} + \alpha \{1 - a_{ij} - a'_{ij}\}) \leq \max_{i,j} a_{ij} + \alpha \{1 - \max_{i,j} a_{ij} - \min_{i,j} a'_{ij}\}$

Hence $\max_{i,j} (a_{ij} + \alpha \{1 - a_{ij} - a'_{ij}\}) \leq \max_{i,j} a_{ij} + \alpha \{1 - \max_{i,j} a_{ij} - \min_{i,j} a'_{ij}\}$

Similarly we can prove that

$$\min_{i,j} \{a'_{ij} + (1 - \alpha) (1 - a_{ij} - a'_{ij})\} \geq \min_{i,j} a'_{ij} + (1 - \alpha) \{1 - \max_{i,j} a_{ij} - \min_{i,j} a'_{ij}\}$$

From the above two inequalities we conclude that $c[D_\alpha(A)] \leq D_\alpha[c(A)]$.

Theorem 3.3:

For any IFM $A, B \in F_{mn}$, the following statements are valid

- (i) $A^{[k]} = U$ (the universal matrix) for some $k = 1, 2, \dots$
- (ii) $A^{(k)} = O$ (the zero matrix) for some $k = 1, 2, \dots$
- (iii) $(A \oplus B)^T = A^T \oplus B^T$
- (iv) $(A \odot B)^T = A^T \odot B^T$

Proof:

It is clear From the Definition 2.6.

Theorem 3.4:

If A and B are two IFMs then we have the following

- (i) $\square[Norm(A \oplus B)] \leq \square[Norm(A)] \oplus \square[Norm(B)]$
- (ii) If $(a_{ij}, a'_{ij}) = (1, 0)$ for atleast one i, j then $Norm[A] = A$
- (iii) If A is reflexive or irreflexive then it is same for $Norm[A]$ also.

Proof:

(i) Let $A = [a_{ij}, a'_{ij}]$ and $B = [b_{ij}, b'_{ij}]$, From the Definition 2.6

$$A \oplus B = [(a_{ij} + b_{ij}) \wedge 1, (a'_{ij} + b'_{ij} - 1) \vee 0], \text{Norm}[A] = \left[\frac{a_{ij}}{\max a_{ij}}, \frac{a'_{ij} - \min a'_{ij}}{1 - \min a'_{ij}} \right]$$

$$\text{Norm}[B] = \left[\frac{b_{ij}}{\max b_{ij}}, \frac{b'_{ij} - \min b'_{ij}}{1 - \min b'_{ij}} \right], \square \text{Norm}[A] = \left[\frac{a_{ij}}{\max a_{ij}}, 1 - \frac{a_{ij}}{\max a_{ij}} \right], \square \text{Norm}[B] = \left[\frac{b_{ij}}{\max b_{ij}}, 1 - \frac{b_{ij}}{\max b_{ij}} \right]$$

$$\text{Norm}[A \oplus B] = \left[\frac{(a_{ij} + b_{ij}) \wedge 1}{\max(a_{ij} + b_{ij}) \wedge 1}, \frac{(a'_{ij} + b'_{ij} - 1) \vee 0 - \min(a'_{ij} + b'_{ij} - 1) \vee 0}{1 - \min(a'_{ij} + b'_{ij} - 1) \vee 0} \right]$$

$$\square \text{Norm}[A \oplus B] = \left[\frac{(a_{ij} + b_{ij}) \wedge 1}{\max(a_{ij} + b_{ij}) \wedge 1}, 1 - \frac{(a_{ij} + b_{ij}) \wedge 1}{\max(a_{ij} + b_{ij}) \wedge 1} \right]$$

$$\text{Now } \square[\text{Norm}(A)] \oplus \square[\text{Norm}(B)] = \left[\left\{ \frac{a_{ij}}{\max a_{ij}} + \frac{b_{ij}}{\max b_{ij}} \right\} \wedge 1, \{1 - \frac{a_{ij}}{\max a_{ij}} + 1 - \frac{b_{ij}}{\max b_{ij}} - 1\} \vee 0 \right]$$

Since $\frac{a_{ij}}{\max a_{ij}} \geq \frac{a_{ij}}{\max(a_{ij} + b_{ij})}$ and $\frac{b_{ij}}{\max b_{ij}} \geq \frac{b_{ij}}{\max(a_{ij} + b_{ij})}$ gives $\frac{a_{ij}}{\max a_{ij}} + \frac{b_{ij}}{\max b_{ij}} \geq \frac{a_{ij} + b_{ij}}{\max(a_{ij} + b_{ij})} \dots \dots \dots 3.5$

$$\left\{ \frac{a_{ij}}{\max a_{ij}} + \frac{b_{ij}}{\max b_{ij}} \right\} \wedge 1 \geq \frac{(a_{ij} + b_{ij}) \wedge 1}{\max(a_{ij} + b_{ij}) \wedge 1} \dots 3.6$$

From equation 3.5 we have

$$1 - \left\{ \frac{a_{ij}}{\max a_{ij}} + \frac{b_{ij}}{\max b_{ij}} \right\} \leq 1 - \frac{a_{ij} + b_{ij}}{\max(a_{ij} + b_{ij})}$$

$$\cdot \left\{ 1 - \frac{a_{ij}}{\max a_{ij}} - \frac{b_{ij}}{\max b_{ij}} \right\} \vee 0 \leq 1 - \frac{(a_{ij} + b_{ij}) \wedge 1}{\max(a_{ij} + b_{ij}) \wedge 1} \dots \dots \dots 3.7$$

From equation 3.6 and 3.7 we prove $\square[\text{Norm}(A \oplus B)] \leq \square[\text{Norm}(A)] \oplus \square[\text{Norm}(B)]$.

(ii) If at least one $(a_{ij}, a'_{ij}) = (1, 0)$ then $\max a_{ij} = 1$ and $\min a'_{ij} = 0$

Therefore from the definition $\text{Norm}[A] = \left[\frac{a_{ij}}{1}, \frac{a'_{ij} - 0}{1 - 0} \right] = [a_{ij}, a'_{ij}] = A$

(iii) Suppose A is reflexive then $A \geq I_n$, that is $(a_{ij}, a'_{ij}) = (1, 0)$ when $i = j$

In this case clearly $\text{Norm}[A] = A$ which is reflexive from (ii)

Suppose A is irreflexive then $(a_{ij}, a'_{ij}) = (0, 1)$ when $i = j$

$$\text{when } i = j, \text{Norm}[A] = \left[\frac{a_{ii}}{\max a_{ij}}, \frac{a'_{ii} - \min a'_{ij}}{1 - \min a'_{ij}} \right] = \left[\frac{0}{\max a_{ij}}, \frac{1 - \min a'_{ij}}{1 - \min a'_{ij}} \right] = [0, 1]$$

Hence $\text{Norm}[A]$ is irreflexive.

4. REFERENCES:

- [1]. Zadeh L.A., Fuzzy Sets, Journal of Information and Control, Vol.8 (1965), 338-353
- [2]. Atanassov K., Intuitionistic Fuzzy Sets, VIITKR's session, Sofia, June 1983.
- [3]. Atanassov K., New Operations defined over the intuitionistic fuzzy sets, Fuzzy Sets and System.
- [4]. Atanassov K., Tcvetkov R., On Lukasiwicz's intuitionistic fuzzy conjunction and Disjuncz, Annals of "Informatics", Union of Scientists in Bulgaria, vol 3 (2010), 90-94.
- [5]. Meenakshi A.R and Gandhimathi, Intuitionistic fuzzy relational equations, Advances in fuzzy mathematics, vol 5(2010), 239-244.
- [6]. Zeshui.Xu., Intuitionistic fuzzy Aggregation and Clustering, Studies in Fuzziness and Soft Computing (2012), vol 279,159-190
- [7]. Zhang.Xu., A new method for ranking intuitionistic fuzzy values and its applications in multi attribute decision making, Fuzzy optim decision making (2012), vol 11,135-146.
- [8]. Zeshui Xu and Ronald R.Yager, Some aggregation operators based on IFS, International journal of general system, 35:4(2006), 417-433.
- [9]. Im Lee E.P and Park S.W., The Determinant of square IFMs, For East Journal of Mathematical Scienes, Vol.3 (5) (2001), 789-796.
- [10]. Sriram. S and Muthuraji.T., New operators for Intuitionistic Fuzzy Matrix, Presented in the International conference on Mathematical Modeling 2012, organized by Maths Department, Annamalai University.



- [11]. Muthuraji.T and Sriram.S., Commutative monoids and monoid homomorphism on Lukasiwicz conjunction and disjunction operators over IFMs., International Journal of Pure and Engg Maths, Vol 3, No 11(2015), 61-73.
- [12]. Ejegwa P.A at el., An overview on IFS, International journal of scientific and Tech Research vol 3, issue 3, March 2014,142-145.
- [13]. Supriya Kumar De, Ranjit Biswas, Akhil Ranjan Roy., Some operators on Intuitionistic fuzzy sets, Fuzzy sets and system 114(2000) 477-484.
- [14]. A.M.Ibrahim, P.A.Ejegwa., Remark on some operators of Intuitionistic Fuzzy Sets, International journal of science and technology, volume 2, No-1, January 2013,94-95.

