



## Common Fixed Point for Weakly Compatible Mappings on Dislocated Metric Spaces

Elida Hoxha

Department of Mathematics; Faculty of Natural Sciences  
University of Tirana; Tirana, Albania  
[hoxhaelida@yahoo.com](mailto:hoxhaelida@yahoo.com)

**Abstract:** In this paper we prove a common fixed point theorem for multi-valued and single-valued mappings in a dislocated metric space. The set of generalized  $\psi$ - $\varphi$  weak contractive mappings considered in this paper contains the family of generalized  $\varphi$ -weak contractive mappings as a proper subset. The theorem use weakly compatibility and  $\psi$ - $\varphi$ -weak contractivity condition. It extends the work of several authors.


**Keywords:** Dislocated metric spaces; common fixed point; generalized  $\psi$ - $\varphi$ -weak contractions; weakly compatible mappings; complete dislocated metric spaces.

### Academic Discipline And Sub-Disciplines

Mathematics, Functional Analysis, Fixed point theory.

### SUBJECT CLASSIFICATION

Fixed point theory.



## Council for Innovative Research

Peer Review Research Publishing System

**Journal:** Journal of Advances in Mathematics

Vol 4, No 1

[editor@cirworld.com](mailto:editor@cirworld.com)

[www.cirworld.com](http://www.cirworld.com), [member.cirworld.com](http://member.cirworld.com)



## 1. Introduction.

The notion of dislocated metric spaces ( $d$ -metric spaces) was introduced by Pascal Hitzler in [8]. The study of common fixed point mapping in dislocated metric space satisfying certain contractive conditions has been at the center of vigorous research activity..

Weakly contractive mappings in Hilbert spaces setting was first introduced by Alber and Guerr-Delabriere [1] in 1997. Rhoades showed that most results of [1] are still true for any Banach spaces and Bae considered these type of multi-valued mappings [2]. Kamran [13], Zhang and Song [17], Beg and Abbas [3], Bose and Roychowdhury considered some generalized versions of these mappings and proved some fixed point theorems.

After Jungck has introduced the notion of compatible and weakly compatible mappings for single valued maps [6] many authors have immediately extended the concepts to multi-valued maps.

The purpose of the present paper is to contribute in this field. We have considered a family of generalized  $\psi$ - $\phi$  weakly contractive mappings, which contains the class of generalized  $\phi$ -weakly contractive mappings and have proved a fixed point theorem for them when they are weakly compatible maps on dislocated metric spaces, which extends the work of several authors ([7], [4],[16],[12], [5])

## 2. Preliminaries.

For convenience we start with the following definitions, lemmas, and theorems.

**Definition 2.1.**[9] Let  $X$  be a non-empty and let  $d: X \times X \rightarrow [0, +\infty[$  be a function, called a distance function if for all  $x, y, z \in X$  satisfies:

$$d1: d(x, x) = 0$$

$$d2: d(x, y) = d(y, x) = 0 \Rightarrow x = y$$

$$d3: d(x, y) = d(y, x)$$

$$d4: d(x, y) \leq d(x, z) + d(z, y)$$

If  $d$  satisfies the condition  $d1 \square \square d4$ , then  $d$  is called a metric on  $X$ . If it satisfies the conditions  $d1$ ,  $d2$  and  $d4$  it is called a quasi-metric space. If  $d$  satisfies conditions  $d2$ ,  $d3$  and  $d4$  it is called a dislocated metric (or simply  $d$ -metric).

A nonempty set  $X$  with  $d$ -metric  $d$ , i. e.,  $\square X, d \square \square$  is called a dislocated metric space.

**Example.** Let be  $X = \mathbb{R}^+$  and the function  $d: X \times X \square \square \square \square \square \square \square \square \square \square$  where  $d \square \square x, y \square \square \square \square \square \square \square \square \square \square$ . Then  $\square X, d \square \square$  is a dislocated metric space.

**Definition 2.2** [9] A sequence  $\{x_n\}_{n \in \mathbb{N}}$  in a  $d$ -metric space  $\square X, d \square \square$  is called a Cauchy sequence if for all  $\square \square \square \square \square \square 0$ ,  $\square \square n_0 \square \square \mathbb{N}$  such that  $\square m, n \square \square n_0$ , we have  $d(x_m, x_n) \square \square \square \square$ .

**Definition 2.3** [9] A sequence in  $d$ -metric space converges with respect to  $d$ , if there exists  $x \in X$  such that  $\lim_{n \rightarrow \infty} d(x_n, x) = 0$ .

In this case  $x$  is called a  $d$ -limit of  $\{x_n\}_{n \in \mathbb{N}}$  and we write  $x_n \rightarrow x$ .

**Definition 2.4** [9] A  $d$ -metric space  $\square X, d \square \square$  is called complete if every Cauchy sequence in it is convergent with respect to  $d$ .

**Lemma 2.5** [9] Limits in a  $d$ -metric space are unique.

**Lemma 2.6.** [12] Let  $(X, d)$  be a  $d$ -metric space. Then

(A) if  $d(x, y) = 0$ , then  $d(x, x) = d(y, y) = 0$

(B) if  $\{x_n\}_{n \in \mathbb{N}}$  is a sequence such that  $\lim_{n \rightarrow \infty} d(x_n, x_{n+1}) = 0$ , then  $\lim_{n \rightarrow \infty} d(x_n, x_n) = \lim_{n \rightarrow \infty} d(x_{n+1}, x_{n+1}) = 0$

(C) if  $x \neq y$ , then  $d(x, y) > 0$

(D)  $d(x, x) \leq \frac{2}{n} \sum_{i=1}^n d(x, x_i)$  holds for all  $x, x_i \in X$ , where  $1 \leq i \leq n$ .

Let  $(X, d)$  be a  $d$ -metric space. We denote the family of all nonempty, bounded subset of  $X$  by  $B(X)$

**Definition 2.7.** Let  $A, B \in B(X)$ , then

$$H(A, B) = \max \left\{ \sup_{x \in A} d(x, B), \sup_{y \in B} d(y, A) \right\}$$



$$d(A, B) = \inf \{d(x, y) : x \in A, y \in B\}$$

$$\delta(A, B) = \sup \{d(x, y) : x \in A, y \in B\}$$

**Definition 2.8.** A sequence  $\{A_n\}$  of nonempty subset of  $X$  is said to be convergent towards a subset  $A$  of  $X$  if

i) each point  $a \in A$  is a limit of a convergent sequence  $\{a_n\}$ ,  $a_n \in A_n$  for  $n \in N$ .

ii) for arbitrary  $\varepsilon > 0$ , there is an integer  $m$  such that for  $n > m$ ,  $A_n \subseteq A_\varepsilon$ , where  $A_\varepsilon = \{x \in X : \exists a \in A / d(x, a) < \varepsilon\}$ ,

$A$  is then said to be the limit of the sequence  $\{A_n\}$ .

**Lemma 2.9.** Let  $\{A_n\}$ ,  $\{B_n\}$  be sequences in  $B(X)$  converging respectively to  $A$  and  $B$  in  $B(X)$ . Then the sequences of numbers  $\{H(A_n, B_n)\}$ ,  $\{d(A_n, B_n)\}$  and  $\{\delta(A_n, B_n)\}$  converge to  $H(A, B)$ ,  $d(A, B)$  and  $\delta(A, B)$ , respectively.

**Lemma 2.10.** Let  $\{A_n\}$  be sequences in  $B(X)$  and  $y \in X$  such that  $\delta(A_n, y) \rightarrow 0$ . Then the sequence  $\{A_n\}$  converges to the set  $\{y\}$  in  $B(X)$ .

**Definition 2.11.** [10] The self maps  $f, g$  of  $X$  are compatible iff  $\lim_{n \rightarrow \infty} d(fgx_n, gfx_n) = 0$ , whenever  $\{x_n\}$  is a sequence in  $X$  such that  $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = t$  for some  $t \in X$ .

Several authors ([11], [15]) extend the Definition 2.3 by introducing the compatibility of set-valued mappings as below.

**Definition 2.12.** The mappings  $F : X \rightarrow B(X)$  and  $f : X \rightarrow X$  are  $\delta$ -compatible iff  $\lim_{n \rightarrow \infty} \delta(fFx_n, Ffx_n) = 0$ ,

whenever  $\{x_n\}$  is a sequence in  $X$  such that  $fFx_n \in B(X)$ ,  $\lim_{n \rightarrow \infty} Fx_n = \{t\}$  and  $\lim_{n \rightarrow \infty} fx_n = t$  for some  $t \in X$ .

**Definition 2.13** [10] The mappings  $F : X \rightarrow B(X)$  and  $f : X \rightarrow X$  are weakly compatible (or sub compatible) if they commute at coincidence points that is  $\{t \in X / Ft = \{ft\}\} \subseteq \{t \in X / Fft = fFt\}$ .

**Definition 2.14.** Two set-valued mappings  $F, G : X \rightarrow B(X)$ , are called generalized  $\varphi$ -weak contractions if there exist a map

$\varphi : [0, +\infty[ \rightarrow [0, +\infty[$  with  $\varphi(t) > 0$  for all  $t > 0$  and  $\varphi(0) = 0$ , such that

$$H(Fx, Gy) \leq M(x, y) - \varphi(M(x, y)) \text{ for all } x, y \in X, \text{ where}$$

$$M(x, y) = \max \left\{ d(x, y), d(x, Fx), d(y, Gy), \frac{d(x, Gy) + d(y, Fx)}{2} \right\}.$$

**Definition 2.15.** Two set-valued mappings  $F, G : X \rightarrow B(X)$ , are called generalized  $\psi$ - $\varphi$ -weak contractions if for all  $x, y \in X$ ,

$$\psi(H(Fx, Gy)) \leq \psi(M(x, y)) - \varphi(M(x, y)),$$

where  $M(x, y) = \max \left\{ d(x, y), d(x, Fx), d(y, Gy), \frac{d(x, Gy) + d(y, Fx)}{2} \right\}$  and  $\psi, \varphi : [0, +\infty[ \rightarrow [0, +\infty[$ ,  $\psi$  is non-decreasing and continuous,  $\varphi$  is lower semi continuous function such that,  $\psi(t) > 0$ ,  $\varphi(t) > 0$  for all  $t > 0$  and  $\psi(0) = \varphi(0) = 0$ .

The aim of this work is to prove a theorem on the common fixed point for two multi-valued and two single valued mappings not necessary continuous, where the multi-valued mappings are generalized  $\psi$ - $\varphi$ -weak contractions on dislocated metric spaces.

### 3. MAIN RESULTS.

**Theorem 3.1.** Let  $F, G : X \rightarrow B(X)$  be mappings, where  $(X, d)$  is a complete dislocated metric space and let  $I, J : X \rightarrow X$  be self mappings. Suppose that

(1)  $\psi(H(Fx, Gy)) \leq \psi(N(x, y)) - \varphi(N(x, y))$ , where

$$N(x, y) = \max \left\{ d(Ix, Jy), d(Ix, Fx), d(Jy, Gy), \frac{d(Ix, Gy) + d(Jy, Fx)}{a} \right\}$$



for all  $x, y \in X$ ,  $3 < a \leq 4$  and  $\psi, \varphi: [0, +\infty[ \rightarrow [0, +\infty[$ ,  $\psi$  is non-decreasing and continuous,  $\varphi$  is lower semi continuous function,  $\psi(t) > 0$ ,  $\varphi(t) > 0$  for all  $t > 0$  and  $\psi(0) = \varphi(0) = 0$ .

(2)  $\cup G(X) \subseteq I(X)$ ,  $\cup F(X) \subseteq J(X)$  and either  $I(X)$  or  $J(X)$  is closed

(3) the pairs of mappings  $\{F, I\}$  and  $\{G, J\}$  are weakly compatible,

then  $F, G, I, J$  have a unique common fixed point  $u \in X$ . Moreover,  $Fu = Gu = \{u\}$ .

**Proof.** Let  $x_0$  be an arbitrary point of  $X$ . By the condition (2) there exists  $x_1 \in X$  such that  $Jx_1 \in Fx_0$ . Furthermore, for this point  $x_1$  we can choose  $x_2 \in X$  such that  $Ix_2 \in Gx_1$  and  $d(Jx_1, Ix_2) \leq H(Fx_0, Gx_1)$

By the condition (1) of Theorem 3.1 we have  $\psi(d(Jx_1, Ix_2)) \leq \psi(H(Fx_0, Gx_1)) \leq \psi(N(x_0, x_1)) - \varphi(N(x_0, x_1)) \leq \psi(N(x_0, x_1))$ .

Continuing this process, we can define inductively the sequence  $\{x_n\}$  as follows:

$$Jx_{2n+1} \in Fx_{2n}, \quad Ix_{2n} \in Gx_{2n-1} \text{ for } n \in \mathbb{N} \text{ and } d(Ix_{2n}, Jx_{2n+1}) \leq H(Fx_{2n}, Gx_{2n-1}).$$

Hence as  $\psi$  is monotonically increasing and (1) we have

$$\psi(d(Ix_{2n}, Jx_{2n+1})) \leq \psi(H(Fx_{2n}, Gx_{2n-1})) \leq \psi(N(x_{2n}, x_{2n-1})) - \varphi(N(x_{2n}, x_{2n-1})) \leq \psi(N(x_{2n}, x_{2n-1})) \quad (4)$$

and  $d(Ix_{2n}, Jx_{2n+1}) \leq N(x_{2n-1}, x_{2n})$ .

$$\begin{aligned} N(x_{2n-1}, x_{2n}) &= \max \left\{ d(Ix_{2n}, Jx_{2n-1}), d(Ix_{2n}, Fx_{2n}), d(Jx_{2n-1}, Gx_{2n-1}), \frac{d(Ix_{2n}, Gx_{2n-1}) + d(Jx_{2n-1}, Fx_{2n})}{a} \right\} \\ &\leq \max \left\{ d(Ix_{2n}, Jx_{2n-1}), d(Ix_{2n}, Jx_{2n+1}), d(Jx_{2n-1}, Ix_{2n}), \frac{d(Ix_{2n}, Ix_{2n}) + d(Jx_{2n-1}, Jx_{2n+1})}{a} \right\} \end{aligned} \quad (5)$$

On the other hand, from lemma 2.6 (D) we have

$$d(Ix_{2n}, Ix_{2n}) \leq d(Ix_{2n}, Jx_{2n-1}) + d(Ix_{2n}, Jx_{2n+1})$$

and by definition 2.1 (d4) we have ,

$$d(Jx_{2n-1}, Jx_{2n+1}) \leq d(Ix_{2n}, Jx_{2n-1}) + d(Ix_{2n}, Jx_{2n+1})$$

So,

$$\frac{d(Ix_{2n}, Ix_{2n}) + d(Jx_{2n-1}, Jx_{2n+1})}{a} \leq \frac{2 [d(Ix_{2n}, Jx_{2n-1}) + d(Ix_{2n}, Jx_{2n+1})]}{a}$$

and for  $3 < a \leq 4$  we have

$$N(x_{2n-1}, x_{2n}) \leq \max \{d(Ix_{2n}, Jx_{2n-1}), d(Ix_{2n}, Jx_{2n+1})\}.$$

Now, if  $\max \{d(Ix_{2n}, Jx_{2n-1}), d(Ix_{2n}, Jx_{2n+1})\} = d(Ix_{2n}, Jx_{2n+1})$ , by (4) we have

$$\psi(d(Ix_{2n}, Jx_{2n+1})) \leq \psi(N(x_{2n}, x_{2n-1})) - \varphi(N(x_{2n}, x_{2n-1})) \leq \psi(d(Ix_{2n}, Jx_{2n+1})) - \varphi(d(Ix_{2n}, Jx_{2n+1})), \text{ a contradiction.}$$

Hence, we have

$$N(x_{2n-1}, x_{2n}) \leq \max \{d(Ix_{2n}, Jx_{2n-1}), d(Ix_{2n}, Jx_{2n+1})\} = d(Ix_{2n}, Jx_{2n-1}),$$

So,

$$d(Ix_{2n}, Jx_{2n+1}) \leq N(x_{2n-1}, x_{2n}) \leq d(Ix_{2n}, Jx_{2n-1}) \quad (6)$$

Also

$$d(Ix_{2n+2}, Jx_{2n+1}) \leq H(Fx_{2n+2}, Gx_{2n+1}) \text{ and}$$

$$\psi(d(Ix_{2n+2}, Jx_{2n+1})) \leq \psi(H(Fx_{2n+2}, Gx_{2n+1})) \leq \psi(N(x_{2n}, x_{2n+1})) - \varphi(N(x_{2n}, x_{2n+1})) \leq \psi(N(x_{2n}, x_{2n+1})) \quad (7)$$

so  $d(Ix_{2n+2}, Jx_{2n+1}) \leq N(x_{2n+1}, x_{2n})$ .



In the same way, we have

$$d(Ix_{2n+2}, Jx_{2n+1}) \leq N(x_{2n}, x_{2n+1}) \leq \max\{d(Ix_{2n}, Jx_{2n+1}), d(Jx_{2n+1}, Ix_{2n+2})\} = d(Ix_{2n}, Jx_{2n+1}) . \tag{8}$$

Let put for convenience,

$$y_k = \begin{cases} Ix_{2n} & k = 2n \\ Jx_{2n-1} & k = 2n - 1 . \end{cases}$$

By (6) and (8), we conclude that  $d(y_{k+1}, y_k) \leq N(y_k, y_{k-1}) \leq d(y_k, y_{k-1})$ , for  $k \in N$ .

It implies that the sequence  $\{d(y_k, y_{k+1})\}$  is monotone non-increasing and bounded below. So there exists  $l \geq 0$  such that

$$\lim_{k \rightarrow \infty} d(y_k, y_{k+1}) = \lim_{k \rightarrow \infty} N(y_k, y_{k+1}) = l .$$

Since  $\psi$  is continuous and  $\phi$  is lower semi-continuous,

$$\psi(l) = \lim_{k \rightarrow \infty} \psi(N(y_k, y_{k+1})) \quad \phi(l) \leq \liminf_{k \rightarrow \infty} \phi(N(y_k, y_{k+1})) .$$

By (4), (7) and the property of the function  $\psi$  and  $\phi$  we conclude that

$$\psi(l) \leq \psi(l) - \phi(l) \text{ i.e. } \phi(l) = 0 \text{ and } l = 0, \text{ and so } \lim_{k \rightarrow \infty} d(y_k, y_{k+1}) = \lim_{k \rightarrow \infty} N(y_k, y_{k+1}) = 0 . \tag{9}$$

Next we show that  $\{y_k\}$  is a Cauchy sequence.

Let,  $C_k = \sup\{d(y_i, y_j) : i, j > k\}$ .

Obviously  $\{C_k\}$  is decreasing. So there exists  $C \geq 0$  such that  $\lim_{k \rightarrow \infty} C_k = C$ .

For every  $p \in N$ , there exist  $k(p), s(p) \in N$  such that  $k(p), s(p) \geq p$  and

$$C_p - \frac{1}{p} \leq d(y_{k(p)}, y_{s(p)}) \leq C_p .$$

So,  $\lim_{p \rightarrow \infty} d(y_{k(p)}, y_{s(p)}) = C$ .

Let us to prove that  $C = 0$ .

**Case 1:** If  $k(p)$  is even and  $s(p)$  is odd, so  $k(p) = 2q$  and  $s(p) = 2t - 1$ , then

$$d(y_{k(p)+1}, y_{s(p)+1}) = d(y_{2q+1}, y_{2t}) = d(Jx_{2q+1}, Ix_{2t}) \leq H(Fx_{2q}, Gx_{2t-1})$$

By (1), we have

$$\psi(d(y_{k(p)+1}, y_{s(p)+1})) = \psi(d(y_{2q+1}, y_{2t})) = \psi(d(Jx_{2q+1}, Ix_{2t})) \leq \psi(H(Fx_{2q}, Gx_{2t-1})) \leq \psi(N(x_{2q}, x_{2t-1})) - \phi(N(x_{2q}, x_{2t-1})) \tag{10}$$

So,

$$\begin{aligned} d(y_{k(p)+1}, y_{s(p)+1}) &\leq N(x_{2q}, x_{2t-1}) = N(y_{k(p)}, y_{s(p)}) \\ N(x_{2q}, x_{2t-1}) &= \max\left\{d(Ix_{2q}, Jx_{2t-1}), d(Ix_{2q}, Fx_{2q}), d(Jx_{2t-1}, Gx_{2t-1}), \frac{d(Ix_{2q}, Gx_{2t-1}) + d(Jx_{2t-1}, Fx_{2q})}{a}\right\} \\ &\leq \max\left\{d(Ix_{2q}, Jx_{2t-1}), d(Ix_{2q}, Jx_{2q+1}), d(Jx_{2t-1}, Ix_{2t}), \frac{d(Ix_{2q}, Ix_{2t}) + d(Jx_{2t-1}, Jx_{2q+1})}{a}\right\} \\ &= \max\left\{d(y_{k(p)}, y_{s(p)}), d(y_{k(p)}, y_{k(p)+1}), d(y_{s(p)}, y_{s(p)+1}), \frac{d(y_{k(p)}, y_{s(p)+1}) + d(y_{s(p)}, y_{k(p)+1})}{a}\right\} \end{aligned} \tag{11}$$

As  $p \rightarrow +\infty$  in inequality (11), we have  $\lim_{p \rightarrow \infty} N(x_{2q}, x_{2t-1}) = \lim_{p \rightarrow \infty} N(y_{k(p)}, y_{s(p)}) = \max\{C, 0, 0, \frac{C+C}{a}\} = C$  .  $\tag{12}$

Taking  $p \rightarrow +\infty$  in inequality (10), since  $\psi$  is continuous and  $\phi$  is lower semi-continuous and (9) and (12) hold, we have  $\psi(C) \leq \psi(C) - \phi(C)$ . Hence  $\phi(C) = 0$  and so  $C = 0$ .



So,  $\lim_{p \rightarrow \infty} d(y_{k(p)}, y_{s(p)}) = 0$ , for  $k(p)$  is even and  $s(p)$  is odd. (13)

Similarly, if  $k(p)$  is odd and  $s(p)$  is even.

**Case 2:** If  $k(p)$  and  $s(p)$  is even, so  $k(p) = 2q$  and  $s(p) = 2t$ , then

$$d(y_{k(p)}, y_{s(p)}) = d(y_{2q}, y_{2t}) = d(Ix_{2q}, Ix_{2t}) \leq d((Ix_{2q}, Jx_{2q+1}) + d(Jx_{2q+1}, Ix_{2t})) = d(y_{k(p)}, y_{k(p)+1}) + d(y_{k(p)+1}, y_{s(p)}),$$

Taking  $p \rightarrow +\infty$  in this inequality, since (9) and (13) hold, we have  $\lim_{p \rightarrow \infty} d(y_{k(p)}, y_{s(p)}) = 0$

Similarly, if  $k(p)$  and  $s(p)$  is odd.

Therefore,  $\{y_k\}$  is a Cauchy sequence.

Since  $(X, d)$  is  $d$ -complete, there exists  $u \in X$  such that  $\lim_{k \rightarrow \infty} y_k = u$ .

This implies  $\lim_{n \rightarrow \infty} Ix_{2n} = \lim_{n \rightarrow \infty} Jx_{2n-1} = u$  and sequences  $\{Fx_{2n}\}, \{Gx_{2n+1}\}$  converge at  $\{u\}$ . (14)

Suppose now that the set  $J(X)$  is closed. Then there is, by condition (2), a point  $v \in X$  such that  $Jv = u$ .

Using (1) we obtain

$$\psi(d(Fx_{2n}, Gv)) \leq \psi(H(Fx_{2n}, Gv)) \leq \psi(N(x_{2n}, v)) - \varphi(N(x_{2n}, v)). \quad (15)$$

$$N(x_{2n}, v) = \max \left\{ d(Ix_{2n}, Jv), d(Ix_{2n}, Fx_{2n}), d(Jv, Gv), \frac{d(Ix_{2n}, Gv) + d(Jv, Fx_{2n})}{a} \right\}$$

$$= \max \left\{ d(Ix_{2n}, u), d(Ix_{2n}, Fx_{2n}), d(u, Gv), \frac{d(Ix_{2n}, Gv) + d(u, Fx_{2n})}{a} \right\}$$

Letting  $n \rightarrow \infty$  in the above inequality and by (14) we get

$$\lim_{n \rightarrow \infty} N(x_{2n}, v) = \max \left\{ d(u, u), d(u, \{u\}), d(u, Gv), \frac{d(u, \{u\}) + d(u, Gv)}{2} \right\} =$$

$$\max \left\{ d(u, u), d(u, Gv), \frac{d(u, Gv) + d(u, Gv) + d(u, Gv)}{a} \right\} = \max \{d(u, u), d(u, Gv)\} \quad (16)$$

Taking  $n \rightarrow \infty$  in (5) and by (14), we have  $0 = \max \{d(u, u), d(u, u)\} = d(u, u)$  and by (16) we have

$$\lim_{n \rightarrow \infty} N(x_{2n}, v) = d(u, Gv), \quad (17)$$

Letting  $n \rightarrow \infty$  in (15) and by (17) we get, so  $\varphi(d(u, Gv)) = 0$  and  $d(u, Gv) = 0$ .

Hence  $Gv = \{u\} = \{Jv\}$ . But the pair of maps  $\{G, J\}$  are weakly compatible, thus  $GJv = JGv$ , i.e.  $Gu = \{Ju\}$ . (18)

Now we show that  $u$  is a fixed point for  $G$  and for  $J$ .

$$N(x_{2n}, u) = \max \left\{ d(Ix_{2n}, Ju), d(Ix_{2n}, Fx_{2n}), d(Ju, Gu), \frac{d(Ix_{2n}, Gu) + d(Ju, Fx_{2n})}{a} \right\}.$$

Letting  $n \rightarrow \infty$  in the above inequality and by (14), (18) and equality  $0 = d(u, u)$  we get

$$\lim_{n \rightarrow \infty} N(x_{2n}, u) = \max \left\{ d(u, Gu), d(u, \{u\}), d(Ju, \{Ju\}), \frac{d(u, Gu) + d(u, Gu)}{a} \right\} = d(u, Gu). \quad (19)$$

By (1) we have  $\psi(d(Jx_{2n+1}, Gu)) \leq \psi(H(Fx_{2n}, Gu)) \leq \psi(N(x_{2n}, u)) - \varphi(N(x_{2n}, u))$

Letting  $n \rightarrow \infty$  in this inequality and by (19) we get  $\psi(d(u, Gu)) \leq \psi(H(u, Gu)) \leq \psi(d(u, Gu)) - \varphi(d(u, Gu))$ , so  $\varphi(d(u, Gu)) = 0$  and  $d(u, Gu) = 0$ .

Thus we have  $Gu = \{u\} = \{Ju\}$  and  $u$  is a fixed point for  $G$  and  $J$ .

Now,  $\cup G(X) \subseteq I(X)$  implies that exists  $w \in X$  such that  $Gu = \{Iw\}$ . Hence  $\{u\} = Gu = \{Ju\} = \{Iw\}$  and  $H(Fw, u) = H(Fw, Gu)$

Using (1) we get

$$\psi(H(Fw, u)) = \psi(H(Fw, Gu)) \leq \psi(N(w, u)) - \varphi(N(w, u))$$



$$\text{and } N(w,u) = \max \left\{ d(Iw, Ju), d(Iw, Fw), d(Ju, Gu), \frac{d(Iw, Gu) + d(Ju, Fw)}{a} \right\} = d(u, Fw).$$

So,  $\psi(H(Fw,u)) \leq \psi(d(Fw,u)) - \varphi(d(Fw,u))$  and  $\varphi(d(Fw,u)) = 0$  and  $d(Fw,u) = 0$  and  $\{u\} = Fw = \{Iw\}$ .

Hence,

$$\{u\} = Gu = \{Ju\} = Fw = \{Iw\}.$$

But the pair of maps  $\{F, I\}$  are weakly compatible, thus  $FIw = IFw$ , i.e.  $Fu = \{Iu\}$ .

Moreover,

$$\psi(d(Fu,u)) \leq \psi(H(Fu,u)) = \psi(H(Fu,Gu)) \leq \psi(N(u,u)) - \varphi(N(u,u))$$

$$\text{and } N(u,u) = \max \left\{ d(Iu, Ju), d(Iu, Fu), d(Ju, Gu), \frac{d(Iu, Gu) + d(Ju, Fu)}{a} \right\} = d(u, Fu).$$

So,  $\psi(d(Fu,u)) \leq \psi(H(Fu,u)) \leq \psi(d(Fu,u)) - \varphi(d(Fu,u))$  and  $\varphi(d(Fu,u)) = 0$  and  $d(Fu,u) = 0$  and

$\{u\} = Gu = \{Ju\} = Fu = \{Iu\}$  i.e.  $u$  is a common fixed point for  $F, G, I, J$ .

Similarly, one can reach the above fact by assuming that  $I(X)$  is closed.

Furthermore, we can prove that  $u$  is unique. In fact, if  $u$  and  $v$  are two common fixed points for  $F, G, I, J$ , then by (1) we have

$$\psi(d(u,v)) \leq \psi(H(u,v)) = \psi(H(Fu, Gv)) \leq \psi(N(u,v)) - \varphi(N(u,v))$$

$$\text{and } N(u,v) = \max \left\{ d(Iu, Jv), d(Iu, Fu), d(Jv, Gv), \frac{d(Iu, Gv) + d(Jv, Fu)}{a} \right\} = d(u,v)$$

So,  $\psi(d(u,v)) \leq \psi(d(u,v)) - \varphi(d(u,v))$  and  $\varphi(d(u,v)) = 0$  and  $d(u,v) = 0$  and  $u=v$ . The proof is complete.

**Remark.** The above theorem extends Theorem 3.1 of [7]. If we take  $\psi(t)=t$  for all  $t \geq 0$  then the above theorem reduces to theorem 3.1. of [7].

**Theorem 3.2.** Let  $F, G : X \rightarrow B(X)$  be mappings, where  $(X, d)$  is a complete dislocated metric space and let  $J : X \rightarrow X$  be self mappings. Suppose that

(1)  $\psi(H(Fx, Gy)) \leq \psi(N(x, y)) - \varphi(N(x, y))$ , where

$$N(x, y) = \max \left\{ d(Jx, Jy), d(Jx, Fx), d(Jy, Gy), \frac{d(Jx, Gy) + d(Jy, Fx)}{a} \right\}$$

for all  $x, y \in X$ ,  $3 < a \leq 4$  and  $\psi, \varphi : [0, +\infty[ \rightarrow [0, +\infty[$ ,  $\psi$  is non-decreasing and continuous,  $\varphi$  is lower semi continuous function,  $\psi(t) > 0$ ,  $\varphi(t) > 0$  for all  $t > 0$  and  $\psi(0) = \varphi(0) = 0$ .

(2)  $\cup G(X) \subseteq J(X)$ ,  $\cup F(X) \subseteq J(X)$  and  $J(X)$  is closed

(3) the pairs of mappings  $\{F, J\}$  and  $\{G, J\}$  are weakly compatible,

then  $F, G, J$  have a unique common fixed point  $u \in X$ .

**Proof.**  $F, G, J$  complete the condition in Theorem 3.1 by taking  $I = J$ . Hence  $F, G, J$  have a common fixed point  $u \in X$

**Remark.** This theorem extends the Ciric and Ume theorem (Theorem 2.1.[5])

**Theorem 3.3.** Let  $(X, d)$  be a complete d-metric space, and let  $F, G : X \rightarrow B(X)$  be two multi-valued mappings such that for all  $x, y \in X$

(1)  $\psi(H(Fx, Gy)) \leq \psi(M(x, y)) - \varphi(M(x, y))$

$$\text{where } M(x, y) = \max \left\{ d(x, y), d(x, Fx), d(y, Gy), \frac{d(x, Gy) + d(y, Fx)}{a} \right\}$$

for all  $x, y \in X$ ,  $3 < a \leq 4$  and  $\psi, \varphi : [0, +\infty[ \rightarrow [0, +\infty[$ ,  $\psi$  is non-decreasing and continuous,  $\varphi$  is lower semi continuous function,  $\psi(t) > 0$ ,  $\varphi(t) > 0$  for all  $t > 0$  and  $\psi(0) = \varphi(0) = 0$ .

then  $F, G$ , have a unique common fixed point  $u \in X$ . Moreover,  $Fu = Gu = \{u\}$ .



**Proof.** By taking  $I = J = I_X$  we have  $\cup G(X) \subseteq I(X) = X$ ,  $\cup F(X) \subseteq J(X) = X$  and  $X$  is closed. Hence  $F, G, I, J$  complete the condition in Theorem 3.1 and  $F, G$  have a common fixed point  $u \in X$ .

**Remark.** This theorem extends the Bose and Roychowdhury theorem (Theorem 51. [4]).

**Theorem 3.4.** Let  $(X, d)$  be a complete d-metric space, and let  $F, G, I, J : X \rightarrow X$  four mappings such that for all  $x, y \in X$

$$(1) \psi(H(Fx, Gy)) \leq \psi(N(x, y)) - \varphi(N(x, y))$$

$$\text{where } N(x, y) = \max \left\{ d(Ix, Jy), d(Ix, Fx), d(Jy, Gy), \frac{d(Ix, Gy) + d(Jy, Fx)}{a} \right\}$$

for all  $x, y \in X$ ,  $3 < a \leq 4$  and  $\psi, \varphi : [0, +\infty[ \rightarrow [0, +\infty[$ ,  $\psi$  is non-decreasing and continuous,  $\varphi$  is lower semi continuous function,  $\psi(t) > 0$ ,  $\varphi(t) > 0$  for all  $t > 0$  and  $\psi(0) = \varphi(0) = 0$ .

(2)  $G(X) \subseteq I(X)$ ,  $F(X) \subseteq J(X)$  and either  $I(X)$  or  $J(X)$  is closed

(3) the pairs of mappings  $\{F, I\}$  and  $\{G, J\}$  are weakly compatible,

then  $F, G, I, J$  have a unique common fixed point  $u \in X$ .

**Proof.** The proof of this theorem is similar to the Theorem 3.1 for  $Gx = \{Gx\}$ ,  $Fx = \{Fx\}$ .

**Remark.** If we take  $\psi(t) = t$  and  $\varphi(t) = (1 - \alpha)t$  for all  $t \geq 0$  then the above theorem reduces to theorem 3.1. of Kumari and Kumar and Sarma theorem (Theorem 3.1. [14]).

**Theorem 3.5.** Let  $(X, d)$  be a complete d-metric space, and let  $F, G : X \rightarrow X$  two mappings such that for all  $x, y \in X$

$$(1) \psi(H(Fx, Gy)) \leq \psi(M(x, y)) - \varphi(M(x, y))$$

$$\text{where } M(x, y) = \max \left\{ d(x, y), d(x, Fx), d(y, Gy), \frac{d(x, Gy) + d(y, Fx)}{a} \right\}$$

for all  $x, y \in X$ ,  $3 < a \leq 4$  and  $\psi, \varphi : [0, +\infty[ \rightarrow [0, +\infty[$ ,  $\psi$  is non-decreasing and continuous,  $\varphi$  is lower semi continuous function,  $\psi(t) > 0$ ,  $\varphi(t) > 0$  for all  $t > 0$  and  $\psi(0) = \varphi(0) = 0$ .

then  $F, G$  have a unique common fixed point  $u \in X$ .

**Proof.** The proof of this theorem is similar to the Theorem 3.1 for  $Gx = \{Gx\}$ ,  $Fx = \{Fx\}$  and  $I = J = I_X$

**Remark.** This theorem extends the Shrivastava and Ansari and Sharma theorem (Theorem 3.7.[16])

**Theorem 3.6.** Let  $(X, d)$  be a complete d-metric spaces and  $F : X \rightarrow X$  a self-map, such that for all  $x, y \in X$

$$(1) \psi(H(Fx, Fy)) \leq \psi(M(x, y)) - \varphi(M(x, y))$$

$$\text{where } M(x, y) = \max \left\{ d(x, y), d(x, Fx), d(y, Fy), \frac{d(x, Fy) + d(y, Fx)}{a} \right\}$$

for all  $x, y \in X$ ,  $3 < a \leq 4$  and  $\psi, \varphi : [0, +\infty[ \rightarrow [0, +\infty[$ ,  $\psi$  is non-decreasing and continuous,  $\varphi$  is lower semi continuous function,  $\psi(t) > 0$ ,  $\varphi(t) > 0$  for all  $t > 0$  and  $\psi(0) = \varphi(0) = 0$ .

then  $F$  have a unique common fixed point  $u \in X$ .

**Proof.** The proof of this theorem is similar to the Theorem 3.1 for  $Fx = Gx = \{Fx\}$  and  $I = J = I_X$ .

**Remark.** Theorem 3.5 extends the Karapinar and Salimi theorem (Theorem 1.9. [12]).

## REFERENCES

- [1] Alber, Y.I and Gueer-Delabriere, S., 1997, New Results in Operator Theory and its Applications. The Israel M. Glazman Memorial Volume, Birkhauser Verlag,. MR 1478463 (98j:47126). zbl 0870.00030.
- [2] Bae, J.S, 2003, Fixed point theorems for weakly contractive multivalued mapps, J.Math. Anal. Appl. 284, 690-697. Mr 1998661 (2004f:54038). zbl 1033.47038.
- [3] Beg, I. and Abbas, M., 2006, Coincidence point and in variant approximation for mappings satisfying generalized weak contractive condition, Fixed Point Theory and Application, Volume 2006, Article ID74503, 1-7. MR0614886 (82m:47038). zbl 0475.47043.





- [4] Bose, R.K. and Roychowdhury, M.K., 2009, Fixed point theorems for generalized weakly contractive mappings, *Surveys in Mathematics and its Applications*, Volume 4 , 215-238.
- [5] Cirici, L. and Ume, J.S., 2006, Some common fixed point theorems for weakly compatible mappings, *J. Math. Anal. Appl.* 314, 488-499.
- [6] Daffer, P.Z. and Kaneko, H., 1995, Fixed points of generalized contractive multi-valued mappings, *Jurnal of Mathematical Analysis and Applications*, vol. 192, no. 2, pp. 655-666.
- [7] Hoxha, E., 2013, Common Fixed Point for Weakly Compatible Generalized  $\varphi$ -weak Contractive Mappings. *Current Trends in Technology and Science*. Volume II, Issue V, 317-322.
- [8] Hitzler, P., 2001, Generalized metrics and topology in logic programming semantics. National University of Ireland, (University College Cork), PH.D.Thesis.
- [9] Hitzler, P. and Seda, A.K., 2000, Dislocated topologies, *J.Electr.Engin.*, 51 (12/S):3:7.
- [10] Jungck, G., 1986, Compatible mappings and common fixed points, *Int. J. Math. Math. Sci.* 9, pp.771-779.
- [11] Jungck, G. & Rhoades, B.E, 1993, Some fixed point theorems for compatible maps, *Int.J.Math.Sci.*16 No.3, pp 417-428.
- [12] Karapinar, E. and Salimi, P., 2013, Dislocated metric space to metric spaces with some fixed point theorems, *Fixed point theory and applications* 2013, 2013:222.
- [13] Kamran, T., 2007, Multivalued  $f$ -weakly Picard mappings, *Nonlinear Analysis*, 67, 2289-2296. MR2331879 (2008 d: 47095) zbl 1128.54024.
- [14] Kumari, P.S., Kumar, V.V. and Sarma, I.R., 2012, Common fixed point theorems on weakly compatible maps on dislocated metric spaces, *Mathematical Sciences* 2012, 6:71, <http://www.iaumath.com/content/6/1/71>.
- [15] Liu, L.S., 1992, On common fixed points of single valued mappings, *J. Qufu Norm. Univ. Nat. Sci. Ed.* 18, No.1, pp.6-10.
- [16] Shrivastava, R., Ansari, Z.K. and Sharma, M., 2012, Some results on fixed points in dislocated and dislocated quasi-metric spaces, *Journal of Advanced Studies in Topology*, Vol 3. No.1, 25-31.
- [17] Zhang, Q. and Song, Y., 2009, Fixed point theory of generalized  $\varphi$ -weak contractions, *Applied Mathematics Letters* 22, 75-78, MR2484285. zbl 1163.47304.

## AUTHOR'S PROFILE



**Prof. Elida Hoxha** (DOB-08/01/1961)

completed her M.Sc. in Mathematics from Tirana University in the year 1984 and completed her Ph.D. from Tirana University in 1997. She has a teaching experience of more than 24 years. Presently she is working as Associate Professor in the Department of Mathematics, Faculty of Natural Science, University of Tirana, Albania.

She is a popular teacher in under graduate and post graduate level. Her subject of teaching is Mathematical Analysis, Topology, Functional Analysis. Besides teaching she is actively engaged in research field. Her research fields of Fixed Point Theory, Fuzzy sets and Fuzzy mappings, Topology.